

A database of threat statuses and life-history traits of Red List species in Flanders (northern Belgium)

Dirk Maes[‡], Dimitri Brosens^{‡,§}, Filiep T'jollyn[‡], Stijn Van Hoey[‡], Frederic Piesschaert[‡], Tim Adriaens[‡], Wouter Dekoninck[‡], Koen Devos[‡], Koen Lock, Thierry Onkelinx[‡], Jo Packet[‡], Jeroen Speybroeck[‡], Arno Thomaes[¶], Koen Van Den Berge[‡], Wouter Van Landuyt[‡], Hugo Verreycken[‡]

[‡] Research Institute for Nature and Forest (INBO), Brussels, Belgium

[§] Belgian Biodiversity Platform, Brussels, Belgium

[¶] Royal Belgian Institute of Natural Sciences, Brussels, Belgium

[¶] Research Institute for Nature and Forest (INBO), Geraardsbergen, Belgium

Corresponding author: Dirk Maes (dirk.maes@inbo.be), Dimitri Brosens (dimitri.brosens@inbo.be)

© Dirk Maes, Dimitri Brosens, Filiep T'jollyn, Stijn Van Hoey, Frederic Piesschaert, Tim Adriaens, Wouter Dekoninck, Koen Devos, Koen Lock, Thierry Onkelinx, Jo Packet, Jeroen Speybroeck, Arno Thomaes, Koen Van Den Berge, Wouter Van Landuyt, Hugo Verreycken



Citation: () . <https://doi.org/>

Abstract

Background

Red Lists estimate the extinction risk of species at global or regional levels and are important instruments in conservation policies. Global Red List assessments are readily available via the IUCN website (<https://www.iucnredlist.org/>) and are regularly updated by (taxonomic) experts. Regional Red Lists, however, are not always easy to find and often use local criteria to assess the local extinction risk of species.

New information

Here, we publish a database with the outcome of 38 Red List assessments in Flanders (northern Belgium) between 1994 and 2018. In total, the database contains 6,224 records of 5,039 unique species pertaining to 24 different taxonomic groups. Using a quality control procedure, we evaluated the criteria used, the number of records, the temporal and spatial distribution of the data and the up-to-dateness of the Red Lists. This way, nineteen Red Lists were approved as being of sufficient high quality (i.e. validated) and nineteen others were not. Once validated, Red Lists are approved by the regional minister of environment and published in the Belgian Official Gazette acquiring legal status. For the validated Red Lists, we additionally compiled (life-history) traits that are applicable to a wide variety of species groups (taxonomic kingdom, biome, biotope, nutrient level, dispersal capacity, lifespan and cuddliness). The publication of this dataset allows comparison of Red List statuses with other European regions and countries and permits analyses about how certain (life-history) traits can explain the Red List status of species. The dataset will be regularly updated by adding new Red List (re)assessments and/or additional (life-history) traits.

Keywords

Red List, Flanders (northern Belgium), life-history traits, IUCN, threatened species, conservation

Introduction

Red Lists are important instruments at both the global and the regional scale (Brooks et al. 2016). They estimate the extinction risk in a given region, usually using standardised and internationally accepted criteria (Mace et al. 2008). Although Red Lists are not compiled to prioritise conservation actions (Lamoreux et al. 2003, Rodrigues et al. 2006), they are often used as an important source for conservation policies (McCarthy et al. 2008) such as species action plans (Fitzpatrick et al. 2007, Laycock et al. 2011)

and reintroduction programmes (IUCN/SSC 2013). Also, consecutive Red Lists allow comparison of changes in Red List categories and, in combination with information on threats, inform effective application of conservation efforts to the species' major threats (Brooke et al. 2008). Yet, national or regional Red Lists are often only available locally and/or in the local language and not always easy to access, which hinders analyses on larger scales (Maes et al. 2019). Although a website with national or regional Red Lists exists (<http://www.nationalredlist.org/>), it is often more convenient for local authorities to manage the information on national or regional Red Lists locally (e.g. <https://www.inbo.be/en/search-flanders-red-lists>) and to publish them on open-access platforms for easier availability.

Here, we publish the results of Red List assessments in Flanders (northern Belgium) between 1994 and 2018. For the species on the validated Red Lists, we also include some general (life-history) traits (taxonomic kingdom, biome, biotope use, nutrient level, lifespan, mobility and cuddliness - cf. Trochet et al. 2014), which allows further analyses on the correlation between the Red List status and the species characteristics (cf. Jeppsson and Forslund 2014).

General description

Purpose: This database publishes the Red List statuses of all species that were assessed in Flanders (northern Belgium) since 1994. First, a literature search was done in both local and scientific publications to gather all Red List assessments ever performed in Flanders. All species present in the Red Lists were compiled in a database with the original taxonomic name and Red List status as published in the original Red List. The data were carefully checked for double entries and for typing errors in the published species names. Since Red List categories were not always in accordance with the presently used IUCN categories, we "translated" the originally published Red List category into IUCN Red List categories. Second, all species names were checked against the GBIF taxonomic backbone (<https://www.gbif.org/species>) to obtain currently traceable species names (including synonyms). Since the Flemish Species Decree of 2009 (Soortenbesluit) came into effect, Red List assessments in Flanders are coordinated by the Research Institute for Nature and Forest (INBO, formerly the Institute for Nature Conservation – IN). The task of institute is to instigate the compilation of new Red Lists but also to perform a control procedure to check whether the Red List is of sufficiently high quality. The quality control procedure consists of checking the number of available historical and recent data, the number of sites that was surveyed in both historical and recent times and the spatial coverage of the different ecological districts of Flanders (Maes et al. 2015). For older Red Lists, this quality control was done *post factum*, while for new Red Lists this was done at the start of the Red List assessment. Red Lists fulfilling the quality control criteria are labelled as "validated Red Lists". Applying this procedure, we were able to validate nineteen out of 38 published Red Lists Table 1, while nineteen others could not be validated Table 2.

Table 1. Download as CSV 

Validated Red Lists in Flanders (n = 19) with the Red List criteria used (Local or IUCN Red List criteria), the year of publication, the reference to the Red List, the reference to the life-history traits and the number of species (nSpecies) included in the Red List.

Taxonomic group	Criteria	Year	Reference	Reference(s) life-history traits	nSpecies
Butterflies (Lepidoptera – Rhopalocera)	Local	1996	Maes and Van Dyck (2001)	Maes et al. (2013)	68
Grasshoppers (Orthoptera)	Local	2000	Decler et al. (2000)	Kleukers et al. (1997)	39
Breeding birds (Aves)	Local	2004	Devos et al. (2004)	Birdlife International 2004, Vermeersch et al. (2004)	211
Dragonflies (Odonata)	Local	2005	De Knijf (2006)	De Knijf (2006); Nederlandse vereniging voor Libellenstudie 2002	66
Vascular plants (Tracheophyta)	Local	2006	Van Landuyt et al. (2006)	Biesbrouck et al. (2001), Ellenberg et al. (1992), Hill et al. (2004), Stieperaere and Fransen (1982)	1154
Carabid beetles (Coleoptera – Carabidae)	Local	2008	Desender et al. (2008b)	Desender et al. (2008a), Homburg et al. (2014), Turin (2000)	382
Butterflies (Lepidoptera – Rhopalocera)	IUCN	2011	Maes et al. (2012)	Maes et al. (2013)	70
Amphibians (Amphibia)	IUCN	2012	Jooris et al. (2012)	Jooris et al. (2013)	16

Reptiles (Reptilia)	IUCN	2012	Jooris et al. (2012)	Jooris et al. (2013)	6
Waterbugs (Hemiptera – Gerromorpha, Nepomorpha)	IUCN	2013	Lock et al. (2013)	Aukema et al. (2002), Stoffelen et al. (2013)	62
Freshwater fishes (Pisces)	IUCN	2014	Verreycken et al. (2014)	Kottelat and Freyhof (2007), Vandellannoote et al. (1998), Van Emmerik and De Nie (2006)	42
Ladybirds (Coleoptera – Coccinellidae)	IUCN	2014	Adriaens et al. (2015)	Adriaens and Maes (2004), Adriaens et al. (2008), Baugnée et al. (2011), Cuppen et al. (2017), Roy et al. (2011)	39
Mammals (Mammalia)	IUCN	2014	Maes et al. (2014)	Verkem et al. (2003)	103
Saproxyllic beetles (Coleoptera – Cetoniidae, Dynastidae, Lucanidae)	IUCN	2015	Thomaes et al. (2015)	Thomaes et al. (2015)	19
Breeding birds (Aves)	IUCN	2016	Devos et al. (2016)	Birdlife International (2004), Vermeersch et al. (2004)	217
Grasshoppers (Orthoptera)	IUCN	2017	Maes et al. (2017)	Kleukers et al. (1997)	52
Hornworts (Anthocerotophyta)	IUCN	2017	Van Landuyt and De Beer (2017)	Hill et al. (2007), Siebel (2005), Siebel and During (2006)	4
Liverworts (Marchantiophyta)	IUCN	2017	Van Landuyt and De Beer (2017)	Hill et al. (2007), Siebel (2005), Siebel and During (2006)	114
Mosses (Bryophyta)	IUCN	2017	Van Landuyt and De Beer (2017)	Hill et al. (2007), Siebel (2005), Siebel and During (2006)	399

Table 2. Download as CSV 

Non-validated Red Lists in Flanders (n = 19) with the criteria used (Expert judgement, Local or IUCN Red List criteria), the year of publication, the reference to the Red List and the number of species (nSpecies) included in the Red List.

Taxonomic group	Criteria	Year	Reference	nSpecies
Mammals (Mammalia)	Expert judgement	1994	Criel (1994)	69
Waterbugs (Hemiptera – Gerromorpha, Nepomorpha)	Expert judgement	1994	Bosmans (1994)	58
Waterbeetles (Coleoptera – Dytiscidae, Gyrinidae, Haliplidae, Noteridae, Paelobiidae)	Expert judgement	1994	Bosmans (1994)	139
Carabid beetles (Coleoptera – Carabidae)	Local	1995	Desender et al. (1995)	368
Amphibians (Amphibia)	Local	1996	Bauwens and Claus (1996)	14
Dragonflies (Odonata)	Local	1996	De Knijf and Anselin (1996)	58
Reptiles (Reptilia)	Local	1996	Bauwens and Claus (1996)	5
Fishes (Pisces)	Local	1998	Vandellannoote and Coeck (1998)	55
Hoverflies (Diptera – Syrphidae)	Local	1998	Meerhaeghe and Grootaert (1998)	265
Spiders (Araneae)	Expert judgement	1998	Maelfait et al. (1998)	604
Breeding birds (Aves)	Expert judgement	1999	Devos and Anselin (1999)	71

Macrofungi (Ascomycota, Basidiomycota)	Local	1999	Walley and Verbeken (1999)	552
Dolichopodid flies (Diptera – Dolichopodidae)	Local	2000	Pollet (2000)	260
Empidid flies (Diptera – Empididae, Hybotidae, Atelestidae, Brachystomatidae)	Local	2001	Grootaert et al. (2001)	259
Waterbugs (Hemiptera – Gerromorpha, Nepomorpha)	Local	2001	Bonte et al. (2001)	58
Ants (Hymenoptera – Formicidae)	Local	2003	Dekoninck et al. (2003)	55
Land snails (Mollusca)	Local	2006	van Loen et al. (2006)	120
Waterbeetles (Coleoptera – Dytiscidae)	Local	2012	Scheers (2012)	106
Grasshoppers (Orthoptera)	IUCN	2013	Lock et al. (2011)	45

Thirteen validated Red Lists have already been approved by the minister of environment and were published in the Belgian Official Gazette and thus acquired legal status (publication numbers 2011035522 and 2013204362; http://www.ejustice.just.fgov.be/doc/rech_n.htm). The procedure to have the six most recently validated Red Lists also approved by the minister has been started (saproxylic beetles, breeding birds, grasshoppers, hornworts, liverworts and mosses – Table 1).

Red Lists are usually published as reports of the Research Institute for Nature and Forest (INBO), but some of them remained unpublished and were only available for internal use (waterbeetles – Bosmans 1994, waterbugs – Bosmans 1994; Bonte et al. 2001, hoverflies – Meerhaeghe and Grootaert 1998, land snails – van Loen et al. 2006). Other Red Lists were published as part of (provisional) distribution atlases (amphibians and reptiles – Bauwens and Claus 1996, freshwater fishes – Vandellannoote and Coeck 1998, grasshoppers – Declerck et al. 2000, ants – Dekoninck et al. 2003, breeding birds – Devos et al. 2004, dragonflies – De Knijf 2006, vascular plants – Van Landuyt et al. 2006), as dissertation projects (waterbeetles – Scheers 2012) or as papers in local (spiders – Maelfait et al. 1998, waterbugs – Lock et al. 2013) or international journals (butterflies – Maes et al. 2012, freshwater fishes – Verreycken et al. 2014, ladybirds – Adriaens et al. 2015). In this database, we bring together all the threat statuses of all the species mentioned in the published and unpublished Red Lists since 1994 in Flanders.

Finally, we added (life-history) traits to the species in the validated Red Lists based on regional sources on the biology and/or ecology of the different species groups (see references in Table 1). The workflow for the compilation of the Red List database in Flanders is given in Fig. 1.




Figure 1.

Workflow for the compilation of the Red List database in Flanders (northern Belgium).

We plan to regularly update the database with newly published Red Lists together with the life-history traits of the assessed species. We also aim to reassess all Red Lists (using IUCN criteria) of taxonomic groups for which only local criteria were applied in the past (e.g. carabid beetles, dragonflies, vascular plants).

Since 1994, 38 Red Lists have been compiled in Flanders. In total, this concerned 6,224 records of 5,039 unique species – in some cases listed in consecutive Red Lists – pertaining to 24 taxonomic groups. Most of the older Red Lists (1994-1999) used expert judgement without quantitative analyses to classify species into Red List categories: mammals (Criel 1994), waterbugs (Bosmans 1994), waterbeetles (Bosmans 1994), spiders (Maelfait et al. 1998) and breeding birds (Devos and Anselin 1999). After the publication of local Red List criteria (Maes et al. 1995), Red Lists were compiled using a combination of rarity and decline (Maes and van Swaay 1997). Since 2011, however, IUCN criteria for regional use (IUCN 2003) were adopted in Flanders (Maes et al.

2011) and since then, all Red Lists were compiled using the IUCN criteria. The main difference between local and IUCN criteria is that IUCN criteria allow species to be classified based on declining trends (criterion A), rarity or small population sizes only (criterion B, C or D), while in the previously used local criteria, a species could only be classified as threatened when it had both a declining (distribution or population) trend and when the species was rare distribution.

Geographic coverage

Description: Flanders covers an area of 13,522 km² and is the northern administrative region of Belgium Fig. 2. The bounding box of Flanders is 50.68N to 51.51N latitude and 2.54E to 5.92E longitude. It represents 45% of the Belgian territory and is largely covered by agricultural land and urban areas. With 481 inhabitants per km², Flanders is one of the most densely populated areas in Europe.



Figure 2.
The location of Flanders (in grey) in NW Europe.

Coordinates: 50.67 and 51.51 Latitude; 5.94 and 2.53 Longitude.

Traits coverage

RLCAsPublished and RLC_IUCN

The Red List category as published in the original Red Lists, mostly in Dutch, are given in Table 3, as well as their translation into the Red List category according to the IUCN Red List categories (IUCN 2003). Depending on the Red List categories used, the original Red List categories can be translated into different IUCN categories, and vice versa.

Table 3. [Download as CSV](#)
Translation of published Red List category name into IUCN Red List categories.

RLCAsPublished	RLC_IUCN	IUCN Category
Achteruitgaand	NT	Near Threatened
Bedreigd	EN	Endangered
Bedreigd	VU	Vulnerable ^a
Bedreigd, maar mate waarin ongekend	DD	Data Deficient
Bedreigd, maar niet gekend in welke mate	DD	Data Deficient
Bijna in gevaar	NT	Near Threatened
Critically endangered	CR	Critically Endangered

Endangered	EN	Endangered
Ernstig bedreigd	CR	Critically Endangered
Geografisch beperkt	NT	Near Threatened
Kwetsbaar	VU	Vulnerable
Least concern	LC	Least Concern
Met uitsterven bedreigd	CR	Critically Endangered
Met verdwijning bedreigd	CR	Critically Endangered
Momenteel niet bedreigd	LC	Least Concern
Momenteel niet in gevaar	LC	Least Concern
Near threatened	NT	Near Threatened
Niet bedreigd	LC	Least Concern
Niet geëvalueerd	NE	Not Evaluated
Niet van toepassing	NA	Not Applicable
Niet-inheemse broedvogel	NA	Not Applicable
Not assessed	NE	Not Evaluated
Onregelmatige broedvogel	NE	Not Evaluated
Onvoldoende data	DD	Data Deficient
Onvoldoende gekend	DD	Data Deficient
Regionaal uitgestorven	RE	Regionally Extinct
Regionally extinct	RE	Regionally Extinct
Sterk bedreigd	EN	Endangered
Uitgestorven	RE	Regionally Extinct
Uitgestorven in Vlaanderen	RE	Regionally Extinct
Vatbaar voor bedreiging	NT	Near Threatened
Verdwenen	RE	Regionally Extinct
Verdwenen uit Vlaanderen en het Brussels Gewest	RE	Regionally Extinct
Vermoedelijk bedreigd	DD	Data Deficient
Vulnerable	VU	Vulnerable
Waarschijnlijk bedreigd	DD	Data Deficient
Zeldzaam	NT	Near Threatened
Zeldzaam (vrij zeldzaam)	NT	Near Threatened
Zeldzaam (zeer zeldzaam)	NT	Near Threatened
Zeldzaam (zeldzaam)	NT	Near Threatened

Criteria

The criteria used to compile the Red Lists (Table 4).

Table 4. [Download as CSV](#) 

Criteria used for the Red List assessments in Flanders.

Criteria	Description
Expert judgement	The Red List was compiled on the basis of expert knowledge and no quantitative criteria were used
Local	Quantitative local criteria were used (Maes et al. 1995)
IUCN	Quantitative IUCN criteria were used (Maes et al. 2011)

Kingdom

The taxonomic kingdom to which a species belongs (Fungi, Invertebrates, Plants, Vertebrates – Table 5).

Table 5. [Download as CSV](#) 

The taxonomic kingdom to which the different species groups belong.

Kingdom	Description
Fungi	Agaricomycetes, Geoglossomycetes, Leotiomycetes, Pezizomycetes, Sordariomycetes
Invertebrates	Ants (Formicidae), Butterflies (Lepidoptera – Rhopalocera), Carabid beetles (Carabidae), Dolichopodid flies (Dolichopodidae), Dragonflies (Odonata), Empidid flies (Empididae, Hybotidae, Atelestidae, Brachystomatidae), Grasshoppers (Orthoptera), Hoverflies (Diptera – Syrphidae), Ladybirds (Coleoptera – Coccinellidae), Molluscs (Mollusca), Saproxylid beetles (Coleoptera – Cetoniidae, Dynastidae, Lucanidae), Spiders (Araneae), Waterbeetles (Coleoptera - Dytiscidae, Gyrinidae, Haliplidae, Noteridae, Paelobiidae), Waterbugs (Hemiptera – Gerromorpha, Nepomorpha)
Plants	Hornworts (Anthocerotophyta), Liverworts (Marchantiophyta), Mosses (Bryophyta), Vascular plants (Tracheophyta)
Vertebrates	Amphibians (Amphibia), Breeding birds (Aves), Freshwater fishes (Pisces), Mammals (Mammalia), Reptilia (Reptilia)

Biome

The biome in which the species occurs (Aquatic, Epiphytic, Marine, Semi-aquatic, Terrestrial – Table 6).

Table 6. [Download as CSV](#) 

Description of the biome to which each species belongs.

Biome	Description
Aquatic	The major part of the life cycle is in water
Epiphytic	Living on trees (only liverworts and mosses)
Marine	At least a part of the life cycle is in the sea
Semi-aquatic	Water is necessary for hunting or breeding
Terrestrial	The major part of the life cycle is on land

Biotope1 and Biotope2

The preferred biotope in which the species occurs in Flanders (northern Belgium) (Table 7). This is based on broad biotope classifications used in the land use map of Flanders (Gobin et al. 2009) and in Corine Land Cover (Version 18.5.1) or Natura2000 habitats (Council Directive 92/43/EEC). For species occurring in two different biotope types, both biotopes are given in the database (Biotope1 and Biotope2).

Table 7. Download as CSV 

The biotope type in which each species occurs.

Biotope	Description
Agriculture	Arable fields, agricultural grasslands
Dunes	Sandy shores and coastal dunes
Eurytopic	No clear biotope preference or occurring in different biotope types
Grasslands	Dry and wet semi-natural grasslands
Heathlands	Dry and wet heathlands, (peat)bogs
Marine	Sea
Marshes	Wetlands, mires
Running waters	Rivers, rivulets
Salt marshes	Littoral sediment
Shrubs	Scrubs
Standing waters	Ponds, lakes, ditches
Urban	Anthropogenic (buildings, gardens, cemeteries, railroads ...), industrial sites
Woodlands	Deciduous, coniferous and mixed woodlands

NutrientLevel

The nutrient level of the biotope in which the species occurs (Eutrophic, Mesotrophic, Oligotrophic – Table 8).

Table 8. Download as CSV 

The nutrient level of the biotope in which each species occurs.

Nutrient level	Description
Eutrophic	The biotope in which the species occurs has a high nutrient level
Mesotrophic	The biotope in which the species occurs has an intermediate nutrient level
Oligotrophic	The biotope in which the species occurs has a low nutrient level

Lifespan

The longevity of the species (Table 9). We arbitrarily choose 3 years to discriminate between longlived and shortlived species.

Table 9. Download as CSV 

The lifespan of each species.

Lifespan	Description
Longlived	The species lives ≥ 3 years
Shortlived	The species lives < 3 years

Mobility

The dispersal capacities of the species (Table 10). We arbitrarily choose 5 kilometers to discriminate between mobile and sedentary species.

Table 10. [Download as CSV](#) 

The dispersal capacity of each species.

Mobility	Description
Mobile	The dispersal capacity of the species is ≥ 5 km
Sedentary	The dispersal capacity of the species is < 5 km

Cuddliness

Whether the species is considered cuddly or not (Stokes 2007) (Table 11). This information is only given for animals (invertebrates and vertebrates).

Table 11. [Download as CSV](#) 

The cuddliness of each species.

Cuddliness	Description
Cuddly	The species is considered cuddly
Non cuddly	The species is considered non-cuddly (spiny, dangerous, venomous, predator)

Temporal coverage

Notes: All Flemish Red Lists compiled between 1994 and 2018.

Usage rights

Use license: Creative Commons Public Domain Waiver (CC-Zero)

Data resources

Data package title: Red List species Flanders database

Number of data sets: 2

Data set name: Non-validated Red Lists in Flanders (Belgium)

Character set: UTF-8

Data format: DWC-A

Description: Summary of the Red List status of the 3,161 species for which only non-validated Red Lists are available in Flanders (northern Belgium).

Data set name: Validated Red Lists in Flanders (Belgium)

Character set: UTF-8

Data format: DWC-A

Description: Summary of the Red List status and (life-history) traits of the 3,063 species for which validated Red Lists are available in Flanders (northern Belgium).

Acknowledgements

We thank all volunteers for collecting distribution data with which the different Red Lists were compiled. Marc Pollet is kindly thanked for very helpful and constructive comments on an earlier version of the manuscript. We also thank Peter Desmet for his help with the publication of the data on GBIF.

Author contributions

DM compiled the data base and FT thoroughly checked all entries in the data base, DB, FP and SVH checked the species names against the GBIF Backbone Taxonomy and DB converted the database into DarwinCore. TA (dragonflies, grasshoppers, ladybirds), WD (carabid beetles), KD (breeding birds), KL (grasshoppers, waterbeetles, waterbugs), DM (butterflies), TO (mammals - bats), JP (molluscs), JS (amphibians and reptiles), AT (saproxylic beetles), KVDB (mammals), WVL (hornworts, liverworts, mosses, vascular plants) and HV (freshwater fishes) provided the life-history traits for the different taxonomic groups. All authors assisted in the writing of the manuscript.

References

- Adriaens T, Maes D (2004) Voorlopige verspreidingsatlas van lieveheersbeestjes in Vlaanderen, resultaten van het lieveheersbeestjesproject van de jeugdbonden. Bertram 2 (1bis): 1-72.
- Adriaens T, San Martin y Gomez G, Maes D (2008) Invasion history, habitat preferences and phenology of the invasive ladybird *Harmonia axyridis* in Belgium. BioControl 53 (1): 69-88. <https://doi.org/10.1007/s10526-007-9137-6>
- Adriaens T, San Martin y Gomez G, Bogaert J, Crevecoeur L, Beuckx JP, Maes D (2015) Testing the applicability of regional IUCN Red List criteria on ladybirds (Coleoptera, Coccinellidae) in Flanders (north Belgium): opportunities for conservation. Insect Conservation and Diversity 8 (5): 404-417. <https://doi.org/10.1111/icad.12124>
- Aukema B, Cuppen JGM, Nieser N, Tempelman D (2002) Verspreidingsatlas Nederlandse wantsen (Hemiptera: Heteroptera). Deel I: Dipsocoromorpha, Nepomorpha, Gerromorpha & Leptopodomorpha. EIS Nederland, Leiden.
- Baugnée JY, Branquart E, Maes D, Segers S (2011) Velddeterminatietabel voor de lieveheersbeestjes van België en Nederland (Chilocorinae, Coccinellinae, Epilachninae & Coccidulinae): herziene druk met larventabel. Jeugdbond voor Natuur en Milieu, Jeunes & Nature asbl i.s.m. het Instituut voor Natuur- en Bosonderzoek, Gent/Wavre/Brussel.
- Bauwens D, Claus K (1996) Verspreiding van amfibieën en reptielen in Vlaanderen. De Wielewaal, Turnhout.
- Biesbrouck B, Es K, Van Landuyt W, Vanhecke L, Hermy M, Van den Bremt P (2001) Een ecologisch register voor hogere planten als instrument voor het natuurbehoud in Vlaanderen. Flo.Wer vzw, Instituut voor Natuurbehoud, Nationale Plantentuin van België, KULeuven, Brussel. [ISBN 90-403-1138-7]
- Bonte D, Vandomme V, Muylaert J, Bosmans R (2001) Een gedocumenteerde Rode Lijst van de water- en oppervlaktewantsen van Vlaanderen [unpublished report]. Universiteit Gent, Gent.
- Bosmans R (1994) Een gedocumenteerde Rode lijst van de waterkevers en waterwantsen van Vlaanderen, met inbegrip van enkele case studies [unpublished report]. Laboratorium voor Dierenecologie, Biogeografie en Natuurbehoud, Gent.

- Brooke MD, Butchart SHM, Garnett ST, Crowley GM, Mantilla-Beniers NB, Stattersfield A (2008) Rates of movement of threatened bird species between IUCN red list Categories and toward extinction. *Conservation Biology* 22 (2): 417-427. URL: [ISI:000254796800024](https://doi.org/10.1002/254796800024)
- Brooks TM, Akcakaya HR, Burgess ND, Butchart SHM, Hilton-Taylor C, Hoffmann M, Juffe-Bignoli D, Kingston N, MacSharry B, Parr M, Perianin L, Regan EC, Rodrigues ASL, Rondinini C, Shennan-Farpon Y, Young BE (2016) Analysing biodiversity and conservation knowledge products to support regional environmental assessments. *Scientific Data* 3 <https://doi.org/10.1038/sdata.2016.7>
- Criel D (1994) Rode lijst van de zoogdieren in Vlaanderen. AMINAL, Brussel. [ISBN D/1994/3241/129]
- Cuppen J, Kalkman VJ, Tacoma-Krist G (2017) Verspreiding, biotoop en fenologie van de Nederlandse lieveheersbeestjes (Coleoptera: Coccinellidae). *Entomologische Berichten* 77 (3): 147-188.
- Declerck K, Devriese H, Hofmans K, Lock K, Barenburg B, Maes D (2000) Voorlopige atlas en "rode lijst" van de sprinkhanen en krekels van België (Insecta, Orthoptera). IN.2000/10. SALTABEL i.s.m. IN en KBIN, Brussel. [ISBN 90-403-0119-0]
- De Knijf G, Anselin A (1996) Een gedocumenteerde Rode lijst van de libellen van Vlaanderen. 4. Instituut voor Natuurbehoud, Brussel.
- De Knijf G (2006) De Rode Lijst van de libellen in Vlaanderen. In: De Knijf G, Anselin A, Goffart P, Tailly M (Eds) De libellen (Odonata) van België: verspreiding - evolutie - habitats. Libellenwerkgroep Gomphus ism Instituut voor Natuur- en Bosonderzoek, Brussel, 16 pp.
- Dekoninck W, Vankerkhoven F, Maelfait JP (2003) Verspreidingsatlas en voorlopige Rode Lijst van de mieren van Vlaanderen. IN.R.2003.7. Instituut voor Natuurbehoud, Brussel.
- Desender K, Maes D, Maelfait JP, Van Kerckvoorde M (1995) Een gedocumenteerde Rode Lijst van de zandloopkevers en loopkevers van Vlaanderen. 1. Instituut voor Natuurbehoud, Brussel.
- Desender K, Dekoninck W, Maes D (2008a) An updated Red List of the ground and tiger beetles (Coleoptera, Carabidae) in Flanders (Belgium). *Bulletin van het Koninklijk Belgisch Instituut voor Natuurwetenschappen, Entomologie* 78: 113-131.
- Desender K, Dekoninck W, Maes D, Crevecoeur L, Dufrêne M, Jacobs M, Lambrechts J, Pollet M, Stassen E, Thys N (2008b) Een nieuwe verspreidingsatlas van de loopkevers en zandloopkevers (Carabidae) in België. INBO.R.2008.13. Instituut voor Natuur- en Bosonderzoek & Koninklijk Belgisch Instituut voor Natuurwetenschappen, Brussel.
- Devos K, Anselin A (1999) Broedvogels. In: Kuijken E (Ed.) Natuurrapport 1999. Toestand van de natuur in Vlaanderen: cijfers voor het beleid. Instituut voor Natuurbehoud, Brussel, 3 pp.
- Devos K, Anselin A, Vermeersch G (2004) Een nieuwe Rode Lijst van de broedvogels in Vlaanderen (versie 2004). In: Vermeersch G, Anselin A, Devos K, Herremans M, Stevens J, Gabriëls J, Van Der Krieken B (Eds) Atlas van de Vlaamse broedvogels 2000-2002. Instituut voor Natuurbehoud, Brussel, 15 pp.
- Devos K, Anselin A, Driessens G, Herremans M, Onkelinx T, Spanoghe G, Stienen EWM, T'Jollyn F, Vermeersch G, Maes D (2016) De IUCN Rode-Lijst van de broedvogels in Vlaanderen (2016). *Natuur.oriolus* 82 (4): 109-122. URL: https://www.natuurpunt.be/sites/default/files/documents/publication/natuur.oriolus_2016-4_de_iucn_rode_lijst_van_de_broedvogels_in_vlaanderen.pdf
- Ellenberg H, Weber HE, Dill R, Wirth V, Werner W, Paulissen D (1992) Zeigerwerte von Pflanzen in Mitteleuropa. Erich Goltze Verlag, Göttingen.
- Fitzpatrick U, Murray TE, Paxton RJ, Brown MJF (2007) Building on IUCN regional red lists to produce lists of species of conservation priority: a model with Irish bees. *Conservation Biology* 21 (5): 1324-1332. <https://doi.org/10.1111/j.1523-1739.2007.00782.x>
- Gobin A, Uljee I, Van Esch L, Engelen G, de Kok J, van der Kwast H, Hens M, Van Daele T, Peymen J, Van Reeth W, Overloop S, Maes F (2009) Landgebruik in Vlaanderen. 2009 (20). Instituut voor Natuur- en Bosonderzoek, Brussel.
- Grootaert P, Pollet M, Maes D (2001) A Red Data Book of Empidid Flies of Flanders (northern Belgium) (Diptera, Empididae s.l.): constraints and possible use in nature conservation. *Journal of Insect Conservation* 5 (2): 117-129. <https://doi.org/10.1023/A:1011315313330>
- Hill MO, Preston CD, Roy DB (2004) PLANTATT. Attributes of British and Irish Plants: Status, Size, Life History, Geography and Habitats. Biological Record Centre. NERC Centre for Ecology and Hydrology, Huntingdon, Cambridgeshire.
- Hill MO, Preston CD, Bosanquet SDS, Roy DB (2007) BRYOATT: attributes of British and Irish mosses, liverworts and hornworts. Centre For Ecology And Hydrology, Monks Wood.
- Homburg K, Homburg N, Schäfer F, Schuldt A, Assmann T (2014) Carabids.org – a dynamic online database of ground beetle species traits (Coleoptera, Carabidae). *Insect Conservation and Diversity* 7 (3): 195-205. <https://doi.org/10.1111/icad.12045>
- I (2003) Guidelines for Application of IUCN Red List Criteria at Regional Levels: Version 3.0. IUCN Species Survival Commission, IUCN, Gland, Switzerland and Cambridge, UK. [ISBN 2-8317-0738-2]
- I (2013) Guidelines for Reintroductions and Other Conservation Translocations. Version 1.0. IUCN Species Survival Commission, Gland, Switzerland. URL: <http://intranet.iucn.org/webfiles/doc/SSC/RedList/RedListGuidelines.pdf>

- International ,B (2004) Birds in Europe: population estimates, trends and conservation status. 12. BirdLife International, Cambridge, UK.
- Jeppsson T, Forslund P (2014) Species' traits explain differences in Red list status and long-term population trends in longhorn beetles. *Animal Conservation* 17 (4): 332-341. <https://doi.org/10.1111/acv.12099>
- Jooris R, Engelen P, Speybroeck J, Lewylle I, Louette G, Bauwens D, Maes D (2012) De IUCN Rode Lijst van de amfibieën en reptielen in Vlaanderen. INBO.R.2012.22. Instituut voor Natuur- en Bosonderzoek, Brussel.
- Jooris R, Engelen P, Speybroeck J, Lewylle I, Louette G, Bauwens D, Maes D (2013) De amfibieën en reptielen van Vlaanderen. Recent verspreiding en toelichting bij de nieuwe Rode Lijst. 2013/6. Natuurpunt Studie, Mechelen. [ISBN 978-90-5718-320-1]
- Kleukers R, van Nieuwerkerken E, Odé B, Willemse L, van Wingerden WKRE (1997) De Sprinkhanen en Krekels van Nederland (Orthoptera). 1. Nationaal Natuurhistorisch Museum, KNNV Uitgeverij & EIS-Nederland, Leiden.
- Kottelat M, Freyhof J (2007) Handbook of European freshwater fishes. Publications Kottelat, Cornol, Switzerland. [ISBN 978-2-8399-0298-4]
- Lamoreux J, Akçakaya HR, Bennun L, Collar NJ, Boitani L, Brackett D, Brautigam A, Brooks TM, de Fonseca GAB, Mittermeier RA, Rylands AB, Gärdenfors U, Hilton-Taylor C, Mace G, Stein BA, Stuart S (2003) Value of the IUCN Red List. *Trends in Ecology & Evolution* 18 (5): 214-215. [https://doi.org/10.1016/S0169-5347\(03\)00090-9](https://doi.org/10.1016/S0169-5347(03)00090-9)
- Laycock HF, Moran D, Smart JCR, Raffaelli DG, White PCL (2011) Evaluating the effectiveness and efficiency of biodiversity conservation spending. *Ecological Economics* 70 (10): 1789-1796. <https://doi.org/10.1016/j.ecolecon.2011.05.002>
- Lock K, Adriaens T, Devriese H, San Martin y Gomez G, Decler K (2011) Updated red lists of grasshoppers and crickets (Orthoptera) in Flanders, Brussels and Wallonia. *Bulletin de la Société royale belge d'Entomologie* 147: 211-225.
- Lock K, Stoffelen E, Vercauteren T, Bosmans R, Adriaens T (2013) Updated Red List of the water bugs of Flanders (Belgium) (Hemiptera : Gerromorpha & Nepomorpha). *Bulletin de la Société royale belge d'Entomologie/Bulletin van de Koninklijke Belgische Vereniging voor Entomologie* 149: 57-63.
- Mace GM, Collar NJ, Gaston KJ, Hilton-Taylor C, Akçakaya HR, Leader-Williams N, Milner-Gulland EJ, Stuart SN (2008) Quantification of Extinction Risk: IUCN's System for Classifying Threatened Species. *Conservation Biology* 22 (6): 1424-1442. <https://doi.org/10.1111/j.1523-1739.2008.01044.x>
- Maelfait JP, Baert L, Janssen M, Alderweireldt M (1998) A Red list for the spiders of Flanders. *Bulletin van het Koninklijk Belgisch Instituut voor Natuurwetenschappen, Entomologie* 68: 131-142.
- Maes D, Maelfait JP, Kuijken E (1995) Rode lijsten: een onmisbaar instrument in het moderne Vlaamse natuurbehoud. *Wielewaal* 61 (5): 149-156.
- Maes D, van Swaay CAM (1997) A new methodology for compiling national Red Lists applied on butterflies (Lepidoptera, Rhopalocera) in Flanders (N.-Belgium) and in The Netherlands. *Journal of Insect Conservation* 1 (2): 113-124. <https://doi.org/10.1023/A:1018435110335>
- Maes D, Van Dyck H (2001) Butterfly diversity loss in Flanders (north Belgium): Europe's worst case scenario? *Biological Conservation* 99 (3): 263-276. [https://doi.org/10.1016/S0006-3207\(00\)00182-8](https://doi.org/10.1016/S0006-3207(00)00182-8)
- Maes D, Decler K, De Bruyn L, Hoffmann M (2011) Nieuwe Rode-Lijstcategorieën en -criteria voor Vlaanderen. Een aanpassing aan de internationale IUCN standaarden. *Natuur.focus* 10 (2): 54-61.
- Maes D, Vanreusel W, Jacobs I, Berwaerts K, Van Dyck H (2012) Applying IUCN Red List criteria at a small regional level: A test case with butterflies in Flanders (north Belgium). *Biological Conservation* 145 (1): 258-266. <https://doi.org/10.1016/j.biocon.2011.11.021>
- Maes D, Vanreusel W, Van Dyck H (2013) Dagvlinders in Vlaanderen: nieuwe kennis voor betere actie. Uitgeverij Lannoo nv, Tielt. [ISBN 978 94 014 0790 8]
- Maes D, Baert K, Boers K, Casaer J, Crevecoeur L, Criel D, Dekeukeleire D, Gouw J, Gyselings R, Haelters J, Herman D, Herremans M, Lefebvre ,, Lefevre A, Onkelinx T, Stuyck J, Thomaes A, Van Den Berge K, Vandendriessche B, Verbeylen G, Vercayie D (2014) De IUCN Rode Lijst van de zoogdieren in Vlaanderen. INBO.R.2014.1828211. Instituut voor Natuur- en Bosonderzoek, Brussel.
- Maes D, Isaac NB, Harrower C, Collen B, van Strien A, Roy DB (2015) The use of opportunistic data for IUCN Red List assessments. *Biological Journal of the Linnean Society* 115 (3): 690-706. <https://doi.org/10.1111/bj.12530>
- Maes D, Adriaens T, Decler K, Foquet B, Foquet R, Lambrechts J, Lock K, Piesschaert F (2017) IUCN Rode Lijst van de sprinkhanen en krekels in Vlaanderen. 2017 (29). Instituut voor Natuur- en Bosonderzoek,, Brussel. <https://doi.org/10.21436/inbor.12800242>
- Maes D, Verovnik R, Wiemers M, Brosens D, Beshkov S, Bonelli S, Buszko J, Cantú Salazar L, Cassar LF, Collins S, Dincă V, Djuric M, Dusej G, Elven H, Franeta F, Garcia Pereira P, Geryak Y, Goffart P, Gór A, Hiermann U, Höttinger H, Huemer P, Jakšić P, John E, Kalivoda H, Kati V, Komac B, Kőrösi A, Kulak AV, Kuussaari M, L'Hoste L, Lelo S, Mestdagh X, Micevski N, Mihut S, Monasterio León Y, Munguira ML, Murray T, Nielsen PS, Ólafsson E, Öunap E, Pamperis L, Pavlíčko A, Pettersson LB, Popov S, Popović M, Ryrholm N, Šašić M, Pöry J, Savenkov N, Settele J, Sielezniew M, Sinev S, Stefanescu C, Švitra G, Tammaru T, Tiitsaar A,

- Tzirkalli E, Tzortzakaki O, van Swaay CAM, Viborg AL, Wynhoff I, Zografou K, Warren MS (2019) Integrating national Red Lists for prioritising conservation actions for European butterflies. *Journal of Insect Conservation* in press (Special Issue Butterfly Conservation 2018). <https://doi.org/10.1007/s10841-019-00127-z>
- McCarthy MA, Thompson CJ, Garnett ST (2008) Optimal investment in conservation of species. *Journal of Applied Ecology* 45 (5): 1428-1435. <https://doi.org/10.1111/j.1365-2664.2008.01521.x>
 - Meerhaeghe A, Grootaert P (1998) Een gedocumenteerde Rode lijst van de zweefvliegen in Vlaanderen [unpublished report]. Instituut voor Natuurbehoud, Brussel.
 - Pollet M (2000) Een gedocumenteerde Rode lijst van de slankpootvliegen (Dolichopodidae) van Vlaanderen. 8. Instituut voor Natuurbehoud, Brussel.
 - Rodrigues ASL, Pilgrim JD, Lamoreux JF, Hoffmann M, Brooks TM (2006) The value of the IUCN Red List for conservation. *Trends in Ecology & Evolution* 21 (2): 71-76. <https://doi.org/10.1016/j.tree.2005.10.010>
 - Roy HE, Brown PMJ, Frost R, Poland R (2011) Ladybirds (Coccinellidae) of Britain and Ireland. Biological Record Centre, Wallingford, UK.
 - Scheers K (2012) Rode lijst en verspreidingsonderzoek van de waterroofkevers (Coleoptera: Dytiscidae) van Vlaanderen. Hogeschool Van Hall Larenstein, Velp.
 - Siebel H, During H (2006) Beknopte Mosflora van Nederland en België. KNNV Uitgeverij, Zeist.
 - Siebel HN (2005) Indicatie waarden van mossen. BLWG
 - Stieperaere H, Franssen K (1982) Standaardlijst van de Belgische vaatplanten, met aanduiding van hun zeldzaamheid en socio-oecologische groep. *Dumortiera* 22: 1-14.
 - Stoffelen E, Henderickx H, Vercauteren T, Lock K, Bosmans R (2013) De water- en oppervlaktewantsen van België (Hemiptera, Heteroptera: Nepomorpha & Gerromorpha). Determinatiesleutels, beschrijvingen en foto's van de volwassen water- en oppervlaktewantsen. Koninklijk Belgisch Instituut voor Natuurwetenschappen, Brussels.
 - Stokes DL (2007) Things we like: human preferences among similar organisms and implication for conservation. *Human Ecology* 35 (3): 361-369. <https://doi.org/10.1007/s10745-006-9056-7>
 - Thomaes A, Drumont A, Crevecoeur L, Maes D (2015) Rode lijst van de saproxyle bladsprietkevers (Lucanidae, Cetoniidae en Dynastidae) in Vlaanderen. INBO.R.2015.7843021. Instituut voor Natuur- en Bosonderzoek, Brussel.
 - Trochet A, Moulherat S, Calvez O, Stevens V, Clobert J, Schmeller D (2014) A database of life-history traits of European amphibians. *Biodiversity Data Journal* 2 <https://doi.org/10.3897/BDJ.2.e4123>
 - Turin H (2000) De Nederlandse loopkevers, verspreiding en oecologie (Coleoptera: Carabidae). Nationaal Natuurhistorisch Museum Naturalis, KNNV Uitgeverij & EIS-Nederland, Leiden. [ISBN 90-5011-136-X]
 - Vandellannoote A, Coeck J (1998) Rode lijst van de inheemse en ingeburgerde zoet- en brakwatersoorten en van de rondbekken in Vlaanderen. In: Vandellannoote A, Yseboodt R, Bruylants B, Verheyen RF, Coeck J, Maes J, Belpaire C, Van Thuyne G, Denayer B, Beyens J, De Charleroy D, Vandenabeele P (Eds) Atlas van de Vlaamse beek- en riviervissen. Water-Energik-vLario, Wijnegem, 5 pp.
 - Vandellannoote A, Yseboodt R, Bruylants B, Verheyen RF, Coeck J, Maes J, Belpaire C, Van Thuyne G, Denayer B, Beyens J, De Charleroy D, Vandenabeele P (1998) Atlas van de Vlaamse beek- en riviervissen. Water-Energik-vLario, Wijnegem.
 - Van Emmerik WAM, De Nie HW (2006) De zoetwatervissen van Nederland. Ecologisch bekeken. Sportvisserij Nederland, Bilthoven.
 - Van Landuyt W, Hoste I, Vanhecke L (2006) Rode Lijst van de vaatplanten van Vlaanderen en het Brussels Hoofdstedelijk Gewest. In: Van Landuyt W, Hoste I, Vanhecke L, Van den Brecht P, Vercruyssen W, De Beer D (Eds) Atlas van de Flora van Vlaanderen en het Brussels Gewest. Instituut voor Natuur- en Bosonderzoek, Nationale Plantentuin van België & Flo.Wer., Brussel, 12 pp.
 - Van Landuyt W, De Beer D (2017) Een Rode Lijst van de houwmossen (Anthocerotophyta), levermossen (Marchantiophyta) en bladmossen (Bryophyta) van Vlaanderen. 2017 (48). Instituut voor Natuur- en Bosonderzoek, Brussel.
 - van Loen H, Jordaens K, Backeljau T (2006) Gedocumenteerde Rode Lijst en Naamlijst van de landslakken van Vlaanderen en Brussel [unpublished report]. Onderzoeksopdracht IN/JPM/97002. Instituut voor Natuur- en Bosonderzoek, Koninklijk Belgisch Instituut voor Natuurwetenschappen, Universiteit Antwerpen, Brussel, Antwerpen.
 - Verkem S, De Maeseneer J, Vandendriessche B, Verbeylen G, Yskout S (2003) Zoogdieren in Vlaanderen. Ecologie en verspreiding van 1987 tot 2002. Natuurpunt Studie & JNM-Zoogdierenwerkgroep, Mechelen.
 - Vermeersch G, Anselin A, Devos K, Herremans M, Stevens J, Gabriëls J, Van Der Krieken B (2004) Atlas van de Vlaamse broedvogels 2000-2002. Instituut voor Natuurbehoud, Brussel. [ISBN 90-403-0215-4]
 - Verreycken H, Belpaire C, Van Thuyne G, Breine J, Buysse D, Coeck J, Mouton A, Stevens M, Vandenneucker T, De Bruyn L, Maes D (2014) An IUCN Red List of lampreys and freshwater fishes in Flanders (north Belgium). *Fisheries Management and Ecology* 21 (2): 122-132. <https://doi.org/10.1111/fme.12052>

- voor Libellenstudie ,NV (2002) De Nederlandse libellen (Odonata). Koninklijke Nederlandse Natuurhistorische Vereniging (KNNV); Krikken,J., Nationaal Natuurhistorisch Museum; European Invertebrate Survey, Utrecht; Leiden. [ISBN 90-5011-154-8]
- Walley R, Verbeken A (1999) Een gedocumenteerde Rode Lijst van enkele groepen paddestoelen (macrofungi) van Vlaanderen . 7. Instituut voor Natuurbehoud, Brussel. [ISBN 90-403-0111-5]