

Quantum Computation and Universal Quantum Computing

Traiko Dinev <traiko.dinev@gmail.com>

July 20, 2019

NOTE: This partially follows Introduction to Quantum Computing, a masters level course at the University of Edinburgh.

NOTE: Note this "summary" is NOT a reproduction of the course materials nor is it copied from the corresponding courses. It was entirely written and typeset from scratch.

License: Creative Commons public license; See README.md of repository

1 Quantum Systems

The state of a binary quantum system is described by a vector (a ket) in 2 dimensions:

$$|0\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad |1\rangle = \begin{pmatrix} 0 \\ 1 \end{pmatrix} \quad (1)$$

(2)

Physically these can be any differentiable states. In digital computing we often use thresholding to distinguish the "1"s from the "0"s. Here we use naturally quantized states such as energy levels, spin or (equivalently) polarization. We can extend this to continuous spaces where the discrete quantum states become continuous and sums become integrals. This is the basis of infinite-dimensional Hilbert spaces, which are discussed at the end of this [todo].

2 Double Slit Experiment and Superposition

[TODO: Motivate why the system collapses when a measurement is performed, etc.] [TODO: Motivate superposition, motivate probabilistic view of QM] [TODO: normalization of wavefunctions]

3 Postulates of Quantum Mechanics in Bra-Ket Notation

System State The state of a system is captured by its wavefunction $\psi(x)$. A wavefunction describes the state of the system as a probability distribution over possible states.

Observables and Wavefunction Collapse Every observable in classical mechanics (e.g. position, velocity, momentum) has a corresponding **operator** in quantum mechanics. The operators correspond to observables. Observables

are matrices and are Hermitian.

The most basic observable will be for given state $|\alpha\rangle$. If a system is in state $|\psi\rangle$ the inner product $\langle\alpha|\psi\rangle$ yields the probability of obtaining $\langle\alpha|$ when measuring in a binary basis. The observable is the matrix that has eigenvalues $|\alpha\rangle$ and $|\alpha\rangle^\top$, i.e. not- $|\alpha\rangle$.

When we perform a measurement, the wavefunction is **collapsed** to one of the underlying states we are measuring. We will only consider orthogonal states in this note. So if we measure in the $|0\rangle, |1\rangle$ basis, we will collapse **every quantum state** to either $|0\rangle$ or $|1\rangle$. Different states only differ in the probability between the basis vectors. *This is the essence of quantum mechanics.*

For example, position vectors are orthogonal states, which means we can measure (infinitely-small) the position of a particle. An operator, the observable, will have for eigenvectors (eigen-”states” in QM) each separate position.

Superposition Given the above observation, we can treat quantum states as a superposition of all possible **orthogonal** eigenstates. We can do all sorts of magic by measuring in clever states. Think of it as de-composing a signal into basic frequencies. *Note: the following is my interpretation.* It’s not that the signal itself **is built this way by nature**, it’s a by-product of how we perceive and measure the world. *If we can only treat an object as there and not there, we can treat it as being there AND not there at the same time with some probability.*

Time-evolution Schrödinger *[TODO: add]* Note this is a wave equation. C.f. ”the” wave equation (*[TODO: add rope wave equation]*):

$$H(t) |\psi(t)\rangle = i\hbar \frac{\partial}{\partial t} |\psi(t)\rangle \quad (3)$$

H is the hamiltonian, providing the total energy in the system. This is what we will have to find ourselves for different systems.

3.1 Example in a discrete system

Let’s assume we have the state:

$$|\psi\rangle = \alpha_0 |\phi_0\rangle + \dots \alpha_{K-1} |\phi_{K-1}\rangle \quad (4)$$

This is a superposition of K different states with varying amplitudes. This corresponds to the state having a distribution according to these amplitudes. Now let’s say we have a device that can measure each of those states ϕ_i . This assumes the states are orthogonal, because otherwise we wouldn’t be able to perfectly distinguish those states. Then our **observable** is:

$$\mathbf{M} = \sum_{j=0}^{K-1} \lambda_j |\phi_j\rangle \langle\phi_j| = \text{diag}(\lambda_1, \dots, \lambda_{K-1})$$

We take the outer product of the vectors, giving us a diagonal matrix over all states. The λ_j correspond to the outcomes, i.e. the **numbers we read off of the measurement device**.

Let’s consider an example. Set $K = 3$ and assume the following quantum state:

$$|\psi\rangle = \frac{1}{\sqrt{2}} |\phi_0\rangle + \frac{1}{\sqrt{2}} |\phi_2\rangle$$

The operator for the four states could be:

$$\mathbf{M} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 2 & 0 \end{pmatrix} =$$

$$1 \times |\phi_0\rangle \langle \phi_0| - 1 \times |\phi_1\rangle \langle \phi_1| + 2 \times |\phi_2\rangle \langle \phi_2|$$

The average measurement result would be $\langle \psi | M | \psi \rangle$:

$$\begin{aligned} & \left(\frac{1}{\sqrt{2}} \langle \phi_0| + \frac{1}{\sqrt{2}} \langle \phi_2| \right) \times 1 \times |\phi_0\rangle \langle \phi_0| - 1 \times |\phi_1\rangle \langle \phi_1| + 2 \times |\phi_2\rangle \langle \phi_2| \times \left(\frac{1}{\sqrt{2}} |\phi_0\rangle + \frac{1}{\sqrt{2}} |\phi_2\rangle \right) \\ &= \frac{1}{2} (|\phi_0\rangle \langle \phi_0| + 2 \times |\phi_2\rangle \langle \phi_2|) = \frac{3}{2} \end{aligned}$$

To understand why this makes sense, consider that 1/2 of the time we will observe $|\phi_0\rangle$, i.e. the number (measurement) 1. The other half we will observe $|\phi_2\rangle$ and 2. Which adds up to the same figure.

3.2 Projective Measurements

3.3 Density Matrices and Mixed Quantum Probabilistic States

NOTE: This is a combination of quantum uncertainty and classical uncertainty (noise)

3.4 Tensors and Tensor Products (Systems of Multiple Quantum States)

4 No-cloning Theorem

Assume there exists an operator U such that:

$$U |\psi\rangle \otimes |0\rangle = |\psi\rangle \otimes |\psi\rangle$$

$$U |\phi\rangle \otimes |0\rangle = |\phi\rangle \otimes |\phi\rangle$$

Then:

$$|00\rangle \xrightarrow{U} |00\rangle$$

$$|10\rangle \xrightarrow{U} |11\rangle$$

Hence:

$$\begin{aligned} \alpha |00\rangle + \beta |10\rangle &\xrightarrow{U} \alpha U |00\rangle + \beta U |10\rangle \\ &= \alpha |00\rangle + \beta |11\rangle \end{aligned}$$

However, at the same time:

$$\alpha |00\rangle + \beta |10\rangle = (\alpha |0\rangle + \beta |1\rangle) \otimes |0\rangle \xrightarrow{U} (\alpha |0\rangle + \beta |1\rangle) \otimes (\alpha |0\rangle + \beta |1\rangle)$$


Which are trivially not the same on the right side, but are the same on the left side. Q.e.d.

5 Entanglement and Bell States

Entanglement is a phenomenon that exists only in quantum systems and it's at the core of quantum computing. Mathematically, consider the following operation on two quantum states:

$$\begin{aligned} |+\rangle &= H |0\rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle) \\ |-\rangle &= H |1\rangle = \frac{1}{\sqrt{2}}(|0\rangle - |1\rangle) \end{aligned}$$

where H is the hadamard gate. These are a superposition of states. Now let's use $|+\rangle$ as a control for a CNOT on a $|0\rangle$. I.e. this circuit:




The resulting quantum state (of both qubits) is:

$$\frac{1}{\sqrt{2}}(|0\rangle + |1\rangle) \text{ CNOT } |0\rangle = \frac{1}{\sqrt{2}}(|00\rangle + |11\rangle)$$

where $|00\rangle = |0\rangle \otimes |1\rangle$ is shorthand for a tensor product. This is known as a **maximally entangled** state. This system cannot be separated into two distinct systems (i.e. two tensor-ed quantum states $|\alpha\rangle \otimes |\beta\rangle$). Another way to look at it is that if these qubits were separated (to Alice and Bob's labs), then measurements depend on each other. If Alice measures $|0\rangle$, then the state collapses to $|00\rangle$ and Bob also measures $|0\rangle$ (and vice-versa).

This is a profound result that holds true in reality. In fact, observation even after the fact causes the system to collapse. Key words are **quantum eraser** and **delayed-choice quantum eraser** experiments.

6 Superdense Coding



$$\begin{aligned} b_1 b_2; & \text{ if } b_1 = 1, \text{ then } Z \\ & \text{ if } b_2 = 1, \text{ then } X \end{aligned}$$