

Retinopathy of prematurity

Jing Chen · Lois E. H. Smith

Received: 2 November 2006 / Accepted: 24 January 2007 / Published online: 27 February 2007
© Springer Science+Business Media B.V. 2007

Abstract Retinopathy of prematurity (ROP) is a common blinding disease in children in the developed world despite current treatment, and is becoming increasingly prevalent in the developing world. ROP progresses in two phases. The first phase begins with delayed retinal vascular growth after birth and partial regression of existing vessels, followed by a second phase of hypoxia-induced pathological vessel growth. Two major risk factors of ROP are the use of oxygen and a decreased gestation period. Excessive oxygen contributes to ROP through regulation of vascular endothelial growth factor (VEGF). Suppression of VEGF by oxygen in phase I of ROP inhibits normal vessel growth, whereas elevated levels of VEGF induced by hypoxia in phase II of ROP precipitate pathological vessel proliferation. Insulin-like growth factor 1 (IGF-1) is a critical non-oxygen-regulated factor in ROP. We have found that serum levels of IGF-1 in premature babies directly correlate with the severity of clinical ROP. IGF-1 acts indirectly as a permissive factor by allowing maximal VEGF stimulation of vessel growth. Lack of IGF-1 in preterm infants prevents normal retinal vascular growth in phase I of ROP, despite the presence of VEGF. As infants mature, rising levels of IGF-1 in phase II of ROP allows VEGF stimulated pathological neovascularization. These findings suggest that restoration of IGF-1 to normal levels might be useful in preventing ROP in preterm infants.

Keywords Hypoxia · IGF-1 · Neovascularization · Prematurity · Retinopathy · VEGF

Introduction

Retinopathy of prematurity (ROP) is a major cause of blindness in children in the developing and developed world despite current surgical treatment in the late-stage of the disease [1]. Formerly known as retrolental fibroplasia, ROP was originally described in the 1940s by Terry who first connected the condition with premature birth [2]. At that time, no treatment for ROP was available. Major advances in ROP treatment came in the 1980s and 1990s, when cryotherapy and laser photocoagulation of the avascular retina were shown to be partially effective in preventing blindness in ROP infants. However, although these ablation treatments can reduce the incidence of blindness by 25% in infants with late-stage disease, the patients often still have poor visual acuity after treatment. Preventive and less destructive therapies for ROP would be much more desirable, and understanding the molecular basis of the disease is important to the development of such medical interventions.

Risk factors of ROP

First observed in the nursery [3, 4] and then supported by animal studies [5, 6], ROP was associated with excessive oxygen use shortly after the initial description of the disease. Controlled supplemental oxygen is now delivered to premature infants to maintain

J. Chen · L. E. H. Smith (✉)
Department of Ophthalmology, Harvard Medical School
and Children's Hospital Boston, 300 Longwood Avenue,
Boston, MA 02115, USA
e-mail: Lois.Smith@childrens.harvard.edu

adequate blood levels of oxygen [7]. However, even with monitored oxygen use, the number of infants with ROP has increased, most likely due to the increased survival of very low birth weight infants [8].

Besides oxygen, other factors that might contribute to ROP have been investigated over the years, including vitamin E and light exposure. Since premature infants have lower vitamin E levels in serum, the role of vitamin E in ROP was considered. However, conflicting results were reported in clinical studies of vitamin E and the efficacy of vitamin E prophylaxis remain inconclusive [9, 10]. Studies on light reduction have concluded that reduction of light exposure does not reduce the progression of ROP in either a mouse model of ROP [11] or in the clinic [12].

At present, oxygen use and gestational age/birth weight of the baby appear to be the major risk factors for ROP [13–15], although recent reports suggest that elevated glucose may also play a role [16–18].

Pathogenesis: two phases of ROP

ROP is a biphasic disease consisting of an initial phase of vessel loss followed by a second phase of vessel proliferation. For a better understanding of the process, one needs to examine the development of retinal vasculature. Retinal vascularization originates from the center of optic disc and progresses radially outwards towards the ora seratta. In the human, retinal blood vessel development is initiated during the fourth month of gestation and reaches the retinal periphery just before birth [19]. Therefore, infants born prematurely have incompletely vascularized retinas with a peripheral avascular zone. The gestational age at birth of the premature infant determines the area of the avascular zone.

In premature infants vascular growth that would normally occur in utero slows or ceases and is accompanied by regression of developed retinal vessels. The relative hyperoxia of the extrauterine environment as well as supplemental oxygen given to premature infants are thought to be responsible for this process. Normally in utero, the blood is only ~70% saturated compared to 100% in full term infants in room air. The normal PaO_2 in utero is 30 mm Hg, while a normal infant breathing room air will have a PaO_2 of 60–100 mm Hg [20]. As the infant matures, the non-vascularized retina becomes increasingly metabolically active and, in the absence of an adequate vascular system, leads to tissue hypoxia. The first phase of ROP occurs from birth to postmenstrual age approximately 30–32 weeks.

The second phase of ROP is characterized by hypoxia-induced retinal neovascularization and begins around 32–34 weeks postmenstrual age. This phase of ROP is similar to the neovascularizations in other proliferative retinopathies such as diabetic retinopathy. New vessels form at the junction between the vascularized retina and the avascular zone of the retina. Over time, this pathological growth of vessels produces a fibrous scar extending from the retina to the vitreous gel and lens. Retraction of this scar tissue can separate the retina from the retinal pigment epithelium, resulting in a retinal detachment and likely blindness.

Animal models of ROP: the murine model of oxygen-induced ROP

Much of our understanding regarding the underlying mechanism of the disease process comes from the use of animal models. Many animals such as mice, rats, kittens, and beagle pups have incompletely vascularized retinas at birth and resemble the immature retinal vascular development of premature infants.

We have developed a mouse model of ROP to study the molecular pathways in the disease as well as to take advantage of the genetic manipulation in mice [21]. Neonatal mice are exposed to 75% oxygen from postnatal day 7 until day 12. When mice are exposed to hyperoxia, vessel regression and the cessation of normal radial vessel growth occurs mimicking the first phase of ROP. The association between oxygen exposure and vaso-obliteration has also been observed in other animal models such as rats, kittens, and beagle pups [5, 6, 22, 23]. The extent of vaso-obliteration can be determined by measuring the non-perfused area in retinal whole mount.

Upon return to room air, the non-perfused portions of retina become hypoxic, thereby inducing the expression of angiogenic factors and resulting in retinal neovascularization. The neovascular phase in the oxygen-induced animal model is similar to the second phase of ROP in humans and, in addition, mimics certain aspects of proliferative diabetic retinopathy. The formation of vascular tufts are measured by quantifying neovascular nuclei extending into the vitreous in cross sections of retina, or by quantifying the area of vascular tufts in retinal whole mounts. The mouse model has proven useful in delineating the molecular changes in both phases of the neovascular eye diseases.

The roles of vascular endothelial growth factor (VEGF) and oxygen in ROP

Excessive oxygen use has been associated with development of ROP since the 1950s. It has become widely accepted that hyperoxia-induced vessel loss results in retinal hypoxia, which stimulates the release of factors that influence blood vessel growth [24]. It became clear that a growth factor or factors regulated by hypoxia and hyperoxia is important in the development of ROP. VEGF was discovered initially as a permeability factor [25] and later shown to be an endothelial cell mitogen [26]. VEGF mRNA increases with hypoxia [27] and plays a key role in tumor [28, 29] and retinal associated angiogenesis [30–32].

The role of VEGF in phase I of ROP

VEGF is essential in the development of the retinal vasculature. During normal retinal development, blood vessels grow from the optic nerve to the periphery. As the neural retina develops anterior to the vasculature, the increased oxygen demand of the developing neural tissue generates a wave of “physiological hypoxia” that precedes vessel growth [33]. In response to the hypoxia, astrocyte expression of VEGF stimulates blood vessel growth that follows the astrocyte template [34]. As new vessels form and retinal hypoxia decreases, VEGF expression and further vascular growth is reduced via a local feedback mechanism [35].

Supplemental oxygen to premature infants, however, interferes with normal VEGF driven vascular development. In phase I of the murine model of ROP, hyperoxia suppresses VEGF expression, resulting in the loss of the physiological wave of VEGF anterior to the growing vascular front [36]. Cessation of normal vessel growth and regression of existing vessels subsequently occurs. The hyperoxia-induced vaso-obliteration is caused by apoptosis of vascular endothelial cells and can be partially prevented by administration of exogenous VEGF or PlGF-1, a VEGFR-1 specific ligand [37–39]. This finding indicates that VEGF signaling through VEGFR-1 is required for survival of the immature retinal vasculature and explains, at least in part, the effect of hyperoxia on normal vessel development in ROP [39].

The role of VEGF in phase II of ROP

In the second phase of ROP, driven by hypoxia, there is a temporal and spatial relationship between VEGF

and proliferative retinopathy. Following oxygen-induced vessel loss and subsequent hypoxia, VEGF expression is increased in retina, resulting in pathological neovascularization [21, 36]. Inhibition of VEGF following the intravitreal injection of either an anti-VEGF antisense oligonucleotide or with a VEGF binding molecule (VEGF receptor/IgG chimera) in this phase significantly decreases the neovascular response [40, 41], indicating that VEGF is a critical factor contributing to retinal neovascularization. Studies with other animal models also support the central role of VEGF in ocular neovascularization [42–46]. These results correspond with the observations from clinics. Elevated VEGF levels are found in the vitreous of patients with retinal neovascularization [31, 47]. In retinal specimens from an ROP patient, VEGF expression was found to occur in a pattern consistent with that found in the mouse model of ROP [46]. Based on these and other studies, an anti-VEGF aptamer as well as an anti-VEGF antibody fragment injected intravitreally is now available to treat neovascularization associated with age-related macular degeneration and is in clinical trials for diabetic eye disease. Clinical trials are planned for evaluation of treatment of the proliferative phase of ROP with anti-VEGF intravitreal treatment.

The roles of growth hormone and insulin-like growth factor (GH/IGF-1) in ROP

Although oxygen, acting in part through VEGF, plays a central role in retinal vessel development and in ROP, it is important to note that other biochemical mediators are also involved in the pathogenesis of retinopathy. Inhibition of VEGF does not completely inhibit hypoxia-induced retinal neovascularization in the second phase of ROP. In the first phase of the disease, although oxygen alone can cause cessation of vessel growth and vaso-obliteration, the results from animal models clearly indicate that clinical ROP is multi-factorial. Despite controlled use of supplemental oxygen, the disease persists, suggesting that other factors related to prematurity itself are also at work.

The fact that prematurity is the most significant risk factor for ROP suggests that factors involved in growth and development are critical. Growth hormone (GH) and insulin-like growth factor-1 (IGF-1), which mediates many of the mitogenic aspects of GH, have been investigated as non-oxygen-regulated factors in this process. GH and IGF-1 have been suspected to play a role in retinal neovascularization since the discovery that pituitary ablation has an ameliorative effect on

proliferative diabetic retinopathy [48–50]. Using the ROP mouse model, we have determined that IGF-1 is critical to both phases of ROP, a fact supported by clinical results.

The role of IGF-1 in phase I of ROP

IGF-1 is critical to phase I of ROP and to the normal development of retinal vessels. The IGFs (IGF-1 and IGF-2) are important in fetal growth and development during all stages of pregnancy [51]. The serum concentration of IGF-1, but not IGF-2, increases with gestational age and correlates with fetal size [52, 53]. IGF-1 levels rise significantly in the third trimester of pregnancy, but after birth decrease due to the loss of IGF-1 provided by the placenta and the amniotic fluid [51].

We hypothesized that IGF-1 is critical to normal retinal vascular development, and that lack of IGF-1 is associated with lack of vascular growth and subsequent proliferative ROP. Results from animal studies show that normal retinal blood vessels grow more slowly in IGF-1 knockout mouse than in wild type controls, a pattern very similar to that seen in premature babies with ROP. In addition, IGF-1 controls maximum VEGF activation of the Akt endothelial cell survival pathway [54]. These findings help to explain how the loss of IGF-1 expression could cause ROP by preventing the normal survival of vascular endothelial cells.

The critical role of IGF in phase I of ROP is also supported by clinical observation from infants with ROP. The mean serum levels of IGF-1 in age-matched premature babies are directly correlated with the severity of clinical ROP [54–57]. IGF-1 appears to be as strong a determinant of risk for ROP as postmenstrual age at birth and birth weight [54, 55]. Low postnatal IGF-1 levels in preterm infants also correlates with brain development and might account for abnormal neural retinal function in ROP [56]. In patients with genetic defects of the GH/IGF-1 axis, very low level of IGF-1 directly causes decreased vascular density as evidenced by fewer vascular branching points [58]. This accumulated evidence suggests that low IGF-1 levels during development are associated with vessel loss and may be contributing to early vessel degeneration in phase I that sets the stage for hypoxia-induced proliferative retinopathy.

Taken together, these findings raise the possibility that early restoration of IGF-1 to uterine levels might prevent the disease by allowing normal retinal vascular development. Since normal vessel development of the retina precludes the development of ROP, the extent

of the later destructive phase of ROP is determined by phase I. If we can prevent the vessel loss in the phase I of ROP, the destructive second phase of neovascularization should not occur. Clinical trials are being planned to supplement IGF-1 and IGFBP-3 to in utero levels in premature infants in an effort to evaluate if restoration of IGF-1 to normal levels can prevent or reduce the severity of ROP.

The role of IGF-1 in phase II of ROP

The first study showing that IGF-1 is important in retinopathy came from work in the proliferative phase of ROP (phase II). In the second phase of ROP, retinal neovascularization is substantially reduced in transgenic mice expressing a GH receptor antagonist or normal mice with a somatostatin analogue that decreases GH release [59]. This inhibition of neovascularization by GH is mediated through inhibition of IGF-1, since systemic administration of IGF-1 completely restores neovascularization in normal mice with decreased GH release. Direct proof of the role of IGF-1 in the proliferative phase of ROP in mice was established using an IGF-1 receptor antagonist, which was found to suppress retinal neovascularization without altering VEGF levels induced in mouse ROP [60].

Other studies have investigated the role of both IGF-1 and insulin in the ROP mouse model with a vascular endothelial cell specific knockout of the IGF-1 receptor or insulin receptor [61]. Both types of transgenic mice showed substantial reduction in retinal neovascularization compared with control. In the mice with the insulin receptor knockout, the reduction of neovascularization was associated with a decrease in VEGF expression. These findings suggest that both insulin and IGF-1 signaling in the vascular endothelium may be involved in the regulation of retinal neovascularization.

The interaction of VEGF and IGF-1 in the evolution of ROP

With GH and IGF-1 inhibition in the proliferative phase of ROP, hypoxia-induced VEGF production is unchanged, indicating that IGF-1 does not directly act through VEGF under these physiological conditions. It was suggested that IGF-1 regulates retinal neovascularization at least in part through control of VEGF activation of p44/42 MAPK, establishing a hierarchical relationship between IGF-1 and VEGF receptors

[54, 60]. IGF allows maximal VEGF stimulation of new vessel growth. Reducing IGF-1 levels inhibit vessel growth despite the presence of VEGF. These studies suggest that IGF-1 serves a permissive function, and VEGF alone might not be sufficient for promoting vigorous retinal angiogenesis.

Based on the understanding of the roles of VEGF and IGF-1 in both phases of the disease, a rationale for the evolution of ROP has emerged (Fig. 1). Retinal vessel growth requires both IGF-1 and VEGF. In premature infants, IGF-1, which is normally supplied by placenta and the amniotic fluid, is at very low levels after birth, since the premature liver cannot replace the loss. Blood vessel growth therefore slows or stops since IGF-1 is needed to allow VEGF signaling for vascular endothelial growth and survival. When supplement oxygen is given to the preterm infants, VEGF itself is suppressed. Thus, both prematurity and oxygen are contributing to the suppression of normal vessel growth and vessel loss in phase I of ROP. As the infant grows and the retina matures without an adequate supply of blood oxygen, it becomes hypoxic and induces increased expression of VEGF. In the mean time, as the infant's organs and systems continue to mature, IGF-1 levels rise, suddenly allowing the now-high-levels of VEGF to stimulate blood vessels growth. The result is the destructive neovascular proliferation in the phase II of ROP, leading to blindness.

Future therapeutic targets

The discovery of the importance of VEGF and IGF-1 in the development of ROP is a step forward in our understanding of the pathogenesis of this disease. These studies suggest a number of ways to intervene medically in the disease process. The use of anti-VEGF therapy is the first medical treatment for neovascularization in age-related macular degeneration and is likely to be useful for proliferative retinopathy. However, prevention of vessel loss will be even more important in the treatment of ROP since the extent of the second destructive phase of ROP is determined by the amount of vessel loss in the first phase. The finding


that late development of ROP is associated with low levels of IGF-1 after premature birth suggests that physiological replacement of IGF-1 to levels found in utero might prevent the disease by allowing normal vascular development. In addition, the use of a specific agonist to VEGFR-1, PIGF-1, might be used early in the disease process to prevent vessel loss without promoting proliferative disease.

The current understanding of ROP pathogenesis also makes clear that timing is critical in any medical intervention, since the two phases of ROP require very different approaches. Inhibition of either VEGF or IGF-1 early after birth can prevent normal blood vessel growth while, at the second phase, might prevent pathological neovascularization. Similarly, providing VEGF or IGF-1 early on might promote vessel growth whereas late supplementation in the neovascular phase could exacerbate the disease. In the fragile neonate, any intervention must be made very carefully to promote normal physiological development of both blood vessels and other tissues.

Investigations of rare diseases with similar retinal vessel defects as ROP might provide insight to the treatment of ROP. Other conditions with incomplete retinal vascularization are Norrie's disease, familial exudative vitreoretinopathy and incontinentia pigmenti. One form of familial exudative vitreoretinopathy is caused by Frizzled-4 gene defect [62]. Both norrin, the protein product of norrie's disease gene, and Frizzled-4, a Wnt receptor gene, cause similar defects in humans and mice with incomplete peripheral retina vascularization and the lack of deep retinal vessels [62–64]. It has been shown recently that the norrin ligand binds to frizzled-4, a Wnt receptor [64, 65] and ectopic norrin restores retinal vessels in Norrie disease mice [66]. These findings raise the possibility of a new pharmacological therapy for eye disease with incomplete vascularization of peripheral retina such as ROP.

There are currently many active areas of research that suggest manipulation of currently approved pharmacological interventions or even dietary interventions may help to prevent the ischemia that results in the destructive aspects of neovascularization in proliferative retinopathy including ROP. Anti-inflam-

Fig. 1 Schematic representation of IGF-I/VEGF control of blood vessel development in ROP. (A) In utero, (B) Premature birth, (C) Retinal maturation, (D) Retinal neovascularization. NI: normal



matory agents may also be important in future preventative treatment of retinal neovascularization. Further studies on disease mechanism and development of strategies to allow normal retinal and brain development may lead to a significant reduction in the incidence of ROP.

References

- Silverman W (1980) Retrolental fibroplasia: a modern parable. Grune and Stratton, New York
- Terry TL (1944) Retrolental fibroplasia in the premature infant: V. Further studies on fibroplastic overgrowth of the persistent tunica vasculosa lentis. *Trans Am Ophthalmol Soc* 42:383–396
- Campbell K (1951) Intensive oxygen therapy as a possible cause of retrolental fibroplasia; a clinical approach. *Med J Aust* 2:48–50
- Patz A, Hoeck LE, De La Cruz E (1952) Studies on the effect of high oxygen administration in retrolental fibroplasia. I nursery observations. *Am J Ophthalmol* 35:1248–1253
- Ashton N, Ward B, Serpell G (1953) Role of oxygen in the genesis of retrolental fibroplasia; a preliminary report. *Br J Ophthalmol* 37:513–520
- Ashton N, Ward B, Serpell G (1954) Effect of oxygen on developing retinal vessels with particular reference to the problem of retrolental fibroplasia. *Br J Ophthalmol* 38:397–432
- Kinsey VE, Arnold HJ, Kalina RE, Stern L, Stahlman M, Odell G, Driscoll JM Jr, Elliott JH, Payne J, Patz A (1977) PaO₂ levels and retrolental fibroplasia: a report of the cooperative study. *Pediatrics* 60:655–668
- Flynn JT (1983) Acute proliferative retrolental fibroplasia: multivariate risk analysis. *Trans Am Ophthalmol Soc* 81:549–591
- Johnson L, Quinn GE, Abbasi S, Gerdes J, Bowen FW, Bhutani V (1995) Severe retinopathy of prematurity in infants with birth weights less than 1250 grams: incidence and outcome of treatment with pharmacologic serum levels of vitamin E in addition to cryotherapy from 1985 to 1991. *J Pediatr* 127:632–639
- Phelps DL, Rosenbaum AL, Isenberg SJ, Leake RD, Dorey FJ (1987) Tocopherol efficacy and safety for preventing retinopathy of prematurity: a randomized, controlled, double-masked trial. *Pediatrics* 79:489–500
- Wesolowski E, Smith LE (1994) Effect of light on oxygen-induced retinopathy in the mouse. *Invest Ophthalmol Vis Sci* 35:112–119
- Reynolds JD, Hardy RJ, Kennedy KA, Spencer R, van Heuven WA, Fielder AR (1998) Lack of efficacy of light reduction in preventing retinopathy of prematurity. Light reduction in retinopathy of prematurity (LIGHT-ROP) cooperative group. *N Engl J Med* 338:1572–1576
- Lutty GA, Chan-Ling T, Phelps DL, Adamis AP, Berns KI, Chan CK, Cole CH, D'Amore PA, Das A, Deng WT, Dobson V, Flynn JT, Friedlander M, Fulton A, Good WV, Grant MB, Hansen R, Hauswirth WW, Hardy RJ, Hinton DR, Hughes S, McLeod DS, Palmer EA, Patz A, Penn JS, Raisler BJ, Repka MX, Saint-Geniez M, Shaw LC, Shima DT, Smith BT, Smith LE, Tahija SG, Tasman W, Trese MT (2006) Proceedings of the third international symposium on retinopathy of prematurity: an update on ROP from the lab to the nursery (November 2003, Anaheim, California). *Mol Vis* 12:532–580
- Smith LE (2003) Pathogenesis of retinopathy of prematurity. *Semin Neonatol* 8:469–473
- Tasman W, Patz A, McNamara JA, Kaiser RS, Trese MT, Smith BT (2006) Retinopathy of prematurity: the life of a lifetime disease. *Am J Ophthalmol* 141:167–174
- Blanco CL, Baillargeon JG, Morrison RL, Gong AK (2006) Hyperglycemia in extremely low birth weight infants in a predominantly hispanic population and related morbidities. *J Perinatol* 26:737–741
- Ertl T, Gyarmati J, Gaal V, Szabo I (2006) Relationship between hyperglycemia and retinopathy of prematurity in very low birth weight infants. *Biol Neonate* 89:56–59
- Garg R, Agthe AG, Donohue PK, Lehmann CU (2003) Hyperglycemia and retinopathy of prematurity in very low birth weight infants. *J Perinatol* 23:186–194
- Roth AM (1977) Retinal vascular development in premature infants. *Am J Ophthalmol* 84:636–640
- Bell EF, Klein JM (1994) Iowa neonatology handbook: pulmonary comments on oxygen toxicity and retinopathy (ROP) in the premature infant. online: <http://www.ui-healthcare.com/depts/med/pediatrics/iowanonatologyhandbook/general/commentsoxygen.html>
- Smith LE, Wesolowski E, McLellan A, Kostyk SK, D'Amato R, Sullivan R, D'Amore PA (1994) Oxygen-induced retinopathy in the mouse. *Invest Ophthalmol Vis Sci* 35:101–111
- Flower RW (1990) Perinatal ocular physiology and ROP in the experimental animal model. *Doc Ophthalmol* 74:153–162
- Penn JS, Tolman BL, Henry MM (1994) Oxygen-induced retinopathy in the rat: relationship of retinal nonperfusion to subsequent neovascularization. *Invest Ophthalmol Vis Sci* 35:3429–3435
- Patz A (1982) Clinical and experimental studies on retinal neovascularization. XXXIX Edward Jackson memorial lecture. *Am J Ophthalmol* 94:715–743
- Senger DR, Galli SJ, Dvorak AM, Perruzzi CA, Harvey VS, Dvorak HF (1983) Tumor cells secrete a vascular permeability factor that promotes accumulation of ascites fluid. *Science* 219:983–985
- Leung DW, Cachianes G, Kuang WJ, Goeddel DV, Ferrara N (1989) Vascular endothelial growth factor is a secreted angiogenic mitogen. *Science* 246:1306–1309
- Plate KH, Breier G, Weich HA, Risau W (1992) Vascular endothelial growth factor is a potential tumour angiogenesis factor in human gliomas in vivo. *Nature* 359:845–848
- Kim KJ, Li B, Winer J, Armanini M, Gillett N, Phillips HS, Ferrara N (1993) Inhibition of vascular endothelial growth factor-induced angiogenesis suppresses tumour growth in vivo. *Nature* 362:841–844
- Shweiki D, Itin A, Soffer D, Keshet E (1992) Vascular endothelial growth factor induced by hypoxia may mediate hypoxia-initiated angiogenesis. *Nature* 359:843–845
- Adamis AP, Shima DT, Yeo KT, Yeo TK, Brown LF, Berse B, D'Amore PA, Folkman J (1993) Synthesis and secretion of vascular permeability factor/vascular endothelial growth factor by human retinal pigment epithelial cells. *Biochem Biophys Res Commun* 193:631–638
- Aiello LP, Avery RL, Arrigg PG, Keyt BA, Jampel HD, Shah ST, Pasquale LR, Thieme H, Iwamoto MA, Park JE et al (1994) Vascular endothelial growth factor in ocular fluid of patients with diabetic retinopathy and other retinal disorders. *N Engl J Med* 331:1480–1487

32. Aiello LP, Northrup JM, Keyt BA, Takagi H, Iwamoto MA (1995a) Hypoxic regulation of vascular endothelial growth factor in retinal cells. *Arch Ophthalmol* 113:1538–1544
33. Chan-Ling T, Gock B, Stone J (1995) The effect of oxygen on vasoformative cell division. Evidence that ‘physiological hypoxia’ is the stimulus for normal retinal vasculogenesis. *Invest Ophthalmol Vis Sci* 36:1201–1214
34. Dorrell MI, Aguilar E, Friedlander M (2002) Retinal vascular development is mediated by endothelial filopodia, a preexisting astrocytic template and specific R-cadherin adhesion. *Invest Ophthalmol Vis Sci* 43:3500–3510
35. Stone J, Itin A, Alon T, Pe’er J, Gnessin H, Chan-Ling T, Keshet E (1995) Development of retinal vasculature is mediated by hypoxia-induced vascular endothelial growth factor (VEGF) expression by neuroglia. *J Neurosci* 15:4738–4747
36. Pierce EA, Avery RL, Foley ED, Aiello LP, Smith LE (1995) Vascular endothelial growth factor/vascular permeability factor expression in a mouse model of retinal neovascularization. *Proc Natl Acad Sci USA* 92:905–909
37. Alon T, Hemo I, Itin A, Pe’er J, Stone J, Keshet E (1995) Vascular endothelial growth factor acts as a survival factor for newly formed retinal vessels and has implications for retinopathy of prematurity. *Nat Med* 1:1024–1028
38. Pierce EA, Foley ED, Smith LE (1996) Regulation of vascular endothelial growth factor by oxygen in a model of retinopathy of prematurity. *Arch Ophthalmol* 114:1219–1228
39. Shih SC, Ju M, Liu N, Smith LE (2003) Selective stimulation of VEGFR-1 prevents oxygen-induced retinal vascular degeneration in retinopathy of prematurity. *J Clin Invest* 112:50–57
40. Aiello LP, Pierce EA, Foley ED, Takagi H, Chen H, Riddle L, Ferrara N, King GL, Smith LE (1995b) Suppression of retinal neovascularization in vivo by inhibition of vascular endothelial growth factor (VEGF) using soluble VEGF-receptor chimeric proteins. *Proc Natl Acad Sci USA* 92:10457–10461
41. Robinson GS, Pierce EA, Rook SL, Foley E, Webb R, Smith LE (1996) Oligodeoxynucleotides inhibit retinal neovascularization in a murine model of proliferative retinopathy. *Proc Natl Acad Sci USA* 93:4851–4856
42. Adamis AP, Shima DT, Tolentino MJ, Gragoudas ES, Ferrara N, Folkman J, D’Amore PA, Miller JW (1996) Inhibition of vascular endothelial growth factor prevents retinal ischemia-associated iris neovascularization in a non-human primate. *Arch Ophthalmol* 114:66–71
43. Donahue ML, Phelps DL, Watkins RH, LoMonaco MB, Horowitz S (1996) Retinal vascular endothelial growth factor (VEGF) mRNA expression is altered in relation to neovascularization in oxygen induced retinopathy. *Curr Eye Res* 15:175–184
44. Miller JW, Adamis AP, Shima DT, D’Amore PA, Moulton RS, O’Reilly MS, Folkman J, Dvorak HF, Brown LF, Berse B et al (1994) Vascular endothelial growth factor/vascular permeability factor is temporally and spatially correlated with ocular angiogenesis in a primate model. *Am J Pathol* 145:574–584
45. Stone J, Chan-Ling T, Pe’er J, Itin A, Gnessin H, Keshet E (1996) Roles of vascular endothelial growth factor and astrocyte degeneration in the genesis of retinopathy of prematurity. *Invest Ophthalmol Vis Sci* 37:290–299
46. Young TL, Anthony DC, Pierce E, Foley E, Smith LE (1997) Histopathology and vascular endothelial growth factor in untreated and diode laser-treated retinopathy of prematurity. *J Aapos* 1:105–110
47. Adamis AP, Miller JW, Bernal MT, D’Amico DJ, Folkman J, Yeo TK, Yeo KT (1994) Increased vascular endothelial growth factor levels in the vitreous of eyes with proliferative diabetic retinopathy. *Am J Ophthalmol* 118:445–450
48. Poulsen JE (1953) Recovery from retinopathy in a case of diabetes with Simmonds’ disease. *Diabetes* 2:7–12
49. Sharp PS, Fallon TJ, Brazier OJ, Sandler L, Joplin GF, Kohner EM (1987) Long-term follow-up of patients who underwent yttrium-90 pituitary implantation for treatment of proliferative diabetic retinopathy. *Diabetologia* 30:199–207
50. Wright AD, Kohner EM, Oakley NW, Hartog M, Joplin GF, Fraser TR (1969) Serum growth hormone levels and the response of diabetic retinopathy to pituitary ablation. *Br Med J* 2:346–348
51. Langford K, Nicolaides K, Miell JP (1998) Maternal and fetal insulin-like growth factors and their binding proteins in the second and third trimesters of human pregnancy. *Hum Reprod* 13:1389–1393
52. Lassarre C, Hardouin S, Daffos F, Forestier F, Frankenne F, Binoux M (1991) Serum insulin-like growth factors and insulin-like growth factor binding proteins in the human fetus. Relationships with growth in normal subjects and in subjects with intrauterine growth retardation. *Pediatr Res* 29:219–225
53. Reece EA, Wiznitzer A, Le E, Homko CJ, Behrman H, Spencer EM (1994) The relation between human fetal growth and fetal blood levels of insulin-like growth factors I and II, their binding proteins, and receptors. *Obstet Gynecol* 84:88–95
54. Hellstrom A, Perruzzi C, Ju M, Engstrom E, Hard AL, Liu JL, Albertsson-Wikland K, Carlsson B, Niklasson A, Sjobell L, LeRoith D, Senger DR, Smith LE (2001) Low IGF-I suppresses VEGF-survival signaling in retinal endothelial cells: direct correlation with clinical retinopathy of prematurity. *Proc Natl Acad Sci USA* 98:5804–5808
55. Hellstrom A, Engstrom E, Hard AL, Albertsson-Wikland K, Carlsson B, Niklasson A, Lofqvist C, Svensson E, Holm S, Ewald U, Holmstrom G, Smith LE (2003) Postnatal serum insulin-like growth factor I deficiency is associated with retinopathy of prematurity and other complications of premature birth. *Pediatrics* 112:1016–1020
56. Lofqvist C, Engstrom E, Sigurdsson J, Hard AL, Niklasson A, Ewald U, Holmstrom G, Smith LE, Hellstrom A (2006) Postnatal head growth deficit among premature infants parallels retinopathy of prematurity and insulin-like growth factor-1 deficit. *Pediatrics* 117:1930–1938
57. Smith LE (2004) Pathogenesis of retinopathy of prematurity. *Growth Horm IGF Res* 14(Suppl A):S140–S144
58. Hellstrom A, Carlsson B, Niklasson A, Segnestam K, Boguszewski M, de Lacerda L, Savage M, Svensson E, Smith L, Weinberger D, Albertsson Wikland K, Laron Z (2002) IGF-I is critical for normal vascularization of the human retina. *J Clin Endocrinol Metab* 87:3413–3416
59. Smith LE, Kopchick JJ, Chen W, Knapp J, Kinose F, Daley D, Foley E, Smith RG, Schaeffer JM (1997) Essential role of growth hormone in ischemia-induced retinal neovascularization. *Science* 276:1706–1709
60. Smith LE, Shen W, Perruzzi C, Soker S, Kinose F, Xu X, Robinson G, Driver S, Bischoff J, Zhang B, Schaeffer JM, Senger DR (1999) Regulation of vascular endothelial growth factor-dependent retinal neovascularization by insulin-like growth factor-1 receptor. *Nat Med* 5:1390–1395
61. Kondo T, Vicent D, Suzuma K, Yanagisawa M, King GL, Holzenberger M, Kahn CR (2003) Knockout of insulin and IGF-1 receptors on vascular endothelial cells protects against retinal neovascularization. *J Clin Invest* 111:1835–1842

62. Robitaille J, MacDonald ML, Kaykas A, Sheldahl LC, Zeisler J, Dube MP, Zhang LH, Singaraja RR, Guernsey DL, Zheng B, Siebert LF, Hoskin-Mott A, Trese MT, Pimstone SN, Shastry BS, Moon RT, Hayden MR, Goldberg YP, Samuels ME (2002) Mutant frizzled-4 disrupts retinal angiogenesis in familial exudative vitreoretinopathy. *Nat Genet* 32:326–330
63. Rehm HL, Zhang DS, Brown MC, Burgess B, Halpin C, Berger W, Morton CC, Corey DP, Chen ZY (2002) Vascular defects and sensorineural deafness in a mouse model of Norrie disease. *J Neurosci* 22:4286–4292
64. Xu Q, Wang Y, Dabdoub A, Smallwood PM, Williams J, Woods C, Kelley MW, Jiang L, Tasman W, Zhang K, Nathans J (2004) Vascular development in the retina and inner ear: control by Norrin and Frizzled-4, a high-affinity ligand–receptor pair. *Cell* 116:883–895
65. Niehrs C (2004) Norrin and frizzled; a new vein for the eye. *Dev Cell* 6:453–454
66. Ohlmann A, Scholz M, Goldwich A, Chauhan BK, Hudl K, Ohlmann AV, Zrenner E, Berger W, Cvekl A, Seeliger MW, Tamm ER (2005) Ectopic norrin induces growth of ocular capillaries and restores normal retinal angiogenesis in Norrie disease mutant mice. *J Neurosci* 25:1701–1710