

P-61 Impacts of salvage chemotherapy after nivolumab therapy (NIVO): A REVIVE substudy

H. Matsuoka¹, Y. Narita², T. Misumi³, Y. Sakamoto⁴, T. Kawakami⁵, H. Tanioka⁶, T. Matsushima⁷, H. Miwa⁸, H. Shoji⁹, A. Ishiguro¹⁰, S. Fushida¹¹, K. Miura¹², T. Yamada¹³, K. Shinozaki¹⁴, T. Mizukami¹⁵, T. Moriwaki¹⁶, S. Mitani¹⁷, M. Nakamura¹⁸, K. Muro², T. Nishina¹⁹

¹Fujita Health University, Toyoake, Japan; ²Department of Clinical Oncology, Aichi Cancer Center Hospital, Nagoya, Japan; ³Department of Data Science, National Cancer Center Hospital East, Kashiwa, Japan; ⁴Department of Medical Oncology, Osaka Citizen Hospital, Osaka, Japan; ⁵Division of Gastrointestinal Oncology, Shizuoka Cancer Center, Sunto-Gun, Japan; ⁶Kawasaki Medical School, Kurashiki, Japan; ⁷Saitama Cancer Center, Komuro, Japan; ⁸Hyogo College of Medicine, Nishinomiya, Japan; ⁹Department of Gastrointestinal Medical Oncology, National Cancer Center Hospital, Tokyo, Japan; ¹⁰Department of Medical Oncology, Teine Keijinkai Hospital, Sapporo, Japan; ¹¹Department of Gastroenterologic Surgery, Kanazawa University Graduate School of Medical Sciences, Kanazawa, Japan; ¹²Department of Internal Medicine, Himeji Red Cross Hospital, Himeji, Japan; ¹³Kanagawa Cancer Center, Yokohama, Japan; ¹⁴Hiroshima Prefectural Hospital, Hiroshima, Japan; ¹⁵Department of Clinical Oncology, St. Marianna University School of Medicine, Kawasaki, Japan; ¹⁶Department of Gastroenterology, Faculty of Medicine, University of Tsukuba, Tsukuba, Japan; ¹⁷Kindai University Faculty of Medicine, Osaka-Sayama, Japan; ¹⁸Department of Gastroenterology, Sapporo City General Hospital, Sapporo, Japan; ¹⁹National Hospital Organization Shikoku Cancer Center, Ehime, Japan

Background: The primary endpoints met in the REVIVE study have been previously reported, demonstrating the chemotherapeutic efficacy after the progression on NIVO therapy in patients with advanced gastric cancer (AGC). Therefore, the current study evaluated the survival from the NIVO therapy initiation in all registered participants of the REVIVE trial.

Methods: The REVIVE trial was a prospective, multicenter, observational study evaluating the efficacy and safety of chemotherapy for NIVO-refractory or NIVO-intolerant patients with AGC (UMIN000032182). We primarily register the patients who underwent NIVO therapy as primary registration, and the patients formally were registered as formal registration. The previously reported main study analyzed data of formally registered patients who underwent chemotherapy with irinotecan, trifluridine/tipiracil hydrochloride, or oxaliplatin combination regimens. In this study, patients who discontinued NIVO therapy for any reason at data cutoff among primarily registered patients were selected. The survival of patients who received the best supportive care (cohort A) was compared to those included in the main study (cohort B).

Results: Of 395 primarily registered patients, 108 patients in cohort A and 199 patients in cohort B were included, respectively. Those receiving other chemotherapeutic regimens (N = 47) or continuing NIVO therapy (N = 38) were excluded. Median overall survival (OS) and time to treatment failure (TTF) were 9.3 (95% confidence interval [CI], 8.3–10.2) and 1.8 (95%CI, 1.6–2.2) months at 234 and 307 events, respectively, from the initiation of NIVO therapy in the whole population. The objective response rate (ORR) and disease control rate (DCR) were 9.1% and 43.0%, respectively. Patients in cohort B had significantly better prognosis in OS (median, 12.2 vs. 4.8 months; hazards ratio [HR], 0.43 [95%CI 0.34–0.57]; $p < 0.01$). However, the difference in short-term efficacies was not observed: ORR, 6.5% vs. 10.6%; DCR, 38.0% vs. 45.7%; and median TTF, 1.9 vs. 1.8 months (HR, 1.06 [95%CI 0.84–1.35]; $p = 0.62$) (cohort A vs. B). The post-progression survival (PPS) from the date of NIVO therapy discontinuation was significantly better in cohort B than in cohort A (median PPS, 8.1 vs. 1.9 months; HR, 0.22 [95%CI 0.17–0.30]; $p < 0.01$). The proportion of patients who received the best supportive care after NIVO therapy was similar, regardless of the effectiveness of NIVO therapy (transition rate: 25% in complete response or partial response [responders, N = 28] and 33% in stable or progressive disease [non-responders, N = 267]). The difference of OS from the initiation of NIVO therapy in responders was not observed between cohorts A and B; however, the OS of cohort B in non-responders was significantly longer than that of cohort A (median OS, 10.8 vs. 4.8 months; HR, 0.60 [95%CI 0.52–0.69]; $p < 0.01$).

Conclusions: Salvage chemotherapy as much as possible after NIVO therapy could improve the AGC prognosis.

Legal entity responsible for the study: Comprehensive Support of Project for Oncology Research (CSPOR) of Public Health Research Foundation.

Funding: ONO PHARMACEUTICAL CO., LTD. Bristol-Myers Squibb K.K.

Disclosures: Y. Narita: Honoraria (self): ONO, BMS, AstraZeneca; Speaker Bureau / Expert testimony: ONO, BMS, AstraZeneca. T. Kawakami: Honoraria (self): Taiho Pharmaceutical, Ono pharmaceutical, Bristol-Myers Squibb, Bayer. K. Shinozaki: Honoraria (self): ONO Pharmaceutical CO., LTD., TAIHO Pharmaceutical Co., Ltd, Eli Lilly Japan K.K. . T. Mizukami: Speaker Bureau / Expert testimony: TAIHO pharmaceutical, Eli Lilly Japan, Ono pharmaceutical, Otsuka Pharmaceutical Factory, Asahi Kasei Pharmaceutical, Merck Biopharma, Takeda pharmaceutical, Chugai pharmaceutical, Bayer, Bristol Myers Squibb; Research grant / Funding (self): TAIHO pharmaceutical, Eli Lilly Japan. T. Moriwaki: Speaker Bureau / Expert testimony: Taiho Pharmaceutical, Chugai Pharma; Research grant / Funding (institution): Taiho Pharmaceutical, Chugai Pharma. S. Mitani: Honoraria (self): Taiho Pharmaceutical Co., Ono Pharmaceutical Co.; Advisory / Consultancy: Chugai Pharmaceutical Co.; Research grant / Funding (institution): Taiho Pharmaceutical Co., Eli Lilly. M. Nakamura: Honoraria (self): Bayer Yakuhin, Bristol-Myers Squibb, Chugai, Daiichi Sankyo, Eli Lilly Japan, Ono, Sanofi, Taiho, Takeda, Yakult Honsha. K. Muro: Honoraria (self): Eli Lilly, Chugai, Takeda, Ono, Taiho, Sanofi, Bristol-Myers Squibb, and Bayer; Advisory / Consultancy: Amgen, AstraZeneca, Ono, and Chugai; Research grant / Funding (institution): Astellas, Amgen, Solasia Pharma, Sanofi, Daiichi Sankyo, Parexel International, Taiho, MSD, Merck Biopharma, Pfizer, Eisai, Novartis, and Ono. T. Nishina: Honoraria

(self): Ono, Bristol Myers Squibb; Research grant / Funding (institution): Ono, Bristol Myers Squibb. All other authors have declared no conflicts of interest.

<https://doi.org/10.1016/j.annonc.2022.04.151>

P-62 Characterization and management of cholangiocarcinoma in a tertiary hospital with a high volume of patients

I. Gallego Jimenez, C. Garcia Grove, F. Tallafigo, L. Limon

Hospital Virgen del Rocío, IBIS, Sevilla, Spain

Background: The objective of this study is to evaluate the epidemiological characteristics, potential risk factors and local/systemic treatments in a tertiary care hospital with a high volume of patients.

Methods: Unicentric retrospective study; The patients were seen in consultation from January 2020 to January 2022, this being the end date of follow-up. Demographic variables related to tumor, treatment, and death were also collected. For descriptive statistics, percentages with frequencies and medians were used. Survival was estimated using Kaplan-Meier plots. The SPSS 20.0 program was used for statistical analysis.

Results: Data from a total of 44 patients were collected. 70.7% men and 29.3% women. All Caucasian. The highest incidence peak was in the decade between 50 and 60 years (59%), with the rest of the cases in the decade between 40 and 50 years. There was no significant history of prior smoking, alcohol consumption, prior pancreatitis, history of cholangitis, or heavy metal exposure. However, in our series two personal antecedents stand out: 1) 48% of the patients were type 2 diabetics and 23% type 1 diabetics and 2) 61.3% of the patients prior to diagnosis had periodontal disease. 90.2% of the histology was adenocarcinoma, 53.7% of the tumors being resectable, the rest presenting advanced disease. In the case of local surgery, partial hepatectomy with lymphadenectomy was the most used (68%), capecitabine the adjuvant treatment of choice in operated patients (82.3%) and the treatment scheme for advanced disease Cisplatin in combination with Gemcitabine (82.9%), obtaining partial response in 54.5%, stabilization in 18.2% and disease progression in 27.3% in the first tumor evaluation performed. NGS was performed on 100% of the patients, finding potentially actionable mutations: KRAS G12C mutation in 2 patients, IDH mutation in 1 patient, and microsatellite instability in 2 patients. Regarding germline mutations, these were performed in 12 patients due to family history, finding mutations related to the ATM gene (2), BRCA2 (1) and p53 (1) in 4 of them.

Conclusions: The oncological results observed in our series are comparable to those described in the literature. Its etiology is uncertain, advancing the age of diagnosis and requiring the study of more predictive and prognostic factors. Our study reaffirms the role of NGS in this disease, with the intention of increasing future treatment options in this pathology.

Editorial acknowledgement: University Hospital Virgen del Rocío, Seville.

Legal entity responsible for the study: The authors.

Funding: Has not received any funding.

Disclosures: All authors have declared no conflicts of interest.

<https://doi.org/10.1016/j.annonc.2022.04.152>

P-63 Efficacy, safety and prognostic subgroups for outcome with regorafenib in patients with refractory metastatic colorectal cancer in the real-world setting: The CORE study

N. Martínez-Lago¹, B. Carnero Lopez², J. de la Cámara Gómez³, F. Vázquez-Rivera⁴, A. Fernández Montes⁵, A. Cousillas Castiñeiras⁶, N. de Dios⁷, R. Varela Ponte⁸, M. Reboredo-Lopez³

¹Complejo Hospitalario Universitario, A Coruña, Spain; ²Hospital Universitario Lucas Augusti, Lugo, Spain; ³University Hospital A Coruña, A Coruña, Spain; ⁴Translational Medical Oncology Group, Oncomet, University Hospital, Santiago de Compostela, Galicia, Spain; ⁵Complejo Universitario Hospitalario de Ourense, Ourense, Spain; ⁶Complejo Universitario Hospitalario de Pontevedra, Pontevedra, Spain; ⁷Complejo Universitario Hospitalario de Pontevedra, Pontevedra, Spain; ⁸Complejo Hospitalario Universitario de Santiago, Santiago de Compostela, A Coruña, Spain

Background: CORRECT trial has shown a survival benefit for regorafenib over placebo in patients with metastatic colorectal cancer (mCRC) that progressed after standard therapies. We evaluated survival, safety and prognostic subgroups in patients treated with regorafenib in a real-life setting.

Methods: Retrospective, multicenter, observational study of patients (pts) with mCRC treated with Regorafenib (REG) after failure to standard therapies as part of routine clinical practice at 7 hospitals from the Galician Research Group on Digestive Tumors (GITuD).

Results: We recorded 130 pts treated with REG between September 2013 to December 2019. Median age was 63 years (range 27-79), 94.6% ECOG PS0-1, 55.4% RASmt and 1.5% BRAFmt, 18% time since initial diagnosis 3 metastatic locations and 75% liver metastases. Prior therapy included a median of 3 lines of treatment (range