

P-54 Phase 2a study of NT-17, a long-acting interleukin-7, plus pembrolizumab: Cohort of subjects with checkpoint inhibitor-naïve advanced MSS-colorectal cancer

R. Kim¹, H. Mamdani², M. Barve³, M. Johnson⁴, I. Sahin¹, S. Kopetz⁵, S. Yang⁶, B. Lee⁶, T. Adebajo⁶, R. Georgevitch⁶, S. Ferrando-Martinez⁶, M. Chaney⁷, J. Fan⁶, A. Naing⁵

¹Moffitt Cancer Center, Tampa, United States; ²Karmanos Cancer Institute, Detroit, United States; ³Mary Crowley Cancer Research, Dallas, United States; ⁴Sarah Cannon Research Institute/Tennessee Oncology, Nashville, United States; ⁵University of Texas MD Anderson Cancer Center, Houston, United States; ⁶NeolmmuneTech, Inc., Rockville, United States; ⁷Merck & Co., Inc., Kenilworth, United States

Background: Checkpoint inhibitor (CPI) monotherapy is ineffective for microsatellite stable colorectal cancer (MSS-CRC). NT-17 (efineptakin alfa) is a long-acting IL-7 that can increase T-cell infiltration in the tumor microenvironment (TME). We hypothesize that NT-17 may create a favorable immune-reactive TME to enhance the efficacy of CPI when combined with pembrolizumab (pembro).

Methods: This is an open-label, phase 2a study in subjects with relapsed/refractory (R/R) tumors, including CPI-naïve R/R MSS-CRC. Subjects were enrolled following Simon's 2-stage minimax design; 17 were enrolled in the first stage, and 8 additional subjects were enrolled for the second stage. Subjects received the recommended-phase-2-dose of NT-17 intramuscularly at 1200 µg/kg every 6 weeks (Q6W) plus pembro 200 mg intravenously Q3W. Preliminary anti-tumor activity based on Overall Response Rate (ORR) was assessed by Response Evaluation Criteria in Solid Tumors (RECIST) v1.1 and iRECIST. Biomarker analyses in peripheral blood and tumor biopsy were performed.

Results: As of 14-Jan-2022, 28 subjects were enrolled in the CPI-naïve R/R MSS-CRC cohort. Median age was 56.0 years [37-81], with ECOG PS 0 (28.6%) and 1 (71.4%). Twenty-three (82%) subjects received ≥ 2 prior therapies. All subjects had metastatic or locally advanced disease at enrollment. The median duration of follow-up was 5.3 months. Among these 25 evaluable subjects ORR and disease control rate (DCR) was 4% (1/25 subjects) and 40% (10/25) per RECIST v1.1; 12% (3/25 subjects) and 44% (11/25) per iRECIST. In addition to 3 subjects with iPR, 11 subjects are still ongoing to follow up responders. Interestingly, in patients with rectal malignancy both ORR and DCR by iRECIST were 50% (2/4). The ORR and DCR by iRECIST were 25% (1/4) and 50% (2/4) in 2L; 17% (1/6) and 33% (2/6) in 3L; and 13% (1/8) and 50% (4/8) in 4L. Among subjects without/with ≥ 2 liver mets, the ORR was 33.3% (2/6) vs 5.3% (1/19); DCR was 83.3% (5/6) vs 31.6% (6/19) and PFS was 11.6 weeks vs 6.1 weeks. All subjects with responses continue on treatment. Treatment-related adverse events (trAEs) occurred in 27 (96.4%) subjects, 12 (42.8%) G1-2 events and 14 (50%) G3 events; 1 (3.6%) G4 and no G5 trAEs were reported. No subjects discontinued from the study due to trAE. One iPR subject with available biopsy data showed an enhanced T-cell infiltration in the TME at week 5.

Conclusions: The chemotherapy-free combination of NT-17 + pembro was well tolerated in heavily pretreated subjects with CPI-naïve R/R MSS-CRC. The encouraging antitumor activity showed that subjects without liver metastasis sites especially benefited from the combination of NT-17 and pembro therapy. Biomarker analyses demonstrated improved peripheral and intratumoral T cell responses. Plan is to enroll 25 more patients to further evaluate efficacy of NT-17 + pembro in CPI-naïve subjects with R/R MSS-CRC.

Clinical trial identification: NCT04332653.

Legal entity responsible for the study: NeolmmuneTech, Inc.

Funding: NeolmmuneTech, Inc.

Disclosures: R. Kim: Advisory / Consultancy: Incyte, Roche, Servier, Eisai, Exelixis; Speaker Bureau / Expert testimony: Incyte, Taiho. M. Johnson: Advisory / Consultancy: Abbvie, Amgen, Astellas, AstraZeneca, Axelia Oncology, Black Diamond, Boehringer Ingelheim, Bristol Myers Squibb, Calithera Biosciences, Checkpoint Therapeutics, CytomX Therapeutics — ALL PAID TO INSTITUTION, Daiichi Sankyo, Ecor1, Editas Medicine, Eisai EMD Serono, G1 Therapeutics, Genentech/Roche, Genmab, Gritstone Oncology, Ideaya Biosciences, iTeos — ALL PAID TO INSTITUTION, Janssen, Lilly, Merck, Mirati Therapeutics, Oncorus, Regeneron Pharmaceuticals, Ribon Therapeutics, Sanofi-Aventis, Turning Point Therapeutics, WindMIL — ALL MONEY PAID TO INSTITUTION; Research grant / Funding (institution): Abbvie, Acta, Adaptimmune, Amgen, Apexigen, Arcus Biosciences, Array BioPharma, Artios Pharma, AstraZeneca, Atreca, BeiGene, BerGenBio, BioAtla, Boehringer Ingelheim, Calithera Biosciences, Checkpoint Therapeutics, Corvus Pharmaceuticals, Curis, CytomX, Daiichi Sankyo, Dracen Pharmaceuticals, Dynavax, Lilly, Elicio Therapeutics, EMD Serono, Erasca, Exelixis, Fate Therapeutics, Genentech/Roche, Genmab, Genocoe Biosciences, GlaxoSmithKline, Gritstone Oncology, Guardant Health, Harpoon, Helsinn Healthcare, Hengrui Therapeutics, Hutchinson Med-iPharma, Ideaya Biosciences, IGM Biosciences, Immunocore, Incyte, Janssen, Jounce Therapeutics, Kadmon Pharmaceuticals, Loxo Oncology, Lycera, Memorial Sloan-Kettering, Merck, Merus, Mirati, Neolmmune Tech, Neovia Oncology, Novartis, Numab Therapeutics, Nuvalent, OncoMed Pharmaceuticals, Pfizer, PMV Pharmaceuticals, RasCal Therapeutics, Regeneron Pharmaceuticals, Relay Therapeutics, Revolution Medicines, Ribon Therapeutics, Rubius Therapeutics, Sanofi, Seven and Eight Biopharmaceuticals / Birdie Biopharmaceuticals, Shattuck Labs, Silicon Therapeutics, Stem CentRx, Syndax Pharmaceuticals, Takeda Pharmaceuticals, Tarveda, TCR2 Therapeutics, Tempest Therapeutics, Tizona Therapeutics, TMUNITY Therapeutics, Turning Point Therapeutics, University of Michigan, Vyriad, WindMIL, Y-mAbs Therapeutics. S. Kopetz: Advisory / Consultancy: Roche, Redx Pharma, Jacobio, Merck, Navire Pharma, Holy Stone, Biocartis, Natera, Karyopharm Therapeutics, Repare Therapeutics, Genentech, Lilly, AstraZeneca/MedImmune, EMD Serono, Daiichi Sankyo, Amal Therapeutics, Boehringer Ingelheim, Bayer Health, Lutris, Pfizer, Amgen, Pierre Fabre, Symphogen, Novartis, Boston Biomedical, Ipsen, HalioDx, Inivata, GSK, Jazz pharmaceuticals Iydon, Xilis, Abbvie, Gilead Sciences, Mirati Therapeutics, Flame Biosciences, Servier, Carina Biotechnology, Bicara Therapeutics, Endeavor BioMedicines, Numab Pharma, Johnson & Johnson/Janssen, Genomic Health,

For; Research grant / Funding (self): Sanofi, Biocartis, Guardant Health, Array BioPharma, Genentech/ Roche, EMD Serono, MedImmune, Novartis, Amgen, Lilly, Daiichi Sankyo; Shareholder / Stockholder / Stock options: MolecularMatch, Frontier Medicines, Lutris, Iydon. S. Yang: Full / Part-time employment: NeolmmuneTech. B. Lee: Shareholder / Stockholder / Stock options: NeolmmuneTech Inc.; Full / Part-time employment: NeolmmuneTech Inc. S. Ferrando-Martinez: Full / Part-time employment: NeolmmuneTech, Inc. M. Chaney: Full / Part-time employment: Merck and Co., Inc. J. Fan: Full / Part-time employment: NeolmmuneTech, Inc. Officer / Board of Directors: Chief Clinical Officer of NeolmmuneTech, Inc. A. Naing: Advisory / Consultancy: Novartis, CytomX Therapeutics, OncoSec, STCube Pharmaceuticals Inc, Kymab, Takeda (I), CSL Behring (I), Horizon Pharma (I), Genome & Company, Immune-Onc, Deka Bioscience, Nouscom; Research grant / Funding (institution): NCI, EMD Serono, MedImmune, Atterocor, Amplimmune, ARMO BioSciences, Karyopharm Therapeutics, Incyte, Novartis, Regeneron, Merck, Bristol Myers Squibb, Pfizer, CytomX Therapeutics, Neon Therapeutics, Calithera Biosciences, TopAlliance BioSciences Inc, Healos, Lilly, Kymab, PsiOxus Therapeutics, Immune Deficiency Foundation (I), Arcus Biosciences, Neo-ImmuneTech, ImmuneOncia, Surface Oncology, Baxalta (I), Jeffrey Modell Foundation (I), Chao Physician-Scientist Awards (I); Travel / Accommodation / Expenses: ARMO BioSciences; Spouse / Financial dependant: Research funding: The Texas Medical Center Digestive Diseases Center, Jeffrey Modell Foundation, Immune Deficiency Foundation, Baxalta US Inc, Chao Physician-Scientist Foundation., Consultant/Advisory board: Takeda, Pharming Healthcare Inc, and Horizon Therapeutics USA, Inc., Ad hoc consultancy speaker: Alfaisal University. All other authors have declared no conflicts of interest.

<https://doi.org/10.1016/j.jannonc.2022.04.144>

P-55 Efficacy and safety data from patients with pre-treated metastatic colorectal cancer receiving trifluridine/tipiracil: Real-world data from the non-interventional TACTIC study

D. Semsek¹, H. Kröning², T. Göhler³, T. Decker⁴, G. Kojuharoff⁵, J. Lipke⁶, E. Moorahrend⁷, F. Hartmann⁸, T. Reisländer⁸, R. de Buhr⁹, M. Frank⁹, C. Hogrefe⁹, N. Marschner⁹, K. Potthoff⁹, I. Schwaner¹⁰

¹Praxis für interdisziplinäre Onkologie, Freiburg im Breisgau, Germany; ²Schwerpunktpraxis Hämatologie/Onkologie, Magdeburg, Germany; ³Onkzentrum Dresden/Freiberg, Dresden, Germany; ⁴Studiencentrum Onkologie Ravensburg, Ravensburg, Germany; ⁵Onkologische Schwerpunktpraxis Darmstadt, Darmstadt, Germany; ⁶Gemeinschaftspraxis für Hämatologie und Onkologie, Dortmund, Germany; ⁷Zentrum für Hämatologie und Onkologie MVZ GmbH, Porta Westfalica, Germany; ⁸SERVIER Deutschland GmbH, Munich, Germany; ⁹OMEDICO AG, Freiburg im Breisgau, Germany; ¹⁰Onkologische Schwerpunktpraxis Kurfürstendamm, Berlin, Germany

Background: In the pivotal phase III RECURSE trial, trifluridine/tipiracil (FTD/TPI) significantly improved overall and progression-free survival (OS, PFS) in patients with pre-treated metastatic colorectal cancer (mCRC) compared to placebo [1]. While randomised controlled trials represent the most reliable method of hypothesis testing, in- & exclusion criteria inevitably impede translation of their results to a real-world patient collective. Omitting restrictive in- & exclusion criteria we challenged the observations from the RECURSE trial on a patient population which more accurately reflects daily clinical practice in Germany.

Methods: In this prospective, multi-centre, open-label, non-interventional study, patients with pre-treated mCRC were treated with oral FTD/TPI (35 mg/m² bid on days 1 – 5 and 8 – 12 of each 28-day cycle). Primary endpoint was OS. Secondary endpoints included PFS and safety. Additionally, 3 subgroups were defined according to a post-hoc analysis of the RECURSE trial [2]: best, good and poor prognostic characteristics (BPC, GPC, PPC). Patients with < 3 metastatic sites at inclusion and ≥18 months from diagnosis to inclusion were considered to have GPC. GPC patients without liver metastasis at inclusion were considered to have BPC. All remaining patients were considered to have PPC.

Results: From June 2018 to August 2021, 307 patients were treated with FTD/TPI (mean treatment duration 3.4 cycles) at 52 German sites. Median age was 67.7 years and 17.0% of patients had an ECOG PS2/3. When focusing on patients with ECOG PS≤1, median OS of patients in the full analysis set (n=243: 8.6 months; 95% CI 7.4 – 9.3) as well as of the defined subgroups (BPC n=54 vs GPC n=147 vs PPC n=96: 16.2 vs 9.8 vs 6.3 months; 95% CI 9.7 – 19.4 vs 8.6 – 11.7 vs 4.5 – 7.8) were in line with results of RECURSE study (all patients n=534: 7.1 months/ BPC n=97 vs GPC n=261 vs PPC n=273: 16.4 vs 9.3 vs 5.3 months) [1] with a longer survival of patients with BPC and GPC compared to PPC. Similar results observed when analysing data from patients with ECOG PS≤3 (all patients n=300: 7.4 months; 95% CI 6.4 – 8.6/ BPC n=65 vs GPC n=176 vs PPC n=124: 13.3 vs 8.9 vs 5.1 months; 95% CI 9.1 – 17.6 vs 7.6 – 9.8 vs 4.4 – 7.0). Median PFS of all patients in the full analysis set was 2.9 months (95% CI 2.8 – 3.3). BPC (n=65) and GPC (n=176) patients were characterised by a longer median PFS compared to PPC (n=124) patients (4.0 vs 3.4 vs 2.6 months; 95% CI 3.3 – 5.3 vs 3.0 – 3.7 vs 2.4 – 2.8). The most frequent TEAEs were anaemia (20.5%), leukopenia (18.6%) and neutropenia (16.9%).

Conclusions: Administration of FTD/TPI to patients with pre-treated mCRC was associated with prolonged survival, delayed progression and a manageable toxicity profile confirming efficacy and safety of FTD/TPI in a real-world population. Independent of other baseline characteristics such as ECOG PS and age, low metastatic burden and indolent disease were factors of good prognosis with regards of OS and PFS. [1]Mayer et al, NEnglJMed2015 [2]Tabernero et al, ESMOOpen2020.