

Results: A total of 102 out of 735 mCRC patients (13.9%) were analyzed. Median age was 57.3 year and 55% were male. The outcomes were as follows: ORR 12%, DCR 39%, and mPFS 4.0 months (CI95% 3.29-5.03). Of note, 28 patients (27%) had a mPFS > 6 months with rechallenge (range from 6.57 to 14.2 months) and constituted the best-responders group. Data of prognostic characteristics and molecular alterations are available for 19 of these patients. The prognostic subgroup classification was as follows: 9 patients (47.4%) PPC, 5 patients (26.3%) GPC, and 5 (26.3%) BPC. Most frequent molecular alterations were: APC (68.4%), TP53 (63%), RAS (47.4%), and PIK3CA (21%). Concomitant APC and TP53 mutations were detected in 47.4% of patients. No associations between prognostic characteristics and molecular alterations were observed.

Conclusions: This study suggests that rechallenge with oxaliplatin can achieve a clinically meaningful mPFS > 6 months in 27% of patients. No enrichment in GPC and BPC was observed. APC, TP53 and RAS detected in this best-responders group are the major tumour genes which are frequently mutated in mCRC. A more extensive molecular analysis should be carried out to better characterise the patients who benefit the most from this treatment strategy.

Legal entity responsible for the study: The author.

Funding: Has not received any funding.

Disclosures: F. Salva: Honoraria (self): Amgen, Merck, Sanofi. R. Dientsmann: Advisory / Consultancy: Roche, Boehringer-Ingelheim; Speaker Bureau / Expert testimony: Roche, MSD, Amgen. J. Tabernero: Honoraria (self): educational collaboration with Immedex, Medscape Education, MJH Life Sciences, PeerView Institute for Medical Education and Physicians Education Resource (PER); Advisory / Consultancy: scientific consultancy role for Array Biopharma, AstraZeneca, Avintia, Bayer, Boehringer Ingelheim, Chugai, Daiichi Sankyo, F. Hoffmann-La Roche Ltd, Genentech Inc, HalioDX SAS, Hutchison MediPharma International, Ikema Oncology, Inspira Inc, IQVIA, Lilly, Menarini, Merck Serono, Merus, MSD, Mirati, Neophore, Novartis, Ona Therapeutics, Orion Biotechnology, Peptomyc, Pfizer, Pierre Fabre, Samsung Bioepis, Sanofi, Seattle Genetics, Scandion Oncology, Servier, Sotio Biotech, Taiho, Tessa Therapeutics and TheraMyc. All other authors have declared no conflicts of interest.

<https://doi.org/10.1016/j.annonc.2022.04.090>

PD-13 Plasma RAS dynamics and efficacy of anti-EGFR rechallenge in patients with RAS/BRAF wild-type metastatic colorectal cancer: REMARRY and PURSUIT trials

Y. Kagawa¹, D. Kotani², H. Bando², N. Takahashi³, Y. Horita⁴, A. Kanazawa⁵, T. Kato⁶, K. Ando⁷, H. Satake⁸, E. Shinozaki⁹, Y. Sunakawa¹⁰, A. Takashima¹¹, K. Yamazaki¹², S. Yuki¹³, H. Nakajima¹⁴, Y. Nakamura¹⁵, M. Wakabayashi¹⁶, H. Taniguchi¹⁶, T. Ohta¹⁷, T. Yoshino¹⁴

¹Department of Gastroenterological Surgery, Osaka General Medical Center, Osaka, Japan; ²Department of Gastroenterology and Gastrointestinal Oncology, National Cancer Center Hospital East, Kashiwa, Japan; ³Department of Gastroenterology, Saitama Cancer Center, Saitama, Japan; ⁴Department of Medical Oncology, Gastroenterological Oncology Saitama Medical University International Medical Center, Saitama city, Japan; ⁵Department of Surgery, Shimane Prefectural Central Hospital, Matsue, Japan; ⁶Department of Surgery, National Hospital Organization, Osaka National Hospital, Osaka, Japan; ⁷Department of Surgery and Science, Graduate School of Medical Science, Kyushu University, Fukuoka, Japan; ⁸Department of Medical Oncology, Kobe City Medical Center General Hospital, Kobe, Japan; ⁹Cancer Institute Hospital of Japanese Foundation for Cancer Research, Ariake, Japan; ¹⁰Department of Clinical Oncology, St. Marianna University School of Medicine, Kawasaki, Japan; ¹¹National Cancer Center Hospital, Tokyo, Japan; ¹²Division of Gastrointestinal Oncology, Shizuoka Cancer Center, Sunto-Gun, Japan; ¹³Department of Gastroenterology and Hepatology, Hokkaido University Hospital, Sapporo, Japan; ¹⁴National Cancer Center Hospital East, Kashiwa, Japan; ¹⁵Clinical Research Support Office, National Cancer Center Hospital East, Kashiwa, Japan; ¹⁶Department of Clinical Oncology, Aichi Cancer Center Hospital, Nagoya, Japan; ¹⁷Department of Clinical Oncology, Kansai Rosai Hospital, Hyogo, Japan

Background: Assessment of plasma RAS (pRAS) mutations in circulating-tumor DNA at 'just before' rechallenge with anti-EGFR monoclonal antibody (EGFR mAb) may predict efficacy for the rechallenge therapy in patients with RAS/BRAF V600E wild-type metastatic colorectal cancer (mCRC). However, the clinical impact of pRAS status at progression on prior EGFR mAb for the rechallenge therapy is unknown. The REMARRY trial is a prospective longitudinal study to investigate the pRAS dynamics, and PURSUIT trial is a phase II trial to investigate the efficacy of EGFR mAb rechallenge in patients with pRAS wild-type just before rechallenge therapy.

Methods: Eligibility criteria of REMARRY included RAS/BRAF V600E wild-type mCRC; ECOG PS 0-1; CR or PR during prior EGFR mAb; and progressed ≤ 2 months from the last administration of EGFR mAb. pRAS status by the BEAMing method (OncoBEAM RAS CRC Kit) was prospectively monitored at timepoints of progression on EGFR mAb and each subsequent therapy. Among participants of the REMARRY, patients who satisfied the following eligibility criteria were enrolled in PURSUIT: pRAS wild-type within 28 days prior to enrollment in PURSUIT; being refractory or intolerant to fluoropyrimidine, oxaliplatin, and irinotecan; and ≥ 4 months of EGFR mAb-free interval. Study treatment was rechallenge with panitumumab 6 mg/kg + irinotecan 150 mg/m²q2wks. Primary endpoint of PURSUIT was a confirmed objective response rate (ORR) according to RECIST v1.1. Biomarker analysis was performed for blood samples after disease progression on prior EGFR mAb, immediately prior to PURSUIT, and after

disease progression in PURSUIT using plasma next generation sequencing (Guardant360). Plasma RAS, BRAF V600E, and EGFR extracellular domain mutations were defined as acquired resistances for EGFR mAb.

Results: Between May 2019 and May 2021, 183 patients were enrolled in REMARRY from 27 institutions, and 50 patients were enrolled in PURSUIT; median age, 68 years; left-sided primary, 44 patients; and prior EGFR mAb, 1st/2nd/≥3rd lines in 28/6/16 patients. Confirmed ORR and disease control rate were 14% (90% CI, 7.8%–23.9%) and 80% (95% CI, 67.0%–88.8%), respectively. In addition, 4 patients showed an unconfirmed PR. Median progression-free survival (PFS) was 3.6 months (95% CI, 3.0–4.7 months). Among 31 patients with biomarker results after disease progression on prior EGFR mAb, ORR occurred in 5 of 21 patients (23.8%) with pRAS/BRAF/EGFR wild-type, whereas no responses occurred in patients whose tumors harbored any pRAS/BRAF/EGFR mutations (0/10) (p=0.092). Median PFS was 4.2 months and 2.8 months in patients without vs with pRAS/BRAF/EGFR mutations, respectively (p=0.06). In terms of pRAS/BRAF/EGFR status immediately prior to PURSUIT, 4 of 22 patients with wild-type responded to the study treatment (ORR, 18.2%), while one patient harboring pKRAS/EGFR co-mutations in 9 patients with any mutations also responded (ORR, 11.1%) (p=0.63). No trend in PFS was observed between the groups; median, 3.1 month and 3.3 months in patients with wild-type and those with any mutations (p=0.62).

Conclusions: Plasma RAS/BRAF/EGFR mutational status after progression on prior EGFR mAb may identify patients with RAS/BRAF V600E wild-type mCRC who could benefit from rechallenge with EGFR mAb.

Clinical trial identification: REMARRY: UMIN000036424 PURSUIT: jRCTs031190096.

Legal entity responsible for the study: The authors.

Funding: REMARRY: Sysmex Corporation PURSUIT: Takeda Pharmaceutical Company Limited.

Disclosures: Y. Kagawa: Speaker Bureau / Expert testimony: Lilly, Taiho, Yakult, MSD, Bayer, Daiichisankyo, Sanofi, Chugai, Ono, Takeda, Merck. D. Kotani: Honoraria (self): Takeda, Chugai. H. Bando: Honoraria (self): Eli Lilly Japan, Taiho pharmaceutical, Ono pharmaceutical; Research grant / Funding (self): Ono pharmaceutical. T. Kato: Honoraria (self): CHUGAI PHARMACEUTICAL CO., LTD, ONO PHARMACEUTICAL CO, Eli Lilly and Company; Honoraria (institution): CHUGAI PHARMACEUTICAL CO., LTD.; Research grant / Funding (self): Ono Pharmaceutical Co; Research grant / Funding (institution): Ono Pharmaceutical Co. H. Satake: Honoraria (self): Ono pharmaceutical co., Ltd., Daiichi Sankyo Co., Ltd., Eli Lilly Japan Co., Ltd., Merck Bio Pharma Co., Ltd., MSD Co., Ltd., Bayer Co., Ltd., Bristol-Myers Squibb Co., Ltd., Chugai Pharmaceutical Co., Ltd., Sanofi Co., Ltd., Taiho Pharmaceutical Co., Ltd., Takeda Co., Ltd. and Yakult Honsha Co., Ltd.; Research grant / Funding (institution): Ono pharmaceutical co., Ltd., Daiichi Sankyo, Takeda Pharmaceutical Co., Ltd., Sanofi, Taiho Pharmaceutical Co., Ltd., E. Shinozaki: Honoraria (self): Merck biopharma. Y. Sunakawa: Honoraria (self): Bristol-Myers Squibb, Chugai Pharm, Eli Lilly Japan; Advisory / Consultancy: Daiichi-Sankyo, Bristol-Myers Squibb, Guardant Health; Research grant / Funding (self): Chugai Pharm, Takeda, Taiho Pharm. K. Yamazaki: Honoraria (self): Chugai Pharma, Takeda, Taiho; Research grant / Funding (institution): Taiho Pharmaceutical. S. Yuki: Honoraria (self): Ono Pharmaceutical Co., Ltd., Merck Biopharma Co., Ltd., Y. Nakamura: Research grant / Funding (institution): Taiho Pharmaceutical, Guardant Health, Chugai Pharmaceutical. H. Taniguchi: Honoraria (self): Takeda, Taiho, Merck Biopharma; Research grant / Funding (institution): Daiichi-Sankyo, Sysmex, Takeda. T. Ohta: Honoraria (self): Takeda Pharmaceutical; Research grant / Funding (self): Takeda Pharmaceutical. T. Yoshino: Honoraria (self): Taiho Pharmaceutical, Chugai Pharmaceutical, Eli Lilly, Merck Biopharma, Bayer Yakuhin, Ono Pharmaceutical and MSD; Research grant / Funding (institution): Ono Pharmaceutical, Sanofi, Daiichi Sankyo, PAREXEL International, Pfizer Japan, Taiho Pharmaceutical, MSD, Amgen, Genomedia, Sysmex, Chugai Pharmaceutical and Nippon Boehringer Ingelheim. All other authors have declared no conflicts of interest.

<https://doi.org/10.1016/j.annonc.2022.04.091>

PD-14 Resectability, conversion, and resection rates with survival according to RAS and BRAF mutations in a prospective metastatic colorectal cancer study (liver-limited subgroup in the RAXO study)

A. Uutela¹, T. Salminen², E. Osterlund³, J. Kononen⁴, R. Kallio⁵, K. Lehtomäki², A. Älgars⁶, A. Lamminmäki⁷, P. Halonen¹, R. Ristimäki⁸, L. Soveri⁹, H. Stedt⁷, E. Heervä¹⁰, A. Nordin¹, A. Ristimäki¹, S. Kytölä¹¹, T. Kuopio¹¹, M. Mäkinen¹², L. Nieminen¹³, J. Sundström⁸, H. Isoniemi¹, P. Osterlund¹⁴

¹Helsinki University Hospital and Helsinki University, Helsinki, Finland; ²Tampere University Hospital and University of Tampere, Tampere, Finland; ³Uppsala University, Department of Immunology, Genetics and Pathology and Helsinki University Hospital, Uppsala, Sweden; ⁴Docrates Hospital and Central Hospital of Central Finland, Helsinki, Finland; ⁵Oulu University Hospital, Oulu, Finland; ⁶Turku University Hospital and University of Turku, Turku, Finland; ⁷Kuopio University Central Hospital and University of Eastern Finland, Kuopio, Finland; ⁸Turku University Central Hospital and Turku University, Turku, Finland; ⁹Helsinki University Hospital and Hyvinkää Hospital & Homecare, Helsinki, Finland; ¹⁰University of Turku and Turku University Hospital, Turku, Finland; ¹¹Central hospital of Central Finland, Jyväskylä, Finland; ¹²Oulu University Hospital and University of Oulu, Oulu, Finland; ¹³Tampere University Hospital, FIMLAB, Tampere, Finland; ¹⁴Tampere, Karolinska and Helsinki Universities and University Hospitals, Tampere, Stockholm, Helsinki, Finland

Background: RAS and BRAF mutations are associated with worse outcomes for patients with colorectal cancer (CRC) liver metastases, but little is known about their effects on resectability, conversion, and resection rates.