



Advancing the global public health agenda for NAFLD: a consensus statement

Jeffrey V. Lazarus 1,2 → Henry E. Mark 2, Quentin M. Anstee 3,4, Juan Pablo Arab 5, Rachel L. Batterham⁶, Laurent Castera 7, Helena Cortez-Pinto 8, Javier Crespo 9, Kenneth Cusi¹⁰, M. Ashworth Dirac¹¹, Sven Francque 12,13, Jacob George 14, Hannes Hagström¹⁵, Terry T.-K. Huang 16, Mona H. Ismail 17, Achim Kautz¹⁸, Shiv Kumar Sarin 19, Rohit Loomba 20,21, Veronica Miller²², Philip N. Newsome²³, Michael Ninburg 24, Ponsiano Ocama²⁵, Vlad Ratziu²⁶, Mary Rinella²⁷, Diana Romero 28, Manuel Romero-Gómez 29, Jörn M. Schattenberg 30, Emmanuel A. Tsochatzis 1,32, Luca Valenti 33,34, Vincent Wai-Sun Wong 35, Yusuf Yilmaz 36,37, Zobair M. Younossi 8, Shira Zelber-Sagi 39,40 and the NAFLD Consensus Consortium *

Abstract | Non-alcoholic fatty liver disease (NAFLD) is a potentially serious liver disease that affects approximately one-quarter of the global adult population, causing a substantial burden of ill health with wide-ranging social and economic implications. It is a multisystem disease and is considered the hepatic component of metabolic syndrome. Unlike other highly prevalent conditions, NAFLD has received little attention from the global public health community. Health system and public health responses to NAFLD have been weak and fragmented, and, despite its pervasiveness, NAFLD is largely unknown outside hepatology and gastroenterology. There is only a nascent global public health movement addressing NAFLD, and the disease is absent from nearly all national and international strategies and policies for non-communicable diseases, including obesity. In this global Delphi study, a multidisciplinary group of experts developed consensus statements and recommendations, which a larger group of collaborators reviewed over three rounds until consensus was achieved. The resulting consensus statements and recommendations address a broad range of topics — from epidemiology, awareness, care and treatment to public health policies and leadership — that have general relevance for policy-makers, health-care practitioners, civil society groups, research institutions and affected populations. These recommendations should provide a strong foundation for a comprehensive public health response to NAFLD.

Non-alcoholic fatty liver disease (NAFLD) is a potentially serious liver condition that, on a societal level, results in substantial health-care costs, economic losses and reduced health-related quality of life (HRQoL)¹⁻⁸. A biologically and clinically heterogeneous disease, NAFLD covers a broad spectrum of histological conditions that increase both hepatic and non-hepatic morbidity and mortality. The majority of people living with NAFLD have isolated steatosis (non-alcoholic fatty liver, NAFL) and a smaller proportion develop non-alcoholic steatohepatitis (NASH), with increasing hepatic fibrosis leading eventually to cirrhosis, liver cancer, end-stage liver disease and death^{9,10}. NASH is a leading cause of progression to cirrhosis and hepatocellular carcinoma^{11,12}, and an increasingly

common indication for liver transplantation^{13,14}. Liver cancer is now the second leading cause of years of life lost among all cancers globally¹⁵.

NAFLD is part of a multisystem disease and is considered the hepatic manifestation of metabolic syndrome ^{16–18}. Although strongly associated with obesity, NAFLD also occurs in individuals with normal weight, especially in Asian populations ^{19,20}. The causes of death in people living with NAFLD vary depending on disease state. Patients with cirrhosis predominantly have liver-related events, whereas those without cirrhosis have vascular events and non-hepatic cancer ²¹. Overall, cardiovascular disease (CVD) is the leading cause of death in patients with NAFLD; other common causes include extrahepatic

™e-mail: Jeffrey.Lazarus@ isglobal.org
https://doi.org/10.1038/s41575-021-00523-4

malignancies, liver-related complications, chronic kidney disease and type 2 diabetes mellitus (T2DM)^{16,17,22–24}.

NAFLD is closely related to other highly prevalent non-communicable diseases (NCDs) with substantial overlap in the public health and health system approaches needed to prevent and manage these conditions. However, NAFLD is currently absent from major global and national NCD strategies and action plans^{25,26}, and efforts to integrate NAFLD into the NCD agenda have been minimal. Despite the scale of the challenge and the human, social and economic implications of the disease, few people outside the fields of hepatology and gastroenterology are familiar with NAFLD, and there is no global public health movement to address the disease.

In this Consensus Statement, a global multidisciplinary group of experts developed consensus statements and recommendations for tackling the burden of NAFLD. The overarching goal was to develop a foundation for comprehensive public health responses to NAFLD and to outline catalytic actions that will move this agenda forwards in the coming years. Using a Delphi-based approach, the Consensus Statement sets out current thinking on NAFLD in areas ranging from epidemiology, awareness, care and treatment to public health policies and leadership (FIG. 1). The consensus statements and recommendations should have broad relevance for policy-makers, health-care practitioners, civil society groups, research institutions and affected populations.

Author addresses

¹Barcelona Institute for Global Health (ISGlobal), Hospital Clínic, University of Barcelona, Barcelona, Spain. ²EASL International Liver Foundation, Geneva, Switzerland. ³Translational & Clinical Research Institute, Faculty of Medical Sciences, Newcastle University, Newcastle upon Tyne, UK.

⁴Newcastle NIHR Biomedical Research Centre, Newcastle upon Tyne Hospitals NHS Trust, Newcastle upon Tyne, UK. ⁵Departamento de Gastroenterologia, Escuela de Medicina, Pontificia Universidad Catolica de Chile, Santiago, Chile.

⁶Centre for Obesity Research, University College London and National Institute of Health Research, UCLH Biomedical Research Centre, London, UK. ⁷Department of Hepatology, Hôpital Beaujon, Université de Paris, Paris, France.

⁸Clínica Universitária de Gastrenterologia, Laboratório de Nutrição, Faculdade de Medicina, Universidade de Lisboa, Lisbon, Portugal.

⁹Gastroenterology and Heptology Unit, Hospital Universitario Marqués de Valdecilla, Santander, IDIVAL, Santander, Spain.

¹⁰Division of Endocrinology, Diabetes & Metabolism, Department of Medicine, Veterans Health Administration and University of Florida, Gainesville, FL, USA.

¹¹Department of Health Metrics Sciences, Department of Family Medicine, University of Washington, Seattle, WA, USA.

¹²Department of Gastroenterology Hepatology, University Hospital Antwerp, Antwerp, Belgium.

¹³Translational Sciences in Inflammation and Immunology TWI2N, Faculty of Medicine and Health Sciences, University of Antwerp, Antwerp, Belgium.

¹⁴Storr Liver Centre, Westmead Institute for Medical Research, Westmead Hospital and University of Sydney, Sydney, New South Wales, Australia.

¹⁵Department of Medicine, Karolinska Institutet, Stockholm, Sweden.

¹⁶Center for Systems and Community Design and NYU-CUNY Prevention Research Center, Graduate School of Public Health and Health Policy, City University of New York, New York, NY, USA.

¹⁷Division of Gastroenterology, Department of Internal Medicine, King Fahd Hospital of the University, College of Medicine, Imam Abdulrahman Bin Faisal University, Dammam, Saudi Arabia.

¹⁸Kautz5, Köln, Germany.

¹⁹Department of Hepatology, Institute of Liver and Biliary Sciences, New Delhi, India.

²⁰Department of Medicine, NAFLD Research Center, La Iolla, CA, USA.

²¹Department of Medicine, University of California San Diego, La Jolla, CA, USA.

²²University of California Berkeley, School of Public Health,
 Forum for Collaborative Research, Washington, DC, USA.
 ²³National Institute for Health Research Birmingham

Biomedical Research Centre at University Hospitals Birmingham NHS Foundation Trust and the University of Birmingham, Birmingham, UK.

²⁴Hepatitis Education Project, Seattle, WA, USA.

²⁵Department of Medicine, College of Health Sciences, Makerere University, Kampala, Uganda.

²⁶Assistance Publique-Hôpitaux de Paris, Hôpital

Pitie-Salpetriere, University of Paris, Paris, France.

²⁷Department of Medicine, Feinberg School of Medicine,
Northwestern University, Chicago, IL, USA.

²⁸Department of Community Health and Social Sciences, CUNY Graduate School of Public Health and Health Policy, New York, NY, USA.

²⁹UCM Digestive Diseases, CIBEREHD and IBIS, Virgen del Rocío University Hospital, University of Seville, Seville, Spain.

 ³⁰Metabolic Liver Research Program, I. Department of Medicine, University Medical Centre Mainz, Mainz, Germany.
 ³¹University College London Institute for Liver and Digestive Health, Royal Free Hospital, London, UK.
 ³²Sheila Sherlock Liver Centre, Royal Free Hospital,

³³Department of Pathophysiology and Transplantation, Università degli Studi di Milano, Milan, Italy.

London, UK.

³⁴Precision Medicine, Department of Transfusion Medicine and Hematology, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy.

³⁵Department of Medicine and Therapeutics, The Chinese University of Hong Kong, Hong Kong, China.

³⁶Department of Gastroenterology, Marmara University School of Medicine, Istanbul, Turkey.

³⁷Liver Research Unit, Institute of Gastroenterology, Marmara University, Istanbul, Turkey.

³⁸Center for Liver Diseases, Inova Medicine, Falls Church, VA, USA.

³⁹University of Haifa, Faculty of Social Welfare and Health Sciences, School of Public Health, Mount Carmel, Haifa, Israel

⁴⁰Department of Gastroenterology, Tel-Aviv Medical Centre, Tel-Aviv, Israel.

*A list of authors and their affiliations appears at the end of the paper.




Fig. 1 | **Development of the NAFLD consensus statements and recommendations. a** | The development process for the development of the non-alcoholic fatty liver disease (NAFLD) consensus statement and recommendations following a Delphi methodology, **b** | Summary of the 26 recommendations stemming from this process. NCDs, non-communicable diseases.

Methods

Expert panel members and topics

A core group of 33 experts (TABLE 1) were identified by the European Association for the Study of the Liver (EASL) International Liver Foundation (EILF) to lead a Delphi study to develop consensus statements and recommendations to advance the NAFLD public health agenda. The chair (J.V.L.) and project coordinator (H.E.M.) led this group of clinicians, researchers, advocates, academics and civil society experts from 16 countries through the development and implementation of the Delphi process. Core group members identified additional experts to be invited to participate in the Delphi consensus-building process. The final panel comprised 218 individuals, including the core group members. The demographic description of the expert panel is summarized in TABLE 2 and its geographical diversity in Supplementary Table 1.

In December 2020, the core group drafted the statements to be used for the initial Delphi survey round and the statements were grouped into the following categories: (1) human and economic burden; (2) awareness; (3) defining and implementing models of care, including (3a) general considerations, (3b) considerations for children and adolescents, and (3c) considerations for low-resource settings; (4) treatment and care; (5) patient and community perspectives; (6) policy strategies and

a whole-of-society approach; and (7) leadership for the NAFLD public health agenda.

Delphi method data collection

The Delphi method design²⁷ consisted of five components of data collection, including a first and second survey round (R1 in January 2021, R2 in February 2021), an online convening of the core group (16 February 2021), a web-based review of draft recommendations (February 2021), and a final survey round (R3 in March 2021). We used the Qualtrics XM platform to develop and distribute the surveys. The data collection periods for each survey round ranged between 1.5 and 3 weeks, allowing for holiday periods. The R1 survey contained 38 draft statements with four-point Likert-type categories for respondents to indicate their level of agreement with the statements (that is, 'Agree'/'Somewhat agree'/'Somewhat disagree'/'Disagree'). In this round, respondents who agreed or somewhat agreed with a statement could provide comments and suggest edits, while those who disagreed or somewhat disagreed could explain why. The R2 survey contained 37 statements and reflected suggestions from R1, including new, revised and merged statements. In the R2 survey, we included text-box summaries of the edits made to each of the statements for respondents to consider as they indicated their level of agreement or

62 | JANUARY 2022 | VOLUME 19

Table 1 | Core group members (n = 33)

Juan Pablo Arab Pontifical Catholic University of Chile Chile Rachel L. Batterham University College London UK Laurent Castera University of Paris France Helena Cortez Pinto University of Lisbon Portugal lavier Crespo University of Lisbon Spain Kenneth Cusi Veterans Health Administration and University USA M. Ashworth Dirac University of Washington USA W. Ashworth Dirac University of Sydney Belgium Jacob George Antwerp University Hospital Belgium Jacob George University of Sydney Australia Jacob George University of Sydney Australia Jacob George University of New York Graduate School of Public Health and Health Policy Mona H. Ismail College of Medicine, Imam Abdulrahman Bin Faisal University King Fahad Hospital of the University, Al-Khobar Achim Kautz Kautz² gUG Shiv Kumar Sarin Institute of Liver and Biliary Science India Jeffrey V. Lazarus (Chair) Barcelona Institute for Global Health (ISGlobal) Spain Rohit Loomba University of California, San Diego USA Henry E. Mark (Project coordinator) EASL International Liver Foundation Switzerland Veronica Miller University of Galifornia, Berkeley USA Phil N. Newsome University of Galifornia, Berkeley USA Phil N. Newsome University of Birmingham UK Michael Ninburg Hepatitis Education Project USA Ponsiano Ocama Makerere University College of Health Sciences Uganda Vlad Ratziu Sorbonne University Feinberg School of Medicine Diana Romero (Non-voting) City University of New York Graduate School of Public Health and Health Policy Many Rinella Northwestern University Feinberg School of Medicine USA Manuel Romero-Gómez Virgen del Rocio University Hospital Spain Jeffrey V. Lazarus Ukla Health Policy Manuel Romero-Gómez Virgen del Rocio University Hospital Spain Jeffrey V. S. Wong The Chinese University of Hong Kong China Jurkey V. Wong Schattenberg University of Milan Italy Viricent W. S. Wong The Chinese University of Hong Kong China Jurkey V. John Health Policy V. Sa	Name	Affiliation	Country/territory where currently based
Rachel L. Batterham University College London UK Laurent Castera University of Paris France Helena Cortez Pinto University of Lisbon Portugal Javier Crespo Universidad de Cantabria Spain Kenneth Cusi Veterans Health Administration and University of Florida M. Ashworth Dirac University of Washington USA Sven Francque Antwerp University Hospital Belgium Jacob George University of Sydney Australia Hannes Hagström Karolinska Institutet Terry TK. Huang City University of New York Graduate School of Public Health and Health Policy Mona H. Ismail College of Medicine, Imam Abdulrahman Bin Faisal University King Fahad Hospital of the University, Al-Khobar Achim Kautz Kautz* gUG Germany Shiv Kumar Sarin Institute of Liver and Biliary Science India Jeffrey V. Lazarus (Chair) Barcelona Institute for Global Health (ISGlobal) Spain Rohit Loomba Henry E. Mark (Project coordinator) Alennic Ray (Project coordinator) EASL International Liver Foundation Switzerland Veronica Miller University of Galifornia, Berkeley USA Phil N. Newsome University of Birmingham UK Michael Ninburg Hepatitis Education Project USA Ponsian Ocama Makerere University Feinberg School of Medicine USA Mary Rinella Northwestern University Feinberg School of Public USA Diana Romero (Non-voting) City University of New York Graduate School of Public USA Health and Health Policy Manuel Romero-Gómez Virgen del Rocio University Hospital Spain Join M. Schattenberg University Medical Center Mainz Germany Ermanuel Tsochatzis UCL Institute for Liver and Digestive Health UK Luca Valenti University of Milan Italy Virincent W. S. Wong The Chinese University Hospital Usa Fundaria University of Hong Kong China Vusa Yilmaz Marmara University Hospital Spain Jurkey Juniversity of Hong Kong China	Quentin M. Anstee	Newcastle University	UK
Laurent Castera University of Paris France Helena Cortez Pinto University of Lisbon Portugal lavier Crespo Universidad de Cantabria Spain Kenneth Cusi Veterans Health Administration and University USA M. Ashworth Dirac University of Washington USA Sven Francque Antwerp University Hospital Belgium lacob George University of Sydney Australia Hannes Hagström Karolinska Institutet Sweden Terry TK. Huang City University of New York Graduate School of Public USA Health and Health Policy Mona H. Ismail College of Medicine, Imam Abdulrahman Bin Faisal University King Fahad Hospital of the University, Al-Khobar Achim Kautz Kautz² gUG Germany Shiv Kumar Sarin Institute of Liver and Biliary Science India leffrey V. Lazarus (Chair) Barcelona Institute for Global Health (ISGlobal) Spain Rohit Loomba University of California, San Diego USA Henry E. Mark (Project coordinator) EASL International Liver Foundation Switzerland Veronica Miller University of Birmingham UK Michael Ninburg Hepatitis Education Project USA Ponsiano Ocama Makerere University College of Health Sciences Uganda Vlad Ratziu Sorbonne University of Sirmingham UK Mary Rinella Northwestern University Feinberg School of Public USA Diana Romero (Non-voting) City University of New York Graduate School of Public USA Diana Romero (Non-voting) Uicy Iniversity of New York Graduate School of Public USA Diana Romero Gomez Virgen del Rocio University Hospital Spain France Mannand Romero-Gómez Virgen del Rocio University Hospital Spain France Furmanuel Tsochatzis UCL Institute for Liver and Digestive Health UK Luca Valenti University of New York Graduate School of Public Virgen del Rocio University of New York Graduate School of Public Virgen del Rocio University of New York Graduate School of Public Virgen del Rocio University Hospital Spain France Furmanuel Tsochatzis Ucl Institute for Liver and Digestive Health UK Luca Valenti University of Milan Health Policy Namara University of Hong Kong China Furshary Primaz Marmara University of Hong Kong China	Juan Pablo Arab	Pontifical Catholic University of Chile	Chile
Helena Cortez Pinto University of Lisbon Spain Kenneth Cusi Veterans Health Administration and University of Florida M. Ashworth Dirac University of Washington USA M. Ashworth Dirac University of Washington USA Sven Francque Antwerp University Hospital Belgium Jacob George University of Sydney Australia Hannes Hagström Karolinska Institutet Sweden Terry TK. Huang City University of New York Graduate School of Public USA Mona H. Ismail College of Medicine, Imam Abdulrahman Bin Faisal University King Fahad Hospital of the University, Al-Khobar Achim Kautz Kautz' gUG Germany Shiv Kumar Sarin Institute of Liver and Biliary Science India Jeffrey V. Lazarus (Chair) Barcelona Institute for Global Health (ISGlobal) Spain Rohit Loomba University of California, San Diego USA Henry E. Mark (Project coordinator) EASL International Liver Foundation Switzerland Veronica Miller University of Birmingham UK Weronica Miller University of Birmingham Wichael Ninburg Hepatitis Education Project USA Makerere University Feinberg School of Medicine USA Diana Romero (Non-voting) City University of New York Graduate School of Public Health and Health Policy Manuel Romero-Gómez Virgen del Rocio University Hospital Spain Wanuel Romero-Gómez University of New York Graduate School of Public Health and Health Policy Manuel Romero-Gómez University of New York Graduate School of Public USA Manuel Romero-Gómez Virgen del Rocio University Hospital Spain Emmanuel Tsochatzis UCL Institute for Liver and Digestive Health University of Milan Italy Virgent W. S. Wong The Chinese University of Hong Kong China Yusuf Yilmaz Marmara University of Hong Kong China Yusuf Yilmaz Marmara University of Health System USA	Rachel L. Batterham	University College London	UK
Javier Crespo Universidad de Cantabria Spain Kenneth Cusi Veterans Health Administration and University Of Florida M. Ashworth Dirac University of Washington USA Sven Francque Antwerp University Hospital Belgium Jacob George University of Sydney Australia Hannes Hagström Karolinska Institutet Sweden Terry TK. Huang City University of New York Graduate School of Public Health and Health Policy Mona H. Ismail College of Medicine, Imam Abdulrahman Bin Faisal University King Fahad Hospital of the University, Al-Khobar Achim Kautz Kautz² gUG Germany Shiv Kumar Sarin Institute of Liver and Biliary Science India Jeffrey V. Lazarus (Chair) Barcelona Institute for Global Health (ISGlobal) Spain Rohit Loomba University of California, San Diego USA Henry E. Mark (Project coordinator) EASL International Liver Foundation Switzerland Veronica Miller University of Birmingham UK Michael Ninburg Hepatitis Education Project USA Phil N. Newsome University of Birmingham UK Michael Ninburg Hepatitis Education Project USA Ponsian Ocama Makerere University Feinberg School of Medicine USA Diana Romero (Non-voting) City University of New York Graduate School of Public Health and Health Policy Manuel Romero-Gómez Virgen del Rocio University Hospital Spain John M. Schattenberg University of New York Graduate School of Public Health and Health Policy Manuel Romero-Gómez Virgen del Rocio University Hospital Spain John M. Schattenberg University Medical Center Mainz Germany Emmanuel Tsochatzis UCL Institute for Liver and Digestive Health Luca Valenti University of Milan Italy Virineat W. S. Wong The Chinese University of Hong Kong China Yusuf Yilmaz Marmara University of Hong Kong China Yusuf Yilmaz Marmara University of Health System USA	Laurent Castera	University of Paris	France
Kenneth Cusi Veterans Health Administration and University of Florida M. Ashworth Dirac University of Washington USA M. Ashworth Dirac University of Washington USA M. Ashworth Dirac University of Sydney Australia Jacob George University of Sydney Australia Hannes Hagström Karolinska Institutet Sweden Terry TK. Huang City University of New York Graduate School of Public Health and Health Policy Mona H. Ismail College of Medicine, Imam Abdulrahman Bin Faisal University King Fahad Hospital of the University, Al-Khobar Achim Kautz Kautz ⁵ gUG Germany Shiv Kumar Sarin Institute of Liver and Biliary Science India Juniversity of California, San Diego USA Henry E. Mark (Project coordinator) EASL International Liver Foundation Weronica Miller University of California, Berkeley USA Phil N. Newsome University of Florinningham UK Michael Ninburg Hepatitis Education Project USA Ponsiano Ocama Makerere University College of Health Sciences Uganda Vlad Ratziu Sorbonne University of New York Graduate School of Public Health and Health Policy Wire and Health Policy USA Manuel Romero-Gómez Virgu ndel Rocío University Hospital Diana Romero (Non-voting) Usch Liuca Valenti University of Milan Vincent W. S. Wong The Chinese University of Hong Kong China Virus Yilmaz Marmara University Inve Health System USA	Helena Cortez Pinto	University of Lisbon	Portugal
of Florida M. Ashworth Dirac University of Washington USA Sven Francque Antwerp University Hospital Acob George University of Sydney Australia Hannes Hagström Karolinska Institutet Sweden Terry TK. Huang City University of New York Graduate School of Public Health and Health Policy Mona H. Ismail College of Medicine, Imam Abdulrahman Bin Faisal University King Fahad Hospital of the University, Al-Khobar Achim Kautz Kautz ⁵ gUG Germany Shiv Kumar Sarin Institute of Liver and Biliary Science India Leffrey V. Lazarus (Chair) Barcelona Institute for Global Health (ISGlobal) Spain Rohit Loomba University of California, San Diego USA Henry E. Mark (Project coordinator) Liver Sity of California, Berkeley USA Woronica Miller University of California, Berkeley USA Phil N. Newsome University of Birmingham UK Michael Ninburg Hepatitis Education Project USA Ponsiano Ocama Makerere University College of Health Sciences Uganda Vlad Ratziu Sorbonne University Mary Rinella Northwestern University Feinberg School of Medicine USA Diana Romero (Non-voting) City University of New York Graduate School of Public Health and Health Policy USA Manuel Romero-Gómez Virgen del Rocio University Hospital Diana Romero (Non-voting) City University of New York Graduate School of Public Health and Health Policy USA Manuel Romero-Gómez Virgen del Rocio University Hospital UK Luca Valenti University of Milan UK UK University of Hong Kong China Yusuf Yilmaz Marmara University of Hong Kong China Yusuf Yilmaz Nova Medicine, Inova Health System USA	Javier Crespo	Universidad de Cantabria	Spain
Sven Francque Antwerp University Hospital Belgium Jacob George University of Sydney Australia Jacob George University of Sydney Australia Hannes Hagström Karolinska Institutet Sweden Terry TK. Huang City University of New York Graduate School of Public Health and Health Policy Mona H. Ismail College of Medicine, Imam Abdulrahman Bin Faisal University King Fahad Hospital of the University, Al-Khobar Achim Kautz Kautz's GUG Germany Shiv Kumar Sarin Institute of Liver and Biliary Science India Jeffrey V. Lazarus (Chair) Barcelona Institute for Global Health (ISGlobal) Spain Rohit Loomba University of California, San Diego USA Henry E. Mark (Project coordinator) EASL International Liver Foundation Switzerland Veronica Miller University of California, Berkeley USA Phil N. Newsome University of Birmingham UK Michael Ninburg Hepatitis Education Project USA Ponsiano Ocama Makerere University College of Health Sciences Uganda Vlad Ratziu Sorbonne University Feinberg School of Medicine Wany Rinella Northwestern University Feinberg School of Medicine Diana Romero (Non-voting) City University of New York Graduate School of Public Health and Health Policy Manuel Romero-Gómez Virgen del Rocío University Hospital Spain Jörn M. Schattenberg University Medical Center Mainz Germany Emmanuel Tsochatzis UCL Institute for Liver and Digestive Health UK Luca Valenti University of Milan Italy Vincent W. S. Wong The Chinese University of Hong Kong China Yusuf Yilmaz Marmara University Tova Health System USA	Kenneth Cusi		USA
Hannes Hagström Karolinska Institutet Sweden Terry TK. Huang City University of New York Graduate School of Public Health and Health Policy Mona H. Ismail College of Medicine, Imam Abdulrahman Bin Faisal University King Fahad Hospital of the University, Al-Khobar Achim Kautz Kautz ⁵ gUG Germany Shiv Kumar Sarin Institute of Liver and Biliary Science India Jeffrey V. Lazarus (Chair) Barcelona Institute for Global Health (ISGlobal) Spain Rohit Loomba University of California, San Diego USA Henry E. Mark (Project coordinator) EASL International Liver Foundation Switzerland Veronica Miller University of Galifornia, Berkeley USA Phil N. Newsome University of Birmingham UK Michael Ninburg Hepatitis Education Project USA Ponsiano Ocama Makerere University College of Health Sciences Uganda Vlad Ratziu Sorbonne University Feinberg School of Medicine Diana Romero (Non-voting) City University of New York Graduate School of Public Health and Health Policy Manuel Romero-Gómez Virgen del Rocío University Hospital Spain Jörn M. Schattenberg University Medical Center Mainz Germany Emmanuel Tsochatzis UCL Institute for Liver and Digestive Health UK Luca Valenti University of Milan Italy Vincent W. S. Wong The Chinese University of Hong Kong China Yusuf Yilmaz Marmara University Inova Health System USA	M. Ashworth Dirac	University of Washington	USA
Hannes Hagström Karolinska Institutet Sweden Terry TK. Huang City University of New York Graduate School of Public Health and Health Policy Mona H. Ismail College of Medicine, Imam Abdulrahman Bin Faisal University King Fahad Hospital of the University, Al-Khobar Achim Kautz Kautz ⁸ gUG Germany Shiv Kumar Sarin Institute of Liver and Biliary Science India leffrey V. Lazarus (Chair) Barcelona Institute for Global Health (ISGlobal) Spain Rohit Loomba University of California, San Diego USA Henry E. Mark (Project coordinator) EASL International Liver Foundation Switzerland Veronica Miller University of California, Berkeley USA Phil N. Newsome University of Birmingham UK Michael Ninburg Hepatitis Education Project USA Ponsiano Ocama Makerere University College of Health Sciences Uganda VIad Ratziu Sorbonne University Feinberg School of Medicine USA Diana Romero (Non-voting) City University of New York Graduate School of Public Health and Health Policy Manuel Romero-Gómez Virgen del Rocío University Hospital Spain Jörn M. Schattenberg University Medical Center Mainz Germany Emmanuel Tsochatzis UCL Institute for Liver and Digestive Health UK Luca Valenti University of Milan Italy Vincent W. S. Wong The Chinese University of Hong Kong China Yusuf Yilmaz Marmara University Turkey Zobair M. Younossi Inova Medicine, Inova Health System USA	Sven Francque	Antwerp University Hospital	Belgium
Terry TK. Huang City University of New York Graduate School of Public Health and Health Policy Mona H. Ismail College of Medicine, Imam Abdulrahman Bin Faisal University King Fahad Hospital of the University, Al-Khobar Achim Kautz Kautz³ gUG Germany Institute of Liver and Biliary Science India Jeffrey V. Lazarus (Chair) Barcelona Institute for Global Health (ISGlobal) Spain Rohit Loomba University of California, San Diego USA Henry E. Mark (Project coordinator) FASL International Liver Foundation Weronica Miller University of California, Berkeley USA Pibli N. Newsome University of Birmingham UK Michael Ninburg Hepatitis Education Project USA Ponsiano Ocama Makerere University College of Health Sciences Uganda Vlad Ratziu Sorbonne University France Mary Rinella Northwestern University Feinberg School of Medicine USA Diana Romero (Non-voting) City University of New York Graduate School of Public Health and Health Policy Manuel Romero-Gómez Virgen del Rocío University Hospital Spain Jörn M. Schattenberg University Medical Center Mainz Emmanuel Tsochatzis UCL Institute for Liver and Digestive Health UK Luca Valenti University of Milan Italy Vincent W. S. Wong The Chinese University of Hong Kong China Yusuf Yilmaz Marmara University Turkey Zobair M. Younossi Inova Medicine, Inova Health System USA	Jacob George	University of Sydney	Australia
Health and Héalth Policy Mona H. Ismail College of Medicine, Imam Abdulrahman Bin Faisal University King Fahad Hospital of the University, Al-Khobar Achim Kautz Kautz ⁵ gUG Germany Institute of Liver and Biliary Science India Jeffrey V. Lazarus (Chair) Barcelona Institute for Global Health (ISGlobal) Spain Rohit Loomba University of California, San Diego USA Henry E. Mark (Project coordinator) EASL International Liver Foundation Switzerland Veronica Miller University of Birmingham UK Michael Ninburg Hepatitis Education Project USA Ponsiano Ocama Makerere University College of Health Sciences Uganda Vlad Ratziu Sorbonne University France Mary Rinella Northwestern University Feinberg School of Medicine USA Diana Romero (Non-voting) City University of New York Graduate School of Public Health and Health Policy Manuel Romero-Gómez Virgen del Rocío University Hospital Jörn M. Schattenberg University Medical Center Mainz Germany Emmanuel Tsochatzis UCL Institute for Liver and Digestive Health UK Luca Valenti University of Milan Vincent W. S. Wong The Chinese University of Hong Kong China Yusuf Yilmaz Marmara University Inova Medicine, Inova Health System USA	Hannes Hagström	Karolinska Institutet	Sweden
University King Fahad Hospital of the University, Al-Khobar Achim Kautz Kautz ⁵ gUG Germany Shiv Kumar Sarin Institute of Liver and Biliary Science India Jeffrey V. Lazarus (Chair) Barcelona Institute for Global Health (ISGlobal) Spain Rohit Loomba University of California, San Diego USA Henry E. Mark (Project coordinator) EASL International Liver Foundation Veronica Miller University of California, Berkeley USA Phil N. Newsome University of Birmingham UK Michael Ninburg Hepatitis Education Project USA Ponsiano Ocama Makerere University College of Health Sciences Uganda Vlad Ratziu Sorbonne University France Mary Rinella Northwestern University Feinberg School of Medicine USA Diana Romero (Non-voting) City University of New York Graduate School of Public Health and Health Policy Manuel Romero-Gómez Virgen del Rocío University Hospital Jörn M. Schattenberg University Medical Center Mainz Germany Emmanuel Tsochatzis UCL Institute for Liver and Digestive Health UK Luca Valenti University of Milan Italy Vincent W. S. Wong The Chinese University of Hong Kong China Yusuf Yilmaz Marmara University Inova Medicine, Inova Health System USA	Terry TK. Huang		USA
Achim Kautz Kautz² gUG Institute of Liver and Biliary Science India Jeffrey V. Lazarus (Chair) Barcelona Institute for Global Health (ISGlobal) Spain Rohit Loomba University of California, San Diego USA Henry E. Mark (Project coordinator) EASL International Liver Foundation Switzerland Veronica Miller University of California, Berkeley USA Phil N. Newsome University of Birmingham UK Michael Ninburg Hepatitis Education Project USA Ponsiano Ocama Makerere University College of Health Sciences Uganda Vlad Ratziu Sorbonne University France Mary Rinella Northwestern University Feinberg School of Medicine USA Diana Romero (Non-voting) City University of New York Graduate School of Public Health and Health Policy Manuel Romero-Gómez Virgen del Rocío University Hospital Spain Jörn M. Schattenberg University Medical Center Mainz Germany Emmanuel Tsochatzis UCL Institute for Liver and Digestive Health UK Luca Valenti University of Milan Vincent W. S. Wong The Chinese University of Hong Kong China Yusuf Yilmaz Marmara University Inova Medicine, Inova Health System USA	Mona H. Ismail		Saudi Arabia
Shiv Kumar Sarin Institute of Liver and Biliary Science India Jeffrey V. Lazarus (Chair) Barcelona Institute for Global Health (ISGlobal) Spain Rohit Loomba University of California, San Diego USA Henry E. Mark (Project coordinator) EASL International Liver Foundation Switzerland Veronica Miller University of California, Berkeley USA Phil N. Newsome University of Birmingham UK Michael Ninburg Hepatitis Education Project USA Ponsiano Ocama Makerere University College of Health Sciences Uganda Vlad Ratziu Sorbonne University Feinberg School of Medicine USA Diana Romero (Non-voting) City University of New York Graduate School of Public Health and Health Policy Manuel Romero-Gómez Virgen del Rocío University Hospital Spain Jörn M. Schattenberg University Medical Center Mainz Germany Emmanuel Tsochatzis UCL Institute for Liver and Digestive Health UK Luca Valenti University of Milan Italy Vincent W. S. Wong The Chinese University Tova Health System USA Zobair M. Younossi Inova Medicine, Inova Health System USA		King Fahad Hospital of the University, Al-Khobar	
leffrey V. Lazarus (Chair) Rohit Loomba University of California, San Diego USA Henry E. Mark (Project coordinator) EASL International Liver Foundation Veronica Miller University of California, Berkeley USA Phil N. Newsome University of Birmingham UK Michael Ninburg Hepatitis Education Project USA Ponsiano Ocama Makerere University College of Health Sciences Uganda Vlad Ratziu Sorbonne University France Mary Rinella Northwestern University Feinberg School of Medicine USA Diana Romero (Non-voting) City University of New York Graduate School of Public Health and Health Policy Manuel Romero-Gómez Virgen del Rocío University Hospital Jörn M. Schattenberg University Medical Center Mainz Emmanuel Tsochatzis UCL Institute for Liver and Digestive Health UK Luca Valenti University of Milan Italy Vincent W. S. Wong The Chinese University of Hong Kong Cushair M. Younossi Inova Medicine, Inova Health System USA	Achim Kautz	Kautz⁵ gUG	Germany
Rohit Loomba University of California, San Diego USA Henry E. Mark (Project coordinator) EASL International Liver Foundation Switzerland Veronica Miller University of California, Berkeley USA Phil N. Newsome University of Birmingham UK Michael Ninburg Hepatitis Education Project USA Ponsiano Ocama Makerere University College of Health Sciences Uganda Vlad Ratziu Sorbonne University Feinberg School of Medicine USA Diana Romero (Non-voting) City University of New York Graduate School of Public Health and Health Policy Manuel Romero-Gómez Virgen del Rocío University Hospital Spain Jörn M. Schattenberg University Medical Center Mainz Germany Emmanuel Tsochatzis UCL Institute for Liver and Digestive Health UK Luca Valenti University of Milan Italy Vincent W. S. Wong The Chinese University of Hong Kong China Yusuf Yilmaz Marmara University Turkey Zobair M. Younossi Inova Medicine, Inova Health System USA	Shiv Kumar Sarin	Institute of Liver and Biliary Science	India
Henry E. Mark (Project coordinator) EASL International Liver Foundation Switzerland Veronica Miller University of California, Berkeley USA Phil N. Newsome University of Birmingham UK Michael Ninburg Hepatitis Education Project USA Ponsiano Ocama Makerere University College of Health Sciences Uganda Vlad Ratziu Sorbonne University Feinberg School of Medicine USA Diana Romero (Non-voting) City University of New York Graduate School of Public Health and Health Policy Manuel Romero-Gómez Virgen del Rocío University Hospital Spain Jörn M. Schattenberg University Medical Center Mainz Germany Emmanuel Tsochatzis UCL Institute for Liver and Digestive Health UK Luca Valenti University of Milan Italy Vincent W. S. Wong The Chinese University of Hong Kong China Yusuf Yilmaz Marmara University Zobair M. Younossi Inova Medicine, Inova Health System USA	Jeffrey V. Lazarus (Chair)	Barcelona Institute for Global Health (ISGlobal)	Spain
Veronica Miller University of California, Berkeley USA Phil N. Newsome University of Birmingham UK Michael Ninburg Hepatitis Education Project USA Ponsiano Ocama Makerere University College of Health Sciences Uganda Vlad Ratziu Sorbonne University France Mary Rinella Northwestern University Feinberg School of Medicine USA Diana Romero (Non-voting) City University of New York Graduate School of Public Health and Health Policy Manuel Romero-Gómez Virgen del Rocío University Hospital Spain Jörn M. Schattenberg University Medical Center Mainz Germany Emmanuel Tsochatzis UCL Institute for Liver and Digestive Health UK Luca Valenti University of Milan Italy Vincent W. S. Wong The Chinese University of Hong Kong China Yusuf Yilmaz Marmara University Inova Medicine, Inova Health System USA	Rohit Loomba	University of California, San Diego	USA
Phil N. Newsome University of Birmingham UK Michael Ninburg Hepatitis Education Project USA Ponsiano Ocama Makerere University College of Health Sciences Uganda Vlad Ratziu Sorbonne University Feinberg School of Medicine USA Mary Rinella Northwestern University Feinberg School of Medicine USA Diana Romero (Non-voting) City University of New York Graduate School of Public Health and Health Policy Manuel Romero-Gómez Virgen del Rocío University Hospital Spain Jörn M. Schattenberg University Medical Center Mainz Germany Emmanuel Tsochatzis UCL Institute for Liver and Digestive Health UK Luca Valenti University of Milan Italy Vincent W. S. Wong The Chinese University of Hong Kong China Yusuf Yilmaz Marmara University Inova Health System USA	Henry E. Mark (Project coordinator)	EASL International Liver Foundation	Switzerland
Michael Ninburg Hepatitis Education Project USA Ponsiano Ocama Makerere University College of Health Sciences Uganda Vlad Ratziu Sorbonne University Feinberg School of Medicine USA Mary Rinella Northwestern University Feinberg School of Medicine USA Diana Romero (Non-voting) City University of New York Graduate School of Public Health and Health Policy Manuel Romero-Gómez Virgen del Rocío University Hospital Spain Jörn M. Schattenberg University Medical Center Mainz Germany Emmanuel Tsochatzis UCL Institute for Liver and Digestive Health UK Luca Valenti University of Milan Italy Vincent W. S. Wong The Chinese University of Hong Kong China Yusuf Yilmaz Marmara University Inova Health System USA	Veronica Miller	University of California, Berkeley	USA
Ponsiano Ocama Makerere University College of Health Sciences Uganda Vlad Ratziu Sorbonne University France Mary Rinella Northwestern University Feinberg School of Medicine USA Diana Romero (Non-voting) City University of New York Graduate School of Public Health and Health Policy Manuel Romero-Gómez Virgen del Rocío University Hospital Spain Jörn M. Schattenberg University Medical Center Mainz Germany Emmanuel Tsochatzis UCL Institute for Liver and Digestive Health UK Luca Valenti University of Milan Italy Vincent W. S. Wong The Chinese University of Hong Kong China Yusuf Yilmaz Marmara University Inova Medicine, Inova Health System USA	Phil N. Newsome	University of Birmingham	UK
Vlad RatziuSorbonne UniversityFranceMary RinellaNorthwestern University Feinberg School of MedicineUSADiana Romero (Non-voting)City University of New York Graduate School of Public Health and Health PolicyUSAManuel Romero-GómezVirgen del Rocío University HospitalSpainJörn M. SchattenbergUniversity Medical Center MainzGermanyEmmanuel TsochatzisUCL Institute for Liver and Digestive HealthUKLuca ValentiUniversity of MilanItalyVincent W. S. WongThe Chinese University of Hong KongChinaYusuf YilmazMarmara UniversityTurkeyZobair M. YounossiInova Medicine, Inova Health SystemUSA	Michael Ninburg	Hepatitis Education Project	USA
Mary Rinella Northwestern University Feinberg School of Medicine USA Diana Romero (Non-voting) City University of New York Graduate School of Public Health and Health Policy Manuel Romero-Gómez Virgen del Rocío University Hospital Jörn M. Schattenberg University Medical Center Mainz Germany Emmanuel Tsochatzis UCL Institute for Liver and Digestive Health UK Luca Valenti University of Milan Italy Vincent W. S. Wong The Chinese University of Hong Kong Yusuf Yilmaz Marmara University Inova Medicine, Inova Health System USA	Ponsiano Ocama	Makerere University College of Health Sciences	Uganda
Diana Romero (Non-voting) City University of New York Graduate School of Public Health and Health Policy Manuel Romero-Gómez Virgen del Rocío University Hospital Jörn M. Schattenberg University Medical Center Mainz Germany Emmanuel Tsochatzis UCL Institute for Liver and Digestive Health UK Luca Valenti University of Milan Italy Vincent W. S. Wong The Chinese University of Hong Kong China Yusuf Yilmaz Marmara University Inova Medicine, Inova Health System USA	Vlad Ratziu	Sorbonne University	France
Health and Health Policy Manuel Romero-Gómez Virgen del Rocío University Hospital Jörn M. Schattenberg University Medical Center Mainz Germany Emmanuel Tsochatzis UCL Institute for Liver and Digestive Health UK Luca Valenti University of Milan Italy Vincent W. S. Wong The Chinese University of Hong Kong China Yusuf Yilmaz Marmara University Inova Medicine, Inova Health System USA	Mary Rinella	Northwestern University Feinberg School of Medicine	USA
Jörn M. Schattenberg University Medical Center Mainz Germany Emmanuel Tsochatzis UCL Institute for Liver and Digestive Health UK Luca Valenti University of Milan Italy Vincent W. S. Wong The Chinese University of Hong Kong China Yusuf Yilmaz Marmara University Turkey Zobair M. Younossi Inova Medicine, Inova Health System USA	Diana Romero (Non-voting)		USA
Emmanuel Tsochatzis UCL Institute for Liver and Digestive Health UK Luca Valenti University of Milan Italy Vincent W. S. Wong The Chinese University of Hong Kong China Yusuf Yilmaz Marmara University Turkey Zobair M. Younossi Inova Medicine, Inova Health System USA	Manuel Romero-Gómez	Virgen del Rocío University Hospital	Spain
Luca Valenti University of Milan Italy Vincent W. S. Wong The Chinese University of Hong Kong China Yusuf Yilmaz Marmara University Turkey Zobair M. Younossi Inova Medicine, Inova Health System USA	Jörn M. Schattenberg	University Medical Center Mainz	Germany
Vincent W. S. Wong The Chinese University of Hong Kong China Yusuf Yilmaz Marmara University Turkey Zobair M. Younossi Inova Medicine, Inova Health System USA	Emmanuel Tsochatzis	UCL Institute for Liver and Digestive Health	UK
Yusuf Yilmaz Marmara University Turkey Zobair M. Younossi Inova Medicine, Inova Health System USA	Luca Valenti	University of Milan	Italy
Zobair M. Younossi Inova Medicine, Inova Health System USA	Vincent W. S. Wong	The Chinese University of Hong Kong	China
,	Yusuf Yilmaz	Marmara University	Turkey
Shira Zolhar Sagi University of Haifa School of Public Health Israel	Zobair M. Younossi	Inova Medicine, Inova Health System	USA
oniversity of Figure 201000 of Fublic Health Islaet	Shira Zelber-Sagi	University of Haifa School of Public Health	Israel

disagreement with the statements. The open-ended comment options were again provided to all respondents except those who agreed with a given statement.

A majority of core group members (27 of 33) participated in the online convening following the R2 survey, which permitted in-depth breakout-group discussions on issues that arose in the first two rounds. This convening was hosted and facilitated by Wilton Park, a UK-based global forum for strategic dialogue. Concurrent with revising the statements for R3, the core group developed a draft set of recommendations

to accompany the consensus statements. Preliminary feedback on these recommendations from the larger expert panel was sought over a 1-week period via a shared Google document. The resulting 26 recommendations were included with the final set of 37 statements in R3. Given fairly high levels of agreement in the previous survey rounds, the consensus statements and recommendations in R3 were presented with a binary ('Agree'/'Disagree') response option. A text box at the end of each of the survey domain sections provided respondents with the option to include final comments.

Finally, we assigned each statement and recommendation a grade to indicate the level of agreement, utilizing a grading system recently used in other Delphi studies^{28,29} in which 'U' denotes unanimous (100%) agreement, 'A' 90–99% agreement, 'B' 78–89% agreement, and 'C' 67–77% agreement.

Findings

Here, we report the final statements and recommendations along with a summary of the broader literature as it relates to them.

Consensus statements and recommendations

Across Delphi rounds there was a consistent increase in consensus for all statements. The mean percentage of 'Agree' responses rose from 80.3% in R1 to 90.9% in R2 and 98.5% in R3. The incorporation of substantive comments from respondents into the statements

Table 2 | Expert panel demographic composition and level of engagement

Characteristic	Values
Total	218
Gender ^a	
Man	67.0% (146)
Woman	30.7% (67)
Prefer not to say/no response	2.3% (5)
Primary sector of employment	
Academic	71.1% (155)
Civil society	4.1% (9)
Public	15.1% (33)
Private	4.6% (10)
Other/no response	5.0% (11)
Primary field of employment	
Health-care provider	21.1% (46)
Clinical research	62.8% (137)
Non-clinical research	3.2% (7)
Advocacy	5.0% (11)
Other/no response	7.8% (17)
Geographical representation	
Countries/regions of origin (n)	89
Countries/regions currently based in (n)	91
Delphi process engagement ^b	
Round 1 survey	87.6% (191)
Round 2 survey	88.1% (192)
World Café core group meeting	81.8% (27)
Round 3 survey	84.9% (185)
Participation in one or more components	218
Mean no. of surveys engaged in	2.05

a*Non-binary' and 'gender diverse' were also included as responses options. bNumbers sum to >218 owing to engagement in multiple components of the Delphi process. increased the level of support in subsequent survey rounds. In the end, there was unanimous agreement with 7 statements and >90% agreement with another 30 statements (TABLE 3; translations for TABLE 3 are available in Arabic, Chinese (Mandarin), French, German, Italian and Spanish in Supplementary Tables 2–7). For the associated recommendations, the mean percentage of agreement for the 26 recommendations was 98%. Three recommendations met with unanimous agreement, 22 others with >90% agreement, and the final one with >80% agreement (TABLE 4; translations for TABLE 4 are available in Arabic, Chinese (Mandarin), French, German, Italian and Spanish in Supplementary Tables 2–7).

The human and economic burden

- Statements 1.1–1.6
- Recommendations 1–3

Epidemiology in adults. The global prevalence of NAFLD among adults is estimated to be 23–25% 30,31. The burden varies between and within regions, with the highest prevalence in the Middle East (32%) and South America (30%) and the lowest in Africa (13%)³¹. Up to 20% of people with NAFLD are affected by NASH 31–33. However, there are few reliable epidemiological estimates disaggregated by fibrosis stage, age, gender and geographical location. The need for resource-intensive procedures to accurately assess and determine disease severity is a barrier to population-based surveillance for NAFLD, as is the variety of diagnostic methods and criteria. The availability of good quality data continues to hinder concerted national and global action on NAFLD.

In most populations, the burden of NAFLD increases proportionally with increases in BMI³⁴, although the condition is also common in individuals without overt metabolic risk factors^{19,35}. In the vast majority of patients, NAFLD emerges in the context of metabolic syndrome, with insulin resistance an important pathophysiological mechanism¹⁷. NAFLD prevalence is higher among patients with T2DM than in the general population, whereas T2DM incidence is higher in patients with NAFLD^{17,24,36,37}. Driven by increasing prevalences of obesity and T2DM and by ageing populations, the global NAFLD burden is projected to grow in the coming decade^{32,33}.

Between 1990 and 2017, global deaths due to cirrhosis increased from 899,000 to 1.32 million while disability-adjusted life years increased from 30.5 million to 41.4 million. During this period, the number of prevalent cases of compensated cirrhosis due to NASH more than doubled, whereas for decompensated cirrhosis the figure more than tripled. With the expansion of prevention and treatment measures for hepatitis B and C, NASH is expected to overtake them soon as the leading cause of cirrhosis³⁸.

Epidemiology in children and adolescents. Epidemiological data on NAFLD in children and adolescents are scarce. There is marked heterogeneity in the findings of available studies, due in part to variations in

Table 3	Consensus s	tatements for	· a NAFLD	public health agenda

Number	Statement	Grade		
1. The hur	nan and economic burden			
1.1	According to current estimates, 20–25% of the global adult population is affected by NAFLD, and an estimated 20% of people with NAFLD will develop NASH. However, robust epidemiological estimates, disaggregated by fibrosis stage, age, gender, risk profile and geographical area, are limited. Incomplete data hinder concerted action at the national and global levels.	A		
1.2	Data from central registries, electronic health-care records or official statistics are available for certain countries and can be useful sources of information. However, differences in reporting, including the use of different administrative codes (for example, the International Classification of Disease (ICD) codes), limit comparability.	A		
1.3	Data on paediatric NAFLD are scarce. Prevalence estimates vary widely, whilst there is limited information on long-term health outcomes in children living with NAFLD. However, available data indicate that NAFLD is an increasing problem in paediatric populations and is especially prevalent in children with obesity.	U		
1.4	A wide range of factors needs to be considered in developing prevention and treatment approaches for NAFLD. These factors extend from metabolic risks, including insulin resistance, to genetic, social and environmental influences that may play a part in the development and progression of the disease.	U		
1.5	NAFLD shares a bidirectional relationship with other metabolic conditions. Addressing NAFLD will likely reduce the prevalence and severity of these conditions.	Α		
1.6	There are both economic and social arguments for taking action on NAFLD. Evidence shows that NAFLD progression is associated with substantial health-care costs, socioeconomic losses and reduced quality of life, most notably in patients with advanced fibrosis and cirrhosis. Early intervention could help reduce the burden of disease, associated health-care costs and economic losses.	U		
2. Awarer	ness and education			
2.1	Communicating about NAFLD and its consequences has proved to be a major challenge for the liver health community.	Α		
2.2	Raising the profile of NAFLD as a public health issue will require clear messages about the condition, its consequences and what action is required. These messages should be tailored to specific audiences, including the liver and gastroenterology communities, primary care providers, specialists from other relevant disciplines, as well as stakeholders such as at-risk groups, the media and policy-makers.	A		
2.3	Primary care providers and diabetes specialists can play a critical part in identifying and referring patients with advanced fibrosis to liver specialists. Raising the awareness of these medical providers would improve their ability to play this part.	А		
3a. Defini	ng and implementing models of care: general considerations			
3.1	Given the broad disease spectrum of NAFLD and the different levels of care required by patients across this spectrum, having clearly defined, context-specific models of care will be important for addressing the disease burden.	Α		
3.2	The majority of people living with NAFLD can be managed in primary care; only patients with advanced disease need referral to a liver specialist. NAFLD care pathways can guide care decisions, including decisions on when to refer a patient to specialist care.	A		
3.3	People living with NAFLD, especially those with advanced fibrosis, commonly require the management of multiple comorbid conditions, including diabetes, obesity and cardiovascular disease.	Α		
3.4	There is limited evidence on the impact of different NAFLD models of care on patient outcomes and cost-effectiveness. The lack of evidence and of investment in implementation research continues to impede the design and delivery of good care in different health-care settings and contexts.	A		
3.5	Fibrosis stage is an important predictor of long-term liver-related outcomes and overall mortality in people living with NAFLD. Evidence of advanced fibrosis is an adequate indicator of a patient's need for referral to specialist liver care.	А		
3.6	Non-invasive tests (NITs) can be effective at excluding advanced fibrosis and the need for further assessment or referral to specialist liver care, especially when combinations of NITs are used sequentially.	А		
3.7	The availability and use of different NITs vary among health-care settings. Non-commercial blood-based scores could be feasibly implemented in most primary and secondary care settings, such as diabetes clinics, if they were more readily available and widely known.	А		
3.8	People living with type 2 diabetes mellitus (T2DM) and/or obesity are recognized as being at high risk for NAFLD-related complications. Collaboration and coordination across the different components of the health-care system will be needed to care for these patients most effectively.	A		
3b. Defini	3b. Defining models of care: considerations for children (younger than 18 years)			
3.9	The natural history of paediatric NAFLD is poorly understood, due to a lack of prospective studies and the complex nature of the disease, including pathologies that are unique to children living with NAFLD. Better data on the natural history, pathophysiology and risk factors for disease progression would improve the care of this population.	U		
3.10	Models of care for children should address all care needs, including the provision of psychological support, and be designed to facilitate the smooth transfer of care from paediatric to adult services.	A		

Table 3 (cont.) Consensus statements for a NAFLD public health agenda	
Number	·	Grade
	uing models of care: considerations for children (younger than 18 years) (cont.)	Grade
3.11	The lack of validated NITs for use in children is a barrier to timely diagnosis and linkage to care.	А
3.12	Available data show that paediatric NAFLD is associated with both hepatic and non-hepatic morbidity and mortality. Children living with NAFLD might benefit from multidisciplinary management approaches tailored to their unique health-care needs.	A
3c. Defin	ing models of care: considerations for low-resource settings	
3.13	In low-resource settings, the availability of diagnostic tools — including NITs — is likely to be limited, especially the more expensive imaging-based tests. Diagnosis in these settings will often require practitioners to make pragmatic choices and resort to low-cost solutions.	A
3.14	NAFLD is not mentioned in the current guidelines from the WHO on the detection, diagnosis and treatment of major non -communicablediseases (NCDs) in primary care in low-resource settings. Inclusion of NAFLD in such guidance would help to improve care for affected populations in these settings.	A
4. Treatn	nent and care	
4.1	Interventions aimed at modifying lifestyle risk factors are the cornerstone of NAFLD treatment. There is some evidence that these interventions can prevent disease progression and can, in some cases, reverse fibrosis, yet more data will help to identity the most effective approaches and how to implement them in clinical practice.	А
4.2	As the number of effective pharmacological treatments for NAFLD increases, programmes aimed at modifying lifestyle risk factors will continue to be a core element of NAFLD disease management.	A
4.3	Access to treatment programmes for NAFLD requires that they be incorporated into relevant national health-care policies and guidelines and be adequately funded. Private and public payers and/or funders have a key part to play in ensuring financial support (for example, reimbursement) for these services.	A
4.4	The invasive nature of liver biopsy, the inherent variability of histological findings and the lack of an alternative validated surrogate for long-term clinical benefit have complicated the development of efficacious treatments for NAFLD.	A
5. Patien	t and community perspectives	
5.1	People living with NAFLD can provide valuable insights into the design and implementation of interventions to safeguard and improve their health. Patients and patient organizations should be actively involved in developing policies and strategies to address NAFLD; however, few such groups currently address NAFLD.	A
5.2	Given that NAFLD is a largely invisible public health issue, high-profile patients can be especially useful in creating awareness and advocating for greater action on prevention and treatment.	Α
5.3	Professional and patient organizations that address NCDs, including T2DM, obesity, heart disease and cancer, can play an important part in raising the profile of NAFLD, including by providing information to at-risk groups.	U
5.4	Stigma can be a major barrier when seeking to address health issues. Liver disease in general is commonly associated with unhealthy alcohol use, while NAFLD is associated with obesity. Both of these associations are with highly stigmatized conditions, and the implications of such stigma need to be acknowledged and addressed when developing prevention and treatment approaches for NAFLD.	A
,	strategies and a societal approach	
6.1	A national strategy for NAFLD is lacking in almost every country in the world, while NAFLD is explicitly mentioned in very few national strategies or clinical guidelines for related conditions such as obesity or diabetes. This fact highlights the extremely low priority the condition has in national health agendas, and the need for a concerted effort to shape and deliver a robust public health response.	U
6.2	Several highly prevalent NCDs share common risk factors — such as unhealthy diets, physical inactivity and unhealthy alcohol consumption — with NAFLD. Policies, fiscal measures and legislation could address many of these diseases in a coordinated, simultaneous way.	A
6.3	Addressing NAFLD will require collective action that spans diverse disciplines and sectors. Existing frameworks such as the United Nations Sustainable Development Goals (SDGs) can usefully inform and guide the development of multi-sectoral efforts to address the direct, underlying and cross-cutting causes of NAFLD.	A
	ship for the NAFLD public health agenda	
7.1	National and regional liver associations, in collaboration with governments and other stakeholders, have a leading role in responding to NAFLD, including in developing public health strategies and guidelines and in collaborating with other disease associations and organizations.	A
7.2	Multilateral organizations such as the WHO also have a key role in shaping and helping lead the response to NAFLD, firstly by recognizing the condition as a major health issue, and secondly by supporting nationally led efforts to deliver public health responses.	A
7.3	Global efforts to expand universal health coverage and ensure that health systems are people-centred provide a useful mechanism for holistically addressing NCDs, including not only NAFLD, but also associated diseases such as diabetes and obesity.	U
	•	

Grading system: U, denotes unanimous (100%) agreement; A, 90–99% agreement; B, 78–89% agreement; and C, 67–77% agreement. NAFLD, non-alcoholic fatty liver disease; NASH, non-alcoholic steatohepatitis; WHO, World Health Organization.

Number	Recommendation	Grade
1. The hun	nan and economic burden	
1	Investment is needed in research that will improve understanding of NAFLD epidemiology, especially in under-studied population groups such as children, and people without overt metabolic risk factors.	A
2	In the absence of population-based and prospective longitudinal studies, alternate research methods should be considered, such as those employing electronic health records.	Д
3	Investment cases should be developed for NAFLD at global, regional and local levels. To support these cases, toolkits should be prepared to provide guidance on obtaining the requisite economic data and communicating the findings to policy-makers, health-care funders and/or payers and other relevant stakeholders.	А
2. Awaren	ess and education	
4	Professional societies and other relevant stakeholders, such as patient organizations, should collaborate on a transparent process to carefully reconsider the nomenclature of fatty liver diseases, with special attention to the benefits of and barriers to changing the name of 'non-alcoholic fatty liver disease'.	А
5	The liver health community should engage health communication experts to jointly develop effective strategies and practical tools to increase awareness in key audiences, including the media and policy-makers.	А
6	The terminology and concept of 'compensated advanced chronic liver disease' should be adopted, as it better reflects the continuum of advanced disease and the increased risk of decompensation than the current usage of fibrosis stages 3 and 4.	В
7	Professional bodies should develop simple knowledge products and educational courses targeting the liver and gastroenterology communities, primary care providers and specialists from other disciplines, as well as at-risk populations, the media and policy-makers. The courses should include medical school and continuing medical education activities.	А
3. Definin	g and implementing models of care	
8	Health-care planners and providers should design and implement locally feasible NAFLD care pathways, utilizing available tests to efficiently determine a patient's care needs and link them to appropriate services.	U
9	Health-care providers — especially primary care providers, diabetes specialists and those caring for people living with obesity — should be equipped with the tools and knowledge needed to support the care of people living with NAFLD. At a minimum, providers should be able to identify which patients require referral to a liver specialist.	A
10	Multidisciplinary care models should form the basis for managing people living with NAFLD, especially those with advanced fibrosis.	U
11	Research should focus on developing more effective and more accurate non-invasive tests (NITs) for risk-stratifying patients — including children — in primary care, and for staging fibrosis and diagnosing NASH in secondary care.	U
12	Implementation research should be undertaken to better understand the barriers to uptake of currently available NITs.	А
13	Active case finding should be considered in population groups at high risk for advanced fibrosis. The specific target populations ought to be determined locally but should include people living with type 2 diabetes mellitus and central adiposity.	A
14	Implementation research is needed to identify the core elements of effective NAFLD care models in different health-care settings — including low-resource settings — and to provide generalizable findings that can inform the development of models of care in different contexts.	A
15	Preventing and treating childhood NAFLD should be a priority, both as a means of improving child health and as a way of reducing the burden of disease in later life.	А
4. Treatme	ent and care	
16	Research should focus on identifying interventions, including lifestyle treatments (for example, diet and physical activity regimens) and pharmacological treatments that can help people living with NAFLD and obesity to achieve and sustain a weight loss of at least 10%.	А
17	Effective structured lifestyle treatment programmes should be made available to people living with NAFLD, especially those who are at high risk of advanced fibrosis and/or rapid fibrosis progression.	Д
18	Currently accepted surrogate histological end points for conditional NASH drug approval should be standardized, with the goal of eventually replacing them with non-invasive diagnostic and surrogate end point biomarkers.	Д

Table 4 (cont.) | Consensus recommendations for a NAFLD public health agenda

Number	Recommendation	Grade	
5. Patient and community perspectives			
19	Medical associations and other stakeholders should support patient groups in meeting the needs of people living with NAFLD. Where possible, NAFLD-specific groups should be formed. Patient groups focused on related conditions — including diabetes and obesity — should be provided with relevant information on NAFLD to share with their members.	А	
20	Patient groups for liver disease and related non-communicable diseases (NCDs) should be involved in the development of clinical practice guidelines for NAFLD. Medical associations should also support these patient groups in developing relevant materials on NAFLD for their members.	А	
6. Policy st	rategies and a societal approach		
21	Efforts to detect, prevent and treat NAFLD should be integrated within a broader package of cost-effective interventions that holistically address NCD risk factors, focusing specifically on unhealthy diets, physical inactivity and unhealthy alcohol consumption.	Α	
22	Global health organizations (including the WHO) and national institutions should incorporate NAFLD into their technical materials on NCDs and include NAFLD among their priority NCDs.	А	
23	The WHO should dedicate a World Health Day (7 April) to liver health to highlight the global prevalence of NAFLD and its significance for public health.	А	
24	The NAFLD prevention agenda should include the creation of healthier, more equitable and sustainable societies as one of its core goals. One way to do that should be to emphasize the SDG targets that are relevant to preventing and treating NAFLD.	Α	
7. Leadership			
25	A global coalition of organizations and individuals should lead the development of a NAFLD public health roadmap and support the global health community in following it.	Α	
26	Medical societies that provide care for any aspect of metabolic syndrome should formally collaborate to address NAFLD, including by jointly developing guidelines, policy briefs and plans of action.	Α	

Grading system: U, denotes unanimous (100%) agreement; A, 90–99% agreement; B, 78–89% agreement; and C, 67–77% agreement. NAFLD, non-alcoholic fatty liver disease; NASH, non-alcoholic steatohepatitis; WHO, World Health Organization.

study settings, the race and ethnicities of the studied populations and the reference methods used to define NAFLD. A 2015 meta-analysis estimated the prevalence of NAFLD in children aged 1–19 years at 7.6% (95% CI 5.5–10.3%), rising to 34.2% (95% CI 27.8–41.2%) in studies conducted in paediatric obesity clinics³⁹. NAFLD prevalence is generally higher in children living with obesity than in those without obesity^{39–44}, but the extent of this relationship is likely to differ by population group⁴⁵. NAFLD is also a public health problem in children and adolescents with normal weight⁴⁴. Driven in part by rising obesity levels, the burden of childhood NAFLD has increased over the past three decades with an estimated annual change of 1.35% (95% CI 1.16–1.54%)⁴⁶.

Understanding the natural history, pathophysiology and phenotypes of childhood and adolescent NAFLD has advanced in the past two decades, including through articulation of clinically relevant subtypes of paediatric NASH⁴⁷⁻⁴⁹. NAFLD in children with T2DM has a unique pathological phenotype, which seems to be more aggressive than the adult form⁵⁰. Further research is still needed to elucidate the pathophysiology, genetics, natural history and responses to treatment in paediatric NAFLD⁴⁷ and therefore inform prevention and management approaches.

There are fewer data on the long-term impact of NAFLD in childhood than of NAFLD developed in later life. A Danish study estimated that for every 1 unit

increase in BMI for-age Z-score between the ages of 7 and 13 years, the risk of cirrhosis increased by 16%⁵¹. Another study in the same age group showed that a 1 unit increase in BMI increased the risk of liver cancer 30 years later by 20-30%52. Weight gain in childhood or late adolescence is associated with a greater risk of NAFLD than weight gain in late adulthood20, although a high BMI in late adolescence increases the risk of severe liver disease in adulthood, independent of alcohol consumption^{53,54}. A study of paediatric and young adult patients with biopsy-confirmed NAFLD in Sweden showed that, compared with matched controls, the patients with NAFLD had substantially higher rates of all-cause, cancer, liver and cardiometabolic-specific mortality⁵⁵. More data on the long-term consequences of childhood NAFLD, including the life-time risk of developing cirrhosis, will help to inform strategies for prevention and management.

Quality of life. NAFLD research has started to explore the effect of the disease on affected populations using patient-reported outcome (PRO) data. PROs enable researchers and clinicians to look beyond clinical and histological outcomes to understand better the full impact of a condition. PROs capture health status from the perspective of the patient, from general quality of life (QoL) and HRQoL to work productivity, fatigue and satisfaction. Such information enables a comprehensive understanding of disease impact at the individual and

Box 1 | The NAFLD name debate

Since the early 2000s, several proposals have been made to change the name non-alcoholic fatty liver disease (NAFLD). The central arguments for change have been that the adjective 'non-alcoholic' is an unhelpful construction, and that other terms would better reflect the metabolic underpinnings of the disease's aetiology¹⁴⁷. In the absence of widespread consensus, however, NAFLD has remained the commonly used nomenclature. In the past 2 years, the term 'metabolic dysfunction-associated fatty liver disease' (MAFLD) has gained traction as a possible replacement. An international group of experts from 22 countries reached consensus on the change to MAFLD^{148,149}, and the proposed change was endorsed by regional liver associations in South America¹⁵⁰ and the Asian Pacific¹¹⁵ as well as by experts in sub-Saharan Africa¹⁵¹ and the Middle East and North Africa¹⁵². However, other experts have expressed concerns about prematurely changing the name without fully considering its broad implications, from diagnostic criteria to trial end points, calling instead for regional liver societies to work together to reach consensus^{153,154}.

societal levels. Several PRO tools have been developed and validated for use in people living with NAFLD⁵⁶⁻⁵⁸. Overall, QoL worsens with disease progression⁵⁹. People living with NAFLD report worse QoL than those living without the disease, people living with NASH report worse QoL than those with NAFL^{60,61}, and patients with cirrhotic NASH report worse HRQoL than patients with non-cirrhotic NASH². The association between disease stage and HRQoL varies among countries and regions³, highlighting the importance of local data. Research should aim to further our understanding of the outcomes most relevant to people living with NAFLD, so that policies and management strategies can be designed to minimize the effects of the disease on those affected.

Economic burden. In addition to the human burden, NAFLD also has wide-ranging economic implications for affected populations and societies at large^{3,4,6-8}, including both direct medical expenses and indirect costs associated with consequences such as loss of work. Most economic costs associated with NAFLD are incurred in the latter stages of the disease^{4,7,8}, providing a good rationale for funding prevention and early intervention efforts. Investment cases should be developed for NAFLD at global, regional and local levels. To support their development, toolkits should be prepared to provide guidance on obtaining the requisite economic data and communicating the findings to policy-makers, health-care funders and payers and other relevant stakeholders.

The vast human and economic impact of NAFLD provides a compelling imperative for action. More and better data on NAFLD, especially in under-studied populations such as children, are needed to advance our understanding of the impact of the disease and to shape health system and public health responses accordingly. Data disaggregated by disease stage, gender, age, ethnicity and geographical area will be critical. In the absence of population-based and longitudinal studies, alternative research methods should be explored. Electronic health records are one potentially valuable resource⁶². The latest efforts to standardize the administrative codes used to record exposures and outcomes for NAFLD will improve the feasibility of such research and facilitate comparisons between study populations⁶³. As our understanding of

the basic science and epidemiology of NAFLD grows, it will also be important to explore the effectiveness of different operational models on patient outcomes and resource utilization.

The lack of data on the human and economic burden of NAFLD not only inhibits our ability to deliver proportionate health system and public health responses, but to raise awareness of the disease and its consequences among key stakeholders, including policy-makers and at-risk groups. As we strive to better understand the epidemiology of NAFLD, the liver health community will also need to consider how to communicate these findings to different target audiences.

Awareness, education and terminology

- Statements 2.1–2.3
- Recommendations 4-7

Despite being the most prevalent liver disease in history, NAFLD remains largely unknown outside hepatology and gastroenterology. Knowledge of NAFLD among general practitioners⁶⁴ and non-liver health specialists is generally poor, with little sense of the scale of the challenge or the potential gravity of the disease⁶⁵. Patients at higher risk of NAFLD, including people with T2DM and other metabolic risk factors, are also unaware of the disease, their susceptibility to developing it or how it interacts with other metabolic conditions^{66–68}. There are limited data on NAFLD awareness amongst the general public, but what is available points to low levels of awareness⁶⁹.

Increasing awareness of NAFLD will require simple, effective messages and non-stigmatizing terminology that describe risk factors and potential consequences of the disease. Such messages need to be targeted to specific audiences, including health-care professionals — especially hepatologists, gastroenterologists, primary care providers and diabetes specialists — policy-makers and the general public. Health communication experts and the media should be enlisted in developing awareness strategies and tools.

In addition, the liver health community needs to agree upon the terminology we use to describe the disease and its consequences. Compensated advanced chronic liver disease (cACLD) is a relatively new term for the early phases of severe chronic liver disease, covering severe fibrosis and compensated cirrhosis⁷⁰. Adopting this term would improve clinical care and research, as cACLD better reflects the continuum of advanced disease and the increased risk of decompensation than the current use of fibrosis stages 3 and 4 (REFS^{70,71}). Long-standing debates about the nomenclature used for fatty liver disease have also gained traction in the past few years, with 'metabolic dysfunction-associated fatty liver disease' (MAFLD) as a possible replacement for NAFLD (BOX 1). We urge the relevant organizations to engage in a thorough process to achieve consensus on the path forwards. The current lack of clarity risks fragmenting and confusing the liver health community, which would undoubtedly impede efforts to bring much needed attention and action to this critical public health issue. Beyond the clinical and scientific considerations, such a process should also address how a name change might facilitate efforts to increase

awareness about the disease in an audience that is as wide as possible.

Defining and implementing models of care

- Statements 3.1-3.12
- Recommendations 8-15

A model of care (MoC) is a setting-specific framework that outlines how patients with a disease are managed along the care cascade. A comprehensive MoC outlines which services are to be provided, where they should be provided and by whom, and how they are to be integrated and coordinated within a health-care system⁷². Clearly defined, context-specific MoCs will be important for managing the burden of NAFLD, and establishing such MoCs should be a key focus for health-care decision-makers and providers. Yet NAFLD MoCs have received little attention to date, with a review published in 2021 identifying only seven published examples of comprehensive MoCs, only one of which addresses children⁷³.

The majority of patients with NAFLD can be managed in primary care. For patients with isolated steatosis or early-stage fibrosis in the primary care setting, management should focus on preventing disease progression and the development or exacerbation of metabolic comorbidities. Patients with advanced fibrosis might require a hepatologist or gastroenterologist to manage the hepatic component of the disease^{74,75}, whereas a smaller proportion will require tertiary care, such as for transplant surgery^{13,14}.

As a multisystem, comorbid disease, people living with NAFLD will often benefit from multidisciplinary care, especially those with advanced fibrosis⁷⁶. Establishment of multidisciplinary teams (MDTs) can be an effective way to manage the diverse clinical needs of people living with NAFLD⁷⁶. There are several published examples of multidisciplinary secondary care clinics for NAFLD⁷⁷⁻⁸¹. Each provides a model of what is feasible and appropriate within a given health-care setting, with the composition and structure of the MDT and the services it provides varying accordingly.

Development of care pathways. The first step in an MoC is to identify each patient's needs, as determined by disease stage and presence of comorbidities, and to link them to appropriate services — a process known as risk stratification. However, diagnosing and staging NAFLD remains challenging, and diagnoses are often incidental to the identification of abnormal liver enzymes or of steatosis through imaging techniques, neither of which provides information on disease severity⁸².

A care pathway is a framework to support decision-making, including deciding when to refer a patient to specialist care. There are several published examples of care pathways for identifying advanced liver disease 80,81,83-87, and some evidence for the cost-effectiveness of these approaches 88-90. Yet formal pathways do not exist in many health-care settings, and non-invasive tests (NITs) are not routinely used in some settings where they might prove beneficial. Although the availability of specific NITs will vary, it is feasible to implement non-commercial blood-based tests in most

primary and secondary care settings. Managing the burden of NAFLD requires developing locally appropriate care pathways and equipping health-care providers with the tools and knowledge to implement them. That is especially true for primary care providers, as many people living with NAFLD will first present in primary care, where the condition is widely under-diagnosed91. Another key setting is diabetes clinics, where the prevalence of advanced disease is higher than in the general population^{18,24}. Care pathways will also ensure that the necessary health-care infrastructure is in place when more effective pharmacological treatments become available. In the case of hepatitis C, such pathways were not adequately in place before all-oral direct-acting antiviral treatment became available, hampering efforts to link people with treatment^{72,92}.

Care pathways for children living with NAFLD should also address the transition from paediatric to adult services to ensure continuity of care. These pathways need to recognize the differences in the clinical management of children and adults and the psychological factors associated with such a transition⁹³.

Testing and screening for NAFLD. Fibrosis stage is a key indicator for long-term liver and non-liver health outcomes in patients with NAFLD⁹⁴. Various NITs have been validated for detecting advanced fibrosis in clinical practice, ranging from blood-based scores to imaging techniques⁹⁵. The performance of these NITs is strongly influenced by pretest probability. In primary care settings where the population prevalence of advanced disease is low, the negative predictive value of NITs for advanced fibrosis is generally high, whereas the positive predictive value is lower^{96,97}. NITs can be especially effective at identifying advanced disease when used in sequential algorithms^{98–101}. There is also some evidence that certain combinations of NITs can identify patients who have fibrosis stage 2 or greater with a high positive predictive value¹⁰². Although several NITs have been investigated for use in paediatric populations, none of them is currently validated for use in routine clinical practice. Initial screening in children generally relies on liver enzymes and ultrasonography, with a biopsy required to definitively diagnose and stage the disease. There is hope that NIT combinations might replace the need for biopsies in paediatric populations in the near future¹⁰³. The development of more efficient and effective NITs for risk-stratifying patients in primary care and diagnosing and staging NASH in secondary care remains a research priority.

There is broad consensus that certain factors, particularly T2DM and obesity, increase the risk of an individual developing NAFLD and of the disease progressing. However, guidance varies on the benefits and cost-effectiveness of active case finding in specific patient groups (BOX 2). Although appropriate targets for active case-finding should be determined with local epidemiology and resources in mind, the expert panel recommends that it include people living with T2DM and those with central adiposity. These approaches should be evaluated for their impact on patient outcomes and for cost-effectiveness.

Box 2 | Guidance on active case-finding in high-risk population groups

Joint guidance from The European Association for the Study of the Liver (EASL), the European Association for the Study of Diabetes (EASD) and the European Association for the Study of Obesity (EASO) recommends screening for non-alcoholic fatty liver disease (NAFLD) in people with obesity, metabolic syndrome and in particular type 2 diabetes mellitus (T2DM)¹¹¹. The Latin American Association for the Study of the Liver (ALEH)¹¹², the Asian Pacific Association for the Study of the Liver (APASL)¹¹⁵ and the Asia–Pacific Working Party on NAFLD¹⁵⁵ all recommend considering screening in certain high-risk populations, including those with obesity and T2DM. The American Diabetes Association recommends screening for non-alcoholic steatohepatitis and advanced fibrosis in patients with elevated liver enzyme levels or hepatic steatosis on ultrasonography¹⁵⁶. By contrast, the American Association for the Study of Liver Diseases (AASLD) does not recommend systematic screening in these groups, given the lack of cost-effectiveness data for such efforts¹¹³.

Primary care interventions. Access to high-quality primary care preventive interventions is critical to reducing the burden of NCDs104, yet there is little evidence for which primary care interventions will optimize patient outcomes for people living with NAFLD. However, the common risk factors for NAFLD, obesity, T2DM and CVD, including an unhealthy diet and physical inactivity¹⁰⁵, argue for integrated chronic disease management approaches. Structured management programmes for other conditions, such as diabetes, can serve as a starting point for more integrated models¹⁰⁶. In low-resource settings, the World Health Organization (WHO) package of essential NCD interventions for primary health care can be used as a basis for integrating NAFLD care into related disease areas, including diabetes management¹⁰⁷. Technological innovation, such as health information exchanges and mHealth (mobile health) applications, can also help facilitate collaboration between patients and providers and the coordination of services within a health-care system by ensuring the timely and accurate flow of information108.

As the liver community leads efforts to improve the life of those with NAFLD, it should prioritize operational research that furthers our understanding of the effect of different MoCs on patient outcomes and of the cost-effectiveness of these approaches in different health-care settings. This research should also address the structural barriers that make coordination and collaboration within health-care systems a challenge and how to effectively engage across disciplines.

Treatment and care

- Statements 4.1-4.4
- Recommendations 16-18

The treatment and care of patients with NAFLD are highly dependent on their disease stage⁷⁵. Interventions aimed at modifying lifestyle risk factors — namely weight, diet and physical activity — and at the management of comorbidities should be the cornerstone of treatment for all patients^{75,109,110}. This priority is emphasized in the clinical management guidelines from regional liver associations¹¹¹⁻¹¹⁵. In patients with more advanced disease, addressing components of metabolic syndrome, liver-related pharmacotherapy and management of cirrhosis-related complications are all important^{75,116}. Even when effective pharmacological treatments for NAFLD become available,

programmes aimed at modifying lifestyle risk factors should continue to be a core element of NAFLD disease management.

There is some evidence that such lifestyle interventions can prevent disease progression and, in some cases, reverse fibrosis 117,118. In persons affected by overweight and obesity, NAFLD lifestyle interventions aim to achieve and sustain a weight loss of around 10%, which is associated with the improvement of liver enzyme levels and histological findings111-113. Behavioural change approaches are most effective when incorporated into a comprehensive, long-term lifestyle modification programme¹¹⁹. Dietary guidance for people living with NAFLD generally centres on the reduction of saturated fats, sugar-sweetened beverages, refined carbohydrates and red and processed meats 120,121. The Mediterranean diet and the Dietary Approaches to Stop Hypertension (DASH) have proven beneficial in some patients by improving liver status, in particular hepatic insulin sensitivity and lipid profile 120,122. Several forms of physical activity — aerobic, resistance or high-intensity intervals — seem to have a beneficial effect on liver fat^{118,123}. Even in the absence of weight loss, exercise can result in a 20–30% reduction in intrahepatic lipid levels^{117,118}. It is important that diet and exercise programmes are tailored to the patient's needs and preferences to support long-term adherence. Further research is needed to identify the interventions, whether lifestyle or pharmacological interventions, that are most effective in helping people living with NAFLD and obesity to achieve and sustain a weight loss of at least 5-10% of initial body weight. Research is also needed to determine how best to implement lifestyle interventions, including research on how different operational models influence long-term adherence and patient outcomes, and on the cost-effectiveness of different approaches. Collaboration between disciplines, including basic, behavioural and clinical sciences and operations researchers, among others, will help to advance our understanding in this area in the years to come.

Making effective structured lifestyle treatment programmes available to people with NAFLD, especially those who are at high risk of advanced fibrosis and/or rapid fibrosis progression, such as people living with NAFLD, obesity and T2DM, should be a priority of the liver health community. Both public and private funders will have a key part to play in ensuring financial support for such services. As a first step, NAFLD needs to be adequately incorporated into relevant national health-care policies and guidelines, something that is currently lacking in most countries¹²⁴.

Although there are currently no pharmacological treatments specifically approved for NAFLD, clinical trials are exploring numerous drug candidates targeting energy intake, energy disposal, lipotoxic liver injury, inflammation and fibrosis¹²⁵. The invasive nature of liver biopsy, the inherent variability of histological findings and the lack of an alternative validated surrogate for long-term clinical benefit have complicated the development of efficacious treatments. As the field moves forwards, it would be helpful to standardize the surrogate histological end points that are currently accepted

for conditional NASH drug approval, with the goal of eventually replacing them with non-invasive diagnostic and surrogate end point biomarkers.

Patient and community perspectives

- Statements 5.1-5.4
- Recommendations 19-20

People with NAFLD can provide valuable insights into the design and delivery of interventions to safeguard and improve their health. Actively engaging people with lived experiences, especially disproportionately affected communities such as certain minority ethnic groups, and considering their perspectives will help ensure that interventions are patient-centred, improving treatment adherence and outcomes ^{126,127}. Developing education materials that are specific to population groups and reflect diverse backgrounds will also improve outreach and engagement efforts.

Patient groups have a critical role in advocating for greater attention to under-served health issues and as a source of information for the affected population (BOX 3). In thinking about developing a patient community for NAFLD, the history of the viral hepatitis movement is instructive. Patient organizations had a pivotal role in that movement, notably through driving World Health Assembly Resolution 67.6, which was adopted in May 2014 (REF. 128). The resolution framed viral hepatitis as a major public health problem that required comprehensive global and national action. The World Hepatitis Alliance was key to this success, providing a platform that united diverse organizations around a common vision. Although there is no global patient alliance for NAFLD, there are platforms within the wider NCD community, such as the NCD Alliance, that liver health organizations can engage with to increase awareness of NAFLD. These efforts can complement the ongoing work of organizations within the liver health community.

One challenge in engaging people living with NAFLD is the fear of stigma. People commonly connect liver disease with unhealthy alcohol use, while NAFLD is often connected with obesity and its associated stereotypes (such as laziness); both of these conditions are highly stigmatized^{28,129–131}. People living with multiple chronic conditions can also experience multiple interacting

Box 3 | The importance of patient groups

Patient organizations are often a primary source of information for affected populations whilst also providing a platform through which patients can engage in developing policies and strategies, including clinical practice guidelines. However, globally, few non-alcoholic fatty liver disease (NAFLD)-specific patient groups exist and there is no truly global platform or coordination mechanism to support local patient organizations. The liver community should further mobilize financial and technical support to help form such groups. We suggest that professional and patient organizations that address associated conditions such as obesity, type 2 diabetes mellitus, heart disease and cancer can also have an important role in disseminating information to people in high-risk groups. Medical associations will need to lead efforts to develop information tools and support the dissemination of these amongst key target groups. As the meaningful engagement of people with lived experiences is becoming more mainstream in the non-communicable disease field, including within the World Health Organization (WHO)¹⁵⁷, the liver health community should take the opportunity to prioritize this issue into the NAFLD agenda.

forms of stigma¹³². The implications of stigma need to be acknowledged and addressed when developing prevention and treatment approaches for NAFLD. High-profile individuals living with NAFLD could have a great impact in creating awareness and advocating for prevention and treatment efforts, as well as in reducing any stigma associated with the condition.

Policy strategies and a societal approach

- Statements 6.1-6.3
- Recommendations 21-24

Despite being a highly prevalent liver disease, little attention has been paid to the policies and strategies needed to prevent, manage and treat NAFLD. A review of 29 European countries examining the existence of policies for NAFLD found large variations in national responses, and none of the countries was prepared to address the challenge^{124,133}. A global study of 102 countries painted a similar picture, highlighting an overall lack of attention to NAFLD in national health agendas; not a single country reported having a written NAFLD strategy. Even in national strategies and clinical guidelines for related conditions such as obesity or T2DM, NAFLD is seldom mentioned. These findings highlight the extremely low priority the condition has in both disease-specific and national health agendas, and the need for a concerted effort to shape and deliver a robust public health response¹³⁴. Fortunately, there is some cause for optimism, with efforts such as the US NASH action plan by the NASH Council and the Global Liver Institute published in December 2020 that provides a model for others to consider¹³⁵.

At a health system level, chronic disease management is driving the reorientation of health systems away from siloed disease-centred models to multidisciplinary patient-centred care^{136,137}. The liver community can, in collaboration with other actors working with metabolic disease management, help lead this process in the coming years for the benefit of not only patients with liver disease but of all people living with NCDs.

At both a public health level and a clinical management level, there is substantial overlap in the measures required to address NAFLD and the other major NCDs. Common risk factors, such as unhealthy diets, physical inactivity and unhealthy alcohol consumption, provide an opportunity for collaborative approaches to improve public health. Policies, fiscal measures and legislation that address common risk factors for NCDs in a coordinated, synergistic way have the potential to create a lasting impact. Yet despite the common approaches needed to address NAFLD and other NCDs, NAFLD is not mentioned by name in the majority of key global or national NCD strategies; most notably, it is absent from the WHO Action Plan on the Prevention and Control of NCDs²⁵. Liver health organizations must engage more effectively with WHO and other national and international organizations to ensure that measures to prevent and treat NAFLD are fully integrated into a broader package of cost-effective interventions that address NCD risk factors holistically. Each year World Health Day is marked on 7 April and the theme changes each year, providing the opportunity to shine a spotlight

on key issues. Dedicating a World Health Day to liver health would provide a platform for NAFLD advocacy and awareness-raising within the global health field and beyond. Such a day would complement existing advocacy and awareness efforts, including International NASH Day, which is marked on 12 June each year.

Complex health issues also require us to rethink systems and go beyond the immediate determinants of a disease to consider the underlying influences and root causes, as well as the multidisciplinary and multisectoral responses needed to address these¹³⁸. The liver health field must look beyond the health sector as it seeks to address the challenges of NAFLD. It can take lessons from other fields such as obesity, in which thinking has evolved during the past two decades beyond a focus on individual-level factors underlying energy imbalances to a consideration of the biological, social, environmental and policy drivers of health behaviours and outcomes¹³⁹ and a systems approach to the ways these drivers interact140,141. Such an approach calls for coordinated actions from all stakeholders to improve policies and practices spanning multiple sectors and to shift social norms on health 142,143.

The NAFLD prevention agenda should therefore include the creation of healthier, more equitable and sustainable societies as part of its vision. Existing frameworks such as the United Nations Sustainable Development Goals (SDGs) can usefully inform and guide the development of multisectoral efforts to address NAFLD. A recently developed SDG framework for NAFLD (available as a preprint) aims to help conceptualize thinking about the design and delivery of such responses¹⁴⁴. As a first step, this framework can be used as a strategic advocacy tool to build the case for closer collaboration within and among sectors.

Leadership for the NAFLD public health agenda

- Statements 7.1-7.3
- Recommendations 25-26

To move the NAFLD public health agenda forwards, national, regional and international liver associations, in collaboration with governments and other stakeholders, will need to lead the way. Multilateral organizations such as WHO also have a key role in shaping and delivering responses to NAFLD, first by recognizing the condition as a public health issue worthy of attention, and second by supporting national public health responses.

Several existing policy levers and movements can support the development of such responses too. For instance, global efforts to expand universal health coverage and ensure that health systems are people-centred provide useful mechanisms for addressing NCDs holistically. In addition, medical societies whose members provide health care for aspects of metabolic syndrome are well positioned to help lead this change. NAFLD should also be incorporated, whenever appropriate, in the development of joint plans of action, guidelines, policy briefs and educational tools, and these efforts should be adequately resourced. Finally, a global coalition of NAFLD stakeholders, both organizations and individuals, should lead the development of a NAFLD public health roadmap and advocate for its adoption

by the global health community. This coalition should actively engage with those outside the liver health space by growing and nurturing a broad network of individuals and organizations with a common vision and goals.

Study strengths and limitations

Although the Delphi method is the right approach for a consensus-building initiative, it is not without challenges and limitations^{145,146}. We employed purposive sampling to select the members of the initial core group and then used core group member recommendations to generate a larger, more diverse expert panel. In doing so, we sought to mitigate concerns about the inherent bias in the purposive sampling of a relatively small group with more broad-based snowball sampling, which resulted in a diverse expert panel of 218 members from 91 countries/territories and all six WHO regions. The variety of backgrounds represented on the panel — including academia, civil society, government, private sector, research, clinical practice and advocacy - strengthened the validity of the consensus statements and recommendations. That said, we understand that conducting the study in English might have limited the composition of the expert panel and therefore the findings.

Delphi studies often involve a combination of in-person convenings for in-depth deliberation and survey rounds for voting. However, in light of the geographical spread of panel members and COVID-19 travel restrictions, we employed alternative modes of group discourse. The core group was convened virtually at two points in the process, while panel members were able to provide written comments on the draft recommendations and the three survey rounds. Although we received and incorporated a large volume of open-ended comments across all four data collection components, we acknowledge that this approach might not have resulted in the same outcomes as those that would have emerged from real-time discussion and resolution of complicated or contentious issues. Conversely, this method gave panel members multiple opportunities to provide open-ended comments in a space without any dominant voices whose presence sometimes inhibits the expression of minority viewpoints during in-person convenings. The combination of in-person feedback (from core group members) and written feedback (from the entire expert panel) might therefore have resulted in more comprehensive contributions overall.

The increasing levels of agreement with the consensus statements across all three survey rounds, together with the high levels of participation (88% in R1 and R2; 85% in R3; 79% in the online meeting), strengthen our confidence in the rigor of the method used and the resultant findings. Expert panel members' ability to include detailed comments on each of the draft statements enabled us to improve them, as reflected in the increasing mean level of agreement with the statements in successive rounds, from 80.3% in R1 to 98.5% in R3. Moreover, the endorsement of the final consensus statements and recommendations presented in TABLES 3 and 4 by 110 organizations in 59 countries/territories (Supplementary Table 8) at the time of publication further testifies to their global relevance.

Conclusions

NAFLD is a highly prevalent disease that poses a major challenge to global public health. In this Consensus Statement, a diverse international group of experts developed and endorsed a set of consensus statements and recommendations that provide needed guidance for the creation and implementation of health system and public health responses that will rise to this challenge. The public health approach that informed the

consensus-building process helped ensure the relevance of these statements and recommendations for a broad group of stakeholders, from researchers and health-care providers to policy-makers and funders. It is now up to the liver health community to lead the development of a roadmap to translate these statements and recommendations into global vision and action.

Published online 27 October 2021

- Golabi, P. et al. Non-alcoholic fatty liver disease (NAFLD) is associated with impairment of health related quality of life (HRQOL). Health Qual. Life Outcomes 14, 18 (2016).
- McSweeney, L. et al. Health-related quality of life and patient-reported outcome measures in NASH-related cirrhosis. JHEP Rep. 2, 100099 (2020).
- O'Hara, J. et al. Cost of non-alcoholic steatohepatitis in Europe and the USA: the GAIN study. JHEP Rep. 2, 100142 (2020).
- Schattenberg, J. M. et al. Disease burden and economic impact of diagnosed non-alcoholic steatohepatitis (NASH) in five European countries in 2018: a cost-of-illness analysis. *Liver Int.* 41, 1227–1242 (2021).
- Younossi, Z. M. et al. The association of histologic and noninvasive tests with adverse clinical and patient-reported outcomes in patients with advanced fibrosis due to nonalcoholic steatohepatitis. *Castroenterology* https://doi.org/10.1053/ j.gastro.2020.12.003 (2020).
- Younossi, Z. M. et al. The economic and clinical burden of nonalcoholic fatty liver disease in the United States and Europe. *Hepatology* 64, 1577–1586 (2016).
- Hagström, H. et al. Health care costs of patients with biopsy-confirmed nonalcoholic fatty liver disease are nearly twice those of matched controls. Clin. Castroenterol. Hepatol. 18, 1592–1599.e8 (2020)
- Gordon, S. C., Fraysse, J., Li, S., Ozbay, A. B. & Wong, R. J. Disease severity is associated with higher healthcare utilization in nonalcoholic steatohepatitis Medicare patients. Am. J. Gastroenterol. 115, 562–574 (2020).
- Bertot, L. C. & Adams, L. A. The natural course of non-alcoholic fatty liver disease. *Int. J. Mol. Sci.* 17, 774 (2016).
- Ekstedt, M., Nasr, P. & Kechagias, S. Natural history of NAFLD/NASH. Curr. Hepatol. Rep. 16, 391–397 (2017)
- Araújó, A. R., Rosso, N., Bedogni, G., Tiribelli, C. & Bellentani, S. Global epidemiology of non-alcoholic fatty liver disease/non-alcoholic steatohepatitis: what we need in the future. Liver Int. 38, 47–51 (2018).
- Kanwal, F. et al. Risk of hepatocellular cancer in patients with non-alcoholic fatty liver disease. Gastroenterology 155, 1828–1837.e2 (2018).
- Haldar, D. et al. Outcomes of liver transplantation for non-alcoholic steatohepatitis: a European Liver Transplant Registry study. J. Hepatol. 71, 313–322 (2019).
- Terrault, N. A. & Pageaux, G.-P. A changing landscape of liver transplantation: king HCV is dethroned, ALD and NAFLD take over! *J. Hepatol.* 69, 767–768 (2018).
- Global Burden of Disease Cancer Collaboration et al. Global, regional, and national cancer incidence, mortality, years of life lost, years lived with disability, and disability-adjusted life-years for 29 cancer groups, 1990 to 2017: a systematic analysis for the Global Burden of Disease Study. *JAMA Oncol.* 5, 1749–1768 (2019).
- Adams, L. A., Anstee, Q. M., Tilg, H. & Targher, G. Non-alcoholic fatty liver disease and its relationship with cardiovascular disease and other extrahepatic diseases. Gut 66, 1138–1153 (2017).
- Anstee, Q. M., Targher, G. & Day, C. P. Progression of NAFLD to diabetes mellitus, cardiovascular disease or cirrhosis. *Nat. Rev. Gastroenterol. Hepatol.* 10, 330–344 (2013).
- Byrne, C. D. & Targher, G. NAFLD: a multisystem disease. J. Hepatol. 62, S47–S64 (2015).
- Albhaisi, S., Chowdhury, A. & Sanyal, A. J. Non-alcoholic fatty liver disease in lean individuals. JHEP Rep.: Innov. Hepatol. 1, 329–341 (2019).

- Younossi, Z. et al. Global burden of NAFLD and NASH: trends, predictions, risk factors and prevention. Nat. Rev. Gastroenterol. Hepatol. 15, 11–20 (2018).
- Vilar-Gomez, E. et al. Fibrosis severity as a determinant of cause-specific mortality in patients with advanced nonalcoholic fatty liver disease: a multi-national cohort study. Gastroenterology 155, 443–457.e17 (2018).
- Angulo, P. et al. Liver fibrosis, but no other histologic features, is asssociated with long-term outcomes of patients with nonalcoholic fatty liver disease. *Gastroenterologu* 149, 389–397.e10 (2015).
- Gastroenterology **149**, 389–397.e10 (2015).

 23. Haflidadottir, S. et al. Long-term follow-up and liver-related death rate in patients with non-alcoholic and alcoholic related fatty liver disease.

 BMC Gastroenterol. **14**
- BMC Castroenterol. 14, 166 (2014).

 24. Park, S. K., Seo, M. H., Shin, H. C. & Ryoo, J. H. Clinical availability of nonalcoholic fatty liver disease as an early predictor of type 2 diabetes mellitus in Korean men: 5-year prospective cohort study. Hepatology 57, 1378–1383 (2013).
- World Health Organization. Global action plan for the prevention and control of noncommunicable diseases 2013–2020. WHO https://apps.who.int/iris/bitstream/ handle/10665/94384/9789241506236_eng.pdf (2013).
- Bennett, J. E. et al. NCD countdown 2030: worldwide trends in non-communicable disease mortality and progress towards Sustainable Development Goal target 3.4. *Lancet* 392, 1072–1088 (2018).
- Rotondi, A. & Gustafson, D. in Gazing into the Oracle: The Delphi Method and its Application to Social Policy and Public Health (eds Adler, M. & Ziglio, E.) 34–55 (Jessica Kingsley, 1996).
- Rubino, F. et al. Joint international consensus statement for ending stigma of obesity. Nat. Med. 26, 485–497 (2020).
- Lazarus, J. V. et al. Consensus statement on the role of health systems in advancing the long-term well-being of people living with HIV. Nat. Commun. 12, 4450 (2021)
- Vos, T. et al. Global burden of 369 diseases and injuries in 204 countries and territories, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet* 396, 1204–1222 (2020).
- Younossi, Z. M. et al. Global epidemiology of nonalcoholic fatty liver disease—meta-analytic assessment of prevalence, incidence, and outcomes. *Hepatology* 64, 73–84 (2016).
- Estes, C. et al. Modeling NAFLD disease burden in China, France, Germany, Italy, Japan, Spain, United Kingdom, and United States for the period 2016-2030. J. Hepatol. 69, 896–904 (2018).
- Estes, C., Razavi, H., Loomba, R., Younossi, Z. & Sanyal, A. J. Modeling the epidemic of nonalcoholic fatty liver disease demonstrates an exponential increase in burden of disease. *Hepatology* 67, 123–133 (2018).
- Younossi, Z. et al. Global perspectives on nonalcoholic fatty liver disease and nonalcoholic steatohepatitis. Hepatology 69, 2672–2682 (2019).
- Ye, Q. et al. Global prevalence, incidence, and outcomes of non-obese or lean non-alcoholic fatty liver disease: a systematic review and meta-analysis. *Lancet Gastroenterol. Hepatol.* 5, 739–752 (2020).
- Bjorkstrom, K., Stal, P., Hultcrantz, R. & Hagstrom, H. Histologic scores for fat and fibrosis associate with development of type 2 diabetes in patients with nonalcoholic fatty liver disease. Clin. Gastroenterol. Hepatol. 15, 1461–1468 (2017).
- Musso, G., Gambino, R., Cassader, M. & Pagano, G. Meta-analysis: natural history of non-alcoholic fatty liver disease (NAFLD) and diagnostic accuracy of non-invasive tests for liver disease severity. *Ann. Med.* 43, 617–649 (2011).

- Sepanlou, S. G. et al. The global, regional, and national burden of cirrhosis by cause in 195 countries and territories, 1990-2017: a systematic analysis for the Global Burden of Disease Study 2017. Lancet Gastroenterol. Hepatol. 5, 245–266 (2020).
- Anderson, E. L. et al. The prevalence of non-alcoholic fatty liver disease in children and adolescents: a systematic review and meta-analysis. *PLoS ONE* 10, e0140908 (2015).
- Jimenez-Rivera, C. et al. Prevalence and risk factors for non-alcoholic fatty liver in children and youth with obesity. BMC Pediatr. 17, 113 (2017).
- Xanthakos, S. A. et al. High prevalence of nonalcoholic fatty liver disease in adolescents undergoing bariatric surgery. *Gastroenterology* 149, 623–634.e8 (2015).
- Yu, E. L. et al. Prevalence of nonalcoholic fatty liver disease in children with obesity. *J. Pediatr.* 207, 64–70 (2019).
- 43. Schwimmer, J. B. et al. Prevalence of fatty liver in children and adolescents. *Pediatrics* **118**, 1388–1393 (2006).
- Yüksel, F. et al. Fatty liver disease in an autopsy series of children and adolescents. *Hippokratia* 16, 61–65 (2012).
- Rocha, R. et al. Nonalcoholic fatty liver disease in asymptomatic Brazilian adolescents. World J. Gastroenterol. 15, 473–477 (2009).
- Zhang, X. et al. Increasing prevalence of NAFLD/NASH among children, adolescents and young adults from 1990 to 2017: a population-based observational study. BMJ Open 11, e042843 (2021).
- Schwimmer, J. B. et al. Histopathology of pediatric nonalcoholic fatty liver disease. *Hepatology* 42, 641–649 (2005).
- Nobili, V. et al. NAFLD in children: new genes, new diagnostic modalities and new drugs. Nat. Rev. Castroenterol. Hepatol. 16, 517–530 (2019).
- Vos, M. B. et al. Factors to consider in development of drugs for pediatric nonalcoholic fatty liver disease. Gastroenterology 157, 1448–1456.e1 (2019).
- Nadeau, K. J. et al. Youth-onset type 2 diabetes consensus report: current status, challenges, and priorities. *Diabetes Care* 39, 1635–1642 (2016).
- Zimmermann, E. et al. Body mass index in s chool-aged children and the risk of routinely diagnosed non-alcoholic fatty liver disease in adulthood: a prospective study based on the Copenhagen School Health Records Register. BMJ Open 5, e006998 (2015).
- 53. Hagström, H., Stål, P., Hultcrantz, R., Hemmingsson, T. & Andreasson, A. Overweight in late adolescence predicts development of severe liver disease later in life: a 39years follow-up study. J. Hepatol. 65, 363–368 (2016).
- Hagström, H., Tynelius, P. & Rasmussen, F. High BMI in late adolescence predicts future severe liver disease and hepatocellular carcinoma: a national, population-based cohort study in 1.2 million men. Gut 67. 1536–1542 (2018).
- Simon, T. G. et al. Non-alcoholic fatty liver disease in children and young adults is associated with increased long-term mortality. J. Hepatol. https://doi.org/ 10.1016/j.jhep.2021.06.034 (2021).
- Younossi, Z. M. et al. A disease-specific quality of life instrument for non-alcoholic fatty liver disease and non-alcoholic steatohepatitis: CLDQ-NAFLD. *Liver Int.* 37, 1209–1218 (2017).
- Doward, L. C. et al. Development of a patient-reported outcome measure for non-alcoholic steatohepatitis (NASH-CHECK): results of a qualitative study. *Patient* 14, 533–543 (2021).

- Twiss, J. et al. Development of a new patient-reported outcome measure for non-alcoholic steatohepatitis: NASH-CHECK [abstract PGI39]. Value Health 20, A638 (2017).
- Huber, Y. et al. Health-related quality of life in nonalcoholic fatty liver disease associates with hepatic inflammation. Clin. Gastroenterol. Hepatol. 17, 2085–2092.e1 (2019).
- David, K. et al. Quality of life in adults with nonalcoholic fatty liver disease: baseline data from the Nonalcoholic Ateatohepatitis Clinical Research Network. *Hepatology* 49, 1904–1912 (2009).
- Kennedy-Martin, T., Bae, J. P., Paczkowski, R. & Freeman, E. Health-related quality of life burden of nonalcoholic steatohepatitis: a robust pragmatic literature review. J. Patient Rep. Outcomes 2, 28 (2017).
- Casey, J. A., Schwartz, B. S., Stewart, W. F. & Adler, N. E. Using electronic health records for population health research: a review of methods and applications. *Annu. Rev. Public Health* 37, 61–81 (2016).
- Hagström, H. et al. Administrative coding in electronic health care record-based research of NAFLD: an expert panel consensus statement. Hepatology https://doi.org/10.1002/hep.31726 (2021).
- Sanyal, A. J. Putting non-alcoholic fatty liver disease on the radar for primary care physicians: how well are we doing? *BMC Med.* 16, 148 (2018).
- Bergqvist, Č. J. et al. Awareness and opinions of non-alcoholic fatty liver disease by hospital specialists. *Intern. Med. J.* 43, 247–253 (2013).
 Alemany-Pages, M. et al. Insights from qualitative
- 66. Alemany-Pagès, M. et al. Insights from qualitative research on NAFLD awareness with a cohort of T2DM patients: time to go public with insulin resistance? BMC Public Health 20, 1142–1142 (2020).
- Wieland, A. C. et al. Low awareness of nonalcoholic fatty liver disease among patients at high metabolic risk. *J. Clin. Gastroenterol.* 49, e6–e10 (2015).
- Goh, G. B. B. et al. Perceptions of non-alcoholic fatty liver disease – an Asian community-based study. *Gastroenterol. Rep.* 4, 131–135 (2016).
- 69. Ghevariya, V. et al. Knowing what's out there: awareness of non-alcoholic fatty liver disease. *Front. Med.* **1**, 4 (2014).
- Augustin, S., Pons, M., Santos, B., Ventura, M. & Genescà, J. in Portal Hypertension VI: Proceedings of the Sixth Baveno Consensus Workshop: Stratifying Risk and Individualizing Care (ed. de Franchis, R.) 39–49 (Springer, 2016).
- de Franchis, R. Expanding consensus in portal hypertension: report of the Baveno VI Consensus Workshop: Stratifying Risk and Individualizing Care for Portal Hypertension. J. Hepatol. 63, 743–752 (2015).
- Lazarus, J. V. et al. We know DAAs work, so now what? Simplifying models of care to enhance the hepatitis C cascade. J. Intern. Med. 286, 503–525 (2019).
- Lazarus, J. V. et al. Defining comprehensive models of care for NAFLD. Nat. Rev. Gastroenterol. Hepatol. https://doi.org/10.1038/s41575-021-00477-7 (2021).
- Grgurevic, I. et al. Natural history of nonalcoholic fatty liver disease: implications for clinical practice and an individualized approach. *Can. J. Gastroenterol. Hepatol.* 2020, 9181368 (2020).
- Dyson, J. K., Anstee, Q. M. & McPherson, S. Non-alcoholic fatty liver disease: a practical approach to treatment. Frontline Gastroenterol. 5, 277–286 (2014).
- Ruissen, M. M., Mak, A. L., Beuers, U., Tushuizen, M. E. & Holleboom, A. G. Non-alcoholic fatty liver disease: a multidisciplinary approach towards a cardiometabolic liver disease. Eur. J. Endocrinol. 183, R57–R73
- Ähmed, M. H., Woodward, C. & Mital, D. Metabolic clinic for individuals with HIV/AIDS: a commitment and vision to the future of HIV services. *Cardiovasc. Endocrinol.* 6, 109–112 (2017).
- Armstrong, M. J. et al. Severe asymptomatic non-alcoholic fatty liver disease in routine diabetes care; a multi-disciplinary team approach to diagnosis and management. *OJM* 107, 33–41 (2014)
- DeVore, S. et al. A multidisciplinary clinical program is effective in stabilizing BMI and reducing transaminase levels in pediatric patients with NAFLD. J. Pediatr. Gastroenterol. Nutr. 57, 119–123 (2013).
- Moolla, A. et al. A multidisciplinary approach to the management of NAFLD is associated with improvement in markers of liver and cardio-metabolic

- health. *Frontline Gastroenterol.* **10**, 337–346 (2019).
- Neilson, L. J. et al. Implementation of a care bundle improves the management of patients with non-alcoholic fatty liver disease. Frontline Gastroenterol. https://doi.org/10.1136/ flgastro-2020-101480 (2021).
- Dyson, J. K., Anstee, Q. M. & McPherson, S. Non-alcoholic fatty liver disease: a practical approach to diagnosis and staging. Front. Gastroenterol. 5, 211–218 (2014).
- 83. Chalmers, J. et al. The development and implementation of a commissioned pathway for the identification and stratification of liver disease in the community. Front. Gastroenterol. 11, 86–92 (2020).
- Dillon, J. F. et al. Intelligent liver function testing (iLFT): a trial of automated diagnosis and staging of liver disease in primary care. *J. Hepatol.* 71, 699–706 (2019).
- Hudson, M. & McPherson, S, on behalf of the North East & North Cumbria Hepatology Network. Liver Network: Guidelines for the Management of Adults with Asymptomatic Liver Function Abnormalities (Northern England Clinical Strategic Networks, 2020).
- Shaheen, A. A. et al. Risk stratification of patients with nonalcoholic fatty liver disease using a case identification pathway in primary care: a cross-sectional study. CMAJ Open 8, E370–E376 (2020)
- Srivastava, A. et al. Prospective evaluation of a primary care referral pathway for patients with non-alcoholic fatty liver disease. *J. Hepatol.* 71, 371–378 (2019).
- Crossan, C. et al. Referral pathways for patients with NAFLD based on non-invasive fibrosis tests: diagnostic accuracy and cost analysis. *Liver Int.* 39, 2052–2060 (2019).
- Srivastava, A. et al. Cost-comparison analysis of FIB-4, ELF and fibroscan in community pathways for non-alcoholic fatty liver disease. *BMC Gastroenterol*.
 19. 122 (2019).
- Tanajewski, L. et al. Economic evaluation of a community-based diagnostic pathway to stratify adults for non-alcoholic fatty liver disease: a Markov model informed by a feasibility study. BMJ Open 7, e015659 (2017).
- Alexander, M. et al. Real-world data reveal a diagnostic gap in non-alcoholic fatty liver disease. BMC Med. 16, 130 (2018).
- Cooke, G. S. et al. Accelerating the elimination of viral hepatitis: a Lancet Gastroenterology & Hepatology Commission. *Lancet Gastroenterol. Hepatol.* 4, 135–184 (2019).
- Mencin, A. A., Loomba, R. & Lavine, J. E. Caring for children with NAFLD and navigating their care into adulthood. *Nat. Rev. Gastroenterol. Hepatol.* 12, 617–628 (2015).
 Taylor, R. S. et al. Association between fibrosis
- Taylor, R. S. et al. Association between fibrosis stage and outcomes of patients with nonalcoholic fatty liver disease: a systematic review and meta-analysis. *Gastroenterology* 158, 1611–1625. e12 (2020).
- Castera, L., Friedrich-Rust, M. & Loomba, R. Noninvasive assessment of liver disease in patients with nonalcoholic fatty liver disease. *Castroenterology* 156, 1264–1281.e4 (2019).
- Vali, Y. et al. Enhanced liver fibrosis test for the non-invasive diagnosis of fibrosis in patients with NAFLD: a systematic review and meta-analysis. J. Hepatol. 73, 252–262 (2020).
- McPherson, S. et al. Age as a confounding factor for the accurate non-invasive diagnosis of advanced NAFLD fibrosis. Am. J. Gastroenterol. 112, 740–751 (2017).
- Anstee, Q. M. et al. Noninvasive tests accurately identify advanced fibrosis due to NASH: baseline data from the STELLAR trials. *Hepatology* 70, 1521–1530 (2019).
- Augustin, S. et al. Identification of patients with advanced fibrosis due to nonalcoholic fatty liver disease: considerations for best practice. J. Gastrointestin Liver Dis. 29, 235–245 (2020).
- Castera, L. Diagnosis of non-alcoholic fatty liver disease/non-alcoholic steatohepatitis: non-invasive tests are enough. *Liver Int.* 38(Suppl 1), 67–70 (2018)
- Majumdar, A., Campos, S., Gurusamy, K., Pinzani, M. & Tsochatzis, E. A. Defining the minimum acceptable diagnostic accuracy of noninvasive fibrosis testing in cirrhosis: a decision analytic modeling study. Hepatology 71, 627–642 (2020).

- 102. Jung, J. et al. MRE combined with FIB-4 (MEFIB) index in detection of candidates for pharmacological treatment of NASH-related fibrosis. *Gut* 70, 1946–1953 (2021).
- 103. Mosca, A., Panera, N., Crudele, A. & Alisi, A. Noninvasive diagnostic tools for pediatric NAFLD: where are we now? Expert Rev. Gastroenterol. Hepatol. 14, 1035–1046 (2020).
- Di Cesare, M. et al. Inequalities in non-communicable diseases and effective responses. *Lancet* 381, 585–597 (2013).
- 105. Peters, R. et al. Common risk factors for major noncommunicable disease, a systematic overview of reviews and commentary: the implied potential for targeted risk reduction. *Ther. Adv. Chronic Dis.* 10, 2040622319880392–2040622319880392 (2019).
- 106. Fuchs, S., Henschke, C., Blümel, M. & Busse, R. Disease management programs for type 2 diabetes in Germany: a systematic literature review evaluating effectiveness. *Dtsch. Arztebl Int.* 111, 453–463 (2014).
- 107. World Health Organization. WHO package of essential noncommunicable (PEN) disease interventions for primary health care. WHO https://www.who.int/ publications/i/item/who-package-of-essentialnoncommunicable-(pen)-disease-interventionsfor-primary-health-care (2020).
- 108. Dixon, B. E., Embi, P. J. & Haggstrom, D. A. Information technologies that facilitate care coordination: provider and patient perspectives. *Transl. Behav. Med.* 8, 522–525 (2018).
- 109. Ando, Y. & Jou, J. H. Nonalcoholic fatty liver disease and recent guideline updates. *Clin. Liver Dis.* 17, 23–28 (2021).
- Leoni, S. et al. Current guidelines for the management of non-alcoholic fatty liver disease: a systematic review with comparative analysis. World J. Gastroenterol. 24, 3361–3373 (2018).
- European Association for the Study of the Liver (EASL); European Association for the Study of Diabetes (EASD); European Association for the Study of Obesity (EASO) EASL-EASD-EASO Clinical Practice Guidelines for the management of non-alcoholic fatty liver disease. J. Hepatol. 64, 1388–1402 (2016).
- 112. Arab, J. P. et al. Latin American Association for the Study of the Liver (ALEH) practice guidance for the diagnosis and treatment of non-alcoholic fatty liver disease. Ann. Hepatol. 19, 674–690 (2020)
- Chalasani, N. et al. The diagnosis and management of nonalcoholic fatty liver disease: practice guidance from the American Association for the Study of Liver Diseases. *Hepatology* 67, 328–357 (2018).
- 114. Chitturi, S. et al. The Asia—Pacific Working Party on Non-alcoholic Fatty Liver Disease guidelines 2017—Part 2: Management and special groups. J. Gastroenterol. Hepatol. 33, 86—98 (2018).
- 115. Eslam, M. et al. The Asian Pacific Association for the Study of the Liver clinical practice guidelines for the diagnosis and management of metabolic associated fatty liver disease. *Hepatol. Int.* 14, 889–919 (2020).
- 116. Muthiah, M. D. & Sanyal, A. J. Current management of non-alcoholic steatohepatitis. *Liver Int.* 40 (Suppl. 1), 89–95 (2020).
- Hallsworth, K. & Adams, L. A. Lifestyle modification in NAFLD/NASH: facts and figures. JHEP Rep. Innov. Hepatol. 1, 468–479 (2019).
- Romero-Gómez, M., Zelber-Sagi, S. & Trenell, M. Treatment of NAFLD with diet, physical activity and exercise. J. Hepatol. 67, 829–846 (2017).
 Jensen, M. D. et al. 2013 AHA/ACC/TOS guideline for
- 119. Jensen, M. D. et al. 2013 AHA/ACC/TOS guideline for the management of overweight and obesity in adults: a report of the American College of Cardiology/ American Heart Association Task Force on Practice Guidelines and The Obesity Society. Circulation 129, 5102-5138 (2014)
- S102–S138 (2014).

 120. Hydes, T. J., Ravi, S., Loomba, R. & Gray, M. E. Evidence-based clinical advice for nutrition and dietary weight loss strategies for the management of NAFLD and NASH. Clin. Mol. Hepatol. 26, 383–400 (2020).

 121. Geurtsen, M. L., Santos, S., Gaillard, R., Felix, J. F.
- 121. Geurtsen, M. L., Santos, S., Gaillard, R., Felix, J. F. & Jaddoe, V. W. V. Associations between intake of sugar-containing beverages in infancy with liver fat accumulation at school age. *Hepatology* 73, 560–570 (2021).
- Berná, G. & Romero-Gomez, M. The role of nutrition in non-alcoholic fatty liver disease: pathophysiology and management. *Liver Int.* 40 (Suppl. 1), 102–108 (2020).

- Hashida, R. et al. Aerobic vs resistance exercise in non-alcoholic fatty liver disease: a systematic review. *J. Hepatol.* 66, 142–152 (2017).
- 124. Lazarus, J. V. et al. A cross-sectional study of the public health response to non-alcoholic fatty liver disease in Europe. J. Hepatol. 72, 14–24 (2020)
- 125. Neuschwander-Tetri, B. A. Therapeutic landscape for NAFLD in 2020. Gastroenterology 158, 1984–1998. e3 (2020).
- 126. Wong, E. L. Y. et al. Patient experience and satisfaction with inpatient service: development of short form survey instrument measuring the core aspect of inpatient experience. PLoS ONE 10, e0122299 (2015)
- 127. Martin, L. R., Williams, S. L., Haskard, K. B. & Dimatteo, M. R. The challenge of patient adherence. *Ther. Clin. Risk Manag.* 1, 189–199 (2005).
- 128. World Health Assembly. Hepatitis. WHO https://apps who.int/gb/ebwha/pdf_files/WHA67/A67_R6-en. pdf?ua=1 (2014).
- 129. Puhl, R. M. & Heuer, C. A. Obesity stigma: important considerations for public health. *Am. J. Public Health* 100, 1019–1028 (2010).
- Room, R. Stigma, social inequality and alcohol and drug use. *Drug Alcohol. Rev.* 24, 143–155 (2005).
- Vaughn-Sandler, V., Sherman, C., Aronsohn, A. & Volk, M. L. Consequences of perceived stigma among patients with cirrhosis. *Dig. Dis. Sci.* 59, 681–686 (2014).
- Himmelstein, M. S. & Puhl, R. M. At multiple fronts: diabetes stigma and weight stigma in adults with type 2 diabetes. *Diabet. Med.* 38, e14387 (2021).
- 133. Lazarus, J. V. et al. European 'NAFLD Preparedness Index'—is Europe ready to meet the challenge of fatty liver disease? *JHEP Rep.* **3**, 100234 (2021).
- 134. Lazarus, J. V. et al. The global non-alcoholic fatty liver disease (NAFLD) preparedness index: are countries ready to tackle the challenge? *Int. Liver Congr.* 75 (Suppl. 2), 23–26 (2021).
- 135. NASH Council & Global Liver Institute. U.S NASH action plan (Global Liver Institute, 2020).
- 136. World Health Organization. Continuity and coordination of care: a practice brief to support implementation of the WHO Framework on integrated people-centred health services (Alliance for Health Policy and Systems Research, 2018).
- Atun, R. et al. Improving responsiveness of health systems to non-communicable diseases. *Lancet* 381, 690–697 (2013).
- 138. Rutter, H. et al. The need for a complex systems model of evidence for public health. *Lancet* **390**, 2602–2604 (2017).
- 139. Kumanyika, S., Jeffery, R. W., Morabia, A., Ritenbaugh, C. & Antipatis, V. J. Obesity prevention: the case for action. *Int. J. Obes. Relat. Metab. Disord.* 26, 425–436 (2002).
- 140. Gortmaker, S. L. et al. Changing the future of obesity: science, policy, and action. *Lancet* 378, 838–847 (2011).
- Huang, T. T., Drewnosksi, A., Kumanyika, S. & Glass, T. A. A systems-oriented multilevel framework for addressing obesity in the 21st century. *Prev. Chronic Dis.* 6, A82 (2009).
- 142. Huang, T. & Ferris, E. in *Childhood Obesity* Ch. 42 (ed. Goran, M. I.) (Taylor and Francis, 2016).
- 143. Huang, T. T. et al. Mobilisation of public support for policy actions to prevent obesity. *Lancet* 385, 2422–2431 (2015).
- 144. Lazarus, J. V. et al. A Sustainable Development Goal framework to guide multisectoral action on NAFLD through a societal approach. Preprint at medRxiv https://doi.org/10.1101/2021.08.02.21261464 (2021).
- 145. Adler, M. & Ziglio, E. Gazing into the Oracle: The Delphi Method and its Application to Social Policy and Public Health (Jessica Kingsley, 1996).
- 146. Grime, M. M. & Wright, G. in *Wiley StatsRef: Statistics Reference Online* 1–6 (Wiley, 2011).
- 147. Targher, G. & Byrne, C. D. From nonalcoholic fatty liver disease to metabolic dysfunction-associated fatty liver disease: is it time for a change of terminology? *Hepatoma Res.* 6, 64 (2020).
- 148. Eslam, M. et al. A new definition for metabolic dysfunction-associated fatty liver disease: an international expert consensus statement. *J. Hepatol.* 73, 202–209 (2020).
- 149. Eslam, M., Sanyal, A. J. & George, J. MAFLD: a consensus-driven proposed nomenclature

- for metabolic associated fatty liver disease. Gastroenterology 158, 1999–2014.e1 (2020).
- 150. Mendez-Sanchez, N. et al. The Latin American Association for the Study of the Liver (ALEH) position statement on the redefinition of fatty liver disease. *Lancet Gastroenterol. Hepatol.* 6, 65–72 (2021).
- Spearman, C. W. et al. The sub-Saharan Africa position statement on the redefinition of fatty liver disease: from NAFLD to MAFLD. J. Hepatol. 74, 1256–1258 (2021).
- 152. Shiha, G. et al. Nomenclature and definition of metabolic-associated fatty liver disease: a consensus from the Middle East and north Africa. *Lancet Gastroenterol. Hepatol.* 6, 57–64 (2021).
- 153. Ratziu, V. et al. The times they are a-changin' (for NAFLD as well). *J. Hepatol.* **73**, 1307–1309 (2020).
- 154. Younossi, Z. M. et al. From NAFLD to MAFLD: implications of a premature change in terminology. *Hepatology* 73, 1194–1198 (2021).
- 155. Wong, V. W. et al. Asia-Pacific Working Party on Non-alcoholic Fatty Liver Disease guidelines 2017–Part 1: definition, risk factors and assessment. J. Gastroenterol. Hepatol. 33, 70–85 (2018).
- 156. American Diabetes Association. Comprehensive medical evaluation and assessment of comorbidities: standards of medical care in diabetes — 2020. *Diabetes Care* 43, \$37–\$47 (2020).
- 157. World Health Organization. Nothing for us, without us: opportunities for meaningful engagement of people living with NCDs: meeting report (WHO, 2021).

Acknowledgements

The European Association for the Study of the Liver (EASL) International Liver Foundation (EILF) acknowledges funding from Intercept Pharmaceutics, as well as Bristol-Myers Squibb and Merck Sharp & Dohme. The authors thank in particular T. White (ISGlobal) and A. Hobbs (CUNY) for providing technical support during implementation of the Delphi methodology, and to N. Lee and the Wilton Park team for facilitating the online convening of the core group on 16 February 2021. Three observers attended the session: M. Sicuro (EILF), Y. Nedelcheva (EASL) and K. Ray (Nature Reviews Gastroenterology & Hepatology). The translations in Supplementary Tables 2–7 were provided by Springer Healthcare, and checked and revised by the authors, and are reproduced in Supplementary information as supplied.

Author contributions

J.V.L. and H.E.M. conceptualized the idea for the study and developed the first draft of the consensus statements. The voting members of the core authorship group (Q.M.A., J.P.A., R.L.B., L.C., H.C.-P., J.C., K.C., M.A.D., S.F., J.G., H.H., T.T.-K.H., M.H.I., A.K., S.K.S., R.L., V.M., P.N.N., M.N., P.O., V.R., M.R., M.R.-G., J.M.S., E.A.T., L.V., V.W.-S.W., Y.Y., Z.M.Y. and S.Z.-S.) provided input on the draft statements ahead of Delphi round 1. D.R. supported the design of the study methodology, managed the Qualtrics XM platform and extracted data following each Delphi round. All members of the NAFLD Consensus Consortium participated in the Delphi study. J.V.L. and H.E.M. reviewed the data between Delphi rounds and revised the consensus statements based on the qualitative feedback. Core group members participated in the World Café discussion (J.V.L., H.E.M., Q.M.A., J.P.A., R.L.B., L.C., H.C.-P., J.C., S.F., H.H., T.T.-K.H., M.H.I., A.K., S.K.S., V.M., P.N.N., M.N., P.O., M.R.-G., J.M.S., E.A.T., L.V., V.W.-S.W., Y.Y., Z.M.Y. and S.Z.-S.) prior to the third and final Delphi round. J.V.L. and H.E.M. drafted the recommendations, which voting members of the authorship group reviewed and provided feedback on. All members of the NAFLD Consensus Consortium participated in at least one Delphi round. H.E.M. and J.V.L. developed the first draft of the manuscript; all authors contributed to and reviewed the full draft of the article, subsequent revisions, and approved the final version for submission.

Competing interests

A.K. declares grants/contracts from Intercept and Novartis and conference support from Intercept. E.A.T. has received personal fees from Intercept, Gilead, Pfizer and Promethera, and honoraria from Intercept, Gilead and Pfizer. E.A.T. is a member of the Governing Board of the European Association for the Study of the Liver. H.C.-P. has received honoraria from Intercept and Novo Nordisk. H.H. has received grants or contracts from Intercept, AstraZeneca, MSD, Pfizer, EchoSens, Stockholm City Council, Swedish Cancer Foundation, Radiumhemmets Forskningsfonder, Julins Fond, Stockholm Inovation Foundation, Skandia Research Foundation, Åke Wiberg Foundation, Bengt Ihre Foundation and Tore Nilson Foundation, participated on an advisory board for BMS and Gilead, and has stock or stock options in Novo Nordisk. H.E.M. has received grants to EASL International Liver

Foundation to support NAFLD activities from Intercept. Genfit, BMS and MSD. J.C. reports consultant and/or speaker and/or participated in clinical trials sponsored by and/or received grants and research support from Gilead Sciences, AbbVie, MSD, Shionogi, Intercept Pharmaceuticals, Janssen Pharmaceuticals, Celgene and Alexion (all outside the submitted work), J.M.S. declares a research grant from Gilead; consulting fees from Boehringer Ingelheim, BMS, Genfit and Gilead, Intercept Pharmaceuticals, Madrigal, Novartis, Novo Nordisk, Nordic Bioscience, Pfizer, Roche, Sanofi, Siemens Healthcare GmbH; speakers bureau from Falk Foundation, and MSD Sharp δ Dohme. J.V.L. reports grants, personal fees and other support from AbbVie, personal fees from CEPHEID, personal fees from Genfit, grants, personal fees and other support from Gilead Sciences, personal fees from GSK, personal fees from Intercept, personal fees from Janssen, and grants and personal fees from MSD, outside the submitted work, L.C. declares: grants from Gilead: speaker's bureau for Abbvie, Echosens, Gilead, Intercept, Novo Nordisk; has been a member of an advisory board for Allergan, Alexion, Echosens, Gilead, Intercept, MSD, Novo Nordisk, Pfizer and Servier. L.V. declares grants or contracts from Gilead; honoraria from MSD, Gilead, AlfaSigma and Abbvie; support for attending conferences from Gilead; participation on advisory or safety boards for Intercept, Pfizer and Gilead. M.N. has received unrestricted research grants from Gilead, Abbie and Merck, and was President of the World Hepatitis Alliance 2018-2020, M.R. has received honoraria from Alnylam. Amgen, AMRA, BMS, Boehringer Ingelheim, Centara, Coheres, Enanta, Galecto, Intercept, Madrigal, NGM Biopharmaceuticals, Novo Nordisk, Pfizer, Fractyl, Gelesis, Siemens, Thetis, Terns, Rivus, 3vBio (Sagimet), 89bio, Novartis, Immuron, Merck and Taiwan J Pharmaceuticals. M.R.-G. reports grants from Intercept, grants from Gilead Sciences, personal fees from Shionogi, personal fees from Alfa Wassermann, personal fees from ProSciento, personal fees from Kaleido, personal fees from Novo Nordisk, personal fees from BMS, personal fees from BMS, personal fees from Allergan, personal fees from Boehringer Ingelheim, personal fees from Zydus, personal fees from Intercept Pharma and personal fees from Gilead Science, outside the submitted work. Q.M.A. declares grant funding from multiple EFPIA partners via the EU IMI2-funded consortium Litmus (Abbvie, Antaros Medical, Allergan/Tobira, AstraZeneca, BMS, Boehringer Ingelheim International, Echosens, Ellegaard Gottingen Minipigs, Eli Lilly & Company, Exalenz Bioscience Ltd., Genfit, Glympse Bio, GlaxoSmithKline, HistoIndex, Intercept Pharma Europe Ltd., iXscient Ltd., Nordic Bioscience, Novartis Pharma AG, Novo Nordisk A/S, One Way Liver Genomics SL, Perspectum Diagnostics, Pfizer Ltd., Resoundant, Sanofi-Aventis Deutschland GmbH, SomaLogic Inc., Takeda Pharmaceuticals International SA.); grants from Glympse Bio; royalties/licences from Elsevier; consulting fees from Allergan, AstraZeneca, Blade Therapeutics, BMS, Cirius Therapeutics, CymaBay, Eli Lilly, Galmed, Genfit, Gilead, HistoIndex, Intercept, Inventiva, Madrigel, Metacrine, NGMBio, Novartis, Novo Nordisk, Pfizer, Poxel Pharma and The Medicines Company; honoraria from MedScape and Fishawack: and participation on safety or advisory boards for North Sea Therapeutics and Medpace. R.L.B. declares grants or contracts from Novo Nordisk and GLWL Research; consulting fees (personal) from Novo Nordisk, Viiv, Boehringer Ingelheim and International Medical Press; has participated on safety or advisory boards for Novo Nordisk and Pfizer; has been an unpaid trustee of the Association for the Study of Obesity, BOMMS Council Member and Trustee of Obesity Empowerment Network UK. R.L. declares grants or contracts from Allergan, AstraZeneca, Boehringer-Ingelheim, Bristol-Myers Squibb, Eli Lilly, Galectin Therapeutics, Galmed Pharmaceuticals, Genfit, Gilead, Intercept, Inventiva, Janssen, Madrigal Pharmaceuticals, Merck, NGM Biopharmaceuticals, Pfizer and Sonic Incytes; consulting fees from Aardvark Therapeutics, Altimmune, Anylam/Regeneron, Amgen, Arrowhead Pharmaceuticals, AstraZeneca, Bristol-Myers Squibb, CohBar, Eli Lilly, Galmed, Gilead, Glympse bio, Hightide, Inipharm, Intercept, Inventiva, Ionis, Janssen Inc., Madrigal, Metacrine, Inc., NGM Biopharmaceuticals, Novartis, Novo Nordisk, Merck, Pfizer, Sagimet, The rate chnologies, 89 bio and Viking The rapeutics; and is co-founder of Liponexus. S.F. declares grants or contracts from Gilead, Roche, Bristol-Myers Squibb and Genfit; consulting fees from Roche, Gilead, Allergan, Abbvie, Bayer, Bristol-Myers Squibb, Boehringer Ingelheim, Merck Sharp & Dohme, Janssen, Actelion, Astellas, Genfit, Inventiva, Intercept, Echosense, Genentech, Novo Nordisk, Novartis, AstraZeneca, Galmed, Promethera, Coherus, Madrigal, Julius Clinical, NGM Bio; and honoraria from Gilead, Genfit, Bayer, Abbvie, Intercept, Allergan. V.M. declares unrestricted contributions to the Liver Forum. V.R. declares grants or contracts from Gilead: consulting fees from Novo Nordisk, Galmed. Madrigal, AstraZeneca, Intercept, Terns, Theratechnologies, NGM, Bristol-Myers Squibb; and support for attending

meetings from Gilead. V.W.-S.W. declares grants or contracts from Gilead (to institution); consulting fees and participation on boards (personal) from 3V-BIO, AbbVie, Allergan, Boehringer Ingelheim, Center for Outcomes Research in Liver Diseases, Echosens, Gilead Sciences, Hanni Pharmaceutical, Intercept, Inventiva, Merck, Novartis, Novo Nordisk, Perspectum Diagnostics, Pfizer, ProSciento, Sagimet Biosciences, TARGET PharmaSolutions and Terns; honoraria (personal) from Abbott, AbbVie, Bristol-Myers Squibb, Echosens, Gilead Sciences; and support for attending meetings or travel (institution) from Abbvie and Gilead. Y.Y. declares research grants from Biocordex and Gilead; honoraria from Gilead, Billim, Pharmactive, Sanovel and Echosens; and participated on advisory boards for Novo Nordisk and Abbvie, J.P.A., K.C., M.A.D., J.G., T.T.-K.H,

M.H.I., P.N.N., P.O., D.R., S.K.S., S.Z.-S. and Z.M.Y. declare no competing interests.

Peer review information

Nature Reviews Gastroenterology & Hepatology thanks Anna Alisi, Rotonya Carr and the other, anonymous, reviewer(s) for their contribution to the peer review of this work.

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Supplementary information

The online version contains supplementary material available at https://doi.org/10.1038/s41575-021-00523-4.

RELATED LINKS

AASLD: https://www.aasld.org/ ALEH: https://alehlatam.org/

American Diabetes Association: https://www.diabetes.org/

APASL: http://apasl.info/ EASD: https://www.easd.org/ EASL: https://easl.eu/ EASO: https://easo.org/ EILF: https://easl-ilf.org/

NCD Alliance: https://ncdalliance.org WHO: https://www.who.int/ Wilton Park: https://www.wiltonpark.org.uk/

© Springer Nature Limited 2021, corrected publication 2021

the NAFLD Consensus Consortium

Jeffrey V. Lazarus^{1,2}, Henry E. Mark², Quentin M. Anstee^{3,4}, Juan Pablo Arab⁵, Rachel L. Batterham⁶, Laurent Castera⁷, Helena Cortez-Pinto⁸, Javier Crespo⁹, Kenneth Cusi¹⁰, M. Ashworth Dirac11, Sven Francque12.13, Jacob George14, Hannes Hagström15, Terry T.-K. Huang16, Mona H. Ismail17, Achim Kautz18, Shiv Kumar Sarin19, Rohit Loomba20.21, Veronica Miller²², Philip N. Newsome²³, Michael Ninburg²⁴, Ponsiano Ocama²⁵, Vlad Ratziu²⁶, Mary Rinella²⁷, Diana Romero²⁸, Manuel Romero-Gómez²⁹, Jörn M. Schattenberg³⁰, Emmanuel A. Tsochatzis^{31,32}, Luca Valenti^{33,34}, Vincent Wai-Sun Wong³⁵, Yusuf Yilmaz^{36,37}, Zobair M. Younossi³⁸, Shira Zelber-Sagi^{39,40}, Fredrik Åberg^{41,42}, Leon Adams⁴³ Maryam Salem Al Khatry⁴⁴, Khalid Al Naamani⁴⁵, Omar Alfaro Murillo⁴⁶, Alina M. Allen⁴⁷, Faisal Alnaser^{48,49}, Saleh A. Alqahtani^{50,51}, Khalid Alswal⁵², Domenico Alvaro⁵³, Raúl J. Andrade^{54,55}, Marco Arrese⁵, Yaw Asante Awuku⁵⁶, Motala Ayesha⁵⁷, Oidov Baatarkhuu⁵⁸, Shokhista Bakieva⁵⁹, Rita Basu⁶⁰, Ramon Bataller⁶¹, Shahinaz Bedri⁶², Emanuele Bosi⁶³, Marc Bourliere⁶⁴, Radan Bruha⁶⁵, Elisabetta Bugianesi⁶⁶, Patrizia Burra⁶⁷, Maria Buti⁶⁹, Christopher D. Byrne⁶⁹, Jose Luis Calleja⁷⁰, Patrizia Carrieri⁷¹, Flloyd Carter⁷², Marlen Ivon Castellanos Fernandez⁷³, Gabriela Castillo-Lopez⁷⁴, Graciela E. Castro-Narro^{75,76}, Henry Lik Yuen Chan⁷⁷, Wah-Kheong Chan⁷⁸, Yoosoo Chang⁷⁹, Massimo Colombo⁸⁰, Kirsten J. Coppell⁸¹, Kathleen Corey⁸², Antonio Craxi⁸³, Donna Cryer⁸⁴, Anuradha Dassanayake⁸⁵, Antonieta de Ascenção Soares Martins⁸⁶, Victor de $Ledinghen^{87}, Stefano\ Del Prato^{88},\ Alessandro\ Demaio^{89},\ Hailemichael\ Desalegn^{90},\ John\ Dillon^{91},\ Ajay\ Duseja^{92},\ Prabhakaran\ Dorairaj^{93},\ Mattias\ Ekstedt^{94},\ Mohamed\ El\ Kassas^{95},\ Alessandro\ Demaio^{89},\ Alessandro\ Demaio^{89},\ Hailemichael\ Desalegn^{90},\ John\ Dillon^{91},\ Ajay\ Duseja^{92},\ Prabhakaran\ Dorairaj^{93},\ Mattias\ Ekstedt^{94},\ Mohamed\ El\ Kassas^{95},\ Alessandro\ Demaio^{89},\ Hailemichael\ Desalegn^{90},\ John\ Dillon^{91},\ Ajay\ Duseja^{92},\ Prabhakaran\ Dorairaj^{93},\ Mattias\ Ekstedt^{94},\ Mohamed\ El\ Kassas^{95},\ Prabhakaran\ Dorairaj^{93},\ Mattias\ Ekstedt^{94},\ Mohamed\ El\ Kassas^{95},\ Prabhakaran\ Dorairaj^{93},\ Prabhakaran\ Dorairaj^{93},\ Mohamed\ El\ Kassas^{95},\ Prabhakaran\ Dorairaj^{93},\ Mohamed\ El\ Kassas^{95},\ Prabhakaran\ Dorairaj^{93},\ Mohamed\ El\ Kassas^{95},\ Prabhakaran\ Dorairaj^{95},\ Prabhakaran\ D$ Osama M. Elsanousi⁹⁶, Gamal Esmat⁹⁷, Jian-Gao Fan⁹⁸, Nathalie Farpour-Lambert⁹⁹, Robert Flisiak¹⁰⁰, Yasser Fouad¹⁰¹, Michael Fuchs^{102,103}, Rino A. Gani¹⁰⁴, Lynn Gerber¹⁰⁵ Hasmik Ghazinyan¹⁰⁶, Liana Gheorghe¹⁰⁷, George Boon-Bee Goh¹⁰⁸, Henning Grønbæk¹⁰⁹, Aghayeva Gulnara¹¹⁰, Saeed Hamid¹¹¹, Vanessa Hebditch¹¹², Ingrid J. Hickman¹¹³, Samantha L. Hocking¹¹⁴, Bela Hunyady¹¹⁵, Ramazan Idilman¹¹⁶, Vasily A. Isakov¹¹⁷, Mohammad H. Jamal¹¹⁸, Peter Jepsen¹⁰⁹, Natacha Jreige Iskandar¹¹⁹, Myeong Jun Song¹²⁰, K. C. Sudhamshu¹²¹, Satoru Kakizaki¹²², George Kalamitsis¹²³, Fasiha Kanwall²⁴, Jia-Horng Kao¹²⁵, Lee Kaplan⁸², Takumi Kawaguchi¹²⁶, Yousef Khader¹²⁷, Seung Up Kim¹²⁸ Nicolas Kodjoh¹²⁹, Ger Koek¹³⁰, Kazuhiko Koike¹³¹, Narcisse Patrice Komas¹³², Marko Korenjak¹³³, Marcelo Kugelmas¹³⁴, Asma Labidi¹³⁵, Naomi F. Lange¹³⁶, Joel E. Lavine¹³⁷, Mariana Lazo¹³⁸, Nancy Lee¹³⁹, Cosmas Rinaldi A. Lesmana¹⁰⁴, Chun-Jen Liu¹²⁵, Michelle T. Long¹⁴⁰, Patricio Lopez-Jaramillo¹⁴¹, Reza Malekzadeh¹⁴², Mamun Al Mahtab¹⁴³, Giulio Marchesini¹⁴⁴, Rui Marinho¹⁴⁸, Sophia E. Martínez Vázquez⁷⁶, Lyudmila Mateva¹⁴⁶, Charles Mbendi Nlombi¹⁴⁷, Pascal Melin¹⁴⁸, Ivana Mikolasevic¹⁴⁹, Tamara Milovanovic¹⁵⁰, Carla Musso¹⁵¹, Atsushi Nakajima¹⁵², Edna Nava¹⁵³, Alexander V. Nersesov¹⁵⁴, Dafina Nikolova¹⁵⁵, Suzanne Norris¹⁵⁶, Katja Novak¹⁵⁷, Jude Oben¹⁵⁸, Janus P. Ong¹⁵⁸ Charles Onyekwere¹⁶⁰, George Papatheodoridis¹⁶¹, Imran Paruk¹⁶², Keyur Patel¹⁶³, M. Paula Macedo¹⁶⁴, Carlos Penha-Gonçalves¹⁶⁵, Marlene Pérez Figueroa¹⁶⁶, Wolf Peter Hofmann¹⁶⁷, Salvatore Petta⁸³, Claudia Pinto Marques Souza de Oliveira¹⁶⁸, Puneet Puri¹⁰², Calvin Q. Pan¹⁶⁹, Marek Rac¹⁷⁰, Johanna Ralston¹⁷¹, Alnoor Ramji¹⁷², Homie Razavi¹⁷³, Mario Reis Alvares-da-Silva¹⁷⁴, Stuart Roberts¹⁷⁵, Michael Roden^{176,177,178}, Tamsin Rose¹⁷⁹, Samir Rouabhia¹⁸⁰, Patrizia Rovere-Querini¹⁸¹, Ian A. Rowe¹⁸², Shakhlo Sadirova⁵⁹ Riina Salupere¹⁸³, Tobokalova Saparbu¹⁸⁴, Raymond Sayegh¹⁸⁵, Giada Sebastiani¹⁸⁶, Yosuke Seki¹⁸⁷, Josefina Selmo¹⁸⁸, Abdel Karim Serme¹⁸⁹, Jonathan E. Shaw¹⁹⁰, Thrivikrama Shenoy¹⁹¹, Nick Sheron¹⁹², Oren Shibolet¹⁹³, Marcelo Silva¹⁹⁴, Igor Skrypnyk¹⁹⁵, Piotr Socha¹⁹⁶, Joan Soriano¹⁹⁷, C. Wendy Spearman¹⁹⁸, Kannan Sridharan¹⁹⁹, Juan José Suárez²⁰⁰, Dhastagir Sultan Sheriff²⁰¹, Ki-Chul Sung²⁰², Mark Swain²⁰³, Frank Tacke²⁰⁴, Shahrad Taheri²⁰⁵, Soek-Siam Tan²⁰⁶, Elliot B. Tapper²⁰⁷, Hannele Yki-Järvinen^{42,208,209}, Maja Thiele^{210,211}, Isaac Thom Shawa²¹², Ieva Tolmane^{213,214}, Esther A. Torres²¹⁵, Michael Trauner²¹⁶, Sombat Treeprasertsuk²¹⁷, Adela Turcanu²¹⁸, Jonas Valantinas²¹⁹, Mette Vesterhus²²⁰, Imam Waked²²¹, Sarah H. Wild²²², Jose Willemse²²³, Robert J. Wong²²⁴, Stavra Xanthakos^{225,226}, Dan Yock Young²²⁷, Ming-Lung Yu²²⁸, Kenneth I. Zheng²²⁹, Mudjat Zeybel²³⁰ and Ming-Hua Zheng²²⁹

41 Transplantation and Liver Surgery, Helsinki University Hospital, Helsinki, Finland. 42 University of Helsinki, Helsinki, Finland. 43 University of Western Australia, Perth, Western Australia, Australia. *Inlantia. **Oniversity of Nestina University of Nestina Current Australia, Pertit, Western Australia, Pertit, Western Australia, Pertit, Western Australia, Australia, Australia, Australia, Australia, Australia, Australia, Pertit, Western Australia, Pertit, Perti Enfermedades Hepáticas y Digestivas (CIBERehd), Madrid, Spain. 56University of Health and Allied Sciences, Ho, Chana. 57University of KwaZulu-Natal, Durban, South Africa. 58Mongolian National University of Medical Sciences, Ulaanbaatar, Mongolia. 59 Research Institute of Virology of the Ministry of Health of the Republic of Uzbekistan, Tashkent, Uzbekistan. 60 University of Virginia, Charlottesville, VA, USA. ⁶¹University of Pittsburgh, Pittsburgh, PA, USA. ⁶²National Public Health Laboratory of Sudan and Ahfad University for Women, Khartoum, Sudan. ⁶³Diabetes Research Institute, San Raffaele Hospital and San Raffaele Vita Salute University, Milan, Italy. ⁶⁴Saint Joseph Hospital Marseille, Marseille, France. ⁶⁵General University Hospital and Charles University, Prague, Czech Republic. 65Department of Medical Sciences, University of Turin, Turin, Italy. 67University of Padua-Italy, Padua, Italy. 68Vall d'Hebron University Hospital, Barcelona, Spain. 69 Southampton National Institute for Health Research Biomedical Research Centre and the Institute of Developmental Sciences, University of Southampton, Southampton, UK. 79 Hospital Spain. "Southampton National institute of nearth research Centre and the Institute of Developmental Sciences, Continuity of Southampton, Southampton, Southampton, Southampton, Southampton, Southampton, Southampton, William Continuity, Ok. "Prospiration of Puerta de Hierro, Majadahonda, Spain. "I'Aix Marseille Univ, Inserm, IRD, SESSTIM, Sciences Economiques & Sociales de la Santé & Traitement de l'Information Médicale, ISSPAM, Marseille, France. "PUniversity of the West Indies, Nassau, Bahamas. "Institute of Gastroenterology, Medical University of Havana, Havana, Cuba. "Huenos Aires, Argentina." Statin American Association for the Study of the Liver (ALEH), Mexico City, Mexico. "6Gastroenterology Department, Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán, Mexico City, Mexico." University of Medicine, University of Malaya, Kuala Lumpur, Malaysia. PCenter for Cohort Studies, Total Healthcare Center, Kangbuk Samsung Hospital, Sungkyunkwan University School of Medicine, Seoul, Republic of Korea. San Raffaele Hospital, Milan, Italy. 81 Department of Medicine, University of Otago, Dunedin, New Zealand. 82 Massachusetts General Hospital and Harvard Medical School, Boston, MA, USA. 83 University of Palermo, Palermo, Italy. ⁸⁴Global Liver Institute, Washington, DC, USA. ⁸⁵University of Kelaniaya, Colombo, Sri Lanka. ⁸⁶University of Cabo Verde, Praia, Cabo Verde. ⁸⁷Hospital Center University of Bordeaux, Bordeaux, France. ⁸⁸University of Pisa, Pisa, Italy. ⁸⁹VicHealth, Melbourne, Victoria, Australia. ⁹⁰St. Paul's Hospital Millennium Medical College, Addis Ababa, Ethiopia. ⁹¹University of Dundee, Dundee, UK. ⁹²Postgraduate Institute of Medical Education and Research, Chandigarh, India. ⁹³World Heart Federation, Geneva, Switzerland. ⁹⁴Department of Health, Medicine and Caring Sciences, Linköping University, Linköping, Sweden. 95Endemic Medicine Department, Faculty of Medicine, Helwan University, Cairo, Egypt. 96The National Ribat University, Khartoum, Sudan. ⁹⁷Cairo University Hospitals, Cairo, Egypt. ⁹⁰Department of Gastroenterology, Xinhua Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai, China. ⁹⁹European Association for the Study of Obesity (EASO), Teddington, UK. ¹⁰⁰Medical University of Bialystok, Bialystok, Poland. ¹⁰¹Minia University, Minia, Egypt. ¹⁰²Virginia Commonwealth University and Central Virginia Veterans Affairs Health Systems, Richmond, VA, USA. ¹⁰³Virginia Commonwealth University, Division of Gastroenterology, Hepatology and Nutrition, Richmond, VA, USA. ¹⁰⁴Hepatobiliary Division, Department of Internal Medicine, Dr. Cipto Mangunkusumo National General Hospital, Universitas Indonesia, Jakarta, Indonesia. 105 Inova Health System, Falls Church, VA, USA. 106Armenian Association for the Study of the Liver (ARASL), Yerevan, Armenia. 107University of Medicine and Pharmacy Carol Davila, Bucharest, Romania. 108Department of Gastroenterology Arthenian Sociation the Study of the Liver (ARASC), relevan, Arthenian. Onliversity of Medicine and Friannacy Carlo Davila, Bucharlest, Rolliadia. Department of Sastroenterology, Singapore General Hospital, Singapore, Singapore, 1ºº Department of Hepatology and Gastroenterology, Aarhus University Hospital, Aarhus, Denmark. 1ºº Azerbaijan Gastroenterologists and Hepatologists Association, Baku, Azerbaijan. 1ºº Agenta Hospital, Brisbane, Queensland, Australia. 1º4 Department of Endocrinology, Royal Prince Alfred Hospital and Central Clinical School, Faculty of Medicine and Health, University of Sydney, New South Wales, Australia. 1º5 Somogy County Kaposi Mór Teaching Hospital, Kaposvár and University of Pécs, Pécs, Hungary. 1º6 Ankara University School of Medicine, Ankara, Turkey. 1º7 Department of Gastroenterology & Hepatology, Federal Research Center of Nutrition, Biotechnology & Food Safety, Moscow, Russia. 1º8 Kuwait University, Kuwait City, Kuwait. 1º9 Horacio Oduber Hospital, Oranjestad, Aruba. ¹²⁰Department of Internal Medicine, College of Medicine, The Catholic University of Korea, Seoul, Republic of Korea. ¹²¹National Academy of Medicine, College of Medicine, The Catholic University of Korea, Seoul, Republic of Korea. ¹²¹National Academy of Medicane, Houston, TX, USA. ¹²⁵National Hospital Organization Takasaki General Medical Center, Takasaki, Japan. ¹²⁵Liver Patients International, Atthens, Greece. ¹²⁴Skaylor College of Medicine, Houston, TX, USA. ¹²⁵National Taiwan University Hospital, Taipei, Taiwan. ¹²⁶Division of Gastroenterology, Department of Medicine, Kurume University School of Medicine, Kurume, Japan. ¹²⁷Jordan University of Science and Technology, Irbid, Jordan. ¹²⁸Yonsei University, Seoul, Republic of Korea. ¹²⁹University of Abomey Calavi, Faculty of Health Sciences, Cotonou, Benin. 130 Department of Internal Medicine, Division of Gastroenterology and Hepatology, Maastricht University Medical Center, Maastricht, Netherlands. 131 University of Tokyo, Tokyo, Japan. 132 Pasteur Institute of Bangui, Bangui, Central African Republic. 135 European Liver Patients Association, Brussels, Belgium. 134 South Denver Gastroenterology, Denver, CO, USA. 135 Gastroenterology "A"

Department, Rabta Hospital, Faculty of Medicine of Tunis, Tunis, Tunisia. ¹⁵⁶University Clinic for Visceral Surgery and Medicine, Inselspital Bern, Bern, Switzerland. ¹⁵⁷Columbia University, New York, NY, USA. ¹³⁸Dornsife School of Public Health, Drexel University, Philadelphia, PA, USA. ¹³⁹Wilton Park, Steyning, UK. ¹⁴⁰Boston University School of Medicine, Boston, MA, USA. ¹⁴¹University of Santander (UDES), Bucaramanga, Colombia. 142 Digestive Disease Research Center, Tehran University of Medical Science, Tehran, Iran. 143 Bangabandhu Sheikh Mujib Medical University, 144 University of Bologna, Bologna, Italy. 145 Lisbon North University Hospital Center, Lisbon, Portugal. 146 University Hospital St. Ivan Rilski, Sofia, Bulgaria. 147 Service of Driaka, Baligladesti. "University of Bologia, Bologia, Italy. "Elsboth North Oniversity Prospital Center, Lisboth, Portugal." "University Prospital St. Ivan Riski, Solia, Bulgaria. "Service of Hepatogastroenterology, Department of Internal Medicine, University Clinic of Kinshasa, University of Kinshasa, Kinshasa, Democratic Republic of the Congo. 148Saint-Dizier Hospital Center, Saint Dizier, France. 149University Hospital Center Rijeka, Rijeka, Croatia. 159Clinical Center of Serbia, Beograd, Serbia. 151Favaloro Foundation, Buenos Aires, Argentia. 151F Castroenterorlepatology, University Ss. Cyfi and Metriodius, Skopje, Republic of North Macedonia. 158 Jamies's Polypian, Dublin, Hedard. 159 University Medical Center Ljubijana, Ljubijana, Slovenia. 158 Guy's and St Thomas' Hospital, London, UK. 159 University of the Philippines College of Medicine, Manila, Philippines. 160 Lagos State University College of Medicine, Lagos, Nigeria. 151 National and Kapodistrian University of Athens, Athens, Greece. 152 University of KwaZulu Natal, Durban, South Africa. 153 University Health Network Toronto, Ontario, Canada. 154 Centro de Estudos de Doenças Crónicas (CEDOC), NOVA Medical School-FCM, Universidade Nova de Lisboa, Lisboa, Portugal. 155 Instituto Gulbenian de Ciencia, Oeiras, Portugal. 156 Dominican Society of Gastroenterology, Santo Domingo, Dominican Republic. 157 Gastroenterology at Bayerischer Platz, Berlin, Germany. 158 University of Sao Paulo School of Medicine, Sao Paulo, Brazil. 159 NYU Langone Health, NYU Grossman School of Medicine, New York, NY, USA. 170 Teaching Hospital Nitra, Nitra, Slovakia. 171 World Obesity Federation, London, UK. 172 University of British Columbia, Vancouver, British Columbia, Canada. 173 Center for Disease Analysis Foundation, Lafayette, LA, USA. 174 School of Medicine, Universidade Federal do Rio Grande do Sul, Porto Alegre, Brazil. 175 Alfred Health, Melbourne, Victoria, Australia. 176 Department of Endocrinology and Diabetology, Medical Faculty and University Hospital, Heinrich-Heine University, Düsseldorf, Germany. 177Institute for Clinical Diabetology, German Diabetes Center, Leibniz Center for Diabetes Research at Heinrich-Heine University, Düsseldorf, Germany. 178German Center for Diabetes Research (DZD), Partner Düsseldorf, Düsseldorf, Germany. 179Friends of Europe, Brussels, Belgium. 180University Hospital Touhami Benflis, Batna, Algeria. 181IRCCS San Raffaele Hospital and University, Milan, Italy. 182University of Leeds, Leeds, UK. 183University of Tartu, Tartu University Hospital, Tartu, Estonia. 184Kyrgyz State Medical Institute for Advanced Training and Retraining, Bishkek, Kyrgyz Republic. 185Hotel Dieu de France and St Joseph University School of Medicine, Beirut, Lebanon. 185McGill University Health Centre, Montreal, Quebec, Canada. 187Weight Loss Bislikek, Krygyz Republic. "Froter Died de France and Scrosepin Oniversity Scribot of Medicalic, Republic. Surgery Center, Votsuya Medical Cube, Tokyo, Japan. ¹⁸⁸Ciudad Sanitaria Dr. Luis E. Aybar, Santo Domingo, Dominican Republic. ¹⁸⁹University of Ouagadougou, Burkina Faso. ¹⁹⁰Baker Heart and Diabetes Institute, Melbourne, Victoria, Australia. ¹⁹¹Sree Gokulam Medical College, Venjarammoodu, Kerala, India. ¹⁹²Kings College London, London, UK. ¹⁹³Tel-Aviv Medical Centre and Tel-Aviv University, Tel-Aviv, Israel. ¹⁹⁴Austral University Hospital, Pilar, Argentina. ¹⁹⁵Poltava State Medical University, Poltava, Ukraine. ¹⁹⁶The Children's Memorial Health Institute, Warsaw, Poland. ¹⁹⁷La Princesa University Hospital, Madrid, Spain. ¹⁹⁸Division of Hepatology, Department of Medicine, Faculty of Health Sciences, University of Cape Town, Cape Town, South Africa. 199Department of Pharmacology & Therapeutics, College of Medicine & Medical Sciences, Arabian Gulf University, Manama, Kingdom of Bahrain. 200Universidad San Cape flown, South Africa. "Pepartment of Pharmacology & Therapeutics, College of Medicine & Medical Sciences, Arabian Guil University, Mahama, Angdom of Barriant. "Ornwerstada Sciences, Arabian Guil University, Mahama, Angdom of Barriant." "Ornwerstada Sciences, Arabian Guil University, Mahama, Angdom of Barriant. "Ornwerstada Sciences, Arabian Guil University, Mahama, Angdom of Barriant." "Ornwerstada Sciences, Arabian Guil University, Seoul, Horgistada Sciences, Arabian Guil University, Mahama, Angdom of Barriant. "Ornwerstada Sciences, Arabian Guil University, Mahama, Angdom of Barriant." "Ornwerstada Sciences, Arabian Guil University, Mahama, Angdom of Barriant. "Ornwerstada Sciences, Arabian Guil University, Mahama, Angdom of Barriant." "Ornwerstada Sciences, Arabian Guil University, Mahama, Angdom of Barriant. "Ornwerstada Sciences, Arabian Guil University, Mahama, Angdom of Barriant. "Ornwerstada Sciences, Arabian Guil University, Mahama, Angdom of Barriant." "Ornwerstada Sciences, Arabian Guil University, Mahama, Angdom of Barriant. "Ornwerstada Sciences, Arabian Guil University, Mahama, Angdom of Barriant. "Ornwerstada Sciences, Arabian Guil University, Mahama, Angdom of Barriant." "Ornwerstada Sciences, Arabian Guil University, Mahama, Angdom of Barriant. "Ornwerstada Sciences, Arabian Guil University, Mahama, Angdom of Barriant. "Ornwerstada Sciences, Arabian Guil University, Mahama, Angdom of Barriant." "Ornwerstada Sciences, Arabian Guil University, Mahama, Angdom of Barriant. "Ornwerstada Sciences, Arabian Guil University, Mahama, Angdom ornwerstada, Angdom ornwerst Riga, Latvia. 214 University of Latvia, Riga, Latvia. 215 University of Puerto Rico School of Medicine, San Juan, Puerto Rico. 216 Medical University of Vienna, Vienna, Austria. 217 King Chulalongkorn Memorial Hospital and Chulalongkorn University, Bangkok, Thailand. 218 Gastroenterology, State University of Medicine and Pharmacy "Nicolae Testemitanu", Chişinău, Republic of Moldova. ²¹⁹Vilnius University Hospital Santaros Clinics, Vilnius, Lithuania. ²²⁹Haraldsplass Deaconess Hospital and University of Bergen, Bergen, Norway, ²²¹National Liver Institute, Shibin Al Kawm, Egypt. ²²²Usher Institute, University of Edinburgh, UK. ²²⁵Dutch Liver Patients Association, Hoogland, Netherlands. ²²⁴Stanford University School of Medicine, Stanford, CA, USA. ²²⁵Cincinnati Children's Hospital Medical Center, Cincinnati, OH, USA. ²²⁶University of Cincinnati College of Medicine, Cincinnati, OH, USA. ²²⁷National University, Singapore, Singapore. 228 Hepatobiliary Section, Department of Internal Medicine, Kaohsiung Medical University Hospital, Kaohsiung Medical University, Kaohsiung, Taiwan. 229 NAFLD Research Center, Department of Hepatology, the First Affiliated Hospital of Wenzhou Medical University, Wenzhou, China. 250 Koç Üniversitesi, Istanbul, Turkey