

### **Tuliskan kesimpulan analisis.**

Penelitian ini menyimpulkan bahwa faktor usia, jenis kelamin, tinggi badan dan riwayat asma berhubungan secara signifikan dengan nilai PEF. Analisis yang digunakan adalah analisis multivariat dengan menghitung nilai masing-masing variabel terhadap PEF. Dari beberapa pemodelan yang dilakukan didapatkan bahwa kombinasi keempat faktor merupakan pemodelan terbaik dengan nilai AIC terendah.

### **Referensi:**

Thomas ET , Guppy M, Straus SE, Bell K, Glasziou, Rate of normal lung function decline in ageing adults: a systematic review of prospective cohort studies. *BMJ Open*. 2019 27;9(6):e028150. doi: 10.1136/bmjopen-2018-028150.

Ji C, Xia Y, Dai H, Zhao Z, Liu T, Tong S, et al. Reference Values and Related Factors for Peak Expiratory Flow in Middle-Aged and Elderly Chinese. *Front Public Health*. 2021 Aug 20;9:706524. doi: 10.3389/fpubh.2021.706524. eCollection 2021.

Ridwan ES, Wiratama BS, Lin M-Y, Hou W-H, Liu MF, Chen C-M, et al. (2021) Peak expiratory flow rate and sarcopenia risk in older Indonesian people: A nationwide survey. *PLoS ONE* 16(2): e0246179. <https://doi.org/10.1371/journal.pone.0246179>

Landi F, Salini S, Zazzara MB, Martone AM, Fabrizi S, Bianchi M, et al. Relationship between pulmonary function and physical performance among community-living people: results from Look-up 7+ study. *J Cachexia Sarcopenia Muscle*. 2020;11(1):38–45.

Kera T, Kawai H, Hirano H, Kojima M, Fujiwara Y, Ihara K, et al. Relationships among peak expiratory flow rate, body composition, physical function, and sarcopenia in community-dwelling older adults. *Aging Clinical and Experimental Research*. 2018;30(4):331–40.