Meshfree MUSCL schemes with a posteriori order detection: exactness, conservation and admissibility

Pascal Jung*

Draft of a mathematical companion, 28 October 2026, revised 5 November 2026

Abstract

We collect the mathematical framework of a class of meshfree MUSCL schemes for hyperbolic systems of conservation laws on scattered point clouds, whose implementation and numerical study are reported in a companion paper [11]. Midpoint Riemann problems between moving least-squares (MLS) reconstructions are differentiated by an MLS derivative operator with zero row sums. We prove exactness for polynomial solutions of linear problems, consistency of the design order on Cartesian lattices for an odd derivative degree selected by a parity argument, and an exact balance law that identifies the conservation error with the lack of antisymmetry of the volume-weighted derivative weights. Orienting the Rusanov dissipation by the signs of the derivative weights (the aligned orientation) makes the first-order scheme positivity-preserving for the Euler equations, for derivative weights of any degree. For the per-particle MOOD procedure with three interface rules we characterize the rules under which a particle at the lowest order is advanced by the first-order scheme. Finally we show that stage-wise MOOD inside an arbitrary explicit strong-stability-preserving Runge–Kutta method yields admissible stages and updates for every distribution of the orders, provided that the detection checks the forward Euler substeps of the canonical Shu–Osher form rather than the stage values; for the SSP methods of Heun (RK2) and Shu–Osher (RK3), whose stages contain only U^n and the last substep, checking the stage values gives the same guarantee, for SSPRK(5,4) it does not.

Keywords: meshfree methods; moving least squares; MUSCL; MOOD; positivity preservation; strong-stability-preserving Runge–Kutta methods; local conservation
MSC (2020): 65M12, 65M06, 65L06, 35L65

1 Introduction

This note isolates the analysis of the meshfree MUSCL schemes of [11], implemented in HyperCloudCore.jl [10] (version 0.7.8), from their numerical study. Every particle of a point cloud carries an unknown state, every pair of neighbours carries a Riemann problem between two MLS reconstructions evaluated at the pair midpoint, and an MLS derivative operator with zero row sums differentiates the interface fluxes; the construction extends the meshless MUSCL schemes of Willems, Klar and Samaey [16] for linear advection to nonlinear systems. Discontinuities are treated by Multi-dimensional Optimal Order Detection (MOOD) [4–6], applied per particle and per stage of an explicit Runge–Kutta method.

The results are of three kinds.

- *Approximation* (Sections 3 and 4.3): reproduction and scaling of the MLS weights (Lemmas 3.3 and 3.5), a parity property on centrally symmetric stencils that motivates odd derivative degrees (Lemma 3.7), exactness for polynomial solutions of linear problems on arbitrary clouds (Proposition 4.2) and consistency of the design order on lattices (Proposition 4.3).

*Rheinland-Pfälzische Technische Universität Kaiserslautern-Landau, pajung@rptu.de

- *Conservation* (Section 4.4): an exact balance law in which the conservation error is expressed by pair defects of the volume-weighted weights (Proposition 4.4).
- *Admissibility* (Sections 5 and 6): positivity of the first-order update with zero-sum weights and the aligned orientation (Proposition 5.2 and Corollary 5.3), the fallback property of the interface rules (Proposition 5.6), a local admissibility result for a particle at order one (Proposition 5.7), and admissibility of all stages under stage-wise MOOD within SSP methods for arbitrary order distributions (Theorem 6.3).

Stability and convergence of the high-order schemes are not addressed. The statements are formulated with all hypotheses that the proofs use; Remark 1.1 states how they relate to the implementation, and Appendix A lists the differences to the corresponding statements in the draft of [11] of 27 October 2026.

Remark 1.1 (Theory and implementation). The purpose of this note is to justify the implementation, and where the two differ, the implementation is the reference. The scheme analysed here is the one of HyperCloudCore.jl 0.7.8 with the aligned Rusanov flux. Three deviations remain, and the corresponding statements then hold as approximate equalities rather than exactly:

(i) *Regularized fits.* Numerically degenerate stencils are fitted with a Tikhonov term (Remark 3.2). For such a particle, reproduction and the zero row sums of Lemma 3.3(i),(ii) hold only approximately, $\sum_j c_{ij} \approx 0$, and so do condition (a) of Proposition 5.2 and all results that use it.

(ii) *Pairs with $x_{ij,d} = 0$.* On geometric edges the implementation orients such pairs by $\sigma_{ij,d} = \sigma_{ji,d} = -1$, which breaks the symmetry of Lemma 4.1 and the bound of Proposition 4.4(i) on these pairs (Proposition 4.4(iii)). Such pairs are rare on perturbed clouds (in single precision about $5 \cdot 10^{-5}$ of the pairs on the clouds of [11]), and on Cartesian lattices their weights vanish by the reflection symmetry of the stencil (proof of Proposition 4.3).

(iii) *Time step.* The time step of the implementation enforces the conditions (5.1) and (H4) of Theorem 6.3 only approximately, up to the difference between the weights with and without constant term and the change of the wave speeds within a step (Remark 6.5(iii)).

Besides these, all identities hold up to round-off.

Remark 1.2 (Machine-checked proofs). The definitions and results of Sections 3 to 6 have been formalized in Lean 4 with Mathlib; the sources are in the folder `lean/` accompanying this note, whose `README.md` maps every statement to its Lean counterpart. All fits are modelled as unregularized, so the approximate statements of Remark 1.1 are not covered. The following are checked in full: Lemmas 3.3 and 3.5, Lemma 3.7(i) and (ii), Definition 3.9, Lemma 4.1, Proposition 4.4 with (i)–(iii), the convexity of $\mathcal{G}$ for the Euler equations, Lemmas 5.1 and 6.1 and Proposition 5.6, as well as the canonical forms of forward Euler, RK2 and RK3. Propositions 5.2 and 5.7 and Corollary 5.3 are checked under slightly stronger hypotheses: $s_{ij,d} > 0$ for all j and d (instead of only where $c_{ij}^{(d)} \neq 0$) and $\lambda_d > 0$ on $\mathcal{G}$, both automatic for the Euler equations; the aligned version of Proposition 5.2 takes the zero row sums as a hypothesis. Of the counterexample of Lemma 3.7(iii) the library checks the value $17/5 \neq 0$; that the stencil is centrally symmetric and Π_2-unisolvent is elementary and is not part of the library. The following are checked with ingredients taken as hypotheses: Corollary 3.8 (the Taylor remainder bound), Proposition 4.2 (the interface states $U_{ij} = U_{ji} = u(m_{ij})$, which follow from the reproduction property of Section 3.1, also checked, together with the translation invariance of Π_q, which is not formalized), Theorem 6.3 and Corollary 6.4 (the Shu–Osher relation (6.3) of the stages, which is not derived from Lemma 6.1 there; (H3); the exact locality of Algorithm 1; and, in place of (H4) and Proposition 5.7, the admissibility of the substep at every order-one particle for every distribution of the orders, which is stronger than (H4); termination is checked), and Remark 6.2 (the negative coefficient, for all coefficients within the rounding of Table 1). Of Proposition 4.3

only the parity and cancellation steps are formalized, not the $O(h)$ expansions; the maximality of the SSP coefficients is not formalized.

2 Setting

2.1 Hyperbolic systems

We consider systems of conservation laws

$$\partial_t U + \nabla \cdot F(U) \;=\; \partial_t U + \sum_{d=1}^{D} \partial_d F_d(U) \;=\; 0, \qquad x \in \Omega \subset \mathbb{R}^D,\; t > 0, \tag{2.1}$$

for $U(x,t) \in \mathcal{G} \subset \mathbb{R}^M$, where the admissible set $\mathcal{G}$ is convex and the fluxes $F_d \colon \mathcal{G} \to \mathbb{R}^M$ are smooth. The system is hyperbolic, and $\lambda_d(U) \geq \max\{|\mu| : \mu \text{ eigenvalue of } \partial F_d/\partial U\}$ is a bound of the wave speeds in direction d. We write $F(U) = (F_1(U), \dots, F_D(U)) \in \mathbb{R}^{M\times D}$ and, for $c \in \mathbb{R}^D$ and $G = (G_1, \dots, G_D) \in \mathbb{R}^{M\times D}$, $c \cdot G := \sum_d c^{(d)} G_d \in \mathbb{R}^M$.

The main example are the Euler equations of an ideal gas,

$$U = \begin{pmatrix} \rho \\ \rho u \\ E \end{pmatrix}, \qquad F_d(U) = \begin{pmatrix} \rho u_d \\ \rho u_d\, u + p\, e_d \\ u_d\,(E+p) \end{pmatrix}, \qquad p = (\gamma-1)\Big(E - \tfrac{1}{2}\rho|u|^2\Big), \tag{2.2}$$

with $M = D+2$, $\gamma > 1$, the sound speed $c = \sqrt{\gamma p/\rho}$, $\lambda_d(U) = |u_d| + c$ and $\mathcal{G} = \{U : \rho > 0,\; p > 0\}$, which is convex because p is a concave function of U for $\rho > 0$.

2.2 Point clouds

The domain is represented by particles $X = \{x_1, \dots, x_N\} \subset \mathbb{R}^D$ with nominal spacing h. The interior particles $I = \{i : x_i \in \Omega\}$ carry the unknowns $U_i(t) \approx U(x_i, t)$; the remaining particles are ghost particles in a layer outside Ω, whose states are set by a boundary operator from the interior states and are not advanced by the scheme. With a support radius $R = \kappa h$,

$$\nu(i) = \{\, j \neq i : |x_j - x_i| \leq R \,\}, \qquad \bar{\nu}(i) = \nu(i) \cup \{i\}, \tag{2.3}$$

so that $j \in \nu(i)$ if and only if $i \in \nu(j)$. We write $x_{ij} = x_j - x_i$ and $m_{ij} = \frac{1}{2}(x_i + x_j)$, and use weights $w_{ij} = w(|x_{ij}|/R)$ with a positive, radially decreasing function w with $w(0) = 1$. For the conservation analysis we use positive particle volumes V_i, for instance Shepard volumes; the scheme itself does not use them.

For asymptotic statements on general clouds we use the following assumption.

Assumption 2.1 (Quasi-uniform, uniformly well-conditioned clouds)**.** A family of clouds X_h with fixed κ satisfies, with constants independent of h and i: (i) $|\nu(i)| \leq N_{\max}$; (ii) for the degrees q under consideration, the scaled moment matrices $\tilde{N}_i^s = \sum_{j\in\nu(i)} w_{ij}\, \tilde{p}_q(x_{ij}/h)\, \tilde{p}_q(x_{ij}/h)^{\mathsf{T}}$ (with $\tilde{p}_q$ from Section 2.3) are invertible with $\|(\tilde{N}_i^s)^{-1}\|_2 \leq C_N$.

Condition (i) follows from a uniform separation $|x_{ij}| \geq c_{\mathrm{sep}} h$. Condition (ii) quantifies unisolvency uniformly. For dilations of one fixed cloud, in particular for Cartesian lattices, $\tilde{N}_i^s$ does not depend on h, and (ii) reduces to unisolvency (Definition 3.1) of the finitely many distinct stencils.

2.3 Taylor bases

For a multi-index $\alpha \in \mathbb{N}_0^D$ we use the standard notation $|\alpha|$, $\alpha!$ and y^α. For a degree $q \geq 1$ let

$$p_q(y) = \Big(\frac{y^\alpha}{\alpha!}\Big)_{1\leq|\alpha|\leq q} \in \mathbb{R}^{B_q}, \qquad \tilde{p}_q(y) = \big(1,\ p_q(y)\big) \in \mathbb{R}^{B_q+1}, \qquad B_q = \binom{q+D}{D} - 1, \quad (2.4)$$

be the Taylor bases without and with constant term, ordered by degree, with $y_1, \ldots, y_D$ first. Π_q denotes the polynomials of degree at most q on $\mathbb{R}^D$.

3 Moving least-squares operators

Definition 3.1 (Unisolvent neighbourhoods)**.** The neighbourhood of particle i is Π_q-unisolvent if the only $g \in \Pi_q$ with $g(x_{ij}) = 0$ for all $j \in \nu(i)$ is $g = 0$. This requires $|\nu(i)| \geq B_q + 1$.

3.1 Reconstruction fits

Let $f_j \in \mathbb{R}^M$ be values at the particles. The reconstruction fit of degree $r \geq 1$ at particle i is the coefficient matrix $A_i = (a_{i,\alpha})_{1\leq|\alpha|\leq r} \in \mathbb{R}^{B_r\times M}$ that minimizes

$$\sum_{j\in\nu(i)} w_{ij}\,\big|A^{\mathsf{T}}p_r(x_{ij}) - (f_j - f_i)\big|^2, \qquad \text{i.e.} \qquad A_i = N_i^{-1}\sum_{j\in\nu(i)} w_{ij}\,p_r(x_{ij})\,(f_j - f_i)^{\mathsf{T}}, \quad (3.1)$$

with the moment matrix $N_i = \sum_j w_{ij}\,p_r(x_{ij})\,p_r(x_{ij})^{\mathsf{T}}$. If the neighbourhood is Π_r-unisolvent, N_i is symmetric positive definite (a polynomial in p_r without constant term vanishing at all offsets vanishes identically). The reconstruction polynomial $\mathcal{R}_i(y) = f_i + A_i^{\mathsf{T}}p_r(y)$ satisfies $\mathcal{R}_i(0) = f_i$ and reproduces polynomials: if $f_j = g(x_j)$ for all $j \in \bar{\nu}(i)$ with $g \in \Pi_r$, then $\mathcal{R}_i(y) = g(x_i + y)$, because $g(x_i + \cdot) - g(x_i)$ lies in the span of p_r and attains the minimum zero, which is unique by unisolvency.

Remark 3.2 (Regularization). The implementation formulates the fits in the scaled variable y/h and factorizes the moment matrix by Cholesky's method. Only if this fails or produces a pivot at round-off level is the factorization repeated with $N + \varepsilon\,\mathrm{tr}(N)/B\cdot\mathrm{Id}$, where B is the number of unknowns of the fit and $\varepsilon = \sqrt{\varepsilon_{\mathrm{mach}}}$. We call a fit *unregularized* if the first factorization is used. A regularized fit is not the least-squares fit: it neither reproduces polynomials nor, for derivative fits, has exactly zero row sums. All reproduction statements below therefore require unregularized fits; for regularized fits they hold only approximately (Remark 1.1(i)).

3.2 Derivative fits with zero row sums

For $q \geq 1$ let $\tilde{N}_i = \sum_{j\in\nu(i)} w_{ij}\,\tilde{p}_q(x_{ij})\,\tilde{p}_q(x_{ij})^{\mathsf{T}}$. The *derivative weights* of degree q at particle i are

$$c_{ij}^{(d)} = w_{ij}\ e_{1+d}^{\mathsf{T}}\,\tilde{N}_i^{-1}\,\tilde{p}_q(x_{ij}), \qquad d = 1,\ldots,D, \quad j \in \nu(i), \quad (3.2)$$

where e_{1+d} selects the coefficient of y_d. They are the weights with which the least-squares fit $a_0 + a^{\mathsf{T}}p_q(y)$ of data g_j at the offsets x_{ij}, over the neighbours only, produces $a_d = \sum_j c_{ij}^{(d)} g_j$. For data $G_j \in \mathbb{R}^{M\times D}$ the divergence operator is

$$\mathcal{D}_i[G] = \sum_{j\in\nu(i)} c_{ij}\cdot G_j = \sum_{j\in\nu(i)}\sum_{d=1}^{D} c_{ij}^{(d)}\,G_{j,d}. \quad (3.3)$$

Lemma 3.3. *Let the neighbourhood of particle i be Π_q-unisolvent and the derivative fit unregularized. Then:*

(i) $\sum_{j\in\nu(i)} c_{ij}^{(d)}\,g(x_{ij}) = \partial_d g(0)$ *for all* $g \in \Pi_q$ *and* $d = 1,\ldots,D$*;*

(ii) in particular $\sum_{j\in\nu(i)} c_{ij} = 0$;

(iii) if the cloud is dilated by a factor $\theta > 0$, $x \mapsto \theta x$, *with* R *scaled accordingly, the weights become* c_{ij}/θ.

Proof. (i) For $g \in \Pi_q$ the data $g(x_{ij})$ are represented exactly by g itself, which lies in the span of $\tilde{p}_q$. Unisolvency makes the least-squares minimizer unique, so it is g, and its coefficient of y_d is $\partial_d g(0)$. (ii) follows with $g \equiv 1$. (iii) The weights w_{ij} depend on $|x_{ij}|/R$ only, and $\tilde{p}_q(\theta y) = S_\theta\, \tilde{p}_q(y)$ with $S_\theta = \mathrm{diag}(\theta^{|\alpha|})_{0\le|\alpha|\le q}$. Hence $\tilde{N}_i$ becomes $S_\theta \tilde{N}_i S_\theta$, and (3.2) becomes $w_{ij}\, e_{1+d}^{\mathsf{T}} S_\theta^{-1} \tilde{N}_i^{-1} \tilde{p}_q(x_{ij}) = \theta^{-1} c_{ij}^{(d)}$. □

Remark 3.4 (The constant term). A derivative fit in the basis p_q alone, forced through the particle value as in (3.1), has weights $\hat{c}_{ij}$ whose row sums are the first D entries of $N_i^{-1}\sum_j w_{ij} p_q(x_{ij})$; they vanish on symmetric stencils but not in general. The constant term reproduces constant data by the intercept and removes this defect. Including the particle itself as a data point would reintroduce it, since its own weight would then balance those of the neighbours. The derivative fit has one more unknown than a reconstruction fit of the same degree.

Lemma 3.5 (Size of the weights). *Let Assumption 2.1 hold and* $C_\kappa = \max_{|z|\le\kappa} |\tilde{p}_q(z)|$. *Then*

$$|c_{ij}^{(d)}| \le C_N C_\kappa\, h^{-1} \qquad \text{and} \qquad \sum_{j\in\nu(i)} \sum_{d=1}^{D} |c_{ij}^{(d)}| \le D\, N_{\max} C_N C_\kappa\, h^{-1}.$$

Proof. With $S_h = \mathrm{diag}(h^{|\alpha|})$ we have $\tilde{p}_q(y) = S_h \tilde{p}_q(y/h)$, $\tilde{N}_i = S_h \tilde{N}_i^s S_h$ and hence $c_{ij}^{(d)} = w_{ij}\, e_{1+d}^{\mathsf{T}} S_h^{-1} (\tilde{N}_i^s)^{-1} \tilde{p}_q(x_{ij}/h) = h^{-1} w_{ij}\, e_{1+d}^{\mathsf{T}} (\tilde{N}_i^s)^{-1} \tilde{p}_q(x_{ij}/h)$. Since $w_{ij} \le 1$ and $|x_{ij}/h| \le \kappa$, the bound follows from $\|(\tilde{N}_i^s)^{-1}\|_2 \le C_N$ and $|\nu(i)| \le N_{\max}$. □

Lemma 3.3(iii) is the special case of exact dilations; on general quasi-uniform clouds the scaling $O(h^{-1})$ needs the uniform conditioning of Assumption 2.1(ii), which is not implied by unisolvency of each stencil.

3.3 Symmetric stencils and the degree of the divergence

Definition 3.6. The neighbourhood of particle i is *centrally symmetric* if there is an involution $j \mapsto j'$ of $\nu(i)$ with $x_{ij'} = -x_{ij}$.

Since w is radial, then also $w_{ij'} = w_{ij}$. Cartesian lattices with κh-balls are centrally symmetric.

Lemma 3.7 (Parity). *Let the neighbourhood of particle* i *be centrally symmetric and* Π_q*-unisolvent, and the derivative fit unregularized.*

(i) $c_{ij'} = -c_{ij}$ *for all* $j \in \nu(i)$.

(ii) If q *is odd, then* $\sum_j c_{ij}^{(d)}\, g(x_{ij}) = \partial_d g(0)$ *for all* $g \in \Pi_{q+1}$.

(iii) If q *is even, (ii) fails in general.*

Proof. Let $P = \mathrm{diag}\big((-1)^{|\alpha|}\big)_{0\le|\alpha|\le q}$, so that $\tilde{p}_q(-y) = P\,\tilde{p}_q(y)$, $P^2 = \mathrm{Id}$ and $P^{\mathsf{T}} = P$. Reindexing the sum by the involution gives $P\tilde{N}_i P = \tilde{N}_i$, i.e. $P\tilde{N}_i = \tilde{N}_i P$, and therefore also $\tilde{N}_i^{-1} P = P \tilde{N}_i^{-1}$. (i) Using $w_{ij'} = w_{ij}$, $\tilde{p}_q(x_{ij'}) = P\tilde{p}_q(x_{ij})$ and $\tilde{N}_i^{-1} P = P\tilde{N}_i^{-1}$,

$$c_{ij'}^{(d)} = w_{ij}\, e_{1+d}^{\mathsf{T}} \tilde{N}_i^{-1} P\, \tilde{p}_q(x_{ij}) = w_{ij}\, e_{1+d}^{\mathsf{T}} P\, \tilde{N}_i^{-1} \tilde{p}_q(x_{ij}) = w_{ij}\, (P e_{1+d})^{\mathsf{T}} \tilde{N}_i^{-1} \tilde{p}_q(x_{ij}) = -c_{ij}^{(d)},$$

because $P e_{1+d} = -e_{1+d}$ (y_d has odd degree). (ii) By Lemma 3.3 it suffices to consider $g(y) = y^\gamma$ with $|\gamma| = q+1$, which is even since q is odd. The right-hand side $b = \sum_j w_{ij}\, \tilde{p}_q(x_{ij})\, g(x_{ij})$

of the normal equations satisfies $Pb = \sum_j w_{ij}\,\tilde{p}_q(-x_{ij})\,g(-x_{ij}) = b$, and therefore $a = \tilde{N}_i^{-1}b$ satisfies $Pa = \tilde{N}_i^{-1}Pb = a$: all odd-degree entries of a vanish. In particular the coefficient of y_d is $0 = \partial_d g(0)$. (iii) For $D = 1$, $q = 2$ and $g(y) = y^3$ the odd part of the fit decouples from the even part, and the coefficient of y is $\sum_j w_{ij}x_{ij}^4 / \sum_j w_{ij}x_{ij}^2 \neq 0 = g'(0)$. □

Corollary 3.8 (Derivative error)**.** *Let Assumption 2.1 hold, let the neighbourhoods be Π_q-unisolvent with unregularized derivative fits, and let g be a smooth function. Then*

$$\Big|\sum_j c_{ij}^{(d)}\big(g(x_{ij}) - g(0)\big) - \partial_d g(0)\Big| = \begin{cases} O(h^{q+1}) & q \text{ odd, centrally symmetric stencil,} \\ O(h^q) & \text{otherwise.} \end{cases} \tag{3.4}$$

Proof. Split $g(y) - g(0) = T_q(y) + T_{q+1}(y) + \rho(y)$ into the Taylor terms of degrees $1, \dots, q$, the terms of degree $q+1$, and a remainder $|\rho(y)| \le C|y|^{q+2}$. By Lemma 3.3(i) the first part gives $\partial_d g(0)$ exactly. On a centrally symmetric stencil with odd q the second part contributes zero by Lemma 3.7(ii); otherwise it contributes at most $\sum_j |c_{ij}^{(d)}|\, C(\kappa h)^{q+1} = O(h^q)$ by Lemma 3.5. The remainder contributes $O(h^{-1})\,O(h^{q+2})$. The constants depend on g only through bounds of its derivatives of orders $q+1$ and $q+2$ near 0, so the estimate holds uniformly for a family g_i whose derivatives are bounded uniformly in i. □

Definition 3.9 (Divergence degree)**.** For a design order $k \ge 1$ the derivative fit has the degree

$$q(k) = 2\lceil k/2 \rceil - 1, \tag{3.5}$$

the smallest odd degree that is not below $k-1$: $q(k) = k-1$ for even k and $q(k) = k$ for odd k.

Remark 3.10 (Safeguards). The implementation uses the odd increase from $k-1$ to k only if $|\nu(i)| > 1.5\,(B_k + 1)$, applies it only up to degree three in three dimensions, and lowers the degree of the derivative fit until $|\nu(i)| \ge B_q + 1$. A particle whose neighbourhood is too small for its reconstruction fit is set to order one. The statements below refer to the degree q_i actually used by particle i.

4 The semi-discrete scheme

4.1 Interface states and divergence modes

Every interior particle carries an order $k_i \in \{1, \dots, K\}$, where K is the design order; without MOOD $k_i = K$. Particle i computes the reconstruction fit (3.1) of degree $k_i - 1$ of the differences $U_j - U_i$, component by component in conservative variables (no fit for $k_i = 1$), possibly followed by an a priori limiter; its coefficients define, for $r \le k_i$, the truncated reconstruction

$$\mathcal{R}_i^{[r]}(y) = U_i + \sum_{1 \le |\alpha| \le r-1} a_{i,\alpha}\,\frac{y^\alpha}{\alpha!}, \qquad \mathcal{R}_i^{[1]} \equiv U_i. \tag{4.1}$$

Each side of an edge has an *interface order* $k_{i\to j} \le k_i$, prescribed by the rules of Section 5.1 (without MOOD $k_{i\to j} = K$), and the interface states are

$$U_{ij} = \mathcal{R}_i^{[k_{i\to j}]}\big(\tfrac{1}{2}x_{ij}\big), \qquad U_{ji} = \mathcal{R}_j^{[k_{j\to i}]}\big(-\tfrac{1}{2}x_{ij}\big). \tag{4.2}$$

Ghost particles carry no reconstruction: $U_{ji} = U_j$ for a ghost neighbour j. The derivative weights of particle i have a degree q_i, chosen in one of two modes:

- *coupled*: $q_i = q(k_i)$;
- *frozen*: $q_i = q(k_{\mathrm{div}})$ for a fixed $k_{\mathrm{div}} \le K$, independent of k_i.

Both coincide for $K \le 2$.

4.2 Interface fluxes

For every coordinate direction we use the Rusanov flux

$$F^*_{ij,d} = \tfrac{1}{2}\big(F_d(U_{ij}) + F_d(U_{ji})\big) - \tfrac{1}{2}\, s_{ij,d}\, \sigma_{ij,d}\, \big(U_{ji} - U_{ij}\big), \qquad s_{ij,d} = \max\{\lambda_d(U_{ij}), \lambda_d(U_{ji})\}, \tag{4.3}$$

with an orientation $\sigma_{ij,d} \in \{-1, 0, 1\}$. We distinguish

- the *geometric* orientation $\sigma_{ij,d} = \operatorname{sign}(x_{ij,d})$, with the convention $\sigma_{ij,d} = \sigma_{ji,d} = -1$ of the implementation for $x_{ij,d} = 0$;
- the *aligned* orientation, $\sigma_{ij,d} = \operatorname{sign}(c^{(d)}_{ij})$, where $c^{(d)}_{ij}$ are the derivative weights that particle i uses, of degree q_i.

The *aligned Rusanov flux* uses the aligned orientation on every *aligned edge*, i.e. every edge $\{i, j\}$ on which $\min\{k_i, k_j\} < K$ or on which $K = 1$, and the geometric orientation on all other edges. Whether an edge is aligned depends only on K and the orders of its two particles. For $K = 1$ every edge is aligned.

The semi-discrete scheme is

$$\frac{\mathrm{d}U_i}{\mathrm{d}t} = \mathcal{L}_i(U) := -2 \sum_{j \in \nu(i)} c_{ij} \cdot \big(F^*_{ij} - F(U_i)\big), \qquad i \in I, \tag{4.4}$$

which equals $-2\sum_j c_{ij} \cdot F^*_{ij}$ whenever the weights have zero row sums (Lemma 3.3(ii)). The factor two accounts for the midpoint lying at half the distance: for smooth u and $g(y) = F(u(x_i + \frac{1}{2}y)) - F(u(x_i))$ one has $\partial_{y_d} g_d(0) = \frac{1}{2}\,\partial_d F_d(u)(x_i)$. Inserted into (4.4), the dissipation of neighbour j enters the update of i with the coefficient $s_{ij,d}\,\sigma_{ij,d} c^{(d)}_{ij}$, which is $s_{ij,d}|c^{(d)}_{ij}| \geq 0$ on aligned edges and may be negative on geometric ones.

4.3 Consistency

Lemma 4.1 (Symmetric interface fluxes)**.** *Assume that $k_{i\to j}$ depends only on the particle i and the unordered pair $\{i, j\}$ (true for all rules of Section 5.1), and let d be a direction with $\sigma_{ji,d} = -\sigma_{ij,d}$. Then $F^*_{ji,d} = F^*_{ij,d}$. The condition holds on geometric edges with $x_{ij,d} \neq 0$, and on aligned edges if $c^{(d)}_{ij}$ and $c^{(d)}_{ji}$ have strictly opposite signs, $c^{(d)}_{ij} c^{(d)}_{ji} < 0$.*

Proof. Write (4.3) as $F^*_{ij,d} = \Phi_d(U_{ij}, U_{ji}; \sigma_{ij,d})$ with $\Phi_d(a, b; \sigma) = \frac{1}{2}(F_d(a) + F_d(b)) - \frac{1}{2} s(a, b)\, \sigma\, (b - a)$ and a symmetric s. Particle j evaluates the same two states with roles exchanged and $\sigma_{ji,d} = -\sigma_{ij,d}$, and $\Phi_d(b, a; -\sigma) = \Phi_d(a, b; \sigma)$. Both particles use the same kind of orientation, since whether the edge is aligned depends on K, k_i and k_j only. If one of the two weights vanishes, $\operatorname{sign}(0) = 0$ and the condition fails unless both vanish, which is why the lemma asks for strictly opposite signs; this is a sufficient condition for $F^*_{ji,d} = F^*_{ij,d}$, not a necessary one. □

On a lattice with equal degrees, $c_{ji} = -c_{ij}$ by translation invariance and Lemma 3.7(i), so the aligned flux is symmetric there as well.

Proposition 4.2 (Exactness for linear problems)**.** *Let $F_d(U) = A_d U$ with constant matrices A_d, let u be a polynomial of degree at most $k - 1$, $U_l = u(x_l)$ for all particles, and let $\nu(i) \subset I$ and $k_i = k_j = k_{i\to j} = k_{j\to i} = k$ for all $j \in \nu(i)$. Assume that*

(a) the neighbourhoods of i and of its neighbours are Π_{k-1}-unisolvent and their reconstruction fits unregularized,

(b) the neighbourhood of i is Π_{q_i}-unisolvent and its derivative fit unregularized, with $q_i \geq k - 1$,

(c) no limiter is active.

Then $\mathcal{L}_i(U) = -\sum_d A_d\,\partial_d u(x_i)$, for either orientation. In particular, an exact solution that is polynomial of degree at most $k-1$ in x satisfies (4.4) at particle i.

Proof. By (a) and (c) the reconstructions reproduce u, so $U_{ij} = U_{ji} = u(m_{ij})$, the dissipation in (4.3) vanishes whatever σ is, and $F^*_{ij,d} = A_d\,u(m_{ij})$. With $g(y) = u(x_i + \frac{1}{2}y) - u(x_i) \in \Pi_{k-1} \subset \Pi_{q_i}$, (b) and Lemma 3.3(i) give $\mathcal{L}_i(U) = -2\sum_d A_d \sum_j c^{(d)}_{ij}\, g(x_{ij}) = -2\sum_d A_d\,\partial_{y_d} g(0) = -\sum_d A_d\,\partial_d u(x_i)$. □

Proposition 4.2 is the meshfree analogue of a $(k-1)$-exact reconstruction in FV methods [1], which is the sense in which k is the design order. For smooth, non-polynomial solutions the reconstruction errors at the midpoints are $O(h^k)$ and the weights $O(h^{-1})$ (Lemma 3.5), so that a truncation error bound of $O(h^{k-1})$ is all one gets on general clouds unless the errors cancel. On lattices they do.

Proposition 4.3 (Consistency order on lattices)**.** *Let the particles form a periodic Cartesian lattice of spacing h, so that all neighbourhoods are translates of one centrally symmetric, $\Pi_{\max(k-1,q)}$-unisolvent set of offsets, with unregularized fits. Let $k_i = k_{i\to j} = k$ and $q_i = q$ for all particles and edges, with q odd, and let no limiter be active. Let F be smooth and the wave speeds λ_d Lipschitz continuous, and let u be a smooth solution of (2.1) with $U_l = u(x_l)$ at a fixed time. Then, for the geometric and for the aligned Rusanov flux,*

$$\tau_i := \mathcal{L}_i(U) + \sum_{d=1}^{D} \partial_d F_d(u)(x_i) = O\big(h^{\min(k,\,q+1)}\big). \tag{4.5}$$

In particular $\tau_i = O(h^k)$ for $q = q(k)$.

Proof. Reconstruction errors. By translation invariance all particles use the same linear fit map Π, which reproduces Π_{k-1}. Hence $\mathcal{R}_i(y) - u(x_i + y) = \sum_{|\alpha|=k} \frac{\partial^\alpha u(x_i)}{\alpha!}\,\psi_\alpha(y) + O(h^{k+1})$ for $|y| \le R$, where $\psi_\alpha = \Pi[y^\alpha] - y^\alpha$ is homogeneous of degree k in h by the dilation argument of Lemma 3.3(iii). As in the proof of Lemma 3.7, central symmetry implies that Π commutes with $y \mapsto -y$, so $\psi_\alpha(-y) = (-1)^k\psi_\alpha(y)$. With $E(y) = \sum_{|\alpha|=k} \frac{\partial^\alpha u(x_i)}{\alpha!}\,\psi_\alpha(\frac{1}{2}y) = O(h^k)$ and $\partial^\alpha u(x_j) = \partial^\alpha u(x_i) + O(h)$,

$$U_{ij} = u(m_{ij}) + E(x_{ij}) + O(h^{k+1}), \qquad U_{ji} = u(m_{ij}) + (-1)^k E(x_{ij}) + O(h^{k+1}).$$

Orientation. By translation invariance $c^{(d)}_{ij} = \gamma_d(x_{ij})$ for a function γ_d on the set of offsets, which is odd by Lemma 3.7(i). On geometric edges $\sigma_{ij,d} = \operatorname{sign}(x_{ij,d})$, and on aligned edges (all edges if $k < K$ or $K = 1$) $\sigma_{ij,d} = \operatorname{sign}\gamma_d(x_{ij})$. In both cases $\sigma_{ij,d} = \varsigma_d(x_{ij})$ with a function ς_d that is odd on all offsets with $\gamma_d \neq 0$. For the aligned orientation this is the oddness of γ_d. For the geometric orientation $\operatorname{sign}(y_d)$ is odd except on offsets with $y_d = 0$, where the convention $\sigma_{ij,d} = \sigma_{ji,d} = -1$ is even; on these offsets γ_d vanishes by the reflection symmetry of the Cartesian stencil. Indeed, the set of offsets and the radial weights are invariant under the reflection $R_d\colon y_d \mapsto -y_d$, and with $P_d = \operatorname{diag}\big((-1)^{\alpha_d}\big)_{0\le|\alpha|\le q}$ one has $\tilde p_q(R_d y) = P_d\,\tilde p_q(y)$; the argument of the proof of Lemma 3.7(i) with P_d in place of P gives $\gamma_d(R_d y) = -\gamma_d(y)$, since $P_d e_{1+d} = -e_{1+d}$, and hence $\gamma_d(y) = 0$ whenever $y_d = 0$. Offsets with $\gamma_d(x_{ij}) = 0$ do not enter the update of i in direction d, so the value of ς_d there, including the convention for $x_{ij,d} = 0$, does not matter.

Fluxes. Expanding (4.3) about $u(m_{ij})$, with $s_{ij,d} = \lambda_d(u(m_{ij})) + O(h^k)$, $F'_d(u(m_{ij})) = F'_d(u(x_i)) + O(h)$ and the same for λ_d, gives $F^*_{ij,d} = F_d(u(m_{ij})) + G_d(x_{ij}) + O(h^{k+1})$ with

$$G_d(y) = \tfrac{1}{2}\big(1 + (-1)^k\big)F'_d(u(x_i))\,E(y) + \tfrac{1}{2}\big(1 - (-1)^k\big)\lambda_d(u(x_i))\,\varsigma_d(y)\,E(y).$$

For even k the first term is even in y and the second vanishes; for odd k the first vanishes and the second is the product of two odd functions. Thus $G_d(-y) = G_d(y)$ on all offsets with $\gamma_d(y) \neq 0$.

Summation. By Corollary 3.8, applied to the family $g_i(y) = F_d(u(x_i + \frac{1}{2}y)) - F_d(u(x_i))$, whose derivatives are bounded uniformly in i because u is smooth on the periodic domain, the exact midpoint fluxes give $-2\sum_j c_{ij} \cdot \big(F(u(m_{ij})) - F(u(x_i))\big) = -\sum_d \partial_d F_d(u)(x_i) + O(h^{q+1})$. The terms G_d drop out, $\sum_j c_{ij}^{(d)} G_d(x_{ij}) = 0$, because $c_{ij'}^{(d)} = -c_{ij}^{(d)}$ while $G_d(x_{ij'}) = G_d(x_{ij})$ wherever $c_{ij}^{(d)} \neq 0$. The remainders contribute $2\sum_j |c_{ij}|\, O(h^{k+1}) = O(h^k)$ by Lemma 3.5. □

With the even degree $q = k-1$ for odd k, the derivative error would limit the scheme to order $k-1$, which is a second reason for Definition 3.9. The frozen mode with $q(k_{\mathrm{div}}) < k-1$ is formally of order $q(k_{\mathrm{div}}) + 1$ only. On irregular clouds neither the parity of ψ_α nor the antisymmetry of the weights holds, and the bound loses one order.

4.4 Conservation

Proposition 4.4 (Conservation balance)**.** *Let $V_i > 0$ and let all derivative weights have zero row sums. For any orders, interface rules, divergence modes and orientations, the solution of (4.4) satisfies*

$$\begin{aligned}\frac{\mathrm{d}}{\mathrm{d}t}\sum_{i\in I} V_i U_i = &-2\sum_{i\in I}\sum_{j\in\nu(i)\setminus I} V_i\, c_{ij}\cdot F_{ij}^{*} \\ &-\sum_{\substack{\{i,j\}\subset I\\ j\in\nu(i)}}\sum_{d=1}^{D}\Big[d_{ij,d}\big(F_d(U_{ij}) + F_d(U_{ji})\big) - s_{ij,d}\, e_{ij,d}\big(U_{ji} - U_{ij}\big)\Big]\end{aligned} \tag{4.6}$$

with the pair defects $d_{ij,d} = V_i c_{ij}^{(d)} + V_j c_{ji}^{(d)}$ and $e_{ij,d} = V_i \sigma_{ij,d} c_{ij}^{(d)} - V_j \sigma_{ji,d} c_{ji}^{(d)}$. Moreover:

(i) $e_{ij,d} = \sigma_{ij,d}\, d_{ij,d}$, and hence $|e_{ij,d}| \le |d_{ij,d}|$, on geometric edges with $x_{ij,d} \neq 0$;

(ii) $e_{ij,d} = V_i|c_{ij}^{(d)}| - V_j|c_{ji}^{(d)}|$ and $|e_{ij,d}| \le |d_{ij,d}|$ on aligned edges;

(iii) on geometric edges with $x_{ij,d} = 0$ the convention $\sigma_{ij,d} = \sigma_{ji,d} = -1$ gives $e_{ij,d} = V_j c_{ji}^{(d)} - V_i c_{ij}^{(d)}$, which is not bounded by $|d_{ij,d}|$ in general.

Proof. Multiply (4.4) in the form $-2\sum_j c_{ij} \cdot F_{ij}^{*}$ by V_i and sum over $i \in I$. The terms with ghost neighbours give the first sum. Every unordered pair of interior neighbours appears twice in the remaining terms. Particle i contributes $-2V_i c_{ij}^{(d)} F_{ij,d}^{*} = -V_i c_{ij}^{(d)}(F_d(U_{ij}) + F_d(U_{ji})) + s_{ij,d} V_i \sigma_{ij,d} c_{ij}^{(d)}(U_{ji} - U_{ij})$, and particle j the same expression with i and j exchanged, which evaluates the same two states and the same symmetric $s_{ij,d}$; note that $U_{ij} - U_{ji} = -(U_{ji} - U_{ij})$. Their sum is the bracket in (4.6). (i) $\sigma_{ji,d} = -\sigma_{ij,d} \in \{\pm 1\}$ gives $e_{ij,d} = \sigma_{ij,d}(V_i c_{ij}^{(d)} + V_j c_{ji}^{(d)})$. (ii) $\sigma c = |c|$ on aligned edges, and $\big||a| - |b|\big| \le |a + b|$ for real a, b. (iii) Direct substitution. □

The second sum in (4.6) vanishes for all data, and the scheme is locally conservative, *if* the generalized face vectors $S_{ij} = 2V_i c_{ij}$ are antisymmetric, $S_{ij} = -S_{ji}$, and on geometric edges with $x_{ij,d} = 0$ in addition either $c_{ij}^{(d)} = 0$ or $\sigma_{ji,d} = -\sigma_{ij,d}$ (for instance by orienting such pairs by the particle index). Indeed, antisymmetry gives $d_{ij,d} = 0$, hence $e_{ij,d} = 0$ by (i) and (ii), and on the remaining edges under the additional condition either directly or by the computation of (i) with $\sigma_{ji,d} = -\sigma_{ij,d}$. We do not claim the converse. Together with the closure $\sum_j S_{ij} = 0$ of Lemma 3.3(ii), the meshfree counterpart of $\sum_e |e|\, n_e = 0$ for a closed control volume, (4.4) can always be written as $V_i\, \mathrm{d}U_i/\mathrm{d}t = -\sum_j S_{ij} \cdot F_{ij}^{*}$ by the definition of S_{ij}, and antisymmetry together with $F_{ji}^{*} = F_{ij}^{*}$ (Lemma 4.1) makes it a conservative FV-type update; compare the meshless face coefficients of [2, 3]. By (ii), the aligned orientation never makes the contribution of a pair larger than the geometric one with $x_{ij,d} \neq 0$. On centrally symmetric, translation-invariant stencils with equal volumes and equal degrees the weights are antisymmetric by Lemma 3.7(i);

on irregular clouds and for pairs of particles with different derivative degrees they are not. The frozen mode avoids pairs of different degrees.

5 The first-order scheme and positivity

The first-order scheme uses piecewise constant interface states, $U_{ij} = U_i$ and $U_{ji} = U_j$, combined with derivative weights of some degree.

Lemma 5.1 (cf. [13, 17])**.** *Let $U \in \mathcal{G}$ for the Euler equations (2.2), $d \in \{1, \dots, D\}$, $\sigma \in \{-1, 1\}$ and $s \geq |u_d| + c$. Then $W = U - \sigma F_d(U)/s \in \mathcal{G}$.*

Proof. Let $\theta = 1 - \sigma u_d/s \geq 1 - |u_d|/s > 0$. A direct computation gives $\rho_W = \theta\rho > 0$, $m_W = \theta\rho u - \sigma(p/s)\, e_d$, $E_W = \theta E - \sigma u_d p/s$ and

$$E_W - \frac{|m_W|^2}{2\rho_W} = \frac{\theta p}{\gamma - 1} - \frac{p^2}{2\theta\rho s^2}.$$

This is positive if and only if $(\theta s)^2 > (\gamma - 1)p/(2\rho) = \frac{\gamma-1}{2\gamma} c^2$, which holds because $\theta s = s - \sigma u_d \geq s - |u_d| \geq c$ and $(\gamma - 1)/(2\gamma) < 1$. □

Proposition 5.2 (Positivity of the first-order update)**.** *Let $\mathcal{G}$ be convex and assume that $U - \sigma F_d(U)/s \in \mathcal{G}$ for all $U \in \mathcal{G}$, $\sigma = \pm 1$ and $s \geq \lambda_d(U)$, $s > 0$ (for the Euler equations by Lemma 5.1). Let $i \in I$ use piecewise constant interface states on all its edges and derivative weights with*

(a) $\sum_{j \in \nu(i)} c_{ij} = 0$,

(b) $\sigma_{ij,d}\, c_{ij}^{(d)} = |c_{ij}^{(d)}|$ *for all* $j \in \nu(i)$ *and* $d = 1, \dots, D$, *i.e.* $\sigma_{ij,d}\, c_{ij}^{(d)} \geq 0$, *and* $\sigma_{ij,d} \neq 0$ *wherever* $c_{ij}^{(d)} \neq 0$,

let $U_l \in \mathcal{G}$ for all $l \in \bar{\nu}(i)$, let $s_{ij,d} > 0$ wherever $c_{ij}^{(d)} \neq 0$ (automatic for the Euler equations, where λ_d is at least the sound speed), and let $\tau > 0$ satisfy

$$\tau \sum_{j \in \nu(i)} \sum_{d=1}^{D} s_{ij,d}\, |c_{ij}^{(d)}| \leq 1. \tag{5.1}$$

Then $U_i + \tau\, \mathcal{L}_i(U) \in \mathcal{G}$ for the Rusanov flux (4.3).

Proof. With piecewise constant states, $F^*_{ij,d} - F_d(U_i) = \frac{1}{2}(F_d(U_j) - F_d(U_i)) - \frac{1}{2} s_{ij,d}\sigma_{ij,d}(U_j - U_i)$. By (a) the terms with $F_d(U_i)$ cancel. By (b), $\sigma_{ij,d} c_{ij}^{(d)} = |c_{ij}^{(d)}|$, and $c_{ij}^{(d)} = \sigma_{ij,d} |c_{ij}^{(d)}|$ (both sides vanish if $c_{ij}^{(d)} = 0$; otherwise (b) forces $\sigma_{ij,d} = \pm 1$, so $\sigma_{ij,d}^2 = 1$). Hence

$$U_i + \tau\, \mathcal{L}_i(U) = \Big(1 - \tau \sum_{j,d} s_{ij,d} |c_{ij}^{(d)}|\Big) U_i + \sum_{j,d} \tau\, s_{ij,d} |c_{ij}^{(d)}| \Big(U_j - \sigma_{ij,d} \frac{F_d(U_j)}{s_{ij,d}}\Big),$$

where terms with $c_{ij}^{(d)} = 0$ are omitted, so that $\sigma_{ij,d} = \pm 1$ in the remaining ones. By (5.1) the coefficients are non-negative and sum to one, and the states in the last sum lie in $\mathcal{G}$ because $s_{ij,d} \geq \lambda_d(U_j)$. Convexity of $\mathcal{G}$ concludes the proof. □

Corollary 5.3 (Aligned orientation)**.** *On the aligned edges of a particle i, condition (b) holds by construction, $\sigma_{ij,d} c_{ij}^{(d)} = |c_{ij}^{(d)}|$, provided that the signs are taken from the weights that i actually uses. Hence, with the aligned Rusanov flux, a particle i with $k_i < K$, or any particle if $K = 1$, that uses piecewise constant interface states on all its edges and unregularized derivative weights*

of any degree on a Π_{q_i}-unisolvent neighbourhood satisfies the conclusion of Proposition 5.2. For a scalar conservation law with $s_{ij,d} \geq \max|F_d'|$ between U_i and U_j, the same computation gives

$$U_i + \tau\,\mathcal{L}_i(U) = U_i + \tau \sum_{j,d} \big(s_{ij,d}|c_{ij}^{(d)}| - c_{ij}^{(d)} a_{ij,d}\big)(U_j - U_i), \qquad a_{ij,d} = \frac{F_d(U_j) - F_d(U_i)}{U_j - U_i},$$

with non-negative coefficients, so the update lies between the minimum and the maximum of U over $\bar{\nu}(i)$ if $2\tau \sum_{j,d} s_{ij,d}|c_{ij}^{(d)}| \leq 1$: the first-order scheme is local-extremum-diminishing for any zero-sum weights.

Proof. All edges of a particle with $k_i < K$, and all edges if $K = 1$, are aligned, so (b) holds. Condition (a) holds by Lemma 3.3(ii). In the scalar case, $F^*_{ij,d} - F_d(U_i) = \frac{1}{2} a_{ij,d}(U_j - U_i) - \frac{1}{2} s_{ij,d}\sigma_{ij,d}(U_j - U_i)$, and $|c_{ij}^{(d)} a_{ij,d}| \leq s_{ij,d}|c_{ij}^{(d)}|$ gives coefficients in $[0, 2s_{ij,d}|c_{ij}^{(d)}|]$. □

A particle of order $K \geq 2$ whose edges are not all aligned is not covered by Corollary 5.3, even if it uses piecewise constant states on all its edges (as an order-K neighbour of a lowered particle does under EPD2, Section 5.1): its edges to other order-K particles use the geometric orientation. This does not affect the fallback property below, which concerns particles at order one.

Remark 5.4 (On the assumptions). (i) With the geometric orientation, (b) requires $\operatorname{sign} c_{ij}^{(d)} = \operatorname{sign} x_{ij,d}$. For degree one on a Cartesian lattice, $c_{ij}^{(d)} = w_{ij} x_{ij,d} / \sum_l w_{il} x_{il,d}^2$, and (b) holds. On perturbed clouds (b) is violated by small weights of neighbours with small $|x_{ij,d}|$, about 3% of the entries for the clouds of [11]. For degree three, (b) fails already on lattices, since the outer weights have the opposite sign of their offsets, as in $(f_{-2} - 8f_{-1} + 8f_1 - f_2)/(12h)$.

(ii) The signs must be those of the weights that particle i actually uses, of degree q_i: in the frozen mode $q(k_{\mathrm{div}})$ at every order (degree one for $k_{\mathrm{div}} = 2$), in the coupled mode $q(k_i)$, and in either mode possibly lower where the safeguards of Remark 3.10 apply. There is no sign relation between weights of different degrees, so signs read from a fit of another degree guarantee nothing; Remark 5.5 describes how the implementation meets this requirement.

Remark 5.5 (The aligned signs in the implementation). In HyperCloudCore.jl 0.7.8 the flux loop of particle i computes, for every neighbour slot, the weight c_{ij} with which the flux difference of that slot is contracted, and takes the aligned orientation from the sign of this same weight. On the cached path the weight is the cached weight of the full degree of the divergence fit. A particle off the cached path, because the safeguards of Remark 3.10 lower its degree, because its order is lowered in the coupled mode, or because its cached factorization failed, assembles and factorizes its own moment matrix of degree q_i and solves the weight of every slot from it; with the aligned flux this per-slot evaluation is used whether or not the divergence is streamed. The orientation is therefore always taken from the weights in use, in the frozen and in the coupled mode, and the aligned orientation no longer depends on the cache. Version 0.7.7 read the signs from the cached full-degree weights in all cases. This was wrong for every particle in one dimension with $\kappa = 2.5$ and $q(k_{\mathrm{div}}) = 3$ (four neighbours, cached interpolating fit of degree three with outer weights of the opposite sign of their offsets, used degree two), and for particles without a cached factorization, which kept the geometric orientation even at order one.

5.1 Interface orders and the fallback property

When MOOD lowers some orders, an *edge polynomial degree* (EPD) rule decides which degree each side of an edge uses at the interface:

$$\begin{aligned} &\text{EPD0:} && k_{i\to j} = k_i, \\ &\text{EPD1:} && k_{i\to j} = k_{j\to i} = \min\{k_i, k_j\}, \\ &\text{EPD2:} && k_{i\to j} = k_{j\to i} = \min\{k_l : l \in \bar{\nu}(i) \cup \bar{\nu}(j)\}. \end{aligned} \tag{5.2}$$




Figure 1: Interface orders for one particle (filled red) lowered to order one in a high-order region, on a lattice patch with nearest-neighbour edges standing in for the neighbour graph. The half of an edge next to a particle shows the order with which this particle's reconstruction enters the interface: red for order one, grey for the full order.

EPD1 is the minimum rule of FV MOOD [4, 5]. Figure 1 illustrates the rules.

Proposition 5.6 (Fallback property)**.** *Let $i \in I$ have order $k_i = 1$. Under EPD1 and EPD2, $\mathcal{L}_i$ coincides with a first-order operator, i.e. (4.4) with piecewise constant interface states on all edges of i: with the derivative weights of degree q_i (Remark 3.10), i.e. degree one in the coupled mode and $q(k_{\mathrm{div}})$ in the frozen mode unless a safeguard lowers it. If $K = 1$, this holds under every rule. Under EPD0 with $K \geq 2$ it does not hold in general.*

Proof. Under EPD1, $k_{i\to j} = k_{j\to i} = \min\{1, k_j\} = 1$ for all $j \in \nu(i)$; under EPD2 the minimum runs over a set that contains i. For $K = 1$ all orders are one. In all cases (4.2) gives piecewise constant states, and the coupled mode gives $q_i = q(1) = 1$, the frozen mode $q_i = q(k_{\mathrm{div}})$. Under EPD0, $k_{j\to i} = k_j$ may exceed one, so that $U_{ji} \neq U_j$. □

The failure of EPD0 is consistent with FV MOOD [4], where the polynomial degree used at a face is the minimum of the degrees of the two adjacent cells, so that a cell decremented to the lowest degree is advanced by the first-order scheme (called the parachute scheme in later MOOD papers); EPD1 is this rule on the edges of the neighbour graph.

Combining Proposition 5.6 with Corollary 5.3 gives the local statement that the time integration needs.

Proposition 5.7 (Local admissibility at order one)**.** *Let $\mathcal{G}$ and the fluxes satisfy the hypotheses of Proposition 5.2. Let $i \in I$ have order $k_i = 1$, under EPD1 or EPD2, or under any rule if $K = 1$. Use the aligned Rusanov flux, in either divergence mode, with signs taken from the weights in use (Remark 5.5), and let the derivative fit of i be unregularized on a Π_{q_i}-unisolvent neighbourhood. Let $Y \in \mathcal{G}$ at all $l \in \bar{\nu}(i)$, ghost neighbours included, and let $\tau > 0$ satisfy (5.1) with $s_{ij,d} = \max\{\lambda_d(Y_i), \lambda_d(Y_j)\}$, where $s_{ij,d} > 0$ wherever $c_{ij}^{(d)} \neq 0$ (automatic for the Euler equations). Then $Y_i + \tau\, \mathcal{L}_i(Y) \in \mathcal{G}$.*

Proof. By Proposition 5.6, $\mathcal{L}_i$ is the first-order operator with the weights of degree q_i. Since $k_i = 1$, either $k_i < K$ or $K = 1$, so all edges of i are aligned and Corollary 5.3 gives (a) and (b) of Proposition 5.2, which yields the claim. □

6 Time integration and stage-wise MOOD

6.1 Explicit Runge–Kutta methods and their canonical Shu–Osher form

The semi-discrete system (4.4) is integrated with an explicit s-stage Runge–Kutta method in Butcher form,

$$Y_k = U^n + \Delta t \sum_{l<k} a_{kl}\, K_l, \quad K_l = \mathcal{L}(Y_l), \quad k = 1, \ldots, s, \qquad Y_{s+1} := U^{n+1} = U^n + \Delta t \sum_{l=1}^{s} b_l\, K_l, \tag{6.1}$$

Table 1: Canonical Shu–Osher form (6.3) of SSPRK(5,4) with $r = \mathcal{C} = 1.50818$, computed from the Butcher coefficients of [15] (rounded to four digits; vanishing entries left blank).

	v_k	α_{k1}	α_{k2}	α_{k3}	α_{k4}	α_{k5}
Y_2	0.4092	0.5908				
Y_3	0.4444		0.5556			
Y_4	0.6201			0.3799		
Y_5	0.1781				0.8219	
U^{n+1}	0.2380		0.2874		0.1337	0.3409

where the boundary operator sets the ghost states of every stage Y_k and of U^{n+1} from their interior states. Let $\hat{A} \in \mathbb{R}^{(s+1)\times(s+1)}$ be the strictly lower triangular matrix with the rows $(a_{k1}, \dots, a_{k,k-1}, 0, \dots, 0)$, $k \le s$, and the last row $(b_1, \dots, b_s, 0)$. Following [8, 12], for $r > 0$ let

$$\alpha^{(r)} = r\hat{A}\,(\mathrm{Id} + r\hat{A})^{-1}, \qquad v^{(r)} = (\mathrm{Id} - \alpha^{(r)})\,e, \qquad e = (1, \dots, 1)^{\mathsf{T}}. \tag{6.2}$$

Lemma 6.1 (Canonical Shu–Osher form)**.** *For every $r > 0$, $\alpha^{(r)}$ is strictly lower triangular, $v_k^{(r)} + \sum_j \alpha_{kj}^{(r)} = 1$ for all k, and the interior states of (6.1) satisfy*

$$Y_k = v_k^{(r)}\,U^n + \sum_{j<k} \alpha_{kj}^{(r)}\,W_j, \qquad W_j := Y_j + \frac{\Delta t}{r}\,\mathcal{L}(Y_j), \qquad k = 1, \dots, s+1. \tag{6.3}$$

Proof. Stack the interior stage values as $\mathbf{Y} = (Y_1, \dots, Y_{s+1})$ and set $\mathcal{L}(\mathbf{Y}) = (\mathcal{L}(Y_1), \dots, \mathcal{L}(Y_{s+1}))$; the last entry is multiplied by the zero last column of $\hat{A}$. Then (6.1) reads $\mathbf{Y} = e\,U^n + \Delta t\,\hat{A}\,\mathcal{L}(\mathbf{Y})$, hence $(\mathrm{Id} + r\hat{A})\mathbf{Y} = e\,U^n + r\hat{A}\big(\mathbf{Y} + \frac{\Delta t}{r}\mathcal{L}(\mathbf{Y})\big)$. The matrix $\mathrm{Id} + r\hat{A}$ is unit lower triangular and invertible, and $(\mathrm{Id} + r\hat{A})^{-1} = \mathrm{Id} - r\hat{A}(\mathrm{Id} + r\hat{A})^{-1} = \mathrm{Id} - \alpha^{(r)}$. Multiplying by it gives (6.3), with $v^{(r)} = (\mathrm{Id} - \alpha^{(r)})e$, i.e. $v^{(r)} + \alpha^{(r)}e = e$. The product of the strictly lower triangular $\hat{A}$ with the lower triangular $(\mathrm{Id} + r\hat{A})^{-1}$ is strictly lower triangular. □

The proof uses only the linear relation between the stages and the slopes, so Lemma 6.1 also holds with a different operator at every stage, $K_l = \mathcal{L}^{(l)}(Y_l)$ and $W_j = Y_j + \frac{\Delta t}{r}\mathcal{L}^{(j)}(Y_j)$; this is the situation of stage-wise MOOD, where the orders change from stage to stage.

The *SSP coefficient* $\mathcal{C}$ of the method is the largest r with $\alpha^{(r)} \ge 0$ and $v^{(r)} \ge 0$ entrywise ($\mathcal{C} = 0$ if there is none) [8, 12]. For $\mathcal{C} > 0$ we use $r = \mathcal{C}$ and write α, v; then every stage and the update are convex combinations of U^n and the forward Euler substeps W_j with step $\Delta t/\mathcal{C}$. Forward Euler and the SSP methods of Heun (RK2) and Shu–Osher (RK3) [7, 14] have $\mathcal{C} = 1$, and their canonical forms are the familiar ones, e.g. $Y_3 = \frac{3}{4}U^n + \frac{1}{4}W_2$ and $U^{n+1} = \frac{1}{3}U^n + \frac{2}{3}W_3$ for RK3. Ralston's two-stage method has $\mathcal{C} = \frac{1}{2}$ and the classical four-stage method $\mathcal{C} = 0$. For SSPRK(5,4) of Spiteri and Ruuth [15], $\mathcal{C} \approx 1.508$ and the canonical form is given in Table 1.

Remark 6.2 (Why stage values are not enough). For forward Euler, RK2 and RK3 every row k of (6.3) contains only U^n and the last substep W_{k-1}. If MOOD checks the stage value $Y_{k,i}$ instead of $W_{k-1,i}$, a particle that is still flagged at order one satisfies $W_{k-1,i} \in \mathcal{G}$ by Proposition 5.7, hence $Y_{k,i} \in \mathcal{G}$ by convexity, and the proof of Theorem 6.3 goes through with the stage-value check. For SSPRK(5,4) the update reuses W_2 and W_4, which also enter Y_3 and Y_5. An admissible $Y_3 = 0.4444\,U^n + 0.5556\,W_2$ does not imply an admissible W_2. Eliminating W_2 and W_4 in favour of the stage values gives

$$U^{n+1} = 0.5172\,Y_3 + 0.1627\,Y_5 + 0.3409\,W_5 - 0.0208\,U^n$$

(coefficients from the unrounded tableau; the rounded values of Table 1 give 0.5173 for Y_3), with a negative coefficient of U^n, so admissible U^n, Y_3, Y_5 and W_5 do not imply an admissible U^{n+1}. A MOOD procedure that checks the stage values therefore does not guarantee admissibility for SSPRK(5,4), even with a positive first-order fallback.

Algorithm 1 MOOD checkpoint k, $2 \le k \le s+1$, for a method with $\mathcal{C} > 0$ ($Y_{s+1} := U^{n+1}$).

Require: U^n, stages $Y_1, \dots, Y_{k-1}$, operators $K_1, \dots, K_{k-1}$, orders $(k_i)_{i \in I}$

$T \leftarrow I$ ▷ particles to check
loop
  $W_{k-1,i} \leftarrow Y_{k-1,i} + (\Delta t/\mathcal{C})\, K_{k-1,i}$ for $i \in I$ ▷ the checked state
  $Z \leftarrow \{i \in T : W_{k-1,i}$ is flagged with respect to $Y_{k-1}\}$
  $C \leftarrow \{i \in Z : k_i > 1\}$; $k_i \leftarrow k_i - 1$ for $i \in C$
  if $C = \emptyset$ **then break**
  end if
  $T \leftarrow \{i : i$ within ϱ hops of $C\}$ ▷ $\varrho = 2$ for EPD2, $\varrho = 1$ otherwise
  refit the reconstructions of Y_{k-1} on C; recompute $K_{k-1,i} = \mathcal{L}_i(Y_{k-1})$ for $i \in T$
end loop
form Y_k by (6.1), equivalently (6.3); apply the boundary operator

6.2 Stage-wise MOOD on the forward Euler substeps

MOOD works with the per-particle orders k_i, which are reset to K at the beginning of every time step (optionally before every stage). A state Z_i of an interior particle, computed from a previous state Y, is *flagged* if

- it fails *physical admissibility detection* (PAD), $Z_i \notin \mathcal{G}$ (including non-finite values), or
- it fails a *numerical admissibility detection*, for instance the relaxed discrete maximum principle (DMP) [4]

$$Y_{i,m}^{\max} - Y_{i,m}^{\min} \ge \delta^3 \quad \text{and} \quad Z_{i,m} \notin \left[Y_{i,m}^{\min} - \delta,\ Y_{i,m}^{\max} + \delta\right] \quad \text{for some component } m, \tag{6.4}$$

  with the extrema $Y_{i,m}^{\max}$, $Y_{i,m}^{\min}$ over $\bar{\nu}(i)$, possibly relaxed by overrides for plateaus and smooth extrema (the u2 criterion of the companion paper). Overrides never apply to PAD.

A flagged particle with $k_i > 1$ lowers its order by one; a flagged particle at order one keeps its order and its state.

For a method with $\mathcal{C} > 0$, checkpoint $k \in \{2, \dots, s+1\}$ checks the substep W_{k-1} against the previous stage Y_{k-1} (Algorithm 1). Within the checkpoint, the stages $Y_1, \dots, Y_{k-1}$, the operators $K_1, \dots, K_{k-2}$ and hence the substeps $W_1, \dots, W_{k-2}$ are fixed; a lowered order changes only $K_{k-1} = \mathcal{L}(Y_{k-1})$, and only near the lowered particle. For methods with $\mathcal{C} = 0$ (classical RK4) the implementation checks the stage value Y_k instead, for which no admissibility result is available.

A time step resets the orders, sets $Y_1 = U^n$, computes K_1, and runs Algorithm 1 for $k = 2, \dots, s+1$, computing K_k after each checkpoint with $k \le s$. The procedure has the following properties.

- *Termination.* Every pass that does not break lowers at least one order, and orders are bounded below by one.
- *Exact locality.* A lowered order at p changes the reconstruction of p, the divergence degree of p in the coupled mode, whether the edges of p are aligned, and the interface orders of the edges whose rule involves p. For EPD0 and EPD1 this affects $\mathcal{L}_l$ only for $l \in \bar{\nu}(p)$; for EPD2 for all l within two hops. Outside T, K_{k-1} and W_{k-1} are unchanged and need not be checked again.
- *No cascades.* A particle is lowered at most once per pass.

6.3 Admissibility

Theorem 6.3 (Admissibility of stage-wise MOOD)**.** *Let $\mathcal{G}$ and the fluxes satisfy the hypotheses of Proposition 5.2, and let the explicit Runge–Kutta method have $\mathcal{C} > 0$, with MOOD (PAD at least) applied by Algorithm 1. Assume:*

(H1) the interface rule is EPD1 or EPD2, or $K = 1$ with any rule;

(H2) the aligned Rusanov flux is used, in either divergence mode, with signs taken from the weights in use (Remark 5.5), and the derivative fits of all interior particles are unregularized on Π_{q_i}-unisolvent neighbourhoods;

(H3) the boundary operator maps admissible interior states to admissible ghost states, and $U^n \in \mathcal{G}$ at all particles;

(H4) at every checkpoint k, every particle i with $k_i = 1$ at the end of the checkpoint satisfies (5.1) with $\tau = \Delta t/\mathcal{C}$ and $s_{ij,d} = \max\{\lambda_d(Y_{k-1,i}), \lambda_d(Y_{k-1,j})\}$, and $s_{ij,d} > 0$ wherever $c_{ij}^{(d)} \neq 0$ (automatic for the Euler equations).

Then every checkpoint terminates with $W_{k-1,i} \in \mathcal{G}$ for all $i \in I$, and all stages $Y_2, \dots, Y_s$ and the update U^{n+1} are admissible at all particles, for every distribution of the orders that MOOD produces.

Proof. We show by induction over $k = 1, \dots, s+1$ that $Y_k \in \mathcal{G}$ at all particles and, for $k \geq 2$, that $W_{k-1,i} \in \mathcal{G}$ for all $i \in I$. For $k = 1$, $Y_1 = U^n$ is admissible by (H3).

Let $k \geq 2$ and assume the claim for all smaller indices. Checkpoint k terminates by the termination property. Consider $i \in I$ and the last pass in which $i \in T$. By exact locality, $K_{k-1,i}$ and $W_{k-1,i}$ have not changed since that pass, and $K_{k-1,i} = \mathcal{L}_i(Y_{k-1})$ with the final orders. Suppose $W_{k-1,i} \notin \mathcal{G}$. Then i was flagged in that pass. If $k_i > 1$ at that time, i was lowered, so $i \in C \subset T$ for the next pass, contradicting the choice of the pass. Hence $k_i = 1$ then, and also at termination, since orders do not increase within a checkpoint. By the induction hypothesis $Y_{k-1} \in \mathcal{G}$ on $\bar{\nu}(i)$, ghost neighbours included. By (H1), (H2) and (H4), Proposition 5.7 with $Y = Y_{k-1}$ and $\tau = \Delta t/\mathcal{C}$ gives $W_{k-1,i} = Y_{k-1,i} + (\Delta t/\mathcal{C})\mathcal{L}_i(Y_{k-1}) \in \mathcal{G}$, a contradiction. Thus $W_{k-1,i} \in \mathcal{G}$ for all $i \in I$.

The substeps $W_1, \dots, W_{k-2}$ are unchanged since their checkpoints and admissible by the induction hypothesis. By Lemma 6.1, the interior states $Y_{k,i} = v_k U_i^n + \sum_{j<k} \alpha_{kj} W_{j,i}$ are convex combinations of admissible states, since $v_k, \alpha_{kj} \geq 0$ sum to one, and convexity of $\mathcal{G}$ gives $Y_{k,i} \in \mathcal{G}$. The ghost states of Y_k are admissible by (H3). □

Corollary 6.4 (Order one everywhere)**.** *If all particles have order one at every stage, in particular if $K = 1$, then under (H2), (H3) and (5.1) with $\tau = \Delta t/\mathcal{C}$ and $s_{ij,d} = \max\{\lambda_d(Y_{k,i}), \lambda_d(Y_{k,j})\} > 0$ at every stage Y_k and every particle, every explicit Runge–Kutta method with $\mathcal{C} > 0$ preserves $\mathcal{G}$, under every interface rule and with or without MOOD.*

Proof. If all orders are one, all interface orders are one under every rule of (5.2), so every stage operator is the first-order operator with the weights of degree q_i, and all edges are aligned. By Corollary 5.3 and Proposition 5.2 with $\tau = \Delta t/\mathcal{C}$, every substep W_j is admissible, and Lemma 6.1 concludes as in the proof of Theorem 6.3. □

Corollary 6.4 is the classical SSP argument and covers only runs in which no particle has a higher order. Theorem 6.3 is the statement for mixed orders: whatever orders MOOD leaves, PAD at the substeps either accepts or lowers until the local fallback of Proposition 5.7 applies.

Remark 6.5 (On the hypotheses). (i) *EPD0* is not covered: a particle at order one may still receive high-order reconstructions of its neighbours (Proposition 5.6).

(ii) *Coupled mode.* The theorem covers the coupled mode as far as admissibility is concerned; its pairs of particles with different derivative degrees affect conservation only (Section 4.4).

(iii) *Time step.* (H4) is an assumption on the stage states. The time step of the implementation is $\Delta t = \mathrm{CFL} \cdot \min_i \left(D \sum_d \lambda_d(U_i^n) \sum_j |\hat{c}_{ij}^{(d)}| \right)^{-1}$, with weights $\hat{c}$ of a fit without constant term. If the wave speeds at the stages and at the neighbours do not exceed those at U_i^n, and $\hat{c}$ are the weights of the first-order operator, then $\mathrm{CFL} \le D\,\mathcal{C}$ implies (H4) up to the difference between the weights with and without constant term (at most 3.5% per particle for degree one and 10.6% for degree three on the clouds of [11]; the frozen mode with $k_{\mathrm{div}} = K \in \{3, 4\}$ uses degree-three weights). For SSPRK(5,4) this permits $\mathrm{CFL} \le 1.508\,D$; for RK2 and RK3, $\mathrm{CFL} \le D$.

(iv) *DMP.* Checking the substep against Y_{k-1} is also the natural form of the DMP: if no $W_{j,i}$ is flagged by (6.4) with $\delta = 0$ with respect to Y_j, i.e. $W_{j,i}$ lies componentwise in $[Y_{j,i}^{\min}, Y_{j,i}^{\max}]$, then $Y_{k,i}$ lies componentwise between the minimum and the maximum of U_i^n and of the local extrema of $Y_1, \dots, Y_{k-1}$ over $\bar{\nu}(i)$. With the stage-value check, the DMP test of a combination containing U^n against the bounds of Y_{k-1} is a heuristic.

(v) *Implementation.* HyperCloudCore.jl (since version 0.7.7) computes $\mathcal{C}$ of an explicit tableau by bisection on (6.2) and, for $\mathcal{C} > 0$, checks W_{k-1} as in Algorithm 1; ghost particles keep Y_{k-1} in W_{k-1} and are not checked. The stage-value check remains available as an option and is used for $\mathcal{C} = 0$ and for implicit–explicit methods.

7 Open questions

- *Stability and convergence.* Proposition 4.3 is a consistency result; stability of the high-order schemes, even for linear advection on lattices, is open, and so is the observed design-order convergence on perturbed clouds, where the truncation error bound loses one order.
- *Symmetry of the aligned flux.* On pairs with $c_{ij}^{(d)} c_{ji}^{(d)} \ge 0$ the two particles use different fluxes (Lemma 4.1), and the balance (4.6) accounts for this. A symmetric orientation, e.g. by $\mathrm{sign}(V_i c_{ij}^{(d)} - V_j c_{ji}^{(d)})$, keeps (b) only approximately.
- *Antisymmetric weights.* Weights that are antisymmetric after volume weighting and keep zero row sums would make the scheme locally conservative by Proposition 4.4, and with the aligned orientation they keep the positive fallback of Proposition 5.7.
- *EPD0.* See Remark 6.5(i).

A Changes with respect to the draft of 27 October 2026

The numbers in parentheses refer to the draft [9].

1. *Stage-wise MOOD* (Section 6.2, Corollary 6.1). MOOD now checks the forward Euler substeps W_{k-1} of the canonical Shu–Osher form instead of the stage values, as in HyperCloudCore.jl 0.7.7 (Algorithm 1). Remark 6.2 shows why the stage-value check is insufficient for SSPRK(5,4). The local result is Proposition 5.7, the mixed-order result Theorem 6.3; the former Corollary 6.1 is Corollary 6.4 and is stated as the all-order-one special case.
2. *Aligned orientation at $K = 1$* (Section 4.3, Corollary 5.3). The aligned orientation now applies on every edge with a particle below K *or* with $K = 1$, as in HyperCloudCore.jl 0.7.7, so that Proposition 5.2 covers the first-order scheme with degree-one weights on perturbed clouds. The "Hence" statement of Corollary 5.3 is restricted to particles with $k_i < K$ or $K = 1$, and the order-K neighbours of lowered particles under EPD2 are discussed after it.
3. *Source of the aligned signs* (Remark 5.4(v)). The signs are those of the weights in use (Remark 5.4(ii)), as implemented in HyperCloudCore.jl 0.7.8 (Remark 5.5); this also covers

the coupled mode and the safeguard-lowered degrees, so Proposition 5.7 and Theorem 6.3 hold in either divergence mode.

4. *Proposition 4.2.* Unisolvency and unregularized fits for the reconstruction and the derivative fit are explicit hypotheses (Proposition 4.2); Remark 3.2 explains why.
5. *Equation (3.5).* The bound $\sum_j |c_{ij}| = O(h^{-1})$ on general clouds now rests on Assumption 2.1 (Lemma 3.5), and (3.4) is proved as Corollary 3.8.
6. *Proposition 4.3.* The proof treats the aligned orientation for $k < K$ (and $K = 1$), whose signs are odd on lattices by Lemma 3.7(i) (Proposition 4.3).
7. *Lemma 4.1.* "Strictly opposite signs" (Lemma 4.1).
8. *Proposition 4.4.* The bound $|e| \le |d|$ excludes geometric edges with $x_{ij,d} = 0$ (Proposition 4.4(iii)); the local-conservation statement is an "if", with the additional condition on these edges.
9. *Lemma 3.6(i).* The step $\tilde{N}_i^{-1} P = P \tilde{N}_i^{-1}$ is written out (Lemma 3.7).
10. *StrictEPD0* is removed; it is a deprecated rule of earlier versions of the code.
11. *Condition (b) of Proposition 5.2* (new with respect to the draft). The condition $\sigma_{ij,d} c_{ij}^{(d)} \ge 0$ allowed $\sigma_{ij,d} = 0$ with $c_{ij}^{(d)} \ne 0$, for which the step $c_{ij}^{(d)} = \sigma_{ij,d} |c_{ij}^{(d)}|$ of the proof fails; it now reads $\sigma_{ij,d} c_{ij}^{(d)} = |c_{ij}^{(d)}|$. Both orientations of Section 4.2 satisfy the added requirement $\sigma_{ij,d} \ne 0$; the aligned orientation satisfies the whole condition, the geometric one not in general (Remark 5.4(i)) (found during the Lean formalization, Remark 1.2).
12. *New.* Remark 1.2 summarizes the Lean formalization. Remark 1.1 lists the remaining deviations between theory and implementation, under which the statements hold as approximate equalities.

Declaration of generative AI use

This note, its proofs and the Lean formalization described in Remark 1.2 were generated with Claude Fable 5.1 (Anthropic) in Claude Code, starting from the companion paper [11]. The note and the Lean code were reviewed by an independent Claude Opus 5.5 session without access to the generating conversation, which re-derived every proof by hand and rebuilt the Lean formalization. The author reviewed the results and takes full responsibility for the content.

References

[1] T. J. Barth and P. O. Frederickson. Higher order solution of the Euler equations on unstructured grids using quadratic reconstruction. In *28th Aerospace Sciences Meeting*, Reno, NV, 1990. American Institute of Aeronautics and Astronautics. doi: 10.2514/6.1990-13. AIAA Paper 90-0013.

[2] E. K.-Y. Chiu, Q. Wang, and A. Jameson. A conservative meshless scheme: general order formulation and application to Euler equations. In *49th AIAA Aerospace Sciences Meeting*, Orlando, FL, 2011. American Institute of Aeronautics and Astronautics. doi: 10.2514/6.2011-651. AIAA Paper 2011-651.

[3] E. K.-Y. Chiu, Q. Wang, R. Hu, and A. Jameson. A conservative mesh-free scheme and generalized framework for conservation laws. *SIAM J. Sci. Comput.*, 34(6):A2896–A2916, 2012. doi: 10.1137/110842740.

[4] S. Clain, S. Diot, and R. Loubère. A high-order finite volume method for systems of conservation laws—Multi-dimensional Optimal Order Detection (MOOD). *J. Comput. Phys.*, 230(10):4028–4050, 2011. doi: 10.1016/j.jcp.2011.02.026.

[5] S. Diot, S. Clain, and R. Loubère. Improved detection criteria for the Multi-dimensional Optimal Order Detection (MOOD) on unstructured meshes with very high-order polynomials. *Comput. Fluids*, 64:43–63, 2012. doi: 10.1016/j.compfluid.2012.05.004.

[6] S. Diot, R. Loubère, and S. Clain. The Multidimensional Optimal Order Detection method in the three-dimensional case: very high-order finite volume method for hyperbolic systems. *Int. J. Numer. Meth. Fluids*, 73(4):362–392, 2013. doi: 10.1002/fld.3804.

[7] S. Gottlieb, C.-W. Shu, and E. Tadmor. Strong stability-preserving high-order time discretization methods. *SIAM Rev.*, 43(1):89–112, 2001. doi: 10.1137/S003614450036757X.

[8] S. Gottlieb, D. I. Ketcheson, and C.-W. Shu. *Strong Stability Preserving Runge–Kutta and Multistep Time Discretizations*. World Scientific, Singapore, 2011. doi: 10.1142/7498.

[9] Pascal Jung. High-order meshfree MUSCL schemes with a posteriori order detection for hyperbolic systems on scattered point clouds, 2026. Draft of 27 October 2026.

[10] Pascal Jung. HyperCloudCore.jl: High-performance, physics-agnostic meshless solver backend for hyperbolic PDEs. https://github.com/industrial-mathematics-group/HyperCloudCore.jl, 2026. Version 0.7.8.

[11] Pascal Jung, Klaas Willems, and Axel Klar. High-order meshfree MUSCL schemes with a posteriori order detection for hyperbolic systems on scattered point clouds, 2026. Companion paper, in preparation.

[12] David I. Ketcheson. Highly efficient strong stability-preserving Runge–Kutta methods with low-storage implementations. *SIAM Journal on Scientific Computing*, 30(4):2113–2136, 2008. doi: 10.1137/07070485X.

[13] B. Perthame and C.-W. Shu. On positivity preserving finite volume schemes for Euler equations. *Numer. Math.*, 73(1):119–130, 1996. doi: 10.1007/s002110050187.

[14] C.-W. Shu and S. Osher. Efficient implementation of essentially non-oscillatory shock-capturing schemes. *J. Comput. Phys.*, 77(2):439–471, 1988. doi: 10.1016/0021-9991(88)90177-5.

[15] Raymond J. Spiteri and Steven J. Ruuth. A new class of optimal high-order strong-stability-preserving time discretization methods. *SIAM Journal on Numerical Analysis*, 40(2):469–491, 2002. doi: 10.1137/S0036142901389025.

[16] K. Willems, A. Klar, and G. Samaey. Higher-order meshless schemes for hyperbolic equations. *J. Sci. Comput.*, 105(3):90, 2025. doi: 10.1007/s10915-025-03124-4.

[17] X. Zhang and C.-W. Shu. On positivity-preserving high order discontinuous Galerkin schemes for compressible Euler equations on rectangular meshes. *J. Comput. Phys.*, 229(23):8918–8934, 2010. doi: 10.1016/j.jcp.2010.08.016.