

Single-layer painome · SL 6
75 genes, persistence 55; 38 genes recovered in multiplex sub-modules
(pie = neuron-type sub-module(s) each gene falls into; grey = none)



Multiplex · the same genes split into 12 neuron-type-specific sub-modules
(bubble = one HiDeF cluster, coloured nodes = SL 6 genes, grey dots = other members)



Note: a multiplex sub-module's spatial term largely reflects its neuron-type layer (both are DRG neuron classes) -- read it as layer identity, not independent evidence.

- A-PEP.CHRNA7_SLC18A3 — Cluster2-46 — Ad HTMRs (q=2.5e-13) — overlap 0.60, q=1.9e-03
- A-PEP.NTRK3_S100A16 — Cluster2-33 — Aß nociceptors (q=6.4e-22) — overlap 0.34, q=1.0e-03
- A-Propr.EPHA3 — Cluster3-76 — no spatial term — overlap 0.83, q=1.1e-03
- A-Propr.HAPLN4 — Cluster2-707 — no spatial term — overlap 1.00, q=3.1e-02
- A-Propr.PCDH8 — Cluster3-228 — Aß SA LTMRs (q=1.8e-02) — overlap 0.80, q=1.2e-02
- ATF3 — Cluster2-51 — PENK+ nociceptors (q=1.7e-10) — overlap 0.64, q=6.0e-06
- Ab-LTMR.CCKAR — Cluster2-77 — Putative C-LTMRs (q=1.4e-10) — overlap 0.88, q=8.8e-06
- Ab-LTMR.NSG2 — Cluster3-69 — Aß SA LTMRs (q=2.4e-08) — overlap 1.00, q=1.7e-05
- C-LTMR.CDH9 — Cluster5-3 — Cold nociceptors (q=1.7e-03) — overlap 1.00, q=2.8e-03
- C-NP.MRGPRX1_MRGPRX4 — Cluster3-56 — Putative C-LTMRs (q=3.6e-02) — overlap 1.00, q=1.4e-06
- C-NP.SST — Cluster4-43 — Pruritogen receptor enriched (q=1.7e-02) — overlap 0.71, q=3.2e-03
- C-PEP.TAC1_CHRNA3 — Cluster3-24 — Putative silent nociceptors (q=2.5e-14) — overlap 0.67, q=1.6e-08
- PPI edge (single-layer / within neuron type)
- cross-cell-type edge (multiplex)
- OPRM1: same gene, different neuron types
- KCNJ6: same gene, different neuron types