

## Question 1 (a)

$$\frac{d^2y}{dx^2} - 10\frac{dy}{dx} + 24y = 0$$

## Question 1 (a)

$$\frac{d^2y}{dx^2} - 10\frac{dy}{dx} + 24y = 0$$

$$\lambda^2 - 10\lambda + 24 = 0$$

## Question 1 (a)

$$\frac{d^2y}{dx^2} - 10\frac{dy}{dx} + 24y = 0$$

$$\lambda^2 - 10\lambda + 24 = 0$$

$$\implies (\lambda - 4)(\lambda - 6) = 0$$

## Question 1 (a)

$$\frac{d^2y}{dx^2} - 10\frac{dy}{dx} + 24y = 0$$

$$\lambda^2 - 10\lambda + 24 = 0$$

$$\implies (\lambda - 4)(\lambda - 6) = 0 \implies \lambda = 4, 6$$

## Question 1 (a)

$$\frac{d^2y}{dx^2} - 10\frac{dy}{dx} + 24y = 0$$

$$\lambda^2 - 10\lambda + 24 = 0$$

$$\implies (\lambda - 4)(\lambda - 6) = 0 \implies \lambda = 4, 6$$

$$y(x) = Ae^{4x} + Be^{6x}$$

## Question 1 (b)

$$\frac{d^2y}{dx^2} - 10\frac{dy}{dx} + 25y = 0$$

## Question 1 (b)

$$\frac{d^2y}{dx^2} - 10\frac{dy}{dx} + 25y = 0$$

$$\lambda^2 - 10\lambda + 25 = 0$$

## Question 1 (b)

$$\frac{d^2y}{dx^2} - 10\frac{dy}{dx} + 25y = 0$$

$$\lambda^2 - 10\lambda + 25 = 0$$

$$\implies (\lambda - 5)^2 = 0$$



## Question 1 (b)

$$\frac{d^2y}{dx^2} - 10\frac{dy}{dx} + 25y = 0$$

$$\lambda^2 - 10\lambda + 25 = 0$$

$$\implies (\lambda - 5)^2 = 0 \implies \lambda = 5 \text{ repeated root}$$

## Question 1 (b)

$$\frac{d^2y}{dx^2} - 10\frac{dy}{dx} + 25y = 0$$

$$\lambda^2 - 10\lambda + 25 = 0$$

$$\implies (\lambda - 5)^2 = 0 \implies \lambda = 5 \text{ repeated root}$$

$$y(x) = Ae^{5x} + Bxe^{5x}$$

## Question 1 (c)

$$\frac{d^2y}{dx^2} - 10\frac{dy}{dx} + 26y = 0$$

## Question 1 (c)

$$\frac{d^2y}{dx^2} - 10\frac{dy}{dx} + 26y = 0$$

$$\lambda^2 - 10\lambda + 26 = 0$$

## Question 1 (c)

$$\frac{d^2y}{dx^2} - 10\frac{dy}{dx} + 26y = 0$$

$$\lambda^2 - 10\lambda + 26 = 0$$

$$\Rightarrow \lambda = \frac{10 \pm \sqrt{100 - 4 \times 26}}{2}$$

## Question 1 (c)

$$\frac{d^2y}{dx^2} - 10\frac{dy}{dx} + 26y = 0$$

$$\lambda^2 - 10\lambda + 26 = 0$$

$$\Rightarrow \lambda = \frac{10 \pm \sqrt{100 - 4 \times 26}}{2} = \frac{10 \pm \sqrt{-4}}{2}$$

## Question 1 (c)

$$\frac{d^2y}{dx^2} - 10\frac{dy}{dx} + 26y = 0$$

$$\lambda^2 - 10\lambda + 26 = 0$$

$$\Rightarrow \lambda = \frac{10 \pm \sqrt{100 - 4 \times 26}}{2} = \frac{10 \pm \sqrt{-4}}{2} = 5 \pm i$$

## Question 1 (c)

$$\frac{d^2y}{dx^2} - 10\frac{dy}{dx} + 26y = 0$$

$$\lambda^2 - 10\lambda + 26 = 0$$

$$\Rightarrow \lambda = \frac{10 \pm \sqrt{100 - 4 \times 26}}{2} = \frac{10 \pm \sqrt{-4}}{2} = 5 \pm i$$

$$y(x) = e^{5x} (A \cos x + B \sin x)$$



## Question 2

$$\frac{d^2y}{dx^2} + 4y = 0$$

## Question 2

$$\frac{d^2y}{dx^2} + 4y = 0$$

$$\lambda^2 + 4 = 0$$

## Question 2

$$\frac{d^2y}{dx^2} + 4y = 0$$

$$\lambda^2 + 4 = 0 \quad \implies \quad \lambda^2 = -4$$

## Question 2

$$\frac{d^2y}{dx^2} + 4y = 0$$

$$\lambda^2 + 4 = 0 \quad \implies \quad \lambda^2 = -4 \quad \implies \quad \lambda = \pm 2i$$

## Question 2

$$\frac{d^2y}{dx^2} + 4y = 0$$

$$\lambda^2 + 4 = 0 \quad \implies \quad \lambda^2 = -4 \quad \implies \quad \lambda = \pm 2i$$

$$y = A \cos(2x) + B \sin(2x)$$

## Question 2

$$\frac{d^2y}{dx^2} + 4y = 0$$

$$\lambda^2 + 4 = 0 \quad \implies \quad \lambda^2 = -4 \quad \implies \quad \lambda = \pm 2i$$

$$y = A \cos(2x) + B \sin(2x)$$

$$\frac{d^2y}{dx^2} + 4y = 0$$

## Question 2

$$\frac{d^2y}{dx^2} + 4y = 0$$

$$\lambda^2 + 4 = 0 \quad \implies \quad \lambda^2 = -4 \quad \implies \quad \lambda = \pm 2i$$

$$y = A \cos(2x) + B \sin(2x)$$

$$\frac{d^2y}{dx^2} + 4y = 0 \quad \implies \quad \frac{d^3y}{dx^3} + 4\frac{dy}{dx} = 0$$

## Question 2

$$\frac{d^2y}{dx^2} + 4y = 0$$

$$\lambda^2 + 4 = 0 \quad \implies \quad \lambda^2 = -4 \quad \implies \quad \lambda = \pm 2i$$

$$y = A \cos(2x) + B \sin(2x)$$

$$\frac{d^2y}{dx^2} + 4y = 0 \quad \implies \quad \frac{d^3y}{dx^3} + 4\frac{dy}{dx} = 0 \quad \implies \quad \frac{d^4y}{dx^4} + 4\frac{d^2y}{dx^2} = 0$$



## Question 2

$$\frac{d^2y}{dx^2} + 4y = 0$$

$$\lambda^2 + 4 = 0 \quad \implies \quad \lambda^2 = -4 \quad \implies \quad \lambda = \pm 2i$$

$$y = A \cos(2x) + B \sin(2x)$$

$$\begin{aligned} \frac{d^2y}{dx^2} + 4y = 0 &\implies \frac{d^3y}{dx^3} + 4\frac{dy}{dx} = 0 \implies \frac{d^4y}{dx^4} + 4\frac{d^2y}{dx^2} = 0 \\ &\implies \frac{d^4y}{dx^4} - 16y = 0 \end{aligned}$$

## Question 2

$$\frac{d^2y}{dx^2} + 4y = 0$$

$$\lambda^2 + 4 = 0 \quad \implies \quad \lambda^2 = -4 \quad \implies \quad \lambda = \pm 2i$$

$$y = A \cos(2x) + B \sin(2x)$$

$$\begin{aligned} \frac{d^2y}{dx^2} + 4y = 0 &\implies \frac{d^3y}{dx^3} + 4\frac{dy}{dx} = 0 \implies \frac{d^4y}{dx^4} + 4\frac{d^2y}{dx^2} = 0 \\ &\implies \frac{d^4y}{dx^4} - 16y = 0 \implies \frac{d^5y}{dx^5} - 16\frac{dy}{dx} = 0 \\ &\implies \frac{d^6y}{dx^6} - 16\frac{d^2y}{dx^2} = 0 \end{aligned}$$

## Question 2

$$\frac{d^2y}{dx^2} + 4y = 0$$

$$\lambda^2 + 4 = 0 \quad \implies \quad \lambda^2 = -4 \quad \implies \quad \lambda = \pm 2i$$

$$y = A \cos(2x) + B \sin(2x)$$

$$\begin{aligned} \frac{d^2y}{dx^2} + 4y = 0 &\implies \frac{d^3y}{dx^3} + 4\frac{dy}{dx} = 0 \implies \frac{d^4y}{dx^4} + 4\frac{d^2y}{dx^2} = 0 \\ &\implies \frac{d^4y}{dx^4} - 16y = 0 \implies \frac{d^5y}{dx^5} - 16\frac{dy}{dx} = 0 \\ &\implies \frac{d^6y}{dx^6} - 16\frac{d^2y}{dx^2} = 0 \implies \frac{d^6y}{dx^6} + 64y = 0 \end{aligned}$$

### Question 3 (a)

$$\frac{d^3y}{dx^3} + 3\frac{d^2y}{dx^2} + 3\frac{dy}{dx} + y = 0$$

### Question 3 (a)

$$\frac{d^3y}{dx^3} + 3\frac{d^2y}{dx^2} + 3\frac{dy}{dx} + y = 0$$

$$\lambda^3 + 3\lambda^2 + 3\lambda + 1 = 0$$

### Question 3 (a)

$$\frac{d^3y}{dx^3} + 3\frac{d^2y}{dx^2} + 3\frac{dy}{dx} + y = 0$$

$$\lambda^3 + 3\lambda^2 + 3\lambda + 1 = 0$$

$$\implies (\lambda + 1)^3 = 0$$

### Question 3 (a)

$$\frac{d^3y}{dx^3} + 3\frac{d^2y}{dx^2} + 3\frac{dy}{dx} + y = 0$$

$$\lambda^3 + 3\lambda^2 + 3\lambda + 1 = 0$$

$$\implies (\lambda + 1)^3 = 0 \implies \lambda = -1 \text{ repeated root}$$

### Question 3 (a)

$$\frac{d^3y}{dx^3} + 3\frac{d^2y}{dx^2} + 3\frac{dy}{dx} + y = 0$$

$$\lambda^3 + 3\lambda^2 + 3\lambda + 1 = 0$$

$$\implies (\lambda + 1)^3 = 0 \implies \lambda = -1 \text{ repeated root}$$

Solutions are

$$y = e^{-x} \quad y = xe^{-x} \quad y = x^2e^{-x}$$



### Question 3 (a)

$$\frac{d^3y}{dx^3} + 3\frac{d^2y}{dx^2} + 3\frac{dy}{dx} + y = 0$$

$$\lambda^3 + 3\lambda^2 + 3\lambda + 1 = 0$$

$$\implies (\lambda + 1)^3 = 0 \implies \lambda = -1 \text{ repeated root}$$

Solutions are

$$y = e^{-x} \quad y = xe^{-x} \quad y = x^2e^{-x}$$

General solution

$$y = e^{-x} (A + Bx + Cx^2)$$

## Question 3 (b)

## Question 3 (b)

Auxiliary equation

$$(\lambda - 1)^4 = 0$$

## Question 3 (b)

Auxiliary equation

$$(\lambda - 1)^4 = 0$$

$$\implies \lambda^4 - 4\lambda^3 + 6\lambda^2 - 4\lambda + 1 = 0$$

## Question 3 (b)

Auxiliary equation

$$(\lambda - 1)^4 = 0$$

$$\implies \lambda^4 - 4\lambda^3 + 6\lambda^2 - 4\lambda + 1 = 0$$

So a suitable differential equation is

$$\frac{d^4 y}{dx^4} - 4\frac{d^3 y}{dx^3} + 6\frac{d^2 y}{dx^2} - 4\frac{dy}{dx} + y = 0$$