



# Some bounds on the zero forcing number of a graph

Michael Gentner, Dieter Rautenbach \*

Institut für Optimierung und Operations Research, Universität Ulm, Ulm, Germany

## ARTICLE INFO

### Article history:

Received 31 August 2016

Received in revised form 2 November 2017

Accepted 15 November 2017

Available online 6 December 2017

### Keywords:

Zero forcing

## ABSTRACT

A set  $Z$  of vertices of a graph  $G$  is a zero forcing set of  $G$  if initially labeling all vertices in  $Z$  with 0 and all remaining vertices of  $G$  with 1, and then, iteratively and as long as possible, changing the label of some vertex  $u$  from 1 to 0 if  $u$  is the only neighbor with label 1 of some vertex with label 0, results in all vertices of  $G$  having label 0. The zero forcing number  $Z(G)$ , defined as the minimum order of a zero forcing set of  $G$ , was proposed as an upper bound of the corank of matrices associated with  $G$ , and was also considered in connection with quantum physics and logic circuits. In view of the computational hardness of the zero forcing number, upper and lower bounds are of interest.

Refining results of Amos, Caro, Davila, and Pepper, we show that  $Z(G) \leq \frac{\Delta-2}{\Delta-1}n$  for a connected graph  $G$  of order  $n$  and maximum degree  $\Delta$  at least 3 if and only if  $G$  does not belong to  $\{K_{\Delta+1}, K_{\Delta,\Delta}, K_{\Delta-1,\Delta}, G_1, G_2\}$ , where  $G_1$  and  $G_2$  are two specific graphs of orders 5 and 7, respectively. For a connected graph  $G$  of order  $n$ , maximum degree 3, and girth at least 5, we show  $Z(G) \leq \frac{n}{2} - \Omega\left(\frac{n}{\log n}\right)$ . Using a probabilistic argument, we show  $Z(G) \leq \left(1 - \frac{H_r}{r} + o\left(\frac{H_r}{r}\right)\right)n$  for an  $r$ -regular graph  $G$  of order  $n$  and girth at least 5, where  $H_r$  is the  $r$ th harmonic number. Finally, we show that  $Z(G) \geq (g-2)(\delta-2)+2$  for a graph  $G$  of girth  $g \in \{5, 6\}$  and minimum degree  $\delta$ , which partially confirms a conjecture of Davila and Kenter.

© 2017 Elsevier B.V. All rights reserved.

## 1. Introduction

We consider graphs that are finite, simple, and undirected, and use standard terminology.

Let  $G$  be a graph. For a vertex  $u$  of  $G$ , let  $N_G(u)$  denote the neighborhood  $\{v \in V(G) : uv \in E(G)\}$  of  $u$  in  $G$ , and let  $N_G[u]$  denote the closed neighborhood  $\{u\} \cup N_G(u)$  of  $u$  in  $G$ .

For a set  $Z$  of vertices of  $G$ , let  $\mathcal{F}(Z)$  be the maximal set of vertices of  $G$  that arises from  $Z$  by iteratively adding vertices that are the unique neighbor outside the current set of some vertex inside the current set. Equivalently,

- $|N_G(w) \setminus \mathcal{F}(Z)| \neq 1$  for every vertex  $w$  in  $\mathcal{F}(Z)$ , and,
- the elements of  $\mathcal{F}(Z) \setminus Z$  have a linear order  $u_1, \dots, u_k$  such that for every index  $i$  in  $\{1, \dots, k\}$ , there is some vertex  $v_i$  in  $Z \cup \{u_j : 1 \leq j \leq i-1\}$  such that  $u_i$  is the only neighbor of  $v_i$  in  $\{u_j : i \leq j \leq k\}$ .

In the latter case, we say that  $v_i$  forces  $u_i$  for  $i \in \{1, \dots, k\}$ , and denote this by  $v_i \rightarrow u_i$ . The sequence  $v_1 \rightarrow u_1, v_2 \rightarrow u_2, \dots, v_k \rightarrow u_k$  is called a forcing sequence for  $Z$ .

The set  $Z$  is a zero forcing set of  $G$  if  $\mathcal{F}(Z)$  equals the vertex set  $V(G)$  of  $G$ . The zero forcing number  $Z(G)$  of  $G$  is the minimum order of a zero forcing set of  $G$ . The zero forcing number was proposed by the AIM Minimum Rank - Special Graphs Work

\* Corresponding author.

E-mail addresses: [michael.gentner@uni-ulm.de](mailto:michael.gentner@uni-ulm.de) (M. Gentner), [dieter.rautenbach@uni-ulm.de](mailto:dieter.rautenbach@uni-ulm.de) (D. Rautenbach).

Group [3,17] as an upper bound on the corank of matrices associated with a given graph. Independently, it was considered in connection with quantum physics [5,7,22] as well as logic circuits [6]. It has already been studied in a number of papers [2,10–12,16,19–21,23] and is computationally hard [1,13].

In the present paper we establish some upper and lower bounds on the zero forcing number.

For a connected graph  $G$  of order  $n$  and maximum degree  $\Delta$  at least 2, Amos et al. [2] prove

$$Z(G) \leq \frac{\Delta}{\Delta+1}n \text{ and} \quad (1)$$

$$Z(G) \leq \frac{\Delta-2}{\Delta-1}n + \frac{2}{\Delta-1}. \quad (2)$$

It was shown that the only extremal graph for (1) is the complete graph  $K_{\Delta+1}$  of order  $\Delta+1$  [14], and that the only extremal graphs for (2) are  $K_{\Delta+1}$ , the complete bipartite graph  $K_{\Delta,\Delta}$  with partite sets of order  $\Delta$ , and the cycle  $C_n$  [14,18].

We characterize the graphs for which the additive term  $\frac{2}{\Delta-1}$  in (2) is not needed. In fact, we believe that (2) can be improved considerably, and, in particular, pose the following conjecture.

**Conjecture 1.** *There are positive reals  $\alpha$  and  $\beta$  with  $\alpha < 1/2$  such that  $Z(G) \leq \alpha n + \beta$  for every connected graph  $G$  of order  $n$  and maximum degree 3.*

As a contribution towards this conjecture, we prove  $Z(G) \leq \frac{n}{2} - \Omega\left(\frac{n}{\log n}\right)$  for a connected graph  $G$  of order  $n$ , maximum degree 3, and girth at least 5. Girão, Mészáros, and Smith [15] constructed graphs that imply that  $\alpha$  is at least  $4/9$ . We present a probabilistic upper bound on the zero forcing number and discuss some of its consequences.

In [11] Davila and Kenter conjecture the lower bound

$$Z(G) \geq (g-2)(\delta-2) + 2 \quad (3)$$

for every graph  $G$  of girth  $g$  at least 3 and minimum degree  $\delta$  at least 2. They observe that for  $g > 6$  and sufficiently large  $\delta$ , the conjecture follows by combining results from [4] and [9]. For  $g = 4$ , that is, for triangle-free graphs, it was shown in [14]. Here, we prove the conjecture for  $g \in \{5, 6\}$ .

## 2. Results

We begin with a simple consequence of (2).

**Proposition 2.** *If  $G$  is a connected graph of order  $n$  and maximum degree  $\Delta$  at least 3 that is distinct from  $K_{\Delta+1}$ , then*

$$Z(G) \leq \frac{\Delta-1}{\Delta}n.$$

**Proof.** If  $n \geq 2\Delta$ , then  $Z(G) \stackrel{(2)}{\leq} \frac{(\Delta-2)n+2}{\Delta-1} \leq \frac{(\Delta-1)n}{\Delta}$ . Now, let  $n < 2\Delta$ . Since  $G$  is not complete, it contains an induced path  $uvw$  of order 3. Since the set  $V(G) \setminus \{v, w\}$  is a zero-forcing set of  $G$ , we obtain  $Z(G) \leq n-2 \leq \frac{(\Delta-1)n}{\Delta}$ , which completes the proof.  $\square$

Our next goal is to characterize the graphs for which the additive term in (2) is not needed.

The following lemma is implicit in the greedy argument in [8].

**Lemma 3.** *Let  $G$  be a connected graph of order  $n$  and maximum degree  $\Delta$  at least 3.*

*If there is some set  $Z_0$  of vertices of  $G$  such that  $|\mathcal{F}(Z_0)| \geq \frac{\Delta-1}{\Delta-2}|Z_0|$ , and  $\mathcal{F}(Z_0)$  induces a subgraph of  $G$  without isolated vertices, then  $Z(G) \leq \frac{\Delta-2}{\Delta-1}n$ .*

**Proof.** If  $\mathcal{F}(Z_0) = V(G)$ , then  $Z_0$  is a zero forcing set of  $G$ , and, hence,  $Z(G) \leq |Z_0| \leq \frac{\Delta-2}{\Delta-1}|\mathcal{F}(Z_0)| = \frac{\Delta-2}{\Delta-1}n$ . Therefore, we may assume that  $Z_i$  is a non-empty set of vertices of  $G$  for some non-negative integer  $i$  such that  $|\mathcal{F}(Z_i)| \geq \frac{\Delta-1}{\Delta-2}|Z_i|$ , the set  $\mathcal{F}(Z_i)$  induces a subgraph of  $G$  without isolated vertices, and  $\mathcal{F}(Z_i)$  is a proper subset of  $V(G)$ .

Because  $G$  is connected, there is a vertex  $v$  in  $\mathcal{F}(Z_i)$  that has at least one neighbor in  $V(G) \setminus \mathcal{F}(Z_i)$  as well as at least one neighbor in  $\mathcal{F}(Z_i)$ . Let  $Z_{i+1}$  arise from  $Z_i$  by adding to  $Z_i$  all but exactly one neighbor of  $v$  in  $V(G) \setminus \mathcal{F}(Z_i)$ , that is,  $|Z_{i+1}| = |Z_i| + p - 1$ , where  $p = |N_G(v) \setminus \mathcal{F}(Z_i)|$ . Since  $N_G(v) \subseteq \mathcal{F}(Z_{i+1})$ , we obtain

$$|\mathcal{F}(Z_{i+1})| \geq |\mathcal{F}(Z_i)| + |N_G(v) \setminus \mathcal{F}(Z_i)| = |\mathcal{F}(Z_i)| + p.$$

By the choice of  $v$ , we have  $1 \leq p \leq \Delta - 1$ , which implies  $p(\Delta - 2) \geq (p - 1)(\Delta - 1)$ . Adding this last inequality to the assumption  $|\mathcal{F}(Z_i)|(\Delta - 2) \geq |Z_i|(\Delta - 1)$  yields

$$(|\mathcal{F}(Z_i)| + p)(\Delta - 2) \geq (|Z_i| + p - 1)(\Delta - 1).$$



Fig. 1. The two specific graphs  $G_1$  and  $G_2$ .

Fig. 2. A section of  $C$ . The vertices in  $\{v_1, v_4, v_6, v_7\}$  belong to  $Z_0$  because the vertices in  $\{v_g, v_3, v_5, v_6\}$  have degree 3. The vertex  $v_g$  belongs to  $Z_0$  regardless of the degree of  $v_{g-1}$ . Note that  $v_1 \rightarrow v_2, v_2 \rightarrow v_3, v_3 \rightarrow v_4, v_4 \rightarrow v_5, v_5 \rightarrow v_6, v_6 \rightarrow v_7, v_7 \rightarrow v_8, v_8 \rightarrow v_9, v_9 \rightarrow v_{10}$ , and  $v_{10} \rightarrow v_g$ .

Now, we obtain

$$\frac{|\mathcal{F}(Z_{i+1})|}{|Z_{i+1}|} = \frac{|\mathcal{F}(Z_i)|}{|Z_i| + p - 1} \geq \frac{|\mathcal{F}(Z_i)| + p}{|Z_i| + p - 1} \geq \frac{\Delta - 1}{\Delta - 2},$$

and, hence,  $|\mathcal{F}(Z_{i+1})| \geq \frac{\Delta-1}{\Delta-2}|Z_{i+1}|$ . Furthermore, by construction, the set  $\mathcal{F}(Z_{i+1})$  induces a subgraph of  $G$  without isolated vertices.

Since  $|\mathcal{F}(Z_{i+1})|$  is strictly larger than  $|\mathcal{F}(Z_i)|$ , repeating this extension as long as  $\mathcal{F}(Z_i)$  is a proper subset of  $V(G)$ , we obtain, after finitely many extensions, a zero forcing set  $Z$  of  $G$  with  $|Z| \leq \frac{\Delta-2}{\Delta-1}|\mathcal{F}(Z)| = \frac{\Delta-2}{\Delta-1}n$ , which completes the proof.  $\square$

**Theorem 4.** If  $G$  is a connected graph of order  $n$  and maximum degree  $\Delta$  at least 3, then

$$Z(G) \leq \frac{\Delta - 2}{\Delta - 1}n \quad (4)$$

if and only if  $G \notin \{K_{\Delta+1}, K_{\Delta,\Delta}, K_{\Delta-1,\Delta}, G_1, G_2\}$ , where  $G_1$  and  $G_2$  are the two specific graphs illustrated in Fig. 1.

**Proof.** The necessity follows easily using  $Z(K_{\Delta+1}) = \Delta, Z(K_{\Delta,\Delta}) = 2\Delta - 2, Z(K_{\Delta-1,\Delta}) = 2\Delta - 3, Z(G_1) = 3$ , and  $Z(G_2) = 5$ . We proceed to the proof of the sufficiency. Therefore, let  $G \notin \{K_{\Delta+1}, K_{\Delta,\Delta}, K_{\Delta-1,\Delta}, G_1, G_2\}$  be as in the statement. In order to derive (4) using Lemma 3, it suffices to exhibit a set  $Z_0$  of vertices of  $G$  such that

$$\frac{|\mathcal{F}(Z_0)|}{|Z_0|} \geq \frac{\Delta - 1}{\Delta - 2}, \text{ and } \mathcal{F}(Z_0) \text{ induces a subgraph of } G \text{ without isolated vertices.} \quad (5)$$

Therefore, suppose that such a set does not exist.

If  $G$  has a vertex  $v$  of degree  $d_G(v)$  at most  $\Delta - 2$ , and  $u$  is a neighbor of  $v$ , then let  $Z_0 = N_G[v] \setminus \{u\}$ . Since  $|Z_0| = d_G(v)$  and  $u \in \mathcal{F}(Z_0)$ , we obtain  $\frac{|\mathcal{F}(Z_0)|}{|Z_0|} \geq \frac{d_G(v)+1}{d_G(v)} \geq \frac{\Delta-1}{\Delta-2}$ , that is, the set  $Z_0$  satisfies (5), which is a contradiction. Hence, we may assume that  $G$  has minimum degree at least  $\Delta - 1$ .

Since  $\Delta - 1 \geq 2$ , the graph  $G$  is not a tree. Let  $C : v_1 \dots v_g v_1$  be a shortest cycle in  $G$ . We consider three cases depending on the girth  $g$  of  $G$ .

**Case 1**  $g \geq 5$ .

Since  $G$  has girth at least 5, no vertex in  $V(G) \setminus V(C)$  has more than one neighbor on  $C$ . If all vertices on  $C$  have degree at least 3, then let  $u_i$  be a neighbor of  $v_i$  in  $V(G) \setminus V(C)$  for every  $i \in \{1, \dots, g\}$ . Let  $Z_0 = \bigcup_{i=1}^g N_G[v_i] \setminus \{u_i\}$ . Since  $|Z_0| \leq (\Delta - 2)g$  and  $u_1, \dots, u_g \in \mathcal{F}(Z_0)$ , we obtain  $\frac{|\mathcal{F}(Z_0)|}{|Z_0|} \geq \frac{|Z_0|+g}{|Z_0|} \geq \frac{\Delta-1}{\Delta-2}$ , that is, the set  $Z_0$  satisfies (5), which is a contradiction. Hence, we may assume that  $C$  contains a vertex of degree 2. Since  $G$  has minimum degree at least  $\Delta - 1 \geq 2$ , this implies  $\Delta = 3$ .

Let  $1 \leq i_1 < i_2 < \dots < i_p \leq g$  be such that  $\{v_{i_j} : 1 \leq j \leq p\}$  is the set of vertices of degree 3 on  $C$ . Since  $G$  is connected and has maximum degree 3, we obtain that  $p$  is at least 1. Possibly renaming vertices, we may assume that  $i_p = g$ . Similarly as above, for  $j \in \{1, \dots, p\}$ , let  $u_{i_j}$  be the neighbor of  $v_{i_j}$  in  $V(G) \setminus V(C)$ .

If  $p \leq g - 2$ , then let  $Z_0 = \{v_g\} \cup \{v_{i_j+1} : 1 \leq j \leq p\}$ , where  $v_{g+1} = v_1$ . See Fig. 2 for an illustration.

Since  $|Z_0| \leq p + 1$  and  $V(C) \cup \{u_{i_j} : 1 \leq j \leq p\} \subseteq \mathcal{F}(Z_0)$ , we obtain  $\frac{|\mathcal{F}(Z_0)|}{|Z_0|} \geq \frac{g+p}{p+1} \geq 2 = \frac{\Delta-1}{\Delta-2}$ , that is, the set  $Z_0$  satisfies (5), which is a contradiction. Hence, we may assume that  $p = g - 1$ , that is,  $C$  contains exactly one vertex, say  $v_1$ , of degree

2. Let  $Z_0 = V(C) \setminus \{v_2\}$ . Since  $|Z_0| = g - 1$  and  $V(C) \cup \{u_{ij} : 1 \leq j \leq p\} \subseteq \mathcal{F}(Z_0)$ , we obtain  $\frac{|\mathcal{F}(Z_0)|}{|Z_0|} \geq \frac{2g-1}{g-1} \geq 2 = \frac{\Delta-1}{\Delta-2}$ , that is, the set  $Z_0$  satisfies (5), which is a contradiction. This completes the proof in this case.

### Case 2 $g = 4$ .

First, we assume that  $d_G(v_1) = 2$ . As noted above, this implies  $\Delta = 3$ . If  $d_G(v_2) = 2$ , then  $Z_0 = \{v_1, v_2\}$  satisfies (5), which is a contradiction. Hence, by symmetry, we may assume that  $d_G(v_2) = d_G(v_4) = 3$ . Let  $Z_0 = \{v_1, v_2, v_3\}$ . If  $v_1$  and  $v_3$  are the only common neighbors of  $v_2$  and  $v_4$ , then  $\mathcal{F}(Z_0)$  contains  $v_4$  as well as the two neighbors of  $v_2$  and  $v_4$  that do not lie on  $C$ . Hence,  $|\mathcal{F}(Z_0)| \geq 6$ , that is, the set  $Z_0$  satisfies (5), which is a contradiction. Hence, we may assume that  $N_G(v_2) = N_G(v_4)$ . Since  $G \neq K_{2,3}$ , we may assume, by symmetry, that  $d_G(v_3) = 3$ . Since  $\mathcal{F}(Z_0)$  contains  $N_G[v_1] \cup N_G[v_2]$  and the neighbor of  $v_3$  that does not lie on  $C$ , we obtain  $|\mathcal{F}(Z_0)| \geq 6$ , that is, the set  $Z_0$  satisfies (5), which is a contradiction. Hence, we may assume, by symmetry, that  $G$  contains no cycle of length 4 that contains a vertex of degree 2. Since  $C$  is a shortest cycle, it is induced. For  $i \in \{1, 2, 3, 4\}$ , let  $u_i$  be a neighbor of  $v_i$  that does not lie on  $C$ .

Next, we assume that  $N_G(v_1) \not\subseteq N_G(v_3)$  and  $N_G(v_2) \not\subseteq N_G(v_4)$ . We may assume that  $u_1 \in N_G(v_1) \setminus N_G(v_3)$  and  $u_2 \in N_G(v_2) \setminus N_G(v_4)$ , which implies that  $u_1, u_2, u_3$ , and  $u_4$  are four distinct vertices. Let  $Z_0 = (N_G[v_1] \cup N_G[v_2] \cup N_G[v_3] \cup N_G[v_4]) \setminus \{u_1, u_2, u_3, u_4\}$ . Clearly,  $|Z_0| \leq 4(\Delta - 2)$ . Since  $v_1 \rightarrow u_1, v_2 \rightarrow u_2, v_3 \rightarrow u_3$ , and  $v_4 \rightarrow u_4$ , we obtain  $u_1, u_2, u_3, u_4 \in \mathcal{F}(Z_0)$ , and, hence,  $\frac{|\mathcal{F}(Z_0)|}{|Z_0|} \geq \frac{|Z_0|+4}{|Z_0|} \geq \frac{4(\Delta-2)+4}{4(\Delta-2)} = \frac{\Delta-1}{\Delta-2}$ , that is, the set  $Z_0$  satisfies (5), which is a contradiction. Hence, we may assume, by symmetry, that  $N_G(v_2) = N_G(v_4)$ .

Next, we assume that  $N_G(v_1) \not\subseteq N_G(v_3)$ . Again, let  $u_1 \in N_G(v_1) \setminus N_G(v_3)$ , and note that  $u_1, u_2$ , and  $u_3$  are distinct. If  $|N_G(v_1) \cup N_G(v_3)| \leq 2\Delta - 3$ , then let  $Z_0 = (N_G[v_1] \cup N_G[v_2] \cup N_G[v_3]) \setminus \{u_1, u_2, u_3\}$ . We obtain  $|Z_0| \leq |N_G(v_1) \cup N_G(v_3)| + |N_G(v_2)| - |\{u_1, u_2, u_3\}| \leq 2\Delta - 3 + \Delta - 3 = 3(\Delta - 2)$ . Since  $v_1 \rightarrow u_1, v_2 \rightarrow u_2$ , and  $v_3 \rightarrow u_3$ , we obtain  $u_1, u_2, u_3 \in \mathcal{F}(Z_0)$ , and, hence,  $\frac{|\mathcal{F}(Z_0)|}{|Z_0|} \geq \frac{|Z_0|+3}{|Z_0|} \geq \frac{3(\Delta-2)+3}{3(\Delta-2)} = \frac{\Delta-1}{\Delta-2}$ , that is, the set  $Z_0$  satisfies (5), which is a contradiction.

Hence, we may assume  $|N_G(v_1) \cup N_G(v_3)| \geq 2\Delta - 2$ , which implies that  $v_1$  and  $v_3$  both have degree  $\Delta$ , and do not have a common neighbor apart from  $v_2$  and  $v_4$ . By symmetry, this implies that every vertex in  $N_G(v_2)$  has degree  $\Delta$ , and that every two vertices in  $N_G(v_2)$  do not have a common neighbor apart from  $v_2$  and  $v_4$ . Let  $w_2 \in N_G(v_2) \setminus \{v_2, v_4\}$ , and let  $Z_0 = (N_G[v_1] \cup N_G[v_2] \cup N_G[v_3] \cup N_G[v_4]) \setminus \{u_1, u_2, u_3, w_2\}$ . We obtain  $|Z_0| \leq 4(\Delta - 2)$ . Since  $v_2 \rightarrow u_2, v_1 \rightarrow u_1, v_3 \rightarrow u_3$ , and  $u_2 \rightarrow w_2$ , we obtain  $u_1, u_2, u_3, w_2 \in \mathcal{F}(Z_0)$ , and, hence,  $\frac{|\mathcal{F}(Z_0)|}{|Z_0|} \geq \frac{\Delta-1}{\Delta-2}$ , that is, the set  $Z_0$  satisfies (5), which is a contradiction. Hence, we may assume that  $N_G(v_1) = N_G(v_3)$ .

If some vertex  $v'_4 \in N_G(v_1)$  is not adjacent to some vertex in  $N_G(v_2)$ , then  $N_G(v'_4) \neq N_G(v_2)$ , and one of the previous cases applies to the cycle  $v_1 v_2 v_3 v'_4 v_1$ . Hence, we may assume that all vertices in  $N_G(v_1)$  are adjacent to all vertices in  $N_G(v_2)$ , which implies that  $G$  contains a complete bipartite subgraph  $H$  with partite sets  $N_G(v_1)$  and  $N_G(v_2)$ . If  $N_G(v) \neq N_G(w)$  for two vertices  $v$  and  $w$  that both either belong to  $N_G(v_1)$  or to  $N_G(v_2)$ , then some previous case applies to a cycle of length 4 containing these two vertices. This implies that  $G$  equals  $H$ , and, hence,  $Z(G) = n - 2$ . Since  $G \notin \{K_{\Delta,\Delta}, K_{\Delta-1,\Delta}\}$ , we obtain  $n \leq 2\Delta - 2$ . This implies  $n - 2 \leq \frac{\Delta-2}{\Delta-1}n$ , and (4) follows, which completes the proof in this case.

### Case 3 $g = 3$ .

First, we assume that  $d_G(v_1) = 2$ . Again, this implies  $\Delta = 3$ . Since  $G$  is connected and has maximum degree 3, we may assume that  $d_G(v_2) = 3$ . This implies that the set  $Z_0 = \{v_1, v_2\}$  satisfies (5), which is a contradiction. Hence, we may assume, by symmetry, that  $G$  contains no triangle that contains a vertex of degree 2. For  $i \in \{1, 2, 3\}$ , let  $u_i$  be a neighbor of  $v_i$  that does not lie on  $C$ .

Next, we assume that  $N_G(v_1) \not\subseteq N_G(v_2) \cup N_G(v_3)$  and  $N_G(v_2) \not\subseteq N_G(v_3)$ . We may assume that  $u_1 \in N_G(v_1) \setminus (N_G(v_2) \cup N_G(v_3))$  and  $u_2 \in N_G(v_2) \setminus N_G(v_3)$ . For  $Z_0 = (N_G[v_1] \cup N_G[v_2] \cup N_G[v_3]) \setminus \{u_1, u_2, u_3\}$ , we obtain  $|Z_0| \leq 3(\Delta - 2)$ . Since  $v_3 \rightarrow u_3, v_2 \rightarrow u_2$ , and  $v_1 \rightarrow u_1$ , we obtain  $u_1, u_2, u_3 \in \mathcal{F}(Z_0)$ , and, hence,  $\frac{|\mathcal{F}(Z_0)|}{|Z_0|} \geq \frac{\Delta-1}{\Delta-2}$ , that is, the set  $Z_0$  satisfies (5), which is a contradiction.

Next, we assume that  $N_G(v_1) \not\subseteq N_G(v_2) \cup N_G(v_3)$  and  $N_G(v_2) = N_G(v_3)$ . We may assume that  $u_1 \in N_G(v_1) \setminus (N_G(v_2) \cup N_G(v_3))$ . If  $|N_G(v_1) \cup N_G(v_2)| \leq 2\Delta - 2$ , then let  $Z_0 = (N_G[v_1] \cup N_G[v_2]) \setminus \{u_1, u_2\}$ . Note that  $|Z_0| \leq 2(\Delta - 2)$ . Since  $v_2 \rightarrow u_2$  and  $v_1 \rightarrow u_1$ , we obtain  $u_1, u_2 \in \mathcal{F}(Z_0)$ , and, hence,  $\frac{|\mathcal{F}(Z_0)|}{|Z_0|} \geq \frac{\Delta-1}{\Delta-2}$ , that is, the set  $Z_0$  satisfies (5), which is a contradiction. Hence,  $|N_G(v_1) \cup N_G(v_2)| \geq 2\Delta - 1$ , which implies that  $v_1, v_2$ , and  $v_3$  all have degree  $\Delta$ , and that  $v_3$  is the only common neighbor of  $v_1$  and  $v_2$ . Let  $A = N_G(v_1) \setminus \{v_2, v_3\}$  and  $B = N_G(v_2) \setminus \{v_1, v_3\}$ . Note that  $|A| = |B| = \Delta - 2$ . If some vertex  $u'_1$  in  $A$  is not adjacent to some vertex  $u'_2$  in  $B$ , then let  $Z_0 = (N_G(v_1) \cup N_G(v_2) \cup N_G(v_3)) \setminus \{u'_1, v_1, v_2\}$ . Note that  $|Z_0| \leq 3(\Delta - 2)$ . Since  $u'_2 \rightarrow v_2, v_2 \rightarrow v_1$ , and  $v_1 \rightarrow u'_1$ , we obtain  $v_1, v_2, u'_1 \in \mathcal{F}(Z_0)$ , and, hence,  $\frac{|\mathcal{F}(Z_0)|}{|Z_0|} \geq \frac{\Delta-1}{\Delta-2}$ , that is, the set  $Z_0$  satisfies (5), which is a contradiction. Hence, every vertex in  $A$  is adjacent to every vertex in  $B$ . Note that  $N_G(u) = \{v_2, v_3\} \cup A$  for every vertex  $u$  in  $B$ , and that every vertex in  $A$  has at most one neighbor outside of  $\{v_1\} \cup B$ .

If some vertex  $u'_1$  in  $A$  has a neighbor  $w_1$  outside of  $\{v_1, v_2, v_3\} \cup A \cup B$ , then let  $Z_0 = (N_G(v_1) \cup N_G(v_2)) \setminus \{u'_1, u_2\}$ . Note that  $|Z_0| = 2\Delta - 3$ . Since  $v_2 \rightarrow u_2, v_1 \rightarrow u'_1$ , and  $u'_1 \rightarrow w_1$ , we obtain  $u'_1, u_2, w_1 \in \mathcal{F}(Z_0)$ , and, hence,

$\frac{|\mathcal{F}(Z_0)|}{|Z_0|} \geq \frac{|Z_0|+3}{|Z_0|} = \frac{2\Delta}{2\Delta-3} \geq \frac{\Delta-1}{\Delta-2}$ , that is, the set  $Z_0$  satisfies (5), which is a contradiction. Hence, no vertex in  $A$  has a neighbor outside of  $\{v_1, v_2, v_3\} \cup A \cup B$ . Note that  $A$  induces a subgraph of  $G$  of maximum degree at most 1.

If  $A$  contains two vertices  $u'_1$  and  $u''_1$  that are not adjacent, then let  $Z_0 = (N_G(v_1) \cup N_G(v_2)) \setminus \{u'_1, v_2, u_2\}$ . Note that  $|Z_0| = 2\Delta - 4$ . Since  $u'_1 \rightarrow u_2, v_3 \rightarrow v_2$ , and  $v_1 \rightarrow u''_1$ , we obtain  $u''_1, v_2, u_2 \in \mathcal{F}(Z_0)$ , and, hence,  $\frac{|\mathcal{F}(Z_0)|}{|Z_0|} \geq \frac{|Z_0|+3}{|Z_0|} = \frac{2\Delta-1}{2\Delta-4} > \frac{\Delta-1}{\Delta-2}$ , that is, the set  $Z_0$  satisfies (5), which is a contradiction. Hence, every two vertices in  $A$  are adjacent.

Since  $G$  has maximum degree  $\Delta$ , and every vertex in  $A$  has degree  $|\{v_1\}| + (|A| - 1) + |B| = 2\Delta - 4$ , we obtain  $\Delta \leq 4$ , which implies the contradiction that  $G$  is either  $G_1$  or  $G_2$ . Hence, we may assume, by symmetry, that  $N_G(v_i) \subseteq N_G(v_j) \cup N_G(v_k)$  for  $\{i, j, k\} = \{1, 2, 3\}$ . Note that this implies  $|N_G[v_1] \cup N_G[v_2] \cup N_G[v_3]| \leq \frac{3}{2}(\Delta - 2) + 3$ .

Recall that  $C$  contains no vertex of degree 2.

If  $\Delta = 3$ , then  $N_G(v_i) \subseteq N_G(v_j) \cup N_G(v_k)$  for  $\{i, j, k\} = \{1, 2, 3\}$  implies that  $u_1 = u_2 = u_3$ , that is, all vertices on  $C$  are adjacent to the same vertex outside of  $C$ , which, again using  $\Delta = 3$ , implies the contradiction that  $G$  is  $K_4$ . Hence,  $\Delta \geq 4$ .

Recall that  $C : v_1 v_2 v_3 v_1$  was an arbitrary triangle.

Now, suppose that  $N_G(v_1) \setminus V(C) = N_G(v_2) \setminus V(C) = N_G(v_3) \setminus V(C)$  for every choice of  $C$ , and choose  $C$  such that  $d_G(v_1)$  is as large as possible. If  $u$  and  $u'$  are two non-adjacent vertices in  $N_G(v_1) \setminus V(C)$ , then, considering the triangle  $C' : v_1 v_2 u v_1$ , we obtain  $N_G(u) \setminus V(C') = N_G(v_1) \setminus V(C')$ , and, since  $u' \in N_G(v_1) \setminus V(C')$ , this implies the contradiction  $u' \in N_G(u)$ . Hence,  $N_G(v_1) \setminus V(C)$  is complete, which implies that  $N_G[v_1]$  is complete. By the choice of  $C$ , no vertex in  $N_G[v_1]$  has more neighbors than  $v_1$ , which, by the connectivity of  $G$ , implies the contradiction that  $G$  is the complete graph  $K_{\Delta+1}$  with vertex set  $N_G[v_1]$ . Hence, we may assume, by symmetry, that the triangle  $C : v_1 v_2 v_3 v_1$  is such that  $N_G(v_1) \setminus N_G(v_2)$  is not empty.

We may assume that  $u_1 \in N_G(v_1) \setminus N_G(v_2)$ . Let  $Z_0 = (N_G(v_1) \cup N_G(v_2)) \setminus \{u_1, u_2\}$ . Note that  $|Z_0| \leq \frac{3}{2}(\Delta - 2) + 1$ . Since  $v_2 \rightarrow u_2$  and  $v_1 \rightarrow u_1$ , we obtain  $u_1, u_2 \in \mathcal{F}(Z_0)$ , and, hence,  $\frac{|\mathcal{F}(Z_0)|}{|Z_0|} \geq \frac{|Z_0|+2}{|Z_0|} \geq \frac{\frac{3}{2}(\Delta-2)+3}{\frac{3}{2}(\Delta-2)+1} \geq \frac{\Delta-1}{\Delta-2}$ , that is, the set  $Z_0$  satisfies (5), which is a contradiction. This completes the proof.  $\square$

While our [Conjecture 1](#) remains widely open, we are able to improve (2) at least by some lower order term for subcubic graphs of girth at least 5.

**Theorem 5.** *If  $G$  is a connected graph of order  $n$ , maximum degree 3, and girth at least 5, then*

$$Z(G) \leq \frac{n}{2} - \frac{n}{24\log_2(n) + 6} + 2.$$

**Proof.** Let  $G$  be as in the statement. We begin with an extension statement similar to [Lemma 3](#).

**Claim 1.** *Let  $Z$  be a set of vertices of  $G$ . Let  $F = \mathcal{F}(Z)$  and  $R = V(G) \setminus F$ .*

*If  $F$  induces a connected subgraph of  $G$  of order at least 3, and  $R$  contains a vertex of degree at least 2, then there is a set  $Z'$  of vertices of  $G$  with  $Z \subseteq Z'$ ,*

- (i)  $|Z' \setminus Z| \leq 2\log_2(n)$ ,
- (ii)  $|\mathcal{F}(Z') \setminus F| \geq 2|Z' \setminus Z| + 1$ ,
- (iii)  $Z \subseteq Z'$ , and  $\mathcal{F}(Z')$  induces a connected subgraph of  $G$ .

**Proof of Claim 1.** Note that a vertex in  $F$  with a neighbor in  $R$  has exactly one neighbor in  $F$  and two neighbors in  $R$ , in particular, such a vertex has degree 3.

A subgraph  $H$  of  $G$  is an *extension subgraph* if it is of one of the following types:

- Type a: A path  $P : v_0 \dots v_k$  with  $v_0 \in F, v_1, \dots, v_k \in R$ , and  $d_G(v_k) = 2$ .
- Type b: A path  $P : v_0 \dots v_k$  with  $v_0 \in F, v_1, \dots, v_k \in R, d_G(v_k) = 1$ , and  $k \geq 2$ .
- Type c: A path  $P : v_0 \dots v_k$  with  $v_0, v_k \in F, v_1, \dots, v_{k-1} \in R$ , and  $k \geq 2$ .
- Type d: A cycle  $C : u_1 \dots u_\ell u_1$  with  $u_1 \in F, u_2, \dots, u_\ell \in R$ .
- Type e: The union of a path  $P : v_0 \dots v_k$  and a cycle  $C : u_1 \dots u_\ell u_1$  with  $v_0 \in F, v_1, \dots, v_k, u_1, \dots, u_\ell \in R, v_k = u_1$ , and  $V(P) \cap V(C) = \{u_1\}$ .

Whenever we refer to some extension subgraph, we use the notation introduced above.

First, we show the existence of a small extension subgraph. Therefore, suppose that  $G$  does not contain an extension subgraph of order at most  $2\log_2(n) + 1$ . Since  $G$  is connected, and  $R$  contains a vertex of degree at least 2, there is a vertex  $v$  in  $F$  that has a neighbor  $u$  in  $R$  such that  $u$  has degree at least 2. Since there is no extension subgraph of order at most  $2\log_2(n) + 1$ , the vertex  $u$  is the root of a perfect binary subtree  $T$  of  $G$  of height  $\lfloor \log_2(n) \rfloor$  with  $V(T) \subseteq R$ . Since  $v$  has a neighbor in  $F$ , we obtain the contradiction  $n \geq 2 + n(T) = 2 + 2^{\lfloor \log_2(n) \rfloor + 1} - 1 > n$ .

Let  $H$  be an extension subgraph such that the order  $n(H)$  of  $H$  is as small as possible, and, subject to this first condition, the number of vertices of  $H$  in  $R$  is as small as possible.

As shown above,  $n(H) \leq 2\log_2(n) + 1$ .

Since  $G$  has girth at least 5, and the set  $F$  contains more than two vertices, the choice of  $H$  easily implies that

- $H$  is an induced subgraph of  $G$ ,
- no vertex in  $R \setminus V(H)$  is adjacent to two vertices of  $H$ ,
- $V(H) \cap R$  contains a vertex of degree less than 3 only if  $H$  has Type a or Type b, in which case  $v_k$  is the only such vertex, and
- every vertex  $v$  in  $V(H) \cap R$  with  $d_H(v) = 2$  and  $d_G(v) = 3$  has a neighbor  $p(v)$  in  $R \setminus V(H)$ .

The violation of any of these conditions leads to an extension subgraph of smaller order or of the same order but less vertices in  $R$ . As observed above, every vertex  $v$  in  $V(H) \cap F$  has exactly two neighbors in  $R$ , and if only one of these two neighbors belongs to  $H$ , then we denote the other neighbor by  $p(v)$ .

Now, we consider the different types.

First, assume that  $H$  has Type (a). Let  $u$  be the neighbor of  $v_k$  distinct from  $v_{k-1}$ . If  $u \in F$ , then the choice of  $H$  implies  $k = 1$ . Let  $Z' = Z \cup \{v_k\}$ , and let  $p(u)$  be the neighbor of  $u$  in  $R$  distinct from  $v_k$ . Since  $|Z' \setminus Z| = 1$ , we obtain (i). Since  $p(u), p(v_0) \in \mathcal{F}(Z') \setminus F$  and  $Z' \setminus Z \subseteq \mathcal{F}(Z') \setminus F$ , we obtain  $|\mathcal{F}(Z') \setminus F| \geq 2|Z' \setminus Z| + 1$ , and, hence, (ii). If  $u \notin F$ , then let  $Z' = Z \cup \{p(v_i) : 0 \leq i \leq k-1\}$ . Since  $|Z' \setminus Z| = k = n(H) - 1 \leq 2\log_2(n)$ , we obtain (i). Since  $v_1, \dots, v_k, u \in \mathcal{F}(Z') \setminus F$  and  $Z' \setminus Z \subseteq \mathcal{F}(Z') \setminus F$ , we obtain  $|\mathcal{F}(Z') \setminus F| \geq 2k + 1 = 2|Z' \setminus Z| + 1$ , and, hence, (ii). Clearly, (iii) holds in both cases.

Next, assume that  $H$  has Type (b). If  $k = 2$ , then let  $Z' = Z \cup \{v_k\}$ . Since  $|Z' \setminus Z| = 1$ , we obtain (i). Since  $p(v_0), p(v_1) \in \mathcal{F}(Z') \setminus F$  and  $Z' \setminus Z \subseteq \mathcal{F}(Z') \setminus F$ , we obtain  $|\mathcal{F}(Z') \setminus F| \geq 2|Z' \setminus Z| + 1$ , and, hence, (ii). If  $k \geq 3$ , then let  $Z' = Z \cup \{v_k\} \cup \{p(v_i) : 0 \leq i \leq k-3\}$ . Since  $|Z' \setminus Z| = k-1 = n(H) - 2 \leq 2\log_2(n)$ , we obtain (i). Since  $v_1, \dots, v_{k-1}, p(v_{k-2}), p(v_{k-1}) \in \mathcal{F}(Z') \setminus F$  and  $Z' \setminus Z \subseteq \mathcal{F}(Z') \setminus F$ , we obtain  $|\mathcal{F}(Z') \setminus F| \geq 2k \geq 2|Z' \setminus Z| + 1$ , and, hence, (ii). Clearly, (iii) holds in both cases.

Next, assume that  $H$  has Type (c). Let  $Z' = Z \cup \{p(v_i) : 0 \leq i \leq k-2\}$ . Since  $|Z' \setminus Z| = k-1 = n(H) - 2 \leq 2\log_2(n)$ , we obtain (i). Since  $v_1, \dots, v_{k-1}, p(v_{k-1}), p(v_k) \in \mathcal{F}(Z') \setminus F$  and  $Z' \setminus Z \subseteq \mathcal{F}(Z') \setminus F$ , we obtain  $|\mathcal{F}(Z') \setminus F| \geq 2k \geq 2|Z' \setminus Z| + 1$ , and, hence, (ii). Clearly, (iii) holds.

Next, assume that  $H$  has Type (d). Note that, since  $G$  has girth at least 5, we have  $\ell \geq 5$ . Let  $Z' = Z \cup \{u_\ell\} \cup \{p(u_j) : 2 \leq j \leq \ell-2\}$ . Since  $|Z' \setminus Z| = \ell-2 = n(H) - 2 \leq 2\log_2(n)$ , we obtain (i). Since  $u_2, \dots, u_{\ell-1}, p(u_{\ell-1}), p(u_\ell) \in \mathcal{F}(Z') \setminus F$  and  $Z' \setminus Z \subseteq \mathcal{F}(Z') \setminus F$ , we obtain  $|\mathcal{F}(Z') \setminus F| \geq 2\ell - 2 \geq 2|Z' \setminus Z| + 1$ , and, hence, (ii). Clearly, (iii) holds.

Finally, assume that  $H$  has Type (e). Let  $Z' = Z \cup \{p(v_i) : 0 \leq i \leq k-1\} \cup \{u_\ell\} \cup \{p(u_j) : 2 \leq j \leq \ell-2\}$ . Since  $|Z' \setminus Z| = k + \ell - 2 = n(H) - 2 \leq 2\log_2(n)$ , we obtain (i). Since  $v_1, \dots, v_k, u_2, \dots, u_{\ell-1}, p(u_{\ell-1}), p(u_\ell) \in \mathcal{F}(Z') \setminus F$  and  $Z' \setminus Z \subseteq \mathcal{F}(Z') \setminus F$ , we obtain  $|\mathcal{F}(Z') \setminus F| \geq 2k + 2\ell - 2 \geq 2|Z' \setminus Z| + 1$ , and, hence, (ii). Clearly, (iii) holds.

This completes the proof of the claim.  $\square$

Since  $G$  has maximum degree 3, we have  $n \geq 4$ , which implies  $\frac{1}{2} - \frac{1}{8\log_2(n)+2} \geq \frac{4}{9}$ . For some vertex  $v$  of degree 3, and some neighbor  $u$  of  $v$ , let  $Z_0 = N_G[v] \setminus \{u\}$ . Since  $|Z_0| = 3$  and  $|\mathcal{F}(Z_0)| \geq 4$ , we obtain

$$\frac{|Z_0| - 2}{|\mathcal{F}(Z_0)|} \leq \frac{1}{2} - \frac{1}{8\log_2(n) + 2}.$$

Clearly,  $\mathcal{F}(Z_0)$  induces a connected subgraph of  $G$  of order at least 3.

Suppose that  $Z$  is a set of vertices of  $G$  that satisfies the hypotheses of Claim 1 such that

$$\frac{|Z| - 2}{|\mathcal{F}(Z)|} \leq \frac{1}{2} - \frac{1}{8\log_2(n) + 2}. \quad (6)$$

By Claim 1, the set  $Z$  can be extended to a set  $Z'$  with the properties stated in Claim 1. In particular,

$$\frac{|Z' \setminus Z|}{|\mathcal{F}(Z') \setminus \mathcal{F}(Z)|} \stackrel{(ii)}{\leq} \frac{|Z' \setminus Z|}{2|Z' \setminus Z| + 1} \stackrel{(i)}{\leq} \frac{2\log_2(n)}{4\log_2(n) + 1} = \frac{1}{2} - \frac{1}{8\log_2(n) + 2},$$

which implies

$$\frac{|Z'| - 2}{|\mathcal{F}(Z')|} = \frac{(|Z| - 2) + |Z' \setminus Z|}{|\mathcal{F}(Z)| + |\mathcal{F}(Z') \setminus \mathcal{F}(Z)|} \leq \frac{1}{2} - \frac{1}{8\log_2(n) + 2}.$$

In view of the set  $Z_0$  defined above, this implies the existence of a set  $Z$  of vertices of  $G$  that satisfies (6) such that  $F = \mathcal{F}(Z)$  induces a connected subgraph of  $G$  of order at least 3, and all vertices in  $R = V(G) \setminus F$  have degree 1. Since  $G$  is connected, and every vertex in  $F$  has at most two neighbors in  $R$ , we obtain  $|R| \leq 2|F|$ . Since  $n = |F| + |R|$ , this implies  $|F| \geq \frac{n}{3}$  and  $|R| \leq \frac{2n}{3}$ . Note that every vertex  $v$  in  $F$  that has a neighbor in  $R$  has exactly two neighbors in  $R$ . Let  $\tilde{Z}$  arise from  $Z$  by adding,



for every such vertex  $v$  in  $F$ , exactly one of its two neighbors in  $R$  to  $Z$ . Clearly,  $\tilde{Z}$  is a zero forcing set of  $G$ , and we obtain

$$\begin{aligned} |\tilde{Z}| - 2 &= (|Z| - 2) + \frac{1}{2}|R| \\ &\stackrel{(6)}{\leq} \left( \frac{1}{2} - \frac{1}{8\log_2(n) + 2} \right) |F| + \frac{1}{2}|R| \\ &\leq \left( \left( \frac{1}{2} - \frac{1}{8\log_2(n) + 2} \right) \cdot \frac{1}{3} + \frac{1}{2} \cdot \frac{2}{3} \right) n \\ &= \left( \frac{1}{2} - \frac{1}{24\log_2(n) + 6} \right) n, \end{aligned}$$

which completes the proof.  $\square$

We proceed to our probabilistic upper bound. For a set  $N$  and a non-negative integer  $i$ , let  $\binom{N}{i}$  be the set of all subsets of  $N$  of order  $i$ .

**Theorem 6.** *If  $G$  is a graph, then*

$$Z(G) \leq \sum_{u \in V(G)} \sum_{i=0}^{d_G(u)} (-1)^i \sum_{I \in \binom{N_G(u)}{i}} \left| \{u\} \cup \bigcup_{v \in I} N_G[v] \right|^{-1}.$$

**Proof.** Let  $u_1, \dots, u_n$  be a linear order of the vertices of  $G$  selected uniformly at random. Let  $Z$  be the set of those vertices  $u_i$  such that  $u_i$  is not the unique neighbor within  $\{u_i, \dots, u_n\}$  of some vertex  $u_j$  with  $j < i$ . Clearly,  $Z$  is a zero forcing set of  $G$ . Hence, by the first moment method,  $Z(G) \leq \mathbb{E}[|Z|]$ .

Let  $u$  be a vertex of  $G$ . For  $v \in N_G(u)$ , let  $A_v$  be the event that  $u$  is the rightmost vertex from  $N_G[v]$  within the linear order  $u_1, \dots, u_n$ , that is, if  $u = u_j$ , then  $i < j$  for every  $i$  in  $\{1, \dots, n\}$  with  $u_i \in N_G[v] \setminus \{u\}$ . The definition of  $Z$  implies

$$\mathbb{P}[u \in Z] = \mathbb{P} \left[ \overline{\bigcup_{v \in N_G(u)} A_v} \right].$$

Let  $N = \{u\} \cup \bigcup_{v \in N_G(u)} N_G[v]$  and  $d = |N|$ . Note that there are  $d!$  linear orders of  $N$ . Furthermore, if  $I$  is a subset of  $N_G(u)$ , then the number of linear orders  $\sigma$  of  $N$  such that  $u$  is the rightmost vertex from  $\{u\} \cup \bigcup_{v \in I} N_G[v]$  within  $\sigma$  is exactly  $\frac{d!}{\left| \{u\} \cup \bigcup_{v \in I} N_G[v] \right|}$ ,

which implies

$$\mathbb{P} \left[ \bigcap_{v \in I} A_v \right] = \frac{\left| \{u\} \cup \bigcup_{v \in I} N_G[v] \right|^{-1}}{d!}.$$

By inclusion–exclusion, we obtain

$$\begin{aligned} \mathbb{P}[u \in Z] &= \mathbb{P} \left[ \overline{\bigcup_{v \in N_G(u)} A_v} \right] \\ &= \sum_{i=0}^{d_G(u)} (-1)^i \sum_{I \in \binom{N_G(u)}{i}} \mathbb{P} \left[ \bigcap_{v \in I} A_v \right] \\ &= \sum_{i=0}^{d_G(u)} (-1)^i \sum_{I \in \binom{N_G(u)}{i}} \left| \{u\} \cup \bigcup_{v \in I} N_G[v] \right|^{-1}. \end{aligned}$$

By linearity of expectation, we have  $\mathbb{E}[|Z|] = \sum_{u \in V(G)} \mathbb{P}[u \in Z]$ , and the desired result follows.  $\square$

Since the bound in Theorem 6 is not very explicit, we derive some more explicit corollaries.

For a positive integer  $r$ , let  $H_r = \sum_{i=1}^r \frac{1}{i}$ . It is known that  $\lim_{r \rightarrow \infty} (H_r - \ln r) \approx 0,577$ .

**Corollary 7.** If  $G$  is a  $r$ -regular graph of order  $n$  and girth at least 5, then

$$Z(G) \leq \left( \prod_{i=1}^r \left( 1 - \frac{1}{ri+1} \right) \right) n = \left( 1 - \frac{H_r}{r} \right) n + O\left( \left( \frac{H_r}{r} \right)^2 \right) n.$$

**Proof.** By Theorem 6, we obtain

$$\begin{aligned} \frac{Z(G)}{n} &\leq \frac{1}{n} \sum_{u \in V(G)} \sum_{i=0}^{d_G(u)} (-1)^i \sum_{I \in \binom{N_G(u)}{i}} \left| \{u\} \cup \bigcup_{v \in I} N_G[v] \right|^{-1} \\ &= \sum_{i=0}^r (-1)^i \binom{r}{i} \frac{1}{ri+1} \quad (\text{using the regularity and the girth condition}) \\ &= \sum_{i=0}^r (-1)^i \binom{r}{i} \int_0^1 x^{ri} dx \\ &= \int_0^1 \sum_{i=0}^r (-1)^i \binom{r}{i} x^{ri} dx \\ &= \int_0^1 (1-x^r)^r dx \quad (\text{using the binomial theorem}) \\ &= \frac{1}{r} \int_0^1 (1-z)^r z^{\frac{1}{r}-1} dz \quad (\text{substituting } z = x^r) \\ &= \frac{1}{r} B\left(r+1, \frac{1}{r}\right) \quad (\text{where } B(\cdot, \cdot) \text{ is the Beta function}) \\ &= \frac{1}{r} \frac{\Gamma(r+1) \Gamma\left(\frac{1}{r}\right)}{\Gamma\left(1+r+\frac{1}{r}\right)} \quad (\text{where } \Gamma(\cdot) \text{ is the Gamma function}) \\ &= \frac{r!}{\left(r+\frac{1}{r}\right)\left(r-1+\frac{1}{r}\right) \dots \left(1+\frac{1}{r}\right)} \quad (\text{using } \Gamma(x+1) = x\Gamma(x)) \\ &= \prod_{i=1}^r \frac{i}{i+\frac{1}{r}} \\ &= \prod_{i=1}^r \left( 1 - \frac{1}{ri+1} \right), \end{aligned}$$

which implies the first stated bound for  $Z(G)$ .

Note that








$$\begin{aligned} \prod_{i=1}^r \left( 1 - \frac{1}{ri+1} \right) &= 1 - \sum_{i=1}^r \frac{1}{ri+1} + \sum_{i=2}^r (-1)^i \sum_{I \in \binom{[r]}{i}} \prod_{j \in I} \frac{1}{rj+1} \\ &= 1 - \left( \sum_{i=1}^r \frac{1}{ri} - \sum_{i=1}^r \frac{1}{ri(ri+1)} \right) + \sum_{i=2}^r (-1)^i \sum_{I \in \binom{[r]}{i}} \prod_{j \in I} \frac{1}{rj+1} \\ &= 1 - \left( \frac{H_r}{r} - \sum_{i=1}^r \frac{1}{ri(ri+1)} \right) + \sum_{i=2}^r (-1)^i \sum_{I \in \binom{[r]}{i}} \prod_{j \in I} \frac{1}{rj+1}. \end{aligned}$$

Since

$$\left| \sum_{i=1}^r \frac{1}{ri(ri+1)} \right| \leq \frac{1}{r^2} \sum_{i=1}^r \frac{1}{i^2} \leq \frac{\pi^2}{6r^2} = O\left( \left( \frac{H_r}{r} \right)^2 \right)$$



**Table 1**The seven possible types of the vertex  $u$ .

Type 1	Type 2	Type 3	Type 4	Type 5	Type 6	Type 7
						
$p_1 = \frac{81}{140}$	$p_2 = \frac{149}{252}$	$p_3 = \frac{5}{8}$	$p_4 = \frac{171}{280}$	$p_5 = \frac{101}{168}$	$p_6 = \frac{269}{420}$	$p_7 = \frac{17}{28}$

and

$$\begin{aligned}
\left| \sum_{i=2}^r (-1)^i \sum_{I \in \binom{[r]}{i}} \prod_{j \in I} \frac{1}{rj+1} \right| &\leq \sum_{i=2}^r \sum_{I \in \binom{[r]}{i}} \prod_{j \in I} \frac{1}{rj} \\
&= \sum_{i=2}^r \frac{1}{r^i} \sum_{I \in \binom{[r]}{i}} \prod_{j \in I} \frac{1}{j} \\
&\leq \sum_{i=2}^r \frac{1}{r^i} \frac{1}{i!} \left( \sum_{j_1=1}^r \frac{1}{j_1} \right) \left( \sum_{j_2=1}^r \frac{1}{j_2} \right) \cdots \left( \sum_{j_i=1}^r \frac{1}{j_i} \right) \\
&= \sum_{i=2}^r \frac{1}{i!} \left( \frac{H_r}{r} \right)^i \\
&\leq \left( \frac{H_r}{r} \right)^2 \sum_{i=2}^r \frac{1}{i!} \\
&\leq e \left( \frac{H_r}{r} \right)^2 \\
&= o \left( \left( \frac{H_r}{r} \right)^2 \right),
\end{aligned}$$

we obtain the second stated bound for  $Z(G)$ .  $\square$ 

Note that

$$\prod_{i=1}^r \left( 1 - \frac{1}{ri+1} \right) = \begin{cases} \frac{81}{140} \approx 0.579 & \text{, for } r = 3, \\ \frac{2048}{3315} \approx 0.618 & \text{, for } r = 4, \text{ and} \\ \frac{15625}{24024} \approx 0.65 & \text{, for } r = 5. \end{cases}$$

In fact, this expression is less than the factor  $\frac{r-2}{r-1}$  from (2) for  $r \geq 4$ .

If  $G$  is a cubic triangle-free graph such that no component of  $G$  is  $K_{3,3}$ , then, for every vertex  $u$  of  $G$ , the subgraph of  $G$  that contains all vertices at distance at most 2 from  $u$  as well as all edges incident with neighbors of  $u$  is of one of the seven types illustrated in Table 1. This defines the type of the vertex  $u$ .

**Corollary 8.** If  $G$  is a cubic triangle-free graph such that no component of  $G$  is  $K_{3,3}$ , and  $G$  has  $n_i$  vertices of type  $i$  for  $i \in \{1, \dots, 7\}$ , then  $Z(G) \leq \sum_{i=1}^7 p_i n_i$ .

**Proof.** This follows immediately from Theorem 6 by calculating the probabilities  $\mathbb{P}[u \in Z]$  considered within the proof of Theorem 6 for the vertices  $u$  of the different types. If  $u$  has type 4 for instance, then  $\mathbb{P}[u \in Z] = 1 - \frac{3}{4} + \frac{1}{5} + \frac{2}{7} - \frac{1}{8} = \frac{171}{280}$ .  $\square$

We proceed to the proof of two further cases of the conjecture of Davila and Kenter.

**Theorem 9.** If  $G$  is a graph of girth  $g$  in  $\{5, 6\}$  and minimum degree  $\delta$  at least 2, then

$$Z(G) \geq (g-2)(\delta-2) + 2.$$

**Proof.** Let  $G$  be as in the statement. Let  $Z$  be a zero forcing set of minimum cardinality. For a contradiction, suppose that  $|Z| \leq (g-2)(\delta-2) + 1$ . For  $\delta = 2$ , this implies that  $Z$  contains exactly one vertex, say  $v_1$ . Since  $G$  has more than one vertex, and  $v_1$  has degree at least 2, no vertex in  $V(G) \setminus Z$  is the unique neighbor of  $v_1$ , which implies a contradiction. Hence,  $\delta \geq 3$ . Since  $g \geq 5$ , the order  $n$  of  $G$  is at least  $1 + \delta + \delta(\delta-1) = \delta^2 + 1$ . Since  $g \in \{5, 6\}$  and  $\delta \geq 3$ , we obtain  $n - |Z| \geq \delta^2 - (g-2)(\delta-2) \geq g-2$ , which implies that a forcing sequence  $S: v_1 \rightarrow u_1, v_2 \rightarrow u_2, \dots, v_k \rightarrow u_k$  satisfies  $k \geq g-2$ . Let  $Z' = \{v_1, \dots, v_{g-2}\}$ . Let  $N = \left( \bigcup_{v \in Z'} N_G(v) \right) \setminus Z'$ . Since  $S$  is a forcing sequence,  $Z' \cup N \subseteq Z \cup \{u_1, \dots, u_{g-2}\}$ , and, hence,

$$\begin{aligned} |N| &= |Z' \cup N| - (g-2) \\ &\leq |Z \cup \{u_1, \dots, u_{g-2}\}| - (g-2) \\ &= |Z| \\ &\leq (g-2)(\delta-2) + 1. \end{aligned}$$

Let  $G' = G[Z']$ . Let  $G'$  have  $\kappa$  components. Note that  $\kappa \leq |Z'| = g-2 \leq 4$ . Let  $m'$  be the number of edges between  $Z'$  and  $N$ . Since  $g > |Z'|$ , the graph  $G'$  is a forest and has exactly  $g - \kappa - 2$  edges. This implies that  $m' \geq (g-2)\delta - 2(g-\kappa-2) \geq |N| + 2\kappa - 1$ . Since  $m' > |N|$ , some vertex in  $N$  has more than one neighbor in  $Z'$ . Since  $g > |Z'| + 1$ , no vertex in  $N$  has two neighbors in the same component of  $G'$ . This implies that  $\kappa \geq 2$ .

First, we assume that  $\kappa = 2$ . If three vertices in  $N$  have neighbors in both components of  $G'$ , then  $G$  has a cycle of length at most  $g-1$ , which is a contradiction. Hence, at most two neighbors in  $N$  have neighbors in both components of  $G'$ , which implies the contradiction  $m' \leq |N| + 2 < |N| + 2\kappa - 1$ .

Next, we assume that  $\kappa = 3$ . If some vertex  $u$  in  $N$  has neighbors in all three components of  $G'$  and another vertex  $u'$  has two neighbors in  $Z'$ , then  $G$  has a cycle of length at most  $g-1$ , which is a contradiction. Similarly, if two distinct vertices in  $N$  have neighbors in the same two components of  $G'$ , then  $G$  has a cycle of length at most  $g-1$ , which is a contradiction. These observations imply the contradiction  $m' \leq |N| + \binom{3}{2} < |N| + 2\kappa - 1$ .

Finally, we assume that  $\kappa = 4$ , which implies that  $g = 6$ , and that  $Z'$  is an independent set. Again, no two distinct vertices in  $N$  have neighbors in the same two components of  $G'$ . If some vertex in  $N$  has neighbors in three components of  $G'$ , then this implies the contradiction  $m' \leq |N| + 5 < |N| + 2\kappa - 1$ . Similarly, if no vertex in  $N$  has neighbors in three components of  $G'$ , then this implies the contradiction  $m' \leq |N| + \binom{4}{2} < |N| + 2\kappa - 1$ .

This final contradiction completes the proof.  $\square$

## Acknowledgment

We thank Henning Bruhn-Fujimoto for fruitful discussion.

## References

- [1] A. Aazami, Hardness Results and Approximation Algorithms for Some Problems on Graphs (Ph.D. thesis), University of Waterloo, 2008.
- [2] D. Amos, Y. Caro, R. Davila, R. Pepper, Upper bounds on the  $k$ -forcing number of a graph, *Discrete Appl. Math.* 181 (2015) 1–10.
- [3] F. Barioli, W. Barrett, S. Butler, S. Cioaba, D. Cvetkovic, S. Fallat, C. Godsil, W. Haemers, L. Hogben, R. Mikkelsen, S. Narayan, O. Pryporova, I. Sciriha, W. So, D. Stevanovic, H. van der Holst, K.V. Meulen, A.W. Wehe, AIM Minimum Rank - Special Graphs Work Group, Zero forcing sets and the minimum rank of graphs, *Linear Algebra Appl.* 428 (2008) 1628–1648.
- [4] F. Barioli, W. Barrett, S. Fallat, H.T. Hall, L. Hogben, B. Shader, P. van den Driessche, H. van der Holst, Parameters related to tree-width, zero forcing, and maximum nullity of a graph, *J. Graph Theory* 72 (2013) 146–177.
- [5] D. Burgarth, V. Giovannetti, Full control by locally induced relaxation, *Phys. Rev. Lett.* 99 (2007) 100501.
- [6] D. Burgarth, V. Giovannetti, L. Hogben, S. Severini, M. Young, Logic circuits from zero forcing, *Nat. Comput.* 14 (2015) 485–490.
- [7] D. Burgarth, K. Maruyama, Indirect Hamiltonian identification through a small gateway, *New J. Phys.* 11 (2009) 103019.
- [8] Y. Caro, R. Pepper, Dynamic approach to  $k$ -forcing, *arXiv:1405.7573v1*.
- [9] L.S. Chandran, C.R. Subramanian, Girth and treewidth, *J. Combin. Theory Ser. B* 93 (2005) 23–32.
- [10] K.B. Chilakamuri, N. Dean, C.X. Kang, E. Yi, Iteration index of a zero forcing set in a graph, *Bull. Inst. Combin. Appl.* 64 (2012) 57–72.
- [11] R. Davila, F. Kenter, Bounds for the Zero-Forcing Number of Graphs with Large Girth, *arXiv:1406.0482v2*.
- [12] C.J. Edholm, L. Hogben, M. Huynh, J. LaGrange, D.D. Row, Vertex and edge spread of the zero forcing number, maximum nullity, and minimum rank of a graph, *Linear Algebra Appl.* 436 (2012) 4352–4372.
- [13] S. Fallat, K. Meagher, B. Yang, On the complexity of the positive semidefinite zero forcing number, *Linear Algebra Appl.* 491 (2016) 101–122.
- [14] M. Gentner, L.D. Penso, D. Rautenbach, U.S. Souza, Extremal values and bounds for the zero forcing number, *Discrete Appl. Math.* 214 (2016) 196–200.
- [15] A. Girão, G. Mészáros, S.G.Z. Smith, Recent results in zero-forcing of graphs, *arXiv:1611.07513v1*.
- [16] L. Hogben, M. Huynh, N. Kingsley, S. Meyer, S. Walker, M. Young, Propagation time for zero forcing on a graph, *Discrete Appl. Math.* 160 (2012) 1994–2005.
- [17] L. Hogben, H. van der Holst, Forbidden minors for the class of graphs  $G$  with  $\zeta(G) \leq 2$ , *Linear Algebra Appl.* 423 (2007) 42–52.
- [18] L. Lu, B. Wu, Z. Tang, Proof of a conjecture on the zero forcing number of a graph, *Discrete Appl. Math.* 213 (2016) 233–237.

- [19] S.A. Meyer, Zero forcing sets and bipartite circulants, *Linear Algebra Appl.* 436 (2012) 888–900.
- [20] D.D. Row, Zero Forcing Number: Results for Computation and Comparison with Other Graph Parameters (Ph.D. thesis), Iowa State University, 2011.
- [21] D.D. Row, A technique for computing the zero forcing number of a graph with a cut-vertex, *Linear Algebra Appl.* 436 (2012) 4423–4432.
- [22] S. Severini, Nondiscriminatory propagation on trees, *J. Phys. A* 41 (2008) 482002.
- [23] M. Trefois, J.-C. Delvenne, Zero forcing number, constrained matchings and strong structural controllability, *Linear Algebra Appl.* 484 (2015) 199–218.