| |
|---|
| Awarding Body:<br><br>Arden University |
| Programme Name:<br><br>Health and Care Management |
| Module Name:<br><br>Health Analytics and Planning |
| Assessment Title:<br><br>Analysis Briefing Paper |
| Student Number:<br><br>[redacted] |
| Tutor Name:<br><br>[redacted] |
| Word Count:<br><br>2087 words |

## Summary

This paper focuses on the effectiveness of UNICEF-supported community-based interventions in Sub-Saharan Africa that seek to reduce malnutrition among children under the age of five. Child malnutrition remains a severe problem in this region, aggravated by poverty, food insecurity, inadequate sanitation, and restricted healthcare access. These circumstances have resulted in significant malnutrition rates, with stunting (an important malnutrition indicator) afflicting 42% of children on average. Stunting rates vary greatly by location, ranging from 19.1% in Senegal to 58.3% in Burundi, highlighting the unique problems faced by different groups.

The paper focuses on major risk factors such as delayed vaccines, a lack of maternal education, and food hardship. UNICEF's efforts address these issues through programs that encourage maternal education, provide timely vaccines, and enhance food access.
The statistical study, including the Pearson coefficient of 0.69, demonstrates a positive relationship between community size and stunting prevalence, showing that bigger communities confront more difficult obstacles in eliminating malnutrition.

The findings emphasize the need for locally customized, resource-intensive techniques. The recommendations include adapting health and nutrition programs to community needs, expanding assistance for bigger, unde rserved regions, focusing on high-risk populations (e.g., males and children with multiple births), and adopting continuous monitoring to measure success. These techniques seek to increase the effectiveness of treatments, reduce malnutrition, and improve child health outcomes throughout the area.

## Table of Contents

**Summary** ..... **2**

**Research Question** ..... **4**

**Introduction** ..... **4**

**Background Information** ..... **5**

**Data Collect Approach/Presentation of Findings** ..... **6**

Data Analysis ..... 6

Statistical analysis ..... 8

**Discussion** ..... **10**

**Conclusion/Recommendation** ..... **11**

**References** ..... **12**

## Research Question

"How effective are community-based interventions in reducing malnutrition among children under five years old in Sub-Saharan Africa, as supported by UNICEF initiatives?"

## Introduction

In today's dynamic health and social care environment, data analytics is pivotal in improving service quality and enhancing decision-making processes. As health systems strive to optimise care delivery, the ability to measure service performance using established quality frameworks is crucial. This essay aims to explore the application of health analytics in addressing malnutrition among children under five years old in Sub-Saharan Africa, focusing on UNICEF-supported community-based interventions. By analysing and evaluating the effectiveness of these interventions, the study seeks to generate actionable insights that inform evidence-based decisions.

Additionally, this essay will delve into the key digital skills and technologies required for effective data analytics and the challenges faced in implementing such initiatives. Through this, the discussion will align with the learning outcomes of evaluating service performance, applying health analytics concepts, and addressing the digital competencies necessary to drive impactful health outcomes. The research will address the question, "How effective are community-based interventions in reducing malnutrition among children under five years old in Sub-Saharan Africa, as supported by UNICEF initiatives?"

## Background Information

Sub-Saharan Africa is a region struggling with a significant public health challenge: child malnutrition. This condition, characterised by inadequate nutrient intake, leading to poor growth and development, is a major contributor to child mortality and morbidity (Akombi et al., 2017). The prevalence of malnutrition in this region is alarmingly high, with millions of children under five years old affected (Akombi et al., 2017).

Poverty, food insecurity, poor sanitation, limited healthcare, and cultural practices are key factors driving child malnutrition in Sub-Saharan Africa (Owolade et al., 2022). Many families struggle to afford nutritious food and healthcare, while dehydration, conflicts, and economic instability intensify food shortages (Owolade et al., 2022). Unsanitary conditions and limited access to clean water increase the risk of disease, further harming children's health (Owolade et al., 2022). The lack of healthcare facilities and trained professionals prevents early diagnosis and treatment. Additionally, certain cultural practices can contribute to malnutrition.

The consequences of child malnutrition are far-reaching. Malnutrition is a major cause of death among children under five, particularly in the context of infectious diseases. It can negatively impact a child's brain development, leading to cognitive delays and poor educational outcomes (Akombi et al., 2017). Malnutrition can also limit a child's growth and development, affecting their physical health and well-being (Saunders & Smith, 2010).

This research seeks to investigate the effectiveness of UNICEF-supported community-based interventions in reducing malnutrition among children under five years old in Sub-Saharan Africa. It aims to evaluate how well these interventions are improving health outcomes and contributing to the reduction of malnutrition, considering various factors such as the sustainability and reach of the programs.

The research addresses the problem of persistent malnutrition among children under five years old in Sub-Saharan Africa, a critical issue that negatively impacts child survival and development. Specifically, the study will focus on assessing whether community-based interventions, as supported by UNICEF, are sufficiently effective in combating malnutrition and identifying potential improvements to enhance their impact.

## Data Collect Approach/Presentation of Findings

### Data Analysis

The table presents the prevalence of stunting and its associated risk factors among children under five in significant Sub-Saharan African countries. Stunting, a critical indicator of chronic malnutrition, varies by region due to factors like maternal education, food insecurity, and vaccine timing (Vaivada et al., 2020). This study emphasises the need for customised interventions. UNICEF-supported community programs aim to enhance maternal education, ensure timely vaccinations, and improve food security, which can reduce stunting and enhance child health outcomes.

| Age | Study Site & Sample Size | Prevalence of Stunting | Risk Factors |
|---|---|---|---|
| **1-60 months** | 33 countries | 41.1% total; Nigeria 39.2%, Ethiopia 50.8%, DRC 43.4%, Burundi highest 58.3%, Gabon lowest 21.0% | Reduced risk with early vaccines; higher risk with delayed vaccines |
| **< 60 months** | Addis Ababa, 5,822 | 19.6% | Higher risk if mother has less education, food insecurity, male child |
| **< 59 months** | 35 countries, 384,747 | West 33.9%, Central 37.8%, East 35.3%, South 26.5% | Higher risk if mother has no education, poorest and rural households |
| **6-59 months** | Ethiopia, 315 | 22.2% (3.8% severely stunted) | Lower risk with literate mother, older children, wealthier households |
| **< 59 months** | 25 SSA countries, 213,889 | - | Higher risk for twins or multiple births |
| **< 59 months** | Uganda, 104 | 38.5% | Lower risk in pastoralist mothers |

| | | | |
|---|---|---|---|
| **< 59 months** | 18 SSA countries, 55,749 | - | Higher risk with HIV+ mothers, male children, multiple births, rural and poor households |
| **6-59 months** | DRC, 3,721 | 35.2% | Higher risk if mother has no say in financial decisions, male children, shorter birth intervals |
| **< 59 months** | Rwanda, 3,594 | 38% | Higher risk in male children, low birthweight, shorter mothers, no parasite drugs during pregnancy |
| **< 59 months** | Cameroon, 5,053; Nigeria, 18,823; DRC, 3,777 | Cameroon 32%, Nigeria 41%, DRC 44.5% | Higher risk for non-widowed single mothers |
| **6-59 months** | Kenya, 1,245 | 47% (23.4% severely stunted) | Higher risk in boys and children aged 36-47 months |
| **6-59 months** | Mozambique, 874 | 37% | Higher risk in current food aid recipients |
| **< 59 months** | 34 SSA countries, 299,065 | Highest in Burundi 54.6%, lowest in Ghana 19.2% | Higher risk if short pregnancy gap; lower risk with longer pregnancy intervals |

Table 1 – Summary of studies and findings on the prevalence and risk factors of stunting among children aged 0–60 months in Sub-Saharan Africa.
Adapted from Quamme et al., (2022).

Stunting prevalence varies greatly across Sub-Saharan Africa, ranging from 58.3% in Burundi to 19.1% in Senegal (Quamme et al., 2022). Delays in vaccines, a lack of maternal education, food insecurity, poverty, and cultural traditions are all contributing causes. Boys, children from multiple births, and those with HIV-positive moms are at a higher risk (Quamme et al., 2022). Stunting is more common in West (33.9%) and Central Africa (37.8%) than in Southern Africa (26.5%), emphasising the importance of targeted, community-based treatments that address local socioeconomic and cultural factors (Kurniawan et al., 2024).

In data analytics, data cleaning is crucial because it eliminates or corrects inaccurate, lacking, or unnecessary data, guaranteeing accurate and trustworthy findings (Hillier, 2023).

Particularly in health and nutrition programs, where precise data informs successful treatments, data cleaning is crucial for correct insights. Analysts can pinpoint particular problems, like vaccination gaps or maternal education requirements, and precisely monitor progress with clean, consistent data. Data mistakes are reduced with the use of programs like Excel and Python, guaranteeing trustworthy insights that promote focused health enhancements, immediate improvements, and efficient resource distribution for addressing issues unique to a community (Hillier, 2023).

## Statistical analysis

This section of the paper will show statistical analysis to identify trends, correlations or regressions. The statistical analysis tools used are mean, median, mode, standard deviation, and Pearson coefficient. The author of this essay designed the following tables, but the information was evidence-based research and will be correctly cited.

| Age (Months) | Country/Region | Sample Size (Population) | Prevalence of Stunting (%) |
|---|---|---|---|
| 1–60 | Ethiopia | 315 | 51 |
| <60 | Rwanda | 3,594 | 38 |
| 6–59 | Kenya | 1,245 | 47 |
| <59 | Cameroon | 5,053 | 32 |

Table 2 – Prevalence of Stunting (%) depending on the country in Sub-Saharan Africa.

Adapted from Quamme et al., (2022). Adapted from Author.

| Prevalence of Stunting | | | | |
|---|---|---|---|---|
| Mean | Median | Mode | Standard Deviation | Pearson Coefficient |
| 42 | 42,5 | #N/A | 7,449832213 | 0,690239375 |

Table 3 – Prevalence of Stunting statistical analysis.

Adapted from Author.

The average prevalence of stunting in the dataset is 42%. This suggests that stunting affects 42% of the population under observation (Almond, 2024).

The median value, or midpoint of the dataset, is 42.5%. This indicates that half of the observations have stunting rates less than 42.5%, while the other half have higher rates. The proximity of the mean and median indicates that the data is fairly symmetrically distributed (Almond, 2024).

The mode is not provided, implying that no single value appears more frequently than others in the collection. This could imply that the data points vary without any recurrence of values (Almond, 2024).

A standard deviation of 7.45% shows a moderate distribution, indicating that stunting prevalence rates vary across the dataset This indicates that the prevalence of stunting varies by 7.45% from the mean.

The Pearson coefficient is a correlation measure that indicates the strength and direction of the relationship between two continuous variables, both measured on the same interval or ratio scale (Kenton, 2024). The Pearson correlation coefficient of 0.69 shows a moderate to strong positive correlation between the variables examined (the probability of stunting and community size). This shows that as one variable grows, the other tends to follow as well, although not perfectly. This means that when the population increases, the prevalence of stunting also increases.

Larger communities may not always have greater stunting rates, but they do present problems that make it more difficult for organisations such as UNICEF to address the issue effectively. These communities demand additional resources, planning, and flexible tactics. However, UNICEF's success depends on adapting interventions to each community's requirements and leveraging local knowledge to overcome these obstacles.

This briefing paper examines the occurrence of stunting among children under the age of five in Sub-Saharan Africa, utilising two data sets: stunting rates by country and statistical analysis of these rates. Descriptive statistics show an average stunting prevalence of 42%, with a standard deviation of 7.45%, indicating slight variation between nations. The Pearson correlation coefficient of 0.69 indicates a favourable association between population size and stunting rates. Using Excel for visualisation, the findings highlight the importance of focused interventions matched to community size and individual local needs to effectively reduce stunting and enhance child health outcomes.

## Discussion

A statistical investigation of stunting prevalence in Sub-Saharan Africa revealed numerous important findings. The average stunting rate among the nations surveyed is 42%, with significant regional differences ranging from 19.1% in Senegal to 58.3% in Burundi. A Pearson correlation coefficient of 0.69 suggests a reasonably strong positive association between community size and stunting prevalence, implying that as population size grows, so do stunting rates. This is a critical result that underscores the difficulties larger communities face in eliminating malnutrition.

The study addressed the research goals by identifying trends in stunting prevalence and risk factors. Delayed immunisation, a lack of maternal education, food insecurity, and poverty all contribute to increased stunting rates, confirming previous studies (Quamme et al., 2022). Boys, children of multiple births, and those with HIV-positive moms are consistently at higher risk, according to recognised research in the field (Vaivada et al., 2020).

These findings are supported by current research, such as UNICEF's community-based programs, which focus on boosting maternal education, raising food security, and guaranteeing timely vaccines to address malnutrition. Previous research by Vaivada et al. (2020) and Kurniawan et al. (2024) has identified similar risk variables and emphasised the significance of tailored, community-driven treatments to effectively alleviate stunting.

However, the research identifies gaps in addressing bigger community needs, such as increased stunting rates, which may necessitate additional resources, planning, and specific interventions. This emphasises the need for localised, flexible methods that adapt to the particular obstacles faced by different locations, which are critical for reducing stunting and improving child health outcomes in Sub-Saharan Africa.

The statistical analysis supported the significance of focused interventions to reduce stunting by providing answers to the research questions and confirming findings from other studies. This critical viewpoint underlines the importance of ongoing, evidence-based efforts to reduce stunting through context-specific interventions that address the complex interaction of malnutrition-causing causes.

## Conclusion/Recommendation

The study's findings highlight the difficulties and efficacy of community-based, UNICEF-supported interventions aimed at lowering under-five-year-old malnutrition in Sub-Saharan Africa. By concentrating on important elements like maternal education, food security, and vaccine timing, the study shows that these interventions have a favourable effect on lowering stunting. The results emphasise the necessity of customised, localised strategies, given the average stunting rate of 42% and regional variances. Stunting prevalence and community size are positively correlated, which emphasises the greater difficulties faced by bigger communities and the need for resource-intensive, adaptable therapies to address their particular requirements.

To increase the impact of these interventions, many recommendations are proposed:

1. Customise nutrition and health programs to local needs, emphasising early immunisation and improving maternal education.
2. Increase funding for bigger communities with high stunting rates, especially those in low-income and rural regions, to enhance access to nutrition and healthcare.
3. To decrease their vulnerability to malnutrition, give priority to interventions for high-risk populations, such as males, children of multiple births, and children whose mothers are HIV positive.
4. Regularly check on progress by monitoring stunting rates and evaluating the success of treatments, enabling adjustments to enhance outcomes.

## References


Akombi, B.J. *et al.* (2017) *Child malnutrition in sub-Saharan africa: A meta-analysis of demographic and Health Surveys (2006-2016)*, *National Library of medicine* . Available at: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5426674/ (Accessed: 12 October 2024).

Almond, N. (2024) *What are mean median mode range? definition, examples & practice question for Primary School*, *Third Space Learning*. Available at: https://thirdspacelearning.com/blog/what-is-mean-median-mode/#:~:text=Frequently%20asked%20mean%2C%20median%2C%20and%20mode%20questions,-What's%20the%20difference&text=Mean%20median%20mode%20and%20range%20all%20calculate%20the%20averages%20of,is%20the%20most%20common%20number. (Accessed: 22 October 2024).

Hargrave, M. (2024) *Standard deviation formula and uses vs. variance*, *Investopedia*. Available at: https://www.investopedia.com/terms/s/standarddeviation.asp (Accessed: 22 October 2024).

Hillier, W. (2023) *What is data cleaning and why does it matter? [how-to]*, *CareerFoundry*. Available at: https://careerfoundry.com/en/blog/data-analytics/what-is-data-cleaning/ (Accessed: 11 November 2024).

Kenton, W. (2024) *What is the Pearson coefficient? definition, benefits, and history*, *Investopedia*. Available at: https://www.investopedia.com/terms/p/pearsoncoefficient.asp (Accessed: 18 October 2024).

Kurniawan, A.L. *et al.* (2024) *Community-based interventions targeting multiple forms of malnutrition among adolescents in low-income and middle-income countries: Protocol for a scoping review*, *BMJ Open*. Available at: https://bmjopen.bmj.com/content/14/3/e078969 (Accessed: 16 October 2024).

Owolade, A.J.-J. *et al.* (2022) *Malnutrition: An underlying health condition faced in Sub Saharan Africa: Challenges and recommendations*, *ScienceDirect*. Available at: https://www.sciencedirect.com/science/article/pii/S2049080122015291#:~:text=Several%20factors%20contribute%20to%20the,fluctuation%20of%20food%20prices%2C%20etc. (Accessed: 12 October 2024).

Quamme, S.H. and Iversen , P.O. (2022) *Prevalence of child stunting in Sub-Saharan africa and its risk factors*, *ScienceDirect*. Available at: https://www.sciencedirect.com/science/article/pii/S2667268522000092#tbl1 (Accessed: 13 October 2024).

Saunders, J. and Smith, T. (2010) *Malnutrition: Causes and consequences*, *National Library of Medicine*. Available at: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4951875/ (Accessed: 12 October 2024).

UNICEF (no date) *Partnerships and governance for Nutrition*, *UNICEF*. Available at: https://www.unicef.org/nutrition/partnerships-and-governance (Accessed: 16 October 2024).

Vaivada , T. *et al.* (2020) *Stunting in childhood: An overview of global burden, trends, determinants, and drivers of decline*, *ScienceDirect*. Available at: https://www.sciencedirect.com/science/article/pii/S0002916522009546?via%3Dihub (Accessed: 16 October 2024).