

Nathan Smith

---

**Abstract**

Supernovae that show evidence of strong shock interaction between their ejecta and pre-existing slower circumstellar material (CSM) constitute an interesting, diverse, and still poorly understood category of explosive transients. The chief reason they are extremely interesting is because they tell us that in a subset of stellar deaths, the progenitor star becomes wildly unstable in the years, decades, or centuries before explosion. This is something that has not been included in standard stellar evolution models but may significantly change the end product and yield of that evolution and complicates our attempts to map SNe to their progenitors. Another reason they are interesting is because CSM interaction is an efficient engine for making bright transients, allowing superluminous transients to arise from normal SN explosion energy, and transients of normal supernova luminosity to arise from sub-energetic explosions or low radioactivity yield. CSM interaction shrouds the fast ejecta in bright shock emission, obscuring our view of the underlying explosion, and the radiation hydrodynamics is challenging to model. The CSM interaction may also be highly nonspherical, perhaps linked to binary interaction in the progenitor system. In some cases, these complications make it difficult to tell the difference between a core-collapse and thermonuclear explosion or to discern between a nonterminal eruption, failed supernova, or weak supernova. Efforts to uncover the physical parameters of individual events and connections to progenitor stars make this a rapidly evolving topic that challenges paradigms of stellar evolution.

---

N. Smith (✉)

Steward Observatory, University of Arizona, Tucson, AZ, USA

e-mail: [nathans@as.arizona.edu](mailto:nathans@as.arizona.edu); [nathans.steward@gmail.com](mailto:nathans.steward@gmail.com)

## Contents

1	Introduction.....	404
2	Basic Physical Picture.....	406
3	A (Probably) More Realistic Physical Picture.....	410
4	Observed Subtypes.....	413
5	Dust Formation in CSM Interaction.....	417
6	Progenitors and Pre-SN Mass Loss.....	418
6.1	CSM Properties.....	418
6.2	Direct Detections.....	421
6.3	Links to Progenitor Types.....	423
7	Closing Comments.....	425
8	Cross-References.....	426
	References.....	426

## 1 Introduction

Type IIn supernovae (SNe IIn hereafter), defined broadly as SNe that exhibit bright and narrow Balmer lines of H in their spectra, were recognized as a distinct class of objects relatively late compared to other SN subtypes (see Schlegel 1990 and Filippenko 1997 for early background). This was spurred largely by observations of the classic SN IIn 1988Z and connecting it to unusual properties in some other SNe. Our understanding of this class and various subclasses has been evolving continuously since then as more SNe are discovered and the diversity of SNe IIn grows. The related class of SNe Ibn, showing narrow lines of He in the optical spectrum (and relatively weak H lines), was clearly identified only 10 years ago following studies of the prototype SN 2006jc (Foley et al. 2007; Pastorello et al. 2007).

Interpreting observations of interacting SNe is fundamentally different from studying “normal” SNe. For our purposes here, “normal” SNe are those cases where the observed emission comes from a photosphere receding (in mass coordinates) back through freely expanding SN ejecta, or radiation from the optically thin SN ejecta at late times that are heated internally by radioactivity. These are the main subtypes of II-P, II-L, IIf, Ib, Ic, and Ic-BL discussed elsewhere in this volume (see paragraph “Cross-References”). SNe IIn and Ibn represent a physically very different scenario, where the usual diagnostics and physical pictures cannot be applied. As a means of introduction to the topic, we begin with two warnings that should be heeded when thinking of SNe IIn and related subtypes of interacting SNe:

Warning 1. The main engine is conversion of kinetic energy to light in a shock at the outside boundary of the SN. It is not shock-deposited energy leaking out of a homologously expanding envelope (as in early phases of a SN II-P), nor is it internal heating from radioactive decay (most SNe at peak and at late times). Instead, the observed radiation comes primarily from

(continued)

a shock crashing into dense CSM which—unfortunately from the point of view of predictive models—makes the situation complicated and malleable. “Malleable” here means that it is hard to calculate predictive models because the shock luminosity can rise and fall somewhat arbitrarily depending on the radial density structure of the CSM. One should not take the absence of normal signatures (broad lines in the SN ejecta photosphere or nebular emission lines) as an indication that these are not true SNe, because CSM interaction can mask these normal signatures. Moreover, radiation from the shock can propagate back into the SN ejecta, changing their appearance.

Warning 2. SNe IIn and related categories are not really a *supernova type* (or more accurately, not an intrinsic explosion type), but an *external phenomenon* associated with CSM interaction. Any type of core-collapse or thermonuclear SN (or for that matter, any non-SN explosive outflow) can appear as a SN IIn or SN Ibn. All that is required is fast ejecta with sufficient energy crashing into slower ejecta with sufficient density. This is a cause of much confusion and uncertainty.

As a consequence of these differences compared to other SNe, we find a very diverse range of properties in interacting SNe. There is not a singular evolutionary path that leads to an interacting SN, nor is there a singular mapping of SNe IIn to a specific progenitor type. For example, we usually associate SNe II-P as coming from moderately massive red supergiants of 8–20  $M_{\odot}$  (Smartt 2009), but such assignments are not so straightforward for SNe IIn. It can be difficult to tell the difference between a SN IIn that arises from the core-collapse of a very massive luminous blue variable (LBV) or LBV-like star and a thermonuclear explosion of a white dwarf with dense CSM. SNe IIn can arise any time there is dense H-rich material sitting around a star that explodes, which might happen in a number of different ways. Exploring this diversity in progenitors as well as in CSM and explosion properties is a main emphasis of current work because it has important implications for mass loss and instability in the late phases of stellar evolution.

The detailed physical picture and time evolution of interacting SNe may also be very different from normal SNe. Instead of a more-or-less spherical photosphere receding through outflowing ejecta, we have radiation from a thin shell, which may start out as an optically thick sphere but may transition to a limb-brightened shell that is transparent in the middle and optically thick at the edges. We may have underlying explosions that would otherwise be too faint to be observed (e.g., explosions producing very low radioactive yield) and that do not contribute to the traditional populations of extragalactic SNe but nevertheless yield bright transients through their CSM interaction. Thus, we should be mindful that some SNe IIn

may be powered by entirely new (or so far observationally unrecognized) classes of explosions other than SNe II-P, SNe II-L, SNe IIb, SNe Ib, and SNe Ic. Much of this is still little explored or uncharted territory.

For this chapter, it is also worth making a distinction of scope. Actually, all SNe are interacting with surrounding material at some level, because space is not a true vacuum and an interaction arises naturally even when fast SN ejecta expand into low-density ISM or their progenitor's normal wind (this eventually forms a standard SN remnant). Whether this can be detected in extragalactic SNe shortly after explosion—and in what waveband it is detected—depends to a large extent on the CSM density. For the purposes of discussion here, we take “Interacting SNe” to mean those with extraordinarily strong CSM interaction that leads to a radiative shock at early times after explosion, producing narrow lines in the visible-wavelength spectrum and enhanced optical continuum luminosity at early times. We exclude otherwise normal SNe that show signs of interaction based on radio or X-ray emission, although it should be kept in mind that these represent the tail of a continuum of CSM density stretching from superluminous SNe, through more traditional SNe II<sub>n</sub>, and on to lower CSM density.

---


## 2 Basic Physical Picture

When a SN explodes inside a dense cocoon of CSM, a strong shock is driven into the CSM, creating a basic structure as depicted in Fig. 1. A forward shock (FS) is driven outward into the CSM, and a corresponding reverse shock (RS) is driven back into the SN ejecta. In the simplest picture, four zones are delineated where heated material can radiate and contribute to the observed spectrum:

- Zone 1. The unshocked CSM outside the forward shock (photoionized)
- Zone 2. The swept-up CSM that has been hit by the forward shock
- Zone 3. The decelerated SN ejecta that have encountered the reverse shock
- Zone 4. The freely expanding SN ejecta

In normal SNe, we only see emission from Zone 4 at visible wavelengths as the photosphere recedes back through the freely expanding ejecta, creating broad emission and absorption lines whose velocities decrease smoothly with time as the emitting layer recedes through the monotonic outflow. Zones 2 and 3 are sometimes detected in X-rays or radio emission if the CSM is dense enough. Zone 1 might be seen in absorption lines, although this is rare.

In an interacting SN, any of these four zones can contribute strongly to the emitted visible-wavelength spectrum. In fact, the continuum photosphere moves through all four zones as time proceeds, and their relative contributions to line emission change with time. The basic picture of this interaction, which can differ widely in detail from case to case, has been described many times in the literature (Chugai 1997, 2001; Chugai and Danziger 1994; Chugai et al. 2004; Dessart et al. 2015; Smith et al. 2008a, 2010a, 2015).



**Fig. 1** Sketch of the basic picture in CSM interacting SNe. Four different zones are noted with numbers: (1) the pre-shock CSM, (2) the shocked CSM, (3) the shocked SN ejecta, and (4) the freely expanding SN ejecta. These zones are divided by boundaries corresponding to the forward shock, the reverse shock, and the contact discontinuity between the shocked CSM and shocked ejecta where material cools, mixes via Rayleigh-Taylor instabilities, and piles up. This is often called the cold dense shell (CDS) in a SN IIn or SN Ibn. The squiggly radial lines are meant as a reminder that X-rays and UV radiation generated in the shock can propagate out to the CSM or inward to the unshocked ejecta, changing the physical state of the gas there. A zoom-in of Zones 2 and 3 is shown at the right. In practice, efficient radiative cooling can cause these two zones to collapse to very thin layers, and mixing can make them merge into one thin clumpy shell. This figure is adapted from Smith et al. (2008a)

At early times, radiation from the shock propagates upstream and causes an ionized precursor. At this stage (sometimes only lasting a day or two, sometimes several weeks or months), the electron-scattering continuum photosphere is actually in the unshocked CSM, obstructing our view of the shock. The spectrum usually appears as a smooth blue continuum with narrow emission lines that have broad Lorentzian wings. These wings are not caused by Doppler motion of shocked gas, but by electron scattering as narrow line photons work their way out of the optically thick CSM.

As the pre-shock CSM density drops at larger radii, eventually the photosphere recedes into Zones 2 and 3, and we begin to see a transformation in the spectrum. This usually occurs around the time of peak luminosity. Strong intermediate-width (few  $10^3 \text{ km s}^{-1}$ ) components of permitted lines appear, sometimes accompanied by P Cygni absorption. Here we are seeing radiation from the post-shock gas. Because of strong radiative cooling (which occurs by definition in an interacting SN), Zones 2 and 3 can collapse into a geometrically very thin layer, and mixing at the contact

discontinuity may cause these two zones to merge. The standard picture is that this cooled gas is very dense and piles up in a “cold dense shell” (Chugai et al. 2004). This gas is continually reheated by the shock and emits strongly in lines like  $H\alpha$  or  $He\ I$  that appear as the strongest intermediate-width lines in the spectra of SNe IIn and SNe Ibn, respectively. This line emission continues to be strong even after the continuum photosphere recedes further, because the CDS is reheated by X-ray and UV radiation from the FS and RS.

After peak, the continuum photosphere may recede back into Zone 4 in the freely expanding SN ejecta. If the CSM interaction is relatively weak or short-lived, an observer might then see a fairly normal SN spectrum with broad P Cygni lines at the end of the photospheric phase or typical signatures of SN ejecta heated by radioactive decay in the nebular phase. However, if CSM interaction is strong and sustained, an observer might never actually see a normal SN ejecta photosphere or nebular phase, and the P Cygni absorption may be filled in. The underlying SN (heated by the original shock deposition or radioactive decay) may fade in a few months, whereas CSM interaction might remain very bright for many months or years. How long this CSM interaction stays bright is determined mainly by the duration and speed of the pre-SN mass loss, not by the explosion. By the time the CSM fades enough to become transparent, the inner SN ejecta may be too faint to detect on their own, and the ongoing CSM interaction may continue to dominate the spectrum at late times. In fact, the fast SN ejecta may be heated predominantly by radiation from the CSM interaction shock that propagates backward into the outermost ejecta, creating a very different time-dependent spectrum than is seen in a normal SN heated from the inside out by radioactive decay. One must bear these differences in mind when interpreting SNe IIn spectra, especially at late phases.

Of course, if the geometry is asymmetric, any of these zones can be seen simultaneously and at different characteristic velocities, potentially making the interpretation quite complicated. Absorption features may or may not be seen, depending on viewing angle. Even if the interaction is spherically symmetric, we might see multiple zones simultaneously because long path lengths through the limb-brightened edges of the CDS may remain optically thick, while material along our line of sight in the CDS has already become transparent. Very strong and non-monotonic changes in density and velocity make this a complicated situation that presents a significant challenge for radiation transfer codes. For the time being, unraveling the various clues in the spectral evolution is something of an art form.

The total luminosity generated by CSM interaction in this picture can be high because a radiative shock is a very efficient engine to convert kinetic energy into visible-wavelength light. Whereas a normal SN II might have a total radiated energy that is 1 % of its kinetic energy, a Type II SN can have a total radiated energy closer to 50 % or more of the total explosion kinetic energy. The luminosity of CSM interaction  $L$  depends essentially on the rate at which CSM mass is entering the forward shock, which of course depends mostly on the progenitor’s mass-loss rate, and is typically expressed as

$$L = \frac{1}{2} w V_{\text{CDS}}^3, \quad (1)$$

where  $w$  is the wind density parameter  $w = 4\pi R^2 \rho$ , or  $w = \dot{M}/V_{\text{CSM}}$ , and  $V_{\text{CDS}}$  is the value for the evolving speed of the CDS. From the intermediate-width lines in optical spectra, we can measure the value of  $V_{\text{CDS}}$  (be careful not to do this at early times when electron scattering in the CSM is determining the line width; at early times the true value of  $V_{\text{CDS}}$  is hidden from the observer, and it may be much higher than when it is seen at later times). We can also measure  $V_{\text{CSM}}$  from the narrow component due to pre-shock gas, especially if a narrow P Cygni profile is observed (note here that higher-resolution spectra with  $R \simeq 4000$  are especially useful). Using the luminosity ( $L$ ) derived from photometry (note that for  $V$  and  $R$  bands, the bolometric correction may be quite small if the apparent temperature is around 6000–7000 K), we can then turn this equation around to estimate the progenitor star's mass-loss rate

$$\dot{M}_{\text{CSM}} = 2 L \frac{V_{\text{CSM}}}{(V_{\text{CDS}})^3} \quad (2)$$

that occurred at some time  $t = t_{\text{exp}}(V_{\text{CDS}}/V_{\text{CSM}})$  in the past, where  $t_{\text{exp}}$  is the time after explosion to which these measurements correspond (careful again, as  $t_{\text{exp}}$  has its own potentially large uncertainties).

This sort of analysis is oversimplified, but it is sufficient to give us a ballpark idea of the sorts of mass-loss rates that are required to produce a Type IIn supernova. In order to make the CSM interaction luminosity high enough that it can compete with the photosphere of a typical SN (say,  $3 \times 10^8 L_{\odot}$ , or  $-16.5$  mag) and with representative values of  $V_{\text{CSM}} = 100 \text{ km s}^{-1}$  and  $V_{\text{CDS}} = 2500 \text{ km s}^{-1}$ , we require a huge mass-loss rate of order  $0.01 M_{\odot} \text{ yr}^{-1}$ . SNe IIn tend to be more luminous than  $-16.5$  mag, or even superluminous, and so the implied mass-loss rates get very big (although remember that  $V_{\text{CDS}}$  is probably faster than  $2500 \text{ km s}^{-1}$  in the early optically thick phases near peak). This mass-loss rate is well beyond the realm of normal stellar winds (Smith 2014). We return to the implications for progenitors later.

For a constant mass-loss rate in the progenitor wind (and hence, a single value of  $w$ ),  $L$  will usually drop with time as the forward shock decelerates. Equation (1) does a pretty good job of describing the visible continuum luminosity as long as the material is optically thick enough to reprocess most or all of the X-ray and UV luminosity into visible continuum radiation. It does not work so well at very early times during the rise or at peak when diffusion is important, and it does not work at later times when the X-rays begin to leak out and the  $\text{H}\alpha$  line emission becomes very strong. At these phases, Eq. (2) will underestimate the required progenitor mass-loss rate if  $L$  is derived from the visible light curve. At later times, one needs to adopt a very uncertain efficiency factor in order to derive physical parameters from observations. The efficiency of converting kinetic energy into visible light can vary widely depending on CSM density (van Marle et al. 2010), CSM composition (H-free gas tends to be more transparent), and geometry, as discussed next.

### 3 A (Probably) More Realistic Physical Picture


Figure 1 is a good place to begin when picturing the physics of CSM interaction, but it may be something of a fantasy. One exceedingly obvious result from 20 years of *Hubble Space Telescope* imaging of circumstellar shells is that they are often not spherically symmetric, and this is especially true of massive stars and eruptive mass loss. Bipolar shells and equatorial disks or rings are common around evolved massive stars.

This has an impact on our interpretation of observed signatures of CSM interaction, because when spherical symmetry is broken, it can affect the qualitative appearance of the spectrum, and it can change line profile shapes. More importantly, it can dramatically change the inferred total energy and mass budgets, and sometimes it can hide evidence of CSM interaction altogether. If one assumes spherical symmetry in an interacting SN that is actually not spherically symmetric, one can be led to astoundingly wrong conclusions. In this case, the “simplest” assumptions may be simply inappropriate.

Figure 2 shows an example of how the scenario in Fig. 1 might be modified if a SN is surrounded by an equatorial disklike distribution of CSM as opposed to a spherical shell (this is borrowed from Smith et al. 2015, where much of this topic is discussed in more detail). A key point is that if the dense CSM is disklike, then strong CSM interaction only occurs in a small fraction of the total solid angle involving the equatorial regions (see Mauerhan et al. 2014; Smith et al. 2014). The vast majority of the SN ejecta (say, 90 % if the disk only intercepts 10 % of the solid angle seen by the SN) can expand unimpeded in the polar directions. Thus, as far as the majority of the SN ejecta are concerned, their kinetic energy is not tapped by CSM interaction, and they would emit a normal broad-lined spectrum. For the unlucky 10 % of the SN ejecta in the equator, their kinetic energy is converted into heat and radiation by the shock. In this way, CSM interaction with a disk can create SNe that are overluminous compared to normal SNe II-P, for example, but not nearly as luminous as the class of superluminous SNe. This may be applicable to many SNe IIn.

**Enveloped, Swallowed, and Hidden CSM Interaction.** There is a nonintuitive but very interesting result that can occur from interaction with asymmetric CSM, which is that *the resulting SN might not show any narrow lines even if much of its luminosity comes from strong CSM interaction*. This can occur in the following way: Imagine that a SN progenitor is surrounded by a dense CSM disk located at a radius 10–20 AU. The initial shock interaction might produce strong narrow lines in the first few days. In the disk, the fast SN ejecta are decelerated, creating a strong shock that produces a moderately high luminosity and bright narrow-line emission from this CSM interaction. However, the remaining 90 % of the SN ejecta expand unimpeded as if there was no CSM interaction. The normal SN ejecta photosphere quickly expands to a radius of 10 s of AU in a few days. At such time, the optically





**Fig. 2** Sketch of the more complicated interaction when the CSM is highly asymmetric; in this example a disklike distribution surrounds the progenitor. **(a)** The initial pre-explosion state is a progenitor with a wind, as well as a dense disk at radii of  $\sim 10$  AU. **(b)** Immediately after explosion, narrow lines may arise either from a photoionized dense wind or from the disk. Strong CSM interaction with the disk occurs immediately and enhances the early luminosity, but the emergent spectrum may be dominated by narrow lines with Lorentzian wings. **(c)** CSM interaction with the disk has slowed the forward shock in the equatorial plane, but the SN ejecta expand relatively unimpeded in the less dense polar regions. After a few days, the normal SN ejecta photosphere may expand so much that it completely envelops the CSM interaction occurring in the disk. The enveloped CSM interaction can now heat the optically thick ejecta from the inside and contribute significantly to the total visible luminosity, even if no narrow lines are seen, and may cause the photosphere to be asymmetric. **(d)** At late times, the SN ejecta photosphere recedes. At this time, the CSM interaction in the disk is exposed again (now with larger radial velocities because it has been accelerated), and intermediate-width and possibly doubled-peaked lines may be seen from the ongoing interaction. This sketch was originally invoked to explain the observed evolution of PTF11iqb (Smith et al. 2015)

thick SN ejecta can completely envelope the equatorial CSM interaction, hiding some or all of the narrow and intermediate-width lines that arise due to CSM interaction. The extra radiated energy generated by the equatorial CSM interaction is now unable to escape freely because it has been swallowed by the opaque fast ejecta, and this radiative energy is instead deposited in the interior of the optically thick SN ejecta and must diffuse out. It will therefore add extra heat to the outer envelope, perhaps mimicking extra deposition of radioactive decay energy. The asymmetric injection of heat in this way will likely make the photosphere asymmetric. Viewed from the outside at a time of roughly a month after explosion, one might then observe a broad-lined SN photosphere with enhanced luminosity and perhaps a longer than normal duration, unusual line profiles, or significant polarization. At this stage, it would be difficult to tell the difference between some small extra amount of radioactive heating or buried CSM interaction.

All hope is not lost, though, because an observer might then be able to deduce that CSM interaction was contributing to the main light curve all along by watching until late times. After the SN recombination photosphere recedes, the enveloped CSM interaction region is exposed once again. When this happens, the dense CSM disk that has been swept up into a cold dense shell (or cold dense torus) and accelerated to a few thousand  $\text{km s}^{-1}$  may now be seen again via strong intermediate-width line emission. Prominent examples of this are SN 1998S, PTF 11iqb, SN 2009ip, and even the SN I Ib 1993J (Leonard et al. 2000; Matheson et al. 2000; Mauerhan et al. 2014; Pozzo et al. 2004; Smith et al. 2014, 2015). This line emission can be much stronger than the nebular emission from the optically thin inner ejecta heated by the dwindling radioactive decay. Moreover, the ongoing CSM interaction can emit strong X-rays that propagate back into the SN ejecta, potentially changing the emergent nebular spectrum.

Perhaps the most important thing to realize is that we now have two different physical explanations for the wide diversity in peak luminosity of SNe IIn, even if all explosions have the same  $10^{51}$  ergs of kinetic energy. The range of luminosity from the strongest CSM interaction in superluminous SNe down to cases with minimal extra luminosity provided by CSM interaction can be explained either by ramping down the density of the progenitor's wind or by increasing the degree of asymmetry in dense CSM. The strongest clues that significant CSM asymmetry is present are from (1) spectropolarimetry (e.g., Mauerhan et al. 2014), (2) asymmetric line profile shapes (although some asymmetry can also be caused by dust or occultation by the SN photosphere itself), or (3) the time evolution of velocities (e.g., seeing very fast speeds in the SN ejecta that persist after the strongest CSM interaction has subsided; Smith et al. 2014). The last point confirms unequivocally that not all the fast SN ejecta participated in the CSM interaction. There may also be some special observed signatures of asymmetry at certain viewing angles (e.g., edge-on) and specific times. If one does recognize such signatures of asymmetry, one must realize that the progenitor mass-loss rate inferred from equation (2) as well as the SN explosion energy are not just lower limits, but likely underestimates by a factor of  $\sim 10$ . Of course, one can also invoke differences in explosion energy, explosion mechanism, SN ejecta mass, and radioactive yield to also contribute to the observed diversity.

## 4 Observed Subtypes

The main physical parameters that determine the observed properties of a typical core-collapse SN are the mass of ejecta, kinetic energy of the explosion (as well as the mass of synthesized radioactive material), and the composition and structure of the star's envelope at the time of explosion. This leads to the wide diversity in observed types of normal ejecta-dominated core-collapse SNe and thermonuclear SNe (II-P, II-L, IIb, Ib, Ic, Ia).

For interacting SNe, we have all these same free parameters of the underlying explosion—as well as the possibility of SNe with low or no radioactive yield and a wider potential range of explosion energy—but to these, we must also add the variable parameters associated with the CSM into which the SN ejecta crash: CSM mass and/or mass-loss rate, radial distribution (speed and ejection time before explosion), CSM composition, and CSM geometry. Given these parameters, it may not be surprising that the class of interacting SNe is extremely diverse and observations are continually uncovering new or apparently unique characteristics. Moreover, a SN can change type depending on when it is seen. An object that looks like a SN IIn in the first few days can morph into a normal SN II-L or SN IIb if it undergoes enveloped/asymmetric CSM interaction and may then return to being a Type IIn at late times, as discussed in the previous section.

There are, however, some emerging trends among interacting SNe. The list below attempts to capture some of these emerging subtypes among interacting SNe that appear to share some common and distinct traits. The reader should be advised, however, that this is still a rapidly developing field, so this is neither a definitive nor a complete list, and moreover, there are observed events that seem to skirt boundaries or overlap fully between different subtypes. Further subdivisions are likely to be clarified with time. The list below includes a descriptive name and some prototypical or representative observed examples that are often mentioned (again, not a complete list).

**Superluminous IIn; compact shell (SN 2006gy).** SN 2006gy was the first superluminous SN to be discovered, and it remains extremely unusual even though several other SLSNe IIn have since been found (see Ofek et al. 2007; Smith et al. 2007, 2010a; Woosley et al. 2007). It had a slow rise to peak ( $\sim 70$  days) and faded within another 150 days or so, which is unlike the other SLSNe IIn that seem to fade very slowly and steadily from peak. Also unusual was that it had strong intermediate-width P Cygni absorption features in its spectra, strong line-blanketing absorption in the blue, and narrow P Cygni features from the CSM. It is thought to have arisen from a relatively compact and opaque CSM shell, where CSM interaction mostly subsided within a year. In this case, the CSM is thought to have arisen from a single eruption that ejected  $\sim 20 M_{\odot}$  about 8 years prior to core-collapse (Smith et al. 2010a). SN 2006gy is one of the best observed SNe IIn and is often discussed, but readers should be aware that it is not at all typical.

**Superluminous IIn; extended shell (SN 2006tf, SN 2010jl, SN 2003ma).** These are basically the superluminous version of the SN 1988Z-like subclass, discussed next. Unlike SN 2006gy, they show strong, smooth blue continua without much line blanketing and little or no P Cygni absorption in their intermediate-width lines. They also tend to fade slowly and steadily, in some cases remaining bright for several years as the shock runs through an extended dense shell (e.g., Fransson et al. 2014; Rest et al. 2011; Smith et al. 2008a).

**Enduring IIn (SN 1988Z, SN 2005ip).** These SNe IIn show a smooth blue continuum superposed with strong narrow and intermediate-width H lines and in some cases even broad components (Aretxaga et al. 1999; Chugai and Danziger 1994; Smith et al. 2009). Some cases, such as SN 2005ip, have strong narrow coronal emission lines (Smith et al. 2009), implying photoionization of dense clumpy CSM by X-rays generated in the shock. They tend to be more luminous than SNe II-P, but not as bright as SLSNe. These objects tend to fade very slowly and show signs of strong CSM interaction for years or even decades after discovery; SN 1988Z is *still* going strong. One can picture these as a stretched-out version of the SLSNe IIn, in the sense that they have somewhat lower luminosities at peak, but they last longer, eventually sweeping through similar amounts of total CSM mass (of order several to 20  $M_{\odot}$ ). While SLSNe have CSM that is so dense that it requires eruptive LBV-like mass loss in the years or decades before explosion, these enduring SNe IIn like SN 1988Z and SN 2005ip can potentially be explained by extreme RSG winds blowing for centuries before core-collapse. If there is such a thing as a “standard” SN IIn, this is probably what most IIn enthusiasts have in mind.

**Transitional IIn (SN 1998S, PTF11iqb, SN 2013cu).** This class of SNe IIn only shows fleeting signatures of CSM interaction that disappear quickly and may be entirely missed if the SN is not discovered early after explosion. SN 1998S was one of the nearest and best studied SNe IIn that helped shape our understanding of SNe IIn, and so it has often been referred to as “prototypical”—but it really is not. SN 1998S was not very luminous, and its spectral signatures of CSM interaction disappeared pretty quickly, transitioning into a broad-lined ejecta-dominated photosphere within a couple weeks (Leonard et al. 2000; Shivvers et al. 2015). This indicated that the total mass of CSM was actually quite modest (of order 0.1  $M_{\odot}$  or so), substantially different from the types noted above. If it had not been caught so early, SN 1998S might not have been classified as a Type IIn at all. An older example of this was SN 1983K (Niemela et al. 1985). More recently, a few other related objects have been found (SN 2013cu & PTF11iqb; Gal-Yam et al. 2014; Smith et al. 2015), which only showed a Type IIn spectrum for the first few days after discovery (and these cases were thought to have been discovered very early, within 1 day or so of explosion). PTF11iqb and SN 2013cu then morphed into broad-lined SNe with spectra similar to SNe II-L or SNe I Ib, respectively. At late times, PTF11iqb showed strong CSM interaction again, very similar to SN 1998S and the Type I Ib SN 1993J. We do not know how common these are. While SNe IIn

represent roughly 8–9 % of core-collapse SNe (Smith et al. 2011a), this statistic does not include SNe that only exhibit Type IIn characteristics for a brief window of time and then morph into other types after a few days; thus, early strong CSM interaction with dense inner winds might be fairly common among the larger population of ccSNe. The corresponding (potentially very important) implication is that many core-collapse SNe may suffer a brief pulse of episodic mass loss shortly before core-collapse. The reason for this is not yet known but is probably related to the final nuclear burning stages and may have important implications for models of core-collapse.

**Late-time interaction in otherwise normal SNe (SN 1993J, SN 1986J).** The overlap with the previous subclass is probably considerable, but it is worth mentioning that some otherwise normal SNe (perhaps appearing normal simply because we missed the early IIn signatures in the first day or so) show particularly strong CSM interaction at late times. Some of the more common cases are SNe I Ib and SNe II-L, for which there are well-studied and famous nearby examples like SN 1993J and SN 1986J (Matheson et al. 2000; Milisavljevic and Fesen 2013). For these, the transition from a SN into a SN remnant is somewhat blurry, and some of these are well-known nearby aging SNe. Because this interaction is most apparent when the SNe have faded after the first year, this phenomenon can only be studied in nearby galaxies. These may be caused by strong RSG winds or by equatorial CSM deposited by binary interaction in objects like SNe I Ib that are more clearly associated with the end products of binary evolution. The reason why this CSM still resides close to the star at the time of death is still unclear and (as for the previous subclass) may hint at some rapid and dramatic changes in the final nuclear burning sequences in a wider fraction of SN progenitors than just standard SNe IIn.

**Delayed onset, slow rise, multi-peaked (SN 2009ip, SN 2010mc, SN 2008iy, SN 1961V, SN 2014C).** In contrast to the previous subclass, some SNe IIn show little or no signs of CSM interaction at first but then rise to the peak of CSM interaction after a delay of months or years. This delay is presumably due to the time it takes for the fastest SN ejecta to catch up to a CSM shell that was ejected a year or more before core-collapse. The underlying SN ejecta photosphere could be relatively faint at first if the progenitor was a more compact BSG, like SN 1987A, or an LBV; the faintness of the initial SN might cause the initial transient to be missed altogether or mistaken as a pre-SN eruption rather than the actual SN, since it might precede the delayed onset of peak luminosity that actually arises when the ejecta overtake the CSM. It must be admitted, however, that from the time of peak onward, these tend to appear as relatively normal SNe IIn; as such, the distinction between this “delayed onset” subclass and other SNe IIn might be artificial and only distinguishable in cases with fortuitous predisccovery or pre-peak data. The amount of delay in the onset of CSM interaction bears important physical information about the elapsed time between the pre-SN eruption and core-collapse. In the case of SN 2009ip, the delay of  $\sim 40$  days made sense, as fast SN ejecta caught up to CSM moving at 10 % the speed, associated with eruptions that were actually observed a

year or so before the SN (Mauerhan et al. 2013a, 2014; Smith et al. 2014). SN 1961V had a light curve consistent with a normal SN II-P for the first  $\sim 100$  days, followed by a brighter peak (Smith et al. 2011b). In SN 2014C, the delayed onset had different chemical properties, where a H-poor stripped-envelope SN crashed into a H-rich shell after a year (Milisavljevic et al. 2015). In some cases, however, the delayed onset is rather extreme and the CSM interaction strong; SN 2008iy appeared to be a relatively normal core-collapse SN in terms of luminosity, which rose to very luminous peak as a Type IIn after 400 days (Miller et al. 2010). SN 1987A might be thought of as an even more extended version of this, where the onset of CSM interaction was delayed for 10 years as the SN ejecta caught up to CSM ejected  $\sim 10^4$  yr earlier. We do not know what fraction of normal SNe have a delayed onset of CSM interaction, since most SNe are not monitored continuously with large telescopes for years or decades after they fade.

**Type IIn-P (SN 1994W, SN 2011ht, the Crab).** This is a distinct subclass of SNe IIn that exhibit IIn spectra throughout their evolution but have light curves with a well-defined plateau drop (Mauerhan et al. 2013b). They are not to be confused with “transitional” SNe IIn/II-P that may have narrow lines at early times and evolve into an otherwise normal SN II-P. In SNe IIn-P, the narrow H lines with strong narrow P Cygni profiles are seen for the duration of their bright phase. The drop from the plateau is quite extreme (several magnitudes), and the late-time luminosities suggest low yields of  $^{56}\text{Ni}$ . These may be the result of  $\sim 10^{50}$  erg electron capture SN explosions with strong CSM interaction (Mauerhan et al. 2013b; Smith 2013). It has been proposed that the Crab Nebula may be the remnant of this type of event, although this remains uncertain (Smith 2013).

**SN IIn impostors (LBVs, SN 2008S-like, etc.).** These transients have narrow H lines in their spectra similar to SNe IIn but are subluminous (fainter than  $M_R \simeq -15.5$  mag) and have slower velocities than normal SNe IIn (see Smith et al. 2011b and references therein). One may argue that the luminosity cut is arbitrary, but the narrow lines without any broad wings do seem distinct from other SNe IIn (most of the time). Some are certainly nonterminal transients akin to LBVs (because they repeat or the star survives), but some cases are not so clear. Although they may not be SNe, they are included here because we are not sure yet—some objects thrown into this bin could be underluminous because they are interacting transients that arise from “failed” SNe (fallback to a black hole), pulsational pair instability SNe, ecSNe, SNe from compact BSG progenitors with low radioactive yield, mergers with compact object companions, or other terminal events.

**Type Ia/Type IIn or Ia-CSM (SN 2002ic, SN 2005gj, PTF11kx).** These are transients that show spectral features indicative of an underlying SN Ia ejecta photosphere, but with strong superposed narrow H lines and additional continuum luminosity (Silverman et al. 2013). Cases with stronger CSM interaction tend to obscure the Type Ia signatures, leading to ongoing controversy about their potential core-collapse or thermonuclear nature (Benetti et al. 2006). Cases with weaker

CSM interaction (PTF11kx) reveal the Type Ia signatures more clearly, allowing them to be identified unambiguously as thermonuclear events. The relatively rare evolutionary circumstance that leads a thermonuclear SN to have a large mass of H-rich CSM is still unclear.

**Type Ibn (SN 2006jc).** These are similar to SNe IIn, except that H lines are weak or absent, so that narrow or intermediate-width He I lines dominate the spectrum ( $H\alpha$  is weaker than He I  $\lambda 5876$ ). Without the assistance of H opacity, these objects tend to fade more quickly than most SNe IIn. The class is quite diverse and includes a range of H line strengths and CSM speeds, including transitional IIn/Ibn cases (SN 2011hw) where H and He I lines have similar strength (Pastorello et al. 2008, 2015; Smith et al. 2012a). It has been hypothesized that the likely progenitors may be Wolf-Rayet stars or LBVs in transition to a WR-like state (Foley et al. 2007; Pastorello et al. 2007; Smith et al. 2012a).

**Type Icn (hypothetical).** A discovery of this class has not yet been reported (as far as the author is aware), but as transient searches continue, there may be cases where a SN interacts with a dense shell of CSM that is both H and He depleted, yielding narrow or intermediate-width lines of CNO elements, for example. With sufficient creativity, one can imagine stellar evolution pathways that might create this; such SNe are clearly rare, and if they never turn up, their absence will provide interesting physical constraints for some binary models.

---

## 5 Dust Formation in CSM Interaction

The formation of dust by SNe may be essential to explain the amount of dust inferred in high-redshift galaxies. SNe with strong CSM interaction provide an avenue for dust formation that is different from normal SNe and possibly much more efficient. In normal SNe, dust forms in the freely expanding ejecta where there is a competition between cooling and low enough temperatures, while the ejecta are also rapidly expanding and achieving lower and lower densities. Even if dust forms efficiently, this ejecta dust might get destroyed when it crosses the reverse shock. In interacting SNe, by contrast, evidence suggests that dust can form very rapidly in the extremely dense, post-shock cooling layers (Zones 2 and 3 in Fig. 1). Moreover, this dust is already behind the shock and may therefore stand a better chance of surviving and contributing to the ISM dust budget.

The first well-studied case of post-shock dust formation was in SN 2006jc, which was a Type Ibn. The classic signs of dust formation were seen starting at only 50 days post-discovery, in an increase in the rate of fading, excess IR emission, and an increasing blueshifted asymmetry in emission-line profiles (Smith et al. 2008b). The last point strongly favored post-shock dust formation, since this was seen in the intermediate-width lines that were emitted from the post-shock zones. The Type Ibn event may have been the result of a Type Ic explosion crashing into a He-rich shell. In this case, one might think that the C-rich ejecta were important in assisting the



efficient formation of dust, as is seen for the post-shock dust formation in colliding-wind WC+O binaries (Gehrz and Hackwell 1974; Williams et al. 1990). However, similar evidence for this same mode of post-shock dust formation has also been seen in several Type IIn events, such as SN 2005ip, SN 1998S, SN 2006tf, SN 2010jl, and others (Fox et al. 2009; Gall et al. 2014; Pozzo et al. 2004; Smith et al. 2008a, 2009, 2012b). It may therefore be the case that enhanced post-shock dust formation is a common outcome of strong CSM interaction, where efficient post-shock cooling causes the forward shock to collapse and become very dense.

## 6 Progenitors and Pre-SN Mass Loss


Basic considerations about powering SNe IIn with CSM interaction, noted above, argue that extremely strong pre-SN mass loss is required. What sorts of progenitor stars can give rise to this dense CSM?

### 6.1 CSM Properties

The first thing to do is to look around us in the nearby universe and ask what sort of observed classes of stars might fit the bill. Figure 3 makes this comparison by plotting the diversity in wind density of interacting SNe, deduced from the pre-shock wind (or shell) expansion speed and the inferred mass-loss rate, as compared to expected mass loss from known types of stars. The shaded and colored regions in this figure show rough parameters for different classes of evolved massive stars that are potential SN progenitors with strong winds. These are taken from the recent review by Smith (2014); see the caption for definitions. In this plot, a given wind density parameter (required for a particular value of a CSM interaction luminosity via Eq. 1) has a diagonal line increasing to the upper right. The solid line shows a typical value for a moderate-luminosity SN IIn, and the dashed line is a typical lower bound for wind densities required to make a SN IIn (although an object can be slightly below this line and still make a Type IIn spectrum if the wind is clumpy or asymmetric).

We can see immediately that the normal classes of evolved stars with strong but relatively steady winds (WR stars, LBV winds, normal YSGs and RSGs, AGB stars; see caption to Fig. 3 for the key to the abbreviations) do not match up to the wind density required for SNe IIn. These could potentially produce SNe that have strong X-ray or radio emission from CSM interaction, but they are not dense enough to slow the forward shock and to cause the forward shock to cool radiatively and collapse into the cool dense shell that is required for intermediate-width H $\alpha$ . The only observed classes of stars that have the high wind densities required are the giant eruptions of LBVs and the most extreme cool hypergiants (the slower winds of extreme RSGs and YHG make their winds have density comparable to the shells in LBV giant eruptions, even though the mass-loss rates are lower). Both classes of stars are not steady winds, but rather, they are dominated by relatively short-lived





**Fig. 3** Plot of mass-loss rate as a function of wind velocity, comparing values for interacting SNe to those of known types of stars. The solid-colored regions correspond to values for various types of evolved massive stars taken from the review about mass loss by Smith (2014), corresponding to AGB and super-AGB stars, red supergiants (RSGs) and extreme RSGs (eRSG), yellow supergiants (YSG), yellow hypergiants (YHG), LBV winds and LBV giant eruptions, binary Roche-lobe overflow (RLOF), luminous WN stars with hydrogen (WNH), and H-free WN and WC Wolf-Rayet (WR) stars. A few individual stars with well-determined very high mass-loss rates are shown with circles (VY CMa, IRC+10420,  $\eta$  Car’s eruptions, and P Cyg’s eruption). Also shown with X’s are some representative examples of SNe IIIn (and one SN Ibn) that have observational estimates of the pre-shock CSM speed from the narrow emission component as measured in moderately high-resolution spectra as well as estimates of the progenitor mass-loss required, taken from the literature; this figure is from a paper in prep. by the author. Note that in some cases, an observationally derived value for the mass of the CSM has been converted to a mass-loss rate with a rough estimate of the time elapsed since ejection

phases of eruptive or episodic mass loss or extremely dense and clumpy winds. In the case of LBVs, these are eruptions that last a few months to a decade. For extreme RSGs and YHGs, these are phases of enhanced mass loss that may persist for centuries to a few thousand years.

From the point of view of their mass loss, LBVs (and to a lesser extent the most extreme cool hypergiants) are good candidates for the progenitors of SNe IIn. This does not, however, mean that these stars are necessarily all poised to explode in the next couple years, nor does it mean that they are the only possible IIn progenitors. The caveat is that the progenitors of SNe IIn undergo strong eruptive or episodic mass loss immediately before core-collapse and may therefore experience significant changes in their internal structure. The stars may have looked very different in the time period before their pre-SN eruptions began. Hence, these stars in their immediate pre-SN state may not actually exist among nearby populations of stars in the Milky Way and Magellanic Clouds right now.

Also note that this is a mass-loss *rate*, not total mass, and note that these are usually lower limits to the required mass-loss rate (the high rate makes high optical depths and, hence, makes an observational determination difficult). In cases where we have estimates of the total mass ejected, one must divide by an assumed timescale to derive a mass-loss rate; for SNe IIn, this timescale is inferred from the relative velocities of the CSM and CDS. Since we are essentially plotting a wind density, it is important to recognize that two different SNe with the same inferred progenitor mass-loss rate might have had that high mass loss lasting for very different amounts of time and, hence, may have very different amounts of total mass ejected shortly before the SN. In some cases, the total mass can be more constraining than the rate. For example, the total mass of  $\sim 20 M_{\odot}$  required for some superluminous SNe IIn rules out any progenitor stars with initial masses below about  $30\text{--}40 M_{\odot}$ , because we need a comparable mass of SN ejecta to provide enough momentum for a long-lasting interaction phase (not to mention mass loss due to winds throughout the life of the star). Also, there are cases like SN 2005ip where the inferred mass-loss rate is on the low end, but this SN had remarkably long-lasting CSM interaction that suggests a total mass of order  $15 M_{\odot}$  of H-rich material (e.g., Smith et al. 2009; Stritzinger et al. 2012), which rules out lower-mass RSGs and super-AGB stars for its progenitor.

Warning 3. One must be somewhat cautious in drawing conclusions about the progenitor based on only the observed pre-shock CSM speed. We generally think of the outflow speed being proportional to the star's escape velocity, so we expect slower outflows from RSGs and YHG, somewhat faster speeds for LBVs and BSGs (few  $10^2 \text{ km s}^{-1}$ ), and very fast outflows for WR stars (few  $10^3 \text{ km s}^{-1}$ ). This is a good guide if the outflow is a relatively steady radiation-driven wind. However, remember that the observed CSM may be the result of eruptive or explosive mass loss driven by a shock wave. If so, it is possible to get a relatively fast outflow even from a bloated RSG; eruption speeds need not match steady wind speeds for various types of stars in Fig. 3. Moreover, the high luminosity of the SN itself can potentially accelerate the pre-shock CSM (essentially a radiatively driven wind from a temporarily much more

(continued)

luminous star), which could also lead to a faster pre-shock CSM speed than expected for a certain type of star, or multiple velocity components in the CSM.

## 6.2 Direct Detections

The types of inferences described above are all still based on rather indirect and circumstantial evidence. A potentially more direct way to link SNe to their progenitors is to detect the progenitor star in pre-explosion imaging data, usually with the *Hubble Space Telescope*, although ground-based imaging has been used for some very nearby cases (see review by Smartt 2009). This method has been used successfully for some other types of SNe, especially SNe II-P, SNe I Ib, and one particularly famous II-pec event that occurred 20 years ago. The progenitor identification can be confirmed after the SN fades, to verify that the progenitor star is gone (as opposed to being a chance alignment, a host cluster, or a companion star).

For interacting SNe, however, the interpretation of pre-SN direct detections can be a little tricky. First, the “direct” detection of the progenitor might actually be a direct detection of a pre-SN eruption and not the quiescent star. This source might indeed fade after the SN, but it is hard to tell if the star is really dead or if the eruption has just subsided and the star returned to its quiescent state. Inferring an initial mass by comparison with stellar evolution tracks is also complicated if the progenitor might be in an outburst rather than in its quiescent state (we do not have much choice here and are lucky if there is even one *HST* image in the archive, but one just needs to be aware of the caveat). A second complication is that some SNe IIn have persistent CSM interaction for years or decades after the SN, and so one might need to wait a very long time before the SN has faded enough to be fainter than the progenitor. Third, a faint progenitor or even a faint upper limit to a progenitor is very inconclusive in terms of its implication for the mass of the star. This is because with interacting SNe, there is, by definition, a large mass of CSM. Thus, we certainly expect some cases where the progenitor was surrounded by a massive dust shell that should obscure the progenitor star’s visible light output. This makes it difficult to place a progenitor detection (or upper limit) on an HR diagram and infer an initial mass if one has only an optical filter.

Despite this ambiguity, there have been a few lucky and important cases that guide our intuition about the progenitors of SNe IIn and SNe Ibn.

**SN 2005gl:** This was a SN IIn where a very luminous progenitor consistent with an LBV star like P Cygni was detected in *HST* imaging, and the SN had an implied mass-loss rate of  $0.01 M_{\odot} \text{ yr}^{-1}$  (Gal-Yam et al. 2007), and this source then faded after the SN (Gal-Yam and Leonard 2009). Moreover, the pre-shock CSM speed of  $420 \text{ km s}^{-1}$  inferred from narrow H lines was suggestive of the fast outflow from an

LBV (Smith et al. 2010a). As noted above, there is some ambiguity as to whether the pre-SN detection was a massive quiescent LBV-like star or a pre-SN eruption caught by the *HST* image. In either case an eruptive LBV-like star is likely, although the implied initial mass may be different. Based on its spectral evolution, SN 2005gl fits into the subclass of “transitional” SNe IIn like SN 1998S and PTF11iqb, which show a Type IIn spectrum at first but later show broad P Cygni lines indicative of a SN photosphere, so this can be taken as an argument that it was most likely a core-collapse SN event.

**SN 1961V:** Long considered an extreme LBV or SN impostor event, recent arguments favor an actual core-collapse SN for the 1961 transient (Kochanek et al. 2011; Smith et al. 2011a). If this was a SN, then it has one of the best documented progenitor detections and progenitor variability among SNe, and it holds the record for the most massive directly detected SN progenitor. The pre-1961 variability suggests a very massive  $\sim 100 M_{\odot}$  blue LBV-like progenitor that was variable before the SN. The source at the SN position is now more than 5.5 mag fainter than this progenitor (a much more dramatic drop than in the case of SN 2005gl), and there is no IR source with a luminosity comparable to the progenitor (Kochanek et al. 2011), so the extremely massive star is likely dead.

**SN 2006jc:** An eruption in 2004 was noted as a possible LBV or SN impostor, and then a SN occurred at the same position 2 years later. The pre-SN outburst had a peak luminosity similar to SN impostors (Pastorello et al. 2007). The SN explosion 2 years later was of Type Ibn with strong narrow He I emission lines (Foley et al. 2007; Pastorello et al. 2007). There is no detection of the quiescent progenitor, but this unusual case implies an LBV-like eruption that occurred in a WR-like progenitor star that was clearly H depleted. The CSM speed was of order  $1000 \text{ km s}^{-1}$ , which is consistent with WR stars, and faster than typical LBVs.

**SN 2011ht:** This belongs to the subclass of SNe IIn-P, which are thought to arise from lower-energy explosions that may be linked to electron capture SNe in 8–10  $M_{\odot}$  super-AGB stars (Mauerhan et al. 2013b; Smith 2013). There is no detection of the quiescent progenitor, although deep upper limits seem to rule out a luminous, blue quiescent star (Mauerhan et al. 2013b; Roming et al. 2012). However, Fraser et al. (2013b) reported the detection of pre-SN eruption activity in archival data. This may be an important demonstration that non-LBVs can have violent pre-SN eruptions as well.

**SN 2010jl:** This was a SLSN IIn with roughly 10  $M_{\odot}$  or more of CSM. Smith et al. (2011c) identified a candidate source at the location of the SN in pre-explosion HST images that suggested either an extremely massive progenitor star or a very young massive-star cluster; in either case it seems likely that the progenitor had an initial mass above 30  $M_{\odot}$ . We are still waiting for this SN to fade to see if the progenitor candidate was actually the progenitor or a massive young cluster/association.

**SN 2009ip:** This source was initially discovered as an LBV-like outburst in 2009 (its namesake) before finally exploding as a much brighter SN in 2012. A quiescent progenitor star was detected in archival HST data, indicating a very massive 50–80  $M_{\odot}$  progenitor (Foley et al. 2011; Smith et al. 2011b). In this case, the

*HST* detection may well have been the quiescent progenitor, rather than an outburst, because much brighter outbursts came later. It showed slow variability consistent with an S Dor LBV-like episode (Smith et al. 2010b), followed by a series of brief LBV-like giant eruptions (Mauerhan et al. 2013a; Pastorello et al. 2013; Smith et al. 2010b). Unlike any other object, we also have detailed, high-quality spectra of the pre-SN eruptions (Foley et al. 2011; Smith et al. 2010b). The presumably final SN explosion of SN 2009ip in 2012 would fall into the “delayed onset” subclass, since at first the fainter transient showed very broad lines indicative of a SN photosphere. Reaching peak, however, it looked like a normal SN IIn, as the fast ejecta crashed into the slow material ejected 1–3 years earlier (Mauerhan et al. 2013a; Smith et al. 2014). A number of detailed studies of the bright 2012 transient have now been published, although there has been some controversy about whether the 2012 event was a core-collapse SN (Mauerhan et al. 2013a; Ofek et al. 2013; Prieto et al. 2013; Smith et al. 2014) or some type of extremely bright nonterminal event (Fraser et al. 2013a; Margutti et al. 2014; Pastorello et al. 2013). More recently, Smith et al. (2014) showed that the object continues to fade, and its late-time emission is consistent with late-time CSM interaction in normal SNe IIn. If SN 2009ip was indeed a SN, it provides a strong case that very massive stars above  $30 M_{\odot}$  may in fact experience core-collapse and explode and that LBV-like stars are linked to some SNe IIn.

### 6.3 Links to Progenitor Types

We will undoubtedly find more examples of direct detections and pre-SN outbursts in the future. One must bear in mind, though, that LBVs and eruptive precursors are relatively easy to detect because they are brighter than any quiescent stars, so these cases do not rule out alternatives such as dust-enshrouded RSGs or faint and hot quiescent stars. From various clues described above such as the CSM mass and mass-loss rate, CSM expansion speed, H-rich composition, and direct detections of progenitors or environments, we can attempt to link certain subclasses of interacting SNe to different possible progenitors. Some associations are more likely than others.

*SLSN IIn (compact and extended shell; SN 2006gy or SN 2010jl-like):* Based on the extreme required masses of ( $10\text{--}20 M_{\odot}$ ) of H-rich CSM, typically expanding at a few hundred  $\text{km s}^{-1}$ , it seems very likely that the progenitors of SLSNe IIn are very massive, eruptive LBV-like stars (see review by Smith 2014). If they are not truly LBVs, then they do a very good impersonation. The simple fact that very massive stars above  $\sim 40 M_{\odot}$  are exploding as H-rich SNe is a challenge to understand, since stellar evolution models predict all those stars to shed their H envelopes at roughly solar metallicity. Most of these have huge mass ejections occurring just a few years or decades before the SN, so the connection to nearby LBVs—some of which have shells that are hundreds or thousands of years old—is not yet clear. LBV-like progenitors are even likely in cases where the progenitor is optically faint, if a pre-SN eruption has obscured the star with a dust cocoon (as is likely to be the case, given the consequent CSM interaction). Since the progenitors must be very massive

stars simply because of the mass budget, the physical mechanism of pulsational pair instability eruptions is a viable candidate for the pre-SN outbursts (Woosley et al. 2007). This is not the case for the remainder of SNe IIn, because they are too common (Smith et al. 2014).

*Normal, enduring SNe IIn (SN 1988Z-like):* Likely progenitor types are LBVs or extreme cool hypergiants (YHGs/eRSGs), based mostly on the required mass-loss rates and observed CSM speeds. For these enduring cases, we require either strong mass loss in several centuries preceding the SN or a large bipolar shell ejected shortly before the SN with a range of ejection speeds (to accommodate CSM interaction over a large range of radii, lasting for a long time). Total CSM masses that exceed  $10 M_{\odot}$  in some cases and integrated radiated energies that exceed  $10^{51}$  ergs (over years) point to relatively massive progenitor stars. Enhanced late-phase RSG mass loss (Yoon and Cantiello 2010) or instabilities in late nuclear burning sequences (Quataert and Shiode 2012; Smith and Arnett 2014) are good candidates for the episodic mass loss.

*Transitional or fleeting SNe IIn (SN 1998S-like):* Likely progenitors are RSGs, YHGs, or BSG/LBVs with less extreme pre-SN bursts of mass loss, confined to a relatively short-duration event preceding core-collapse (i.e., within a few years preceding the SN). These objects also seem to require highly asymmetric CSM interaction, to allow the expanding SN photosphere to completely or mostly envelope the disk of CSM interaction that emits narrow lines (as discussed in Sect. 3). For this reason, there is a strong suspicion that close binary interaction plays a role in their pre-SN episodic mass loss, although it must still be linked somehow to the final nuclear burning sequences (see Smith and Arnett 2014 regarding the role of close binaries in this scenario).

*SNe IIn-P (SN 2011ht-like):* The favored scenario for these events involves an intermediate-mass progenitor in a super-AGB phase that suffers a sub-energetic ( $10^{50}$  erg) electron-capture SN. This is based on the low  $^{56}\text{Ni}$  yield, the deep absorption features that imply more-or-less spherical symmetry, and the energy/mass budgets inferred (Mauerhan et al. 2013b; Smith 2013). However, observations cannot yet confidently rule out a more massive progenitor that suffers fallback to a black hole, yielding a smaller ejecta mass and very low  $^{56}\text{Ni}$  yield (although in the case of SN 2011ht, progenitor upper limits seem to argue against this higher-mass interpretation). This type of SN also fits the bill for SN 1054 and the Crab Nebula (Smith 2013), whose abundances and kinetic energy seem to point to an electron capture SN from an intermediate-mass star (Nomoto et al. 1982). If these are ecSNe, then the pre-SN episodic mass loss might be related to nuclear flashes in the advanced degenerate core-burning sequences.

*SN IIn impostors:* This group of putatively non-SN transients may be quite diverse, and it may include transients that have narrow lines because they are powered largely by CSM interaction (and weaker explosions than core-collapse SNe) and other transients that have narrow lines because they have slow winds/outflows. This may include LBVs, super AGB stars, binaries with a compact object, stellar mergers, pre-SN nuclear burning instabilities, failed SNe, or all of the above. Any massive supergiant star enshrouded in dust or with strong binary interaction that

suffers an instability is a viable candidate. The theoretical mechanisms for this class of outbursts are still very poorly understood, and there is probably considerable overlap with pre-SN outbursts that lead to SNe IIn and Ibn.

*SNe Ibn:* The most likely progenitors are massive WR stars that for unknown reasons undergo pre-SN outbursts. This is interesting, because there is no known precedent for an observed eruption in a H-deficient massive star.

*SNe Ia/IIn or Ia-CSM:* If these really are thermonuclear SNe Ia (some cases seem clear; others are still debated in the literature), then the exploding progenitors are white dwarfs that have arisen from initial masses below roughly  $8 M_{\odot}$ . In this case, a single degenerate system is clearly required to supply the large mass of H-rich CSM (several  $M_{\odot}$  in some cases). There is, as yet, no viable explanation for the sudden ejection of a large mass of H (by the companion) shortly preceding the thermonuclear explosion of the WD.

---

## 7 Closing Comments

Observations of interacting SNe present one of the most interesting challenges to our understanding of the end phases of evolution for massive stars. What makes these stars explode before they explode? Computational resources are only beginning to meet the needs of the complex problem of simulating convection and nuclear burning coupled with stellar structure and turbulence in these final phases (Arnett and Meakin 2011; Meakin and Arnett 2007). The fact that  $\sim 10\%$  of core-collapse events are preceded by some major reorganization of the stellar structure and energy budget tells us that we have been missing something important, which might be a key ingredient for understanding core-collapse SNe more generally. It will be important to try and understand if this 10% corresponds to the most extreme manifestation of a wider instability (e.g., if all stars undergo some instability in the final burning sequences, only the most extreme cases lead to detectably violent mass loss and SNe IIn) or if SNe IIn are the outcome of special circumstances in a particular evolution channel (i.e., interacting binaries within a certain mass and orbital period range). There is still very little information available on any trends with metallicity, although this is always good to investigate when mass loss plays a critical role.

Recent years have seen something of a paradigm shift in massive star studies. Formerly standard or straightforward assumptions about the simplicity of single-star evolution are giving way to more complicated scenarios as astronomers grudgingly admit that binary stars not only exist but are common and influential (e.g., Sana et al. 2012). This may be especially true among transient sources and stellar deaths that seem otherwise peculiar or difficult to understand. Given the very high multiplicity fraction among massive stars, binary interaction should probably not be considered a last resort or the refuge for an uncreative theorist but, rather, a default assumption.

Whether they are binaries or not, all SNe IIn require some major shift in stellar structure and mass loss before the SN. The synchronization with core-collapse gives a strong implication that something wild is happening in the latest sequences of



nuclear burning. Those cases where a star changes its structure as a result of these nuclear burning instabilities (i.e., an inflated envelope) in a close binary system seem promising for inducing sudden eruptive behavior for various reasons (Smith and Arnett 2014). Single stars may also be able to induce their own violent mass loss in the couple years preceding core-collapse due to energy transported to the envelope from Ne and O burning zones (Quataert and Shiode 2012). However, we do not yet have a good explanation in single-star evolution for the strong mass loss that leads to the “enduring” class of SNe IIn or some of the more extreme cases of “delayed onset” of CSM interaction, where the strong interaction that lasts for years after explosion suggests very strong mass loss for *decades or centuries* before core-collapse. If binary interaction is an important ingredient, then this sort of interaction before a SN might make asymmetric CSM very common, suggesting that we should be paying close attention to possible observed signatures of asymmetry.

---

## 8 Cross-References

- ▶ [Dust and Molecular Formation in Supernovae](#)
- ▶ [Electron Capture Supernovae from Super Asymptotic Giant Branch Stars](#)
- ▶ [Interacting Supernovae: Spectra and Light Curves](#)
- ▶ [Observational and Physical Classification of Supernovae](#)
- ▶ [Supernovae from Massive Stars](#)
- ▶ [Supernova of 1054 and its Remnant, the Crab Nebula](#)
- ▶ [Supernova Progenitors Observed with \*HST\*](#)
- ▶ [The Physics of Supernova 1987A](#)
- ▶ [The Progenitor of SN 1987A](#)
- ▶ [Thermal and Non-thermal Emission from Circumstellar Interaction](#)
- ▶ [The Supernova – Supernova Remnant Connection](#)

**Acknowledgements** My attempt to understand interacting SNe and their connections to massive stars has benefitted greatly from conversations with numerous people but especially Dave Arnett, Matteo Cantiello, Nick Chugai, Selma de Mink, Ori Fox, Morgan Fraser, Dan Kasen, Jon Mauerhan, Stan Owocki, Jose Prieto, Eliot Quataert, Jorick Vink, and Stan Woosley. While drafting this chapter, I received support from NSF grants AST-1312221 and AST-1515559.

---

## References

- Arctxaga I, Benetti S, Terlevich RJ, Fabian AC, Cappellaro E, Turatto M et al (1999) SN 1988Z: spectro-photometric catalogue and energy estimates. MNRAS 309:343
- Arnett WD, Meakin C (2011) Turbulent cells in stars: fluctuations in kinetic energy and luminosity. ApJ 741:33
- Benetti S, Cappellaro E, Turatto M, Tautenberger S, Haratyunyan A, Valenti S (2006) Supernova 2002ic: the collapse of a stripped-envelope, massive star in a dense medium? ApJ 653:L129
- Chugai NN (1997) Supernovae in dense winds. Ap&SS 252:225
- Chugai NN (2001) Broad emission lines from the opaque electron-scattering environment of SN 1998S. MNRAS 326:1448



- Chugai NN, Danziger IJ (1994) SN 1988Z: low-mass ejecta colliding with the clumpy wind? *MNRAS* 268:173
- Chugai NN, Blinnikov SI, Cumming RJ, Lundqvist P, Bragaglia A, Filippenko AV et al (2004) The Type IIn supernova 1994W: evidence for the explosive ejection of a circumstellar envelope. *MNRAS* 352:1213
- Dessart L, Audi E, Hillier DJ (2015) Numerical simulations of superluminous supernovae of Type IIn. *MNRAS* 449:4304
- Filippenko AV (1997) Optical spectra of supernovae. *Annu Rev Astron Astrophys* 35:309
- Foley RJ, Smith N, Ganeshalingam M, Li W, Chornock R, Filippenko AV (2007) SN 2006jc: A Wolf-Rayet star exploding in a dense He-rich circumstellar medium. *ApJL* 657:L105
- Foley RJ, Berger E, Fox O, Levesque EM, Challis PJ, Ivans II et al (2011) The diversity of massive star outbursts. I. Observations of SN 2009ip, UGC 2773 OT2009-1, and their progenitors. *ApJ* 732:32
- Fox O, Skrutskie MF, Chevalier RA, Kanneganti S, Park C, Wilson J et al (2009) Near-infrared photometry of the Type IIn SN 2005ip: the case for dust condensation. *ApJ* 691:650
- Fransson C, Ergon M, Challis PJ, Chevalier RA, France K, Kirshner RP et al (2014) High-density circumstellar interaction in the luminous Type IIn SN 2010jl: the first 1100 days. *ApJ* 797:118
- Fraser M, Inserra C, Jerkstrand A, Kotak R, Pignata G, Benetti S et al (2013a) SN 2009ip à la PESSTO: no evidence for core collapse yet. *MNRAS* 433:1312
- Fraser M, Magee M, Kotak R, Smartt SJ, Smith KW, Polshaw J et al (2013b) Detection of an outburst one year prior to the explosion of SN 2011ht. *ApJ* 779:L8
- Gall C, Hjorth J, Watson D, Dwek E, Maund JR, Fox O et al (2014) Rapid formation of large dust grains in the luminous supernova 2010jl. *Nature* 511:326
- Gal-Yam A, Leonard DC (2009) A massive hypergiant star as the progenitor of the supernova SN 2005gl. *Nature* 458:865
- Gal-Yam A, Leonard DC, Fox DB, Cenko SB, Soderberg AM, Moon DS et al (2007) On the progenitor of SN 2005gl and the nature of Type IIn supernovae. *ApJ* 656:372
- Gal-Yam A, Arcavi I, Ofek EO, Ben-Ami S, Cenko SB, Kasliwal MM et al (2014) A Wolf-Rayet-like progenitor of SN? 2013cu from spectral observations of a stellar wind. *Nature* 509:471
- Gehrz RD, Hackwell JA (1974) Circumstellar dust emission from WC9 stars. *ApJ* 194:619
- Kochanek CS, Szczygiel DM, Stanek KZ (2011) The supernova impostor impostor SN 1961V: Spitzer shows that Zwicky was right (again). *ApJ* 737:76
- Leonard DC, Filippenko AV, Barth AJ, Matheson T (2000) Evidence for asphericity in the Type IIn supernova SN 1998S. *ApJ* 536:239
- Margutti R, Milisavljevic D, Soderberg AM (2014) A Panchromatic view of the restless SN 2009ip reveals the explosive ejection of a massive star envelope. *ApJ* 780:21
- Matheson T, Filippenko AV, Barth AJ, Ho LC, Leonard DC, Bershadsky MA et al (2000) Optical spectroscopy of supernova 1993J during its first 2500 days. *AJ* 120:1487
- Mauerhan JC, Smith N, Filippenko AV, Blanchard KB, Blanchard PK, Casper CFE et al (2013a) The unprecedented 2012 outburst of SN 2009ip: a luminous blue variable star becomes a true supernova. *MNRAS* 430:1801
- Mauerhan JC, Smith N, Silverman JM, Filippenko AV, Morgan AN, Cenko SB et al (2013b) SN 2011ht: confirming a class of interacting supernovae with plateau light curves (Type IIn-P). *MNRAS* 431:2599
- Mauerhan JC, Williams GG, Smith N, Smith PS, Filippenko AV, Hoffman JL et al (2014) Multi-epoch spectropolarimetry of SN 2009ip: direct evidence for aspherical circumstellar material. *MNRAS* 442:1166
- Meakin C, Arnett WD (2007) Turbulent convection in stellar interiors. I. Hydrodynamic simulation. *ApJ* 667:448
- Milisavljevic D, Fesen RA (2013) A detailed kinematic map of cassiopeia A's optical main shell and outer high-velocity ejecta. *ApJ* 772:134

- Miller A, Silverman JM, Butler NR, Bloom JS, Chornock R, Filippenko AV et al (2010) SN 2008iy: an unusual Type II<sub>n</sub> supernova with an enduring 400-d rise time. *MNRAS* 404:305
- Niemela VS, Ruiz MT, Phillips MM (1985) The supernova 1983k in NGC 4699 – Clues to the nature of Type II progenitors. *ApJ* 289:52
- Nomoto K, Sugimoto D, Sparks WM, Fesen RA, Gull TR, Miyaji S (1982) The Crab Nebula's progenitor. *Nature* 299:803
- Ofek EO, Cameron PB, Kasliwal MM, Gal-Yam A, Rau A, Kulkarni SR et al (2007) SN 2006gy: an extremely luminous supernova in the galaxy NGC 1260. *ApJ* 659:L13
- Ofek EO, Lin L, Kouveliotou C, Younes G, Gogus E, Kasliwal MM, Cao Y (2013) SN 2009ip: constraints on the progenitor mass-loss rate. *ApJ* 768:47
- Pastorello A, Smartt SJ, Mattila S, Eldridge JJ, Young D, Itagaki K et al (2007) A giant outburst two years before the core-collapse of a massive star. *Nature* 447:829
- Pastorello A, Mattila S, Zampieri L, Della Valle M, Smartt SJ, Valenti S, Agnoletto I et al (2008) Massive stars exploding in a He-rich circumstellar medium – I. Type Ib<sub>n</sub> (SN 2006jc-like) events. *MNRAS* 389:113
- Pastorello A, Cappellaro E, Inserra C, Smartt SJ, Pignata G, Benetti S et al (2013) Interacting supernovae and supernova impostors: SN 2009ip, is this the end? *ApJ* 767:1
- Pastorello A, Benetti S, Brown PJ, Tsvetkov DY, Inserra C, Taubenberger S et al (2015) Massive stars exploding in a He-rich circumstellar medium – IV. Transitional Type Ib<sub>n</sub> supernovae. *MNRAS* 449:1921
- Pozzo M, Meikle WPS, Fassia A, Geballe T, Lundqvist P, Chugai NN et al (2004) On the source of the late-time infrared luminosity of SN 1998S and other Type II supernovae. *MNRAS* 352:457
- Prieto JL, Brimacombe J, Drake AJ, Howerton S (2013) The 2012 rise of the remarkable Type II<sub>n</sub> SN 2009ip. *ApJL* 763:L27
- Quataert E, Shiode J (2012) Wave-driven mass loss in the last year of stellar evolution: setting the stage for the most luminous core-collapse supernovae. *MNRAS* 423:L92
- Rest A, Foley RJ, Gezari S, Narayan G, Draine B, Olsen K et al (2011) Pushing the boundaries of conventional core-collapse supernovae: the extremely energetic supernova SN 2003 ma. *ApJ* 729:88
- Roming PWA, Pritchard TA, Prieto JL, Kochanek CS, Fryer CL, Davidson K et al (2012) The unusual temporal and spectral evolution of the TYPE II<sub>n</sub> supernova 2011ht. *ApJ* 751:92
- Sana H, de Mink SE, de Koter A, Langer N, Evans CJ, Giele M et al (2012) Binary interaction dominates the evolution of massive stars. *Science* 337:444
- Schlegel EM (1990) A new subclass of Type II supernovae? *MNRAS* 244:269
- Shivvers I, Mauerhan JC, Leonard DC, Filippenko AV, Fox OD (2015) Early emission from the Type II<sub>n</sub> supernova 1998s at high resolution. *ApJ* 806:213
- Silverman JM, Nugent PE, Gal-Yam A, Sullivan M, Howell DA et al (2013) Type Ia supernovae strongly interacting with their circumstellar medium. *ApJ Suppl* 207:3
- Smartt SJ (2009) Progenitors of core-collapse supernovae. *ARA&A* 47:63
- Smith N (2013) The Crab Nebula and the class of Type II<sub>n</sub>-P supernovae caused by sub-energetic electron-capture explosions. *MNRAS* 434:102
- Smith N (2014) Mass loss: its effect on the evolution and fate of high-mass stars. *ARA&A* 52:487
- Smith N, Arnett WD (2014) Preparing for an explosion: hydrodynamic instabilities and turbulence in presupernovae. *ApJ* 785:82
- Smith N, Li W, Foley RJ, Wheeler JC, Pooley D, Chornock R et al (2007) SN 2006gy: Discovery of the most luminous supernova ever recorded, powered by the death of an extremely massive star like  $\eta$  carinae. *ApJ* 666:1116
- Smith N, Chornock R, Li W, Ganeshalingam M, Silverman JM, Foley RJ et al (2008a) SN 2006tf: Precursor eruptions and the optically thick regime of extremely luminous Type II<sub>n</sub> supernovae. *ApJ* 686:467
- Smith N, Foley RJ, Filippenko AV (2008b) Dust formation and He II  $\lambda$ 4686 emission in the dense shell of the peculiar Type Ib supernova 2006jc. *ApJ* 680:568

- Smith N, Silverman JM, Chornock R, Filippenko AV, Wang X et al (2009) Coronal lines and dust formation in SN 2005ip: not the brightest, but the hottest Type IIn supernova. *ApJ* 695:1334
- Smith N, Chornock R, Silverman JM, Filippenko AV, Foley RJ (2010a) Spectral evolution of the extraordinary Type IIn supernova 2006gy. *ApJ* 709:856
- Smith N, Miller A, Li W, Filippenko AV, Silverman JM, Howard AW et al (2010b) Discovery of precursor luminous blue variable outbursts in two recent optical transients: the fitfully variable missing links UGC 2773-OT and SN 2009ip. *AJ* 139:1451
- Smith N, Li W, Filippenko AV, Chornock R (2011a) Observed fractions of core-collapse supernova types and initial masses of their single and binary progenitor stars. *MNRAS* 412:1522
- Smith N, Li W, Silverman JM, Ganeshalingam M, Filippenko AV (2011b) Luminous blue variable eruptions and related transients: diversity of progenitors and outburst properties. *MNRAS* 415:733
- Smith N, Mauerhan JC, Silverman JM, Ganeshalingam M, Filippenko AV, Cenko SB et al (2012a) SN 2011hw: helium-rich circumstellar gas and the luminous blue variable to Wolf-Rayet transition in supernova progenitors. *MNRAS* 426:1905
- Smith N, Silverman JM, Filippenko AV, Cooper MC, Matheson T, Bian F et al (2012b) Systematic blueshift of line profiles in the Type IIn supernova 2010jl: evidence for post-Shock dust formation? *AJ* 143:17
- Smith N, Mauerhan JC, Prieto JL (2014) SN 2009ip and SN 2010mc: core-collapse Type IIn supernovae arising from blue supergiants. *MNRAS* 438:1191
- Smith N, Mauerhan JC, Cenko SB, Kasliwal MM, Silverman JM, Filippenko AV et al (2015) PTF11iqb: cool supergiant mass-loss that bridges the gap between Type IIn and normal supernovae. *MNRAS* 449:1876
- Stritzinger M, Taddia F, Fransson C, Fox OD, Morrell N, Phillips MM et al (2012) Multi-wavelength observations of the enduring Type IIn supernovae 2005ip and 2006jd. *ApJ* 756:173
- van Marle AJ, Smith N, Owocki SP, van Veelen B (2010) Numerical models of collisions between core-collapse supernovae and circumstellar shells. *MNRAS* 407:2305
- Williams PM, van der Hucht KA, Pollock AMT, Florkowski DR, van der Woerd H, Wamsteker WM (1990) Multi-frequency variations of the Wolf-Rayet system HD 193793. I – infrared, X-ray and radio observations. *MNRAS* 243:662
- Woosley SE, Blinnikov S, Heger A (2007) Pulsational pair instability as an explanation for the most luminous supernovae. *Nature* 450:390
- Yoon SC, Cantiello M (2010) Evolution of massive stars with pulsation-driven superwinds during the red supergiant phase. *ApJ* 717:L62