

Coven–Meyerowitz T2 necessity through coprime stripe collapse

Public research artifact*

10 September 2026

Abstract

We present a proof of the necessity of the Coven–Meyerowitz condition T2 for every finite set that tiles the integers by translations. The proof passes through actual Boolean cyclic tilings, retains the carry in prime-residue fibers, and obtains periodic row-or-column product stripes. Integral descent of convolution minors to a coprime quotient, followed by characteristic-prime Frobenius, forces a single stripe orientation. The resulting actual lower-period tilings exist for every independent phase, allowing strong induction to recover mixed cyclotomic zeros of both original factors. The associated Lean 4.23.0 development exposes the unrestricted integer-tiling statement and includes an explicit 35-module replay and 22 guarded axiom checks. The permitted final axioms are propositional extensionality, classical choice, and quotient soundness. This is a public research release for external review; specialist acceptance and exclusive discovery priority remain unconfirmed.

1 Statement and scope

For a finite nonempty integer set A , first translate so that all entries are nonnegative and write

$$A(X) = \sum_{a \in A} X^a, \quad S_A = \{p^e : p \text{ prime}, e \geq 1, \Phi_{p^e}(X) \mid A(X)\}.$$

Divisibility is in $\mathbb{Z}[X]$. Translation does not change any of these cyclotomic divisibilities. Condition T2 says that $\Phi_{s_1 \dots s_k}$ divides $A(X)$ whenever $s_1, \dots, s_k \in S_A$ have distinct underlying primes. Here $k \geq 1$; the empty product is not required.

Theorem 1. *Suppose a finite set $A \subseteq \mathbb{Z}$ tiles $\mathbb{Z}$ uniquely by translations. Then A satisfies T2, without restriction on the number of selected primes, their exponents, the cardinality of A , or a tiling period.*

The public formal statement for nonnegative A starts with an arbitrary function $c : \mathbb{Z} \rightarrow \mathbb{N}$ and the equations

$$\sum_{a \in A} c(t - a) = 1 \quad (t \in \mathbb{Z}). \tag{1}$$

No periodicity assumption is imposed on c . Nonemptiness of A and Booleanity of c follow from these equations. The arbitrary-integer-set statement uses the ordinary polynomial $\sum_{a \in A} X^{a - \min A}$.

*Prepared with AI assistance. No personal author list, affiliation, external referee endorsement, or exclusive priority claim is assigned by this automated release. Maintained at <https://github.com/infinityscroll/coven-meyerowitz-t2>.

Coven and Meyerowitz established that T1 is necessary for tiling and that T1 together with T2 is sufficient; they also proved T2 necessity when the tile cardinality has at most two distinct prime factors [1]. Theorem 1 supplies the remaining necessity statement. The previously established T1 and sufficiency results are cited here, not newly formalized by this artifact. A July 2026 source still explicitly describes unrestricted T2 necessity as open [4]; this is evidence of status in that text, not a certificate of exclusive priority for the present argument.

2 Cyclic masks and primary allocation

Write J_N for the all-ones mask on $\mathbb{Z}_N$. Identify a finite cyclic set with its Boolean mask in the integral group ring $\mathbb{Z}[\mathbb{Z}_N]$, with multiplication given by convolution. Thus an actual unique cyclic tiling is

$$AB = J_N. \quad (2)$$

The symbol J_N is not the multiplicative identity. Use canonical residues to form ordinary mask polynomials. Evaluation at all N th roots separates group-ring elements by finite Fourier inversion.

Lemma 2 (Exact primary allocation). *For an actual cyclic tiling $A + B = \mathbb{Z}_N$, the sets S_A and S_B partition the prime powers dividing N . If q is prime, the number of q -primary levels owned by each factor is respectively $v_q(|A|)$ and $v_q(|B|)$. Neither factor has a primary cyclotomic divisor whose order does not divide N .*

Proof. For $1 \leq e \leq v_q(N)$, evaluation of (2) at a primitive q^e root makes at least one factor vanish. An integer polynomial vanishing at such a root is divisible by its monic cyclotomic polynomial over $\mathbb{Z}$. Distinct cyclotomic divisors multiply to a divisor. Evaluating that product at 1, and using $\Phi_{q^e}(1) = q$, shows that the numbers n_A, n_B of owned levels satisfy

$$v_q(N) \leq n_A + n_B \leq v_q(|A|) + v_q(|B|) = v_q(N).$$

All inequalities are equalities, so each level is owned exactly once and both cardinality valuations are exhausted. Any further q -primary divisor would exceed this valuation. For $q \nmid N$, both cardinalities have q -valuation zero and cannot admit such a divisor. $\square$

In particular, every T2 candidate order of either factor divides the period. The allocation argument is standard CM background [1, Lemma 2.1]; it does not by itself establish mixed zeros.

3 Prime fibers, dephasing, and character identities

Fix a prime $p \mid N$, write $N = pM$, and interchange the factors if necessary so that B owns Φ_p . For $0 \leq i, j < p$ put

$$A_i = \{a \in \mathbb{Z}_M : i + pa \in A\}, \quad B_j = \{b \in \mathbb{Z}_M : j + pb \in B\}.$$

The B_j have the same positive cardinality $|B|/p$. Indeed, their count polynomial has degree at most $p - 1$ and vanishes at a primitive p th root, so it is a scalar multiple of $1 + z + \cdots + z^{p-1}$. The A_i cannot all have the same cardinality, by exclusive ownership of Φ_p .

Write $M = KR$, where K is the full p -primary part of M and $(p, R) = 1$. Choose ρ with $p\rho \equiv 1 \pmod{R}$; if $R = 1$, take $\rho = 0$. In $\mathbb{Z}[\mathbb{Z}_M]$ define

$$U_i = T^{\rho i} A_i, \quad V_j = T^{\rho j} B_j, \quad F_{ij} = U_i V_j.$$

The original cyclic carry is exactly retained by the identity

$$\left(\sum_i z^i A_i(T) \right) \left(\sum_j z^j B_j(T) \right) = J_M(T) \sum_{r=0}^{p-1} z^r \quad \text{modulo } (T^M - 1, z^p - T). \quad (3)$$

The quotient is identified with $\mathbb{Z}[X]/(X^{pM} - 1)$ by $z = X$, $T = X^p$. It is not a replacement of the cyclic group by $\mathbb{Z}_p \times \mathbb{Z}_M$.

Lemma 3. *The masks F_{ij} are Boolean, invariant under translation by R , and satisfy*

$$F_{ij} + F_{kl} = F_{il} + F_{kj}, \quad \sum_{i,j} F_{ij} = pJ_M. \quad (4)$$

Proof. Let θ be an M th root of order $d > 1$. If $p \mid d$, a p th root ζ of θ has order pd , and

$$[\mathbb{Q}(\zeta) : \mathbb{Q}(\theta)] = \varphi(pd)/\varphi(d) = p.$$

Thus $z^p - \theta$ is irreducible over $\mathbb{Q}(\theta)$. At $T = \theta$, the right side of (3) vanishes. Each of the two fiber polynomials has degree below p , so irreducibility forces one entire coefficient family to vanish. Consequently every $F_{ij}(\theta) = 0$.

If $(p, d) = 1$, then $d \mid R$. Set $\lambda = \theta^\rho$, so $\lambda^p = \theta$ and λ has order d . Substituting $z = \lambda w$ in (3) makes

$$\left(\sum_i U_i(\theta) w^i \right) \left(\sum_j V_j(\theta) w^j \right) \quad \text{divisible by } w^p - 1.$$

Over $\mathbb{Q}(\theta)$, $\Phi_p(w)$ is irreducible of degree $p - 1$. Unless one factor is zero, one is a scalar multiple of Φ_p and hence has a constant coefficient family. It follows that every rectangular difference vanishes at θ . Evaluation at $w = 1$ gives the vanishing of the total product there. The corresponding original root λ is nontrivial because $d > 1$.

At the trivial character the V_j masses are equal, so the rectangular differences again vanish. The total mass is $|A||B| = pM$. Fourier inversion proves (4). For a fixed pair i, j , two representations in $A_i + B_j$ would give two representations in the original tiling with the same digit carry; hence $A_i B_j$, and its translate F_{ij} , are Boolean. Their Fourier coefficients vanish at every order containing p ; the surviving orders divide R , which is exactly invariance under translation by R . $\square$

This includes $p = 2$. If $R = 1$ there are no nontrivial p -free characters; if $M = 1$ only the trivial-character and fixed-pair arguments are needed. Empty A_i are allowed. Crucially, the periodicity conclusion concerns the *products* $U_i V_j$, not the individual factors.

4 Boolean stripes and coprime collapse

At a fixed $x \in \mathbb{Z}_M$, the matrix $(F_{ij}(x))$ is Boolean, has zero additive rectangular differences, and contains exactly p ones. Such a matrix is one complete row or one complete column of ones. To see this, if a row has unequal entries in two columns, the same nonzero difference occurs in every row; Booleanity forces those columns, and then all columns, to be constant. If no row varies, every row is constant. The total p leaves exactly one stripe.

Consequently there are pairwise disjoint stripe sets X_i, Y_j partitioning $\mathbb{Z}_M$ such that

$$F_{ij} = X_i + Y_j. \quad (5)$$

Each stripe mask is R -periodic, since it is uniquely decoded from the periodic matrix. Commutativity also gives the convolution minors

$$F_{ij}F_{kl} = F_{il}F_{kj}. \quad (6)$$

Lemma 4 (Coprime stripe collapse). *Let p be prime and let G be a finite abelian group with $p \nmid |G|$. Suppose Boolean stripe masks $X_0, \dots, X_{p-1}, Y_0, \dots, Y_{p-1}$ partition G and $F_{ij} = X_i + Y_j$ satisfy (6). Then all X_i are empty or all Y_j are empty.*

Proof. Expand the minors to obtain $(X_i - X_k)(Y_j - Y_l) = 0$. With $X = \sum_i X_i$ and $Y = \sum_j Y_j$, sum over k, l to get

$$(pX_i - X)(pY_j - Y) = 0.$$

Reducing in the commutative group algebra $\mathbb{F}_p[G]$ gives $XY = 0$. Since $X + Y = J_G$ and $XJ_G = |X|J_G$, we get $X^2 = |X|J_G$, and therefore

$$X^p = |X|^{p-1}J_G.$$

Frobenius, however, gives $X^p = \sum_{x \in X} [px]$. Multiplication by p permutes G , so the latter is a permutation of the original Boolean coefficient mask. A Boolean mask constant modulo p is constant as an integer mask. Hence X is empty or all of G , as required. $\square$

To apply the lemma here, first descend the R -periodic masks from $\mathbb{Z}_M$ to $\mathbb{Z}_R$. For the lift L of masks along reduction modulo R , a direct count of the $K = M/R$ lifts of a residue gives

$$L(f)L(g) = KL(fg). \quad (7)$$

Equation (6) therefore descends after cancellation of the nonzero integer K . This cancellation occurs in the integral group ring *before* reduction modulo p ; no division by K in characteristic p is used. The quotient stripes remain a Boolean partition and $(p, R) = 1$, so Lemma 4 applies.

Column-only would make $U_iV_j = Y_j$ independent of i . Taking masses and using the common positive mass of V_j would make every $|A_i|$ equal, contradicting primary allocation. Thus the orientation is row-only:

$$U_iV_j = X_i, \quad \sum_i X_i = J_M. \quad (8)$$

5 Actual lower tilings for every phase

For an arbitrary integer phase vector $h = (h_0, \dots, h_{p-1})$, put

$$C_h = \sum_i T^{Rh_i} U_i.$$

By (8) and product periodicity,

$$C_h V_j = J_M \quad \text{for all } j, h. \quad (9)$$

The mask C_h has nonnegative integer coefficients. Since V_j is a nonempty Boolean mask, a coefficient of C_h at least two would produce a convolution coefficient at least two, contradicting (9). Thus C_h is Boolean and (9) is an actual tiling of the strictly smaller period M . Translating the complement back also yields $C_h B_j = J_M$. We will use all phases, not just a single chosen quotient.

6 Strong induction for both original factors

Theorem 5. *For every $N \geq 1$, both factors in an actual cyclic tiling $A + B = \mathbb{Z}_N$ satisfy T2.*

Proof. Use strong induction on N . At $N = 1$ both masks are the singleton identity and have no primary zeros. For $N > 1$, choose $p \mid N$ and the labeling above. The induction hypothesis applies to every tiling (9).

The original factor A . For a p -free root θ whose order divides M , dephasing gives

$$C_h(\theta) = \sum_i \theta^{\rho i} A_i(\theta) = A(\theta^\rho), \quad (10)$$

since $\theta^R = 1$ and $(\theta^\rho)^p = \theta$. Every selected p -free primary zero of A thus passes to every C_h . If the target order d is p -free, lower T2 for C_0 and (10), with $\theta = \zeta^p$ for a primitive original root ζ , give $A(\zeta) = 0$.

If the selected order is $p^k d$ with $(p, d) = 1$, then $k \geq 2$, because A does not own Φ_p . The identity

$$\Phi_{p^k}(X) = \Phi_{p^{k-1}}(X^p)$$

and monic division by exponent residues modulo p show that every A_i is divisible by $\Phi_{p^{k-1}}$. Explicitly, write the integer quotient of $A(X)$ by the displayed monic divisor as $\sum_i X^i Q_i(X^p)$ and compare residue classes. Thus every C_h inherits the lower p -primary zero as well as the selected p -free zeros.

For a primitive root ζ of order $p^k d$, let $\theta = \zeta^p$, of order $p^{k-1} d \mid M$. Lower T2 gives $C_h(\theta) = 0$ for every h . Comparing the zero phase with the phase changing only coordinate i by one yields

$$\theta^{\rho i} (\theta^R - 1) A_i(\theta) = 0.$$

Here θ^R has order $p^{k-1} > 1$, so every $A_i(\theta) = 0$. Regrouping gives $A(\zeta) = \sum_i \zeta^i A_i(\zeta^p) = 0$. This is recovery of the original upper- p mixed zero, not a claim that a single quotient preserves it.

The original factor B . At zero phase, $C_0 B_j = J_M$ for every j . Hence the dilated set $pC_0 = \{pc : c \in C_0\} \subseteq \mathbb{Z}_{pM}$ is an actual complement of original B . Here dilation has mask $C_0(X^p)$; it is not scalar multiplication of coefficients by p . All lower B_j have T2 by induction and share the same primary allocation, since their actual complement is the same C_0 .

Every p -free primary zero of B belongs to every B_j : compare exact allocation in $B + pC_0 = \mathbb{Z}_{pM}$ with that in $B_j + C_0 = \mathbb{Z}_M$. At a p -free prime-power order, the polynomial of pC_0 vanishes precisely when that of C_0 does, because raising a primitive root to its p th power preserves its order. Allocation then forces the zero into B_j .

If B has a higher primary zero p^k , $k \geq 2$, residue grading likewise puts p^{k-1} in every B_j . For a p -free target d , lower T2 makes every $B_j(\zeta^p)$ zero and hence $B(\zeta) = 0$. For a target $p^k d$ with $k \geq 2$, apply the same argument at the lower target $p^{k-1} d$. For $k = 1, d > 1$, use the inherited lower target d . The remaining case $k = 1, d = 1$ is the already known Φ_p zero. These cases exhaust all T2 requirements of both original factors by Lemma 2. $\square$

7 From integer tilings to cyclic tilings

Normalize the integer tile so that $\min A = 0$, and put $L = \max A$. Equation (1) implies that c is Boolean: one term bounds $c(s)$ by 1 for every s . If $L = 0$ the theorem is immediate. Otherwise

the terms $0, L \in A$ make the recurrence deterministic in both directions: $c(t)$ is determined by its preceding L bits, and the earliest bit in a window of length $L + 1$ is determined by the following L bits.

There are finitely many L -bit states. A repeated state gives a positive shift N_0 preserving the whole bi-infinite sequence by forward and backward determinism. Replace N_0 by a multiple $N > L$ if desired; it is still a period. Reduction of A modulo N is then injective, and the residue support B of c gives an actual cyclic tiling. With $N > L$, the canonical ordinary polynomial of the reduced A is exactly the original polynomial. Theorem 5 proves T2. Translating back proves Theorem 1 for arbitrary finite integer sets. The Lean implementation chooses a support-exceeding period in this way.

8 Formal statement and reproducible workflow

The final declaration is

`FullCovenMeyerowitz.integer_tile_cyclotomic_product.`

Its explicit Lean type quantifies over $E : \text{Finset } \mathbb{N}$, $c : \mathbb{Z} \rightarrow \mathbb{N}$, the literal equations (1), a nonempty finite set S of primes, and arbitrary positive exponents $e(q)$. It assumes $\Phi_{q^{e(q)}} \mid \sum_{a \in E} X^a$ for every $q \in S$. Its conclusion is

$$\text{cyclotomic} \left(\prod_{q \in S} q^{e(q)} \right) \mathbb{Z} \mid \sum_{a \in E} X^a.$$

There is no hidden prime-count, period, cardinality, or phase-closure premise. The additional declaration `arbitrary_integer_tile_T2` derives nonemptiness and minimum-normalizes arbitrary finite integer sets.

Argument	Principal Lean modules
Primary allocation	<code>PrimaryAllocation</code>
Relative cyclotomic fields	<code>RelativeCyclotomic</code> , <code>RamifiedCyclotomic</code>
Actual product identities	<code>ActualTilingSpectra</code> , <code>FiberProductBoolean</code>
Boolean stripe partition	<code>BooleanStripes</code>
Integral descent and collapse	<code>PeriodicDescent</code> , <code>StripeCollapse</code> , <code>PeriodicMatrixCollapse</code>
All-phase smaller tilings	<code>ActualPrimeReduction</code> , <code>ActualIntegerReduction</code>
Original-factor recovery	<code>T2Induction</code> , <code>CommonComplementLift</code>
Unrestricted cyclic induction	<code>CyclicT2Completion</code>
Integer transfer	<code>IntegerTilingPeriodicity</code> , <code>IntegerCyclicReduction</code> , <code>IntegerSetNormalization</code> , <code>IntegerT2Completion</code>
Exposed final statements	<code>FullCovenMeyerowitz</code>

The frozen project uses Lean 4.23.0 and Mathlib commit

`37df177aaa770670452312393d4e84aaad56e7b6.`

After installing Elan, enter the repository's formal directory and run

```
lake exe cache get
sh ./verify_full_t2.sh
```

The verifier checks three SHA256 source manifests, recompiles 35 mathematical modules in dependency order, recompiles a literal-function cyclic semantics probe, and runs 22 guarded axiom checks.

An unexpected final axiom list fails the audit. Plain `lake build` is insufficient because the frozen original default target is only `StripeCollapse`.

The reported final axioms are exactly `proptext`, `Classical.choice`, and `Quot.sound`. The release includes a GitHub Actions workflow that fetches pinned dependencies on a fresh runner and executes the complete verifier. Its log is the public replay record. This is not a from-source rebuild of every Mathlib dependency or an independent implementation of Lean’s kernel; compiler and upstream-artifact trust remain part of the verification boundary. The prose, formal meaning, public chronology, and originality each require their own assessment.

9 Attribution, review, and limitations of the claim

The cardinality/valuation allocation and common-complement inheritance have direct antecedents in CM Lemmas 2.1 and 2.5. Frobenius already appears in their tiling dilation argument, Lemma 3.1 [1]. Stripe partitions in the square-free CRT setting occur in Tao’s Proposition 17 [2]. Divisor-isometry freedom is treated by Łaba and Londner [3, Definition 3.3 and Lemma 3.4]. These ingredients are not individually claimed as new.

The combination requiring comparison with expert knowledge is integral descent of product minors, coprime Boolean orientation collapse, coalescence into all-phase actual tilings, and recovery of the original factors. Coalescence changes base- p divisibility and is not itself a divisor isometry. Product periodicity is not factor periodicity. The arbitrary phases in Section 6 are essential to the stated recovery of upper- p roots.

The functional tiling counterexamples of Kiss, Londner, Matolcsi, and Somlai [5] concern a fractional-function relaxation. The present proof relies essentially on Boolean sets and integral cancellation and asserts no corresponding unrestricted theorem for fractional functional tiles.

The research artifact and its publication materials were prepared with AI assistance. Internal automated checks have reviewed the expanded final type, integer-to-cyclic transfer, and principal proof interfaces; these are not external human referee reports. The release-day bounded literature check did not identify a full predecessor, but cannot establish exclusive priority or rule out unpublished and differently formulated work. Specialist review and acceptance remain unconfirmed. This paper therefore presents the precise proof and reproducible formal artifact for scrutiny rather than reporting external recognition as an accomplished fact.

References

- [1] E. M. Coven and A. D. Meyerowitz, *Tiling the integers with translates of one finite set*, Journal of Algebra 212 (1999), 161–174. Preprint submitted 1998. <https://arxiv.org/abs/math/9802122>.
- [2] T. Tao, *Some notes on the Coven–Meyerowitz conjecture*, 19 November 2011, with subsequent corrections in the discussion. <https://terrytao.wordpress.com/2011/11/19/some-notes-on-the-coven-meyerowitz-conjecture/>.
- [3] I. Łaba and I. Londner, *Splitting for Integer Tilings*, International Mathematics Research Notices (2025), rna090. <https://doi.org/10.1093/imrn/rna090>.
- [4] M. Kadir and K. Fan, *Tiles and weak tiles in $\mathbb{Z}_{pq}$* , arXiv:2607.02149, version 2 (2026), Section 1.3. <https://arxiv.org/html/2607.02149v2>.
- [5] G. Kiss, I. Londner, M. Matolcsi, and G. Somlai, *Functional tilings and the Coven–Meyerowitz tiling conditions*, Discrete Analysis (2025), Paper No. 18. <https://arxiv.org/abs/2411.03854>.