

Mathematics Extended Essay on the topic of Erdős' Probabilistic method

Candidate Number:

May 25, 2026

Word count: TO UPDATE

Contents

1	Introduction	2
2	Scale Free Networks and Graph Theory	2
3	The Probabilistic Method	2
4	Predicting clique sizes in Scale Free Networks	2
5	Building Scale Free Networks	3
6	Comparing Results	3
7	Conclusion	3

1 Introduction

- A brief introduction - My motivation to explore this topic

To briefly help the reader to understand this essay better, below are a few definitions which might come in handy throughout: An Object is a

2 Scale Free Networks and Graph Theory

- Describing the concept of graphs - Defining terms - Applying the idea to scale free networks

One predominant branch of Mathematics which is often ignored is called combinatorics. It focuses on the idea of counting, which is applied in order to find a result but can also define a final answer. I will talk more about that concept using an example below. Part of combinatorics is the concept of graphs which is related to graph theory which explores the relationship between objects. These should not be confused with a graph on a cartesian plane. These graphs allow the mathematical analysis of networks and connections, like streets, mobile networks, or social media networks. From now on the use of the word graph will refer to the mathematical graph which makes part of discrete mathematics, i.e. a network of nodes connected by edges.

For clarity, I will define the following terms, as they are different around the world: A node is the same as a vertex and is the undivisible point at which one or more edges meet. An edge is the connection between nodes. The degree of distribution I am only making use of undirected graphs, meaning that the edges don't represent a direction in the connection between nodes (as would be required when representing one-way streets for example)

This extended essay will investigate scale free networks which are representative of many real-world networks because of their "Hubs". A scale free network gives a power law relation between degree of

3 The Probabilistic Method

- Describing Probabilistic Notation - Applying using simple a simple Ramsey number example ($R(3,3) = 6$)

As we discussed above, graph theory is computationally difficult to explore. When trying to generate every possible combination of words

4 Predicting clique sizes in Scale Free Networks

- Setting out the problem: we want to find specific clique sizes in a scale free network with 4 hubs - Using the Chung-Lu Random Graph Model, we will describe how to statistically generate a graph, with different probabilities of nodes and edges occurring. - Using the first moment we will explain the concept of

5 Building Scale Free Networks

- Using the help of AI to code a python script, I aim to generate a statistically meaningful number of Scale-Free Networks. - By analysing the sizes of clique's present, I should be able to find similar results to my theoretical/probabilistic approach

6 Comparing Results

- Comparing results and making sure theory aligns with empirical data

7 Conclusion

- Conclude on the uses of my results