

Assignment-based Subjective Questions

Question 1. From your analysis of the categorical variables from the dataset, what could you infer about their effect on the dependent variable? (Do not edit)

Total Marks: 3 marks (Do not edit)

Answer: <Your answer for Question 1 goes below this line> (Do not edit)

I have done analysis on categorical columns using the boxplot and bar plot. Below are the few points we can infer from the visualization –

- Fall season seems to have attracted more booking.
- And, in each season the booking count has increased drastically from 2018 to 2019.
- Most of the bookings has been done during the month of may, june, july, aug, sep and oct. Trend increased starting of the year till mid of the year and then it started decreasing as we approached the end of year.
- Clear weather attracted more booking which seems obvious.
- Thu, Fri, Sat and Sun have more number of bookings as compared to the start of the week.
- When it's not holiday, booking seems to be less in number which seems reasonable as on holidays, people may want to spend time at home and enjoy with family

Question 2. Why is it important to use **drop_first=True** during dummy variable creation? (Do not edit)

Total Marks: 2 marks (Do not edit)

Answer: <Your answer for Question 2 goes below this line> (Do not edit)

drop_first = True is important to use, as it helps in reducing the extra column created during dummy variable creation. Hence it reduces the correlations created among dummy variables. Syntax - drop_first: bool, default False, which implies whether to get k-1 dummies out of k categorical levels by removing the first level.

Question 3. Looking at the pair-plot among the numerical variables, which one has the highest correlation with the target variable? (Do not edit)

Total Marks: 1 mark (Do not edit)

Answer: <Your answer for Question 3 goes below this line> (Do not edit)

'temp' variable has the highest correlation with the target variable.

Question 4. How did you validate the assumptions of Linear Regression after building the model on the training set? (Do not edit)

Total Marks: 3 marks (Do not edit)

Answer: <Your answer for Question 4 goes below this line> (Do not edit)

I have validated the assumption of Linear Regression Model based on below 5 assumptions -

- Normality of error terms o Error terms should be normally distributed
- Multicollinearity check o There should be insignificant multicollinearity among variables.
- Linear relationship validation o Linearity should be visible among variables
- Homoscedasticity o There should be no visible pattern in residual values.
- Independence of residuals o No auto-correlation

Question 5. Based on the final model, which are the top 3 features contributing significantly towards explaining the demand of the shared bikes? (Do not edit)

Below are the top 3 features contributing significantly towards explaining the demand of the shared bikes – temp winter sep

Total Marks: 2 marks (Do not edit)

Answer: <Your answer for Question 5 goes below this line> (Do not edit)

General Subjective Questions

Question 6. Explain the linear regression algorithm in detail. (Do not edit)

Total Marks: 4 marks (Do not edit)

Answer: Please write your answer below this line. (Do not edit)

<Your answer for Question 6 goes here>

Linear regression may be defined as the statistical model that analyses the linear relationship between a dependent variable with given set of independent variables. Linear relationship between variables means that when the value of one or more independent variables will change (increase or decrease), the value of dependent variable will also change accordingly (increase or decrease). Mathematically the relationship can be represented with the help of following equation – $Y = mX + c$ Here, Y is the dependent variable we are trying to predict. X is the independent variable we are using to make predictions. m is the slope of the regression line which represents the effect X has on Y c is a constant, known as the Y-intercept. If $X = 0$, Y would be equal to c. Furthermore, the linear relationship can be positive or negative in nature as explained below– o Positive Linear Relationship: A linear relationship will be called positive if both independent and dependent variable increases. It can be understood with the help of following graph

Assumptions - The following are some assumptions about dataset that is made by Linear Regression model – o Multi-collinearity – o Linear regression model assumes that there is very little or no multi-collinearity in the data. Basically, multi-collinearity occurs when the independent variables or features have dependency in them. o Auto-correlation – o Another assumption Linear regression model assumes is that there is very little or no auto-correlation in the data. Basically, auto-correlation occurs when there is dependency between residual errors. o Relationship between variables – o Linear regression model assumes that the relationship between response and feature variables must be linear. o Normality of error terms

Question 7. Explain the Anscombe's quartet in detail. (Do not edit)

Total Marks: 3 marks (Do not edit)

Answer: Please write your answer below this line. (Do not edit)

<Your answer for Question 7 goes here>

Anscombe's Quartet was developed by statistician Francis Anscombe. It comprises four datasets, each containing eleven (x, y) pairs. The essential thing to note about these datasets is that they share the same descriptive statistics. But things change completely, and I must emphasize

COMPLETELY, when they are graphed. Each graph tells a different story irrespective of their similar summary statistics. The summary statistics show that the means and the variances were identical for x and y across the groups: • Mean of x is 9 and mean of y is 7.50 for each dataset. • Similarly, the variance of x is 11 and variance of y is 4.13 for each dataset • The correlation coefficient (how strong a relationship is between two variables) between x and y is 0.816 for each dataset When we plot these four datasets on an x/y coordinate plane, we can observe that they show the same regression lines as well but each dataset is telling a different story

Question 8. What is Pearson's R? (Do not edit)

Total Marks: 3 marks (Do not edit)

Answer: Please write your answer below this line. (Do not edit)

<Your answer for Question 8 goes here>

Pearson's r is a numerical summary of the strength of the linear association between the variables. If the variables tend to go up and down together, the correlation coefficient will be positive. If the variables tend to go up and down in opposition with low values of one variable associated with high values of the other, the correlation coefficient will be negative. The Pearson correlation coefficient, r, can take a range of values from +1 to -1. A value of 0 indicates that there is no association between the two variables. A value greater than 0 indicates a positive association; that is, as the value of one variable increases, so does the value of the other variable. A value less than 0 indicates a negative association; that is, as the value of one variable increases, the value of the other variable decreases.

Question 9. What is scaling? Why is scaling performed? What is the difference between normalized scaling and standardized scaling? (Do not edit)

Total Marks: 3 marks (Do not edit)

Answer: Please write your answer below this line. (Do not edit)

<Your answer for Question 9 goes here>

Feature Scaling is a technique to standardize the independent features present in the data in a fixed range. It is performed during the data pre-processing to handle highly varying magnitudes or values or units. If feature scaling is not done, then a machine learning algorithm tends to weigh greater values, higher and consider smaller values as the lower values, regardless of the unit of the values. Example: If an algorithm is not using feature scaling method then it can consider the value 3000 meter to be greater than 5 km but that's actually not true and in this case, the algorithm will give wrong predictions. So, we use Feature Scaling to bring all values to same magnitudes and thus, tackle this issue.

Question 10. You might have observed that sometimes the value of VIF is infinite. Why does this happen? (Do not edit)

Total Marks: 3 marks (Do not edit)

Answer: Please write your answer below this line. (Do not edit)

<Your answer for Question 10 goes here>

If there is perfect correlation, then $VIF = \infty$. A large value of VIF indicates that there is a correlation between the variables. If the VIF is 4, this means that the variance of the model coefficient is inflated by a factor of 4 due to the presence of multicollinearity. When the value of VIF is infinite it shows a perfect correlation between two independent variables. In the case of perfect correlation, we get $R^2 = 1$, which leads to $1/(1-R^2) = \infty$. To solve this we need to drop one of the variables from the dataset which is causing this perfect multicollinearity.

Question 11. What is a Q-Q plot? Explain the use and importance of a Q-Q plot in linear regression. (Do not edit)

Total Marks: 3 marks (Do not edit)

Answer: Please write your answer below this line. (Do not edit)

<Your answer for Question 11 goes here>

The quantile-quantile (q-q) plot is a graphical technique for determining if two data sets come from populations with a common distribution. Use of Q-Q plot: A q-q plot is a plot of the quantiles of the first data set against the quantiles of the second dataset. By a quantile, we mean the fraction (or percent) of points below the given value. That is, the 0.3 (or 30%) quantile is the point at which 30% percent of the data fall below and 70% fall above that value. A 45-degree reference line is also plotted. If the two sets come from a population with the same distribution, the points should fall approximately along this reference line. The greater the departure from this reference line, the greater the evidence
