

Extraction of tabled statistical results with tableParser

Appendix A: Example tables

This PDF file may be processed with the tableParser R package. See: [/cran/tableParser](#)

- To convert the tables within this document into R character matrices run:

```
table2matrix(`tableExamples.pdf`)
```

- To convert the reported results in each table into a text string run:

```
table2text('tableExamples.pdf')
```

- To extract the statistical standard results as data frame object with recalculated p-values run:

```
table2stats(`tableExamples.pdf`)
```

See the documentation of each function for further options.

Some blind text

Nam quis enim. Quisque ornare dui a tortor. Fusce consequat lacus pellentesque metus. Duis euismod. Duis non quam. Maecenas vitae dolor in ipsum auctor vehicula. Vivamus nec nibh eget wisi varius pulvinar. Cras a lacus. Etiam et massa. Donec in nisl sit amet dui imperdiet vestibulum. Duis porttitor nibh id eros.

Single-way matrix example

Column A	Column B	Column C
1	3	A
2	4	B

Multi-way matrix example

	Column A	Column B
Row A	1	A
Row B	2	B

Nam quis enim. Quisque ornare dui a tortor. Fusce consequat lacus pellentesque metus. Duis euismod. Duis non quam. Maecenas vitae dolor in ipsum auctor vehicula. Vivamus nec nibh eget wisi varius pulvinar. Cras a lacus. Etiam et massa. Donec in nisl sit amet dui imperdiet vestibulum. Duis porttitor nibh id eros.

Table 1: An example of tabled descriptive statistics

Variable	Category	Frequency (%)	Mean (SD) Income
Age	20–25	78 (39.79)	1,345 (250)
	26–31	65 (33.16)	1,825 (535)
	32 and older	53 (27.05)	2,315 (930)
Gender	Men	97 (49.49)	1,965 (550)
	Women	99 (50.51)	1,685 (350)

Nam quis enim. Quisque ornare dui a tortor. Fusce consequat lacus pellentesque metus. Duis euismod. Duis non quam. Maecenas vitae dolor in ipsum auctor vehicula. Vivamus nec nibh eget wisi varius pulvinar. Cras a lacus. Etiam et massa. Donec in nisl sit amet dui imperdiet vestibulum. Duis porttitor nibh id eros.

Table 2: An example of tabled descriptive statistics and correlations with table notes

Variables	1	2	3
Mean	45 (11.2)	5,200 (21,123)	26.8 (5.2)
Median	40	3,200	25
1. Gender	1	.38**	.04
2. Income		1	.21*
3. Ext.			.87

N=125, * p<.05, ** p<.01

Numbers in brackets represent standard deviations.

Ext.: Extraversion

Nam quis enim. Quisque ornare dui a tortor. Fusce consequat lacus pellentesque metus. Duis euismod. Duis non quam. Maecenas vitae dolor in ipsum auctor vehicula. Vivamus nec nibh eget wisi varius pulvinar. Cras a lacus. Etiam et massa. Donec in nisl sit amet dui imperdiet vestibulum. Duis porttitor nibh id eros.

Table 3: An example of a multi model table with table notes

Variables	Job performance		
	Model 1	Model 2	Model 3
Income	-0.26 (.16)	-0.39 (.21)	-0.36** (.1)
Extraversion		0.12 ⁺ (.6)	0.16*** (.03)
Income * Extraversion			-0.18** (.1)
R2	0.07	0.22	0.31
ΔR2	0.07	0.15	0.9
ΔF	2.16	21.76**	14.37*

+ p=.075; * p<.05; ** p<.01; *** p<.001

Numbers in brackets represent standard errors.

Nam quis enim. Quisque ornare dui a tortor. Fusce consequat lacus pellentesque metus. Duis euismod. Duis non quam. Maecenas vitae dolor in ipsum auctor vehicula. Vivamus nec nibh eget wisi varius pulvinar. Cras a lacus. Etiam et massa. Donec in nisl sit amet dui imperdiet vestibulum. Duis porttitor nibh id eros.

Table 4: An example of an ANOVA table with table notes

Variable	SSq	df	MSq	F	P(>F)
Factor A	12	2	3	9.09	.00
Factor B	4.5	1	4.5	6.82	.01
Factor A * B	3	2	1.5	2.27	.12
Residuals	20	30	0.66		

Total	39.5	35	1.13		
-------	------	----	------	--	--

Italic values are $p < .05$.

Bold values are significant with $p < .01$.

Nam quis enim. Quisque ornare dui a tortor. Fusce consequat lacus pellentesque metus. Duis euismod. Duis non quam. Maecenas vitae dolor in ipsum auctor vehicula. Vivamus nec nibh eget wisi varius pulvinar. Cras a lacus. Etiam et massa. Donec in nisl sit amet dui imperdiet vestibulum. Duis porttitor nibh id eros.