

FYS4150: Project 2

Ingrid A V Holm

September 23, 2016

Unitary and orthogonal transformations (a)

A unitary matrix U has the properties:

$$U^{-1} = U^\dagger \quad (1)$$

An orthogonal matrix (the corresponding real matrix) has the properties:

$$U^T = U^{-1} \quad (2)$$

Where U^\dagger is for matrices in Hilbert space (complex matrices). It represents the transpose of the complex conjugate, i.e. $U^\dagger = (U^*)^T$.

Assume we have an orthogonal basis for the n -dimensional space v_i , where $v_j^T v_i = \delta_{ij}$:

$$\mathbf{v}_i = \begin{bmatrix} v_{i1} \\ \cdots \\ \cdots \\ v_{in} \end{bmatrix}, U\mathbf{v}_i = \mathbf{w}_i$$

Under orthogonal transformations, both the dot product and orthogonality of vectors is conserved:

$$\mathbf{w}_i^T \mathbf{w}_j = (U\mathbf{v}_i)^T (U\mathbf{v}_j) = \mathbf{v}_i^T U^T U \mathbf{v}_j = \mathbf{v}_i^T U^{-1} U \mathbf{v}_j = \mathbf{v}_i^T \mathbf{v}_j = \delta_{ij}$$

Jacobi's rotation method (b)

We want to solve the eigenvalue problem for a tridiagonal matrix:

$$\begin{pmatrix} d_0 & e_0 & 0 & \dots & 0 & 0 \\ e_1 & d_1 & e_1 & \dots & 0 & 0 \\ 0 & e_2 & d_2 & e_2 & \dots & 0 \\ \vdots & & & \ddots & & 0 \\ 0 & \dots & \dots & 0 & e_N & d_N \end{pmatrix} \begin{pmatrix} u_0 \\ u_1 \\ \vdots \\ u_{N-1} \\ u_N \end{pmatrix} = \lambda \begin{pmatrix} u_0 \\ u_1 \\ \vdots \\ u_{N-1} \\ u_N \end{pmatrix} \quad (3)$$

We want to use Jacobi's rotation method. We define an orthogonal matrix S to operate on our matrix A :

$$B = S^T A S \quad (4)$$

With all elements in S equal to zero, except:

$$s_{kk} = s_{ll} = \cos \theta, \quad s_{kl} = -s_{lk} = s \sin \theta, \quad s_{ii} = 1 \quad ; \quad i \neq k \quad i \neq l$$

<p>1. Find $\max(a_{kl})$.</p> <p>2. Compute the rotation parameters $t = \tan \theta$, $c = \cos \theta$, $s = \sin \theta$:</p> $\tau = \cot \theta = \frac{a_{ll} - a_{kk}}{2a_{kl}} \quad (5)$ $t = -\tau \pm \sqrt{1 + \tau^2} \quad (6)$ $c = \frac{1}{1 + t^2}, \quad s = tc \quad (7)$	<p>3. Calculate the elements of B:</p> $b_{ii} = a_{ii}, \quad i \neq k \quad i \neq l$ $b_{ik} = a_{ik}c - a_{il}s, \quad i \neq k \quad i \neq l$ $b_{il} = a_{il}c + a_{ik}s, \quad i \neq k \quad i \neq l$ $b_{kk} = a_{kk}c^2 - 2a_{kl}c * s + a_{ll}s^2$ $b_{ll} = a_{ll}c^2 + 2a_{kl}c * s + a_{kk}s^2$ $b_{kl} = (a_{kk} - a_{ll})c * s + a_{kl}(c^2 - s^2)$ <p>4. If $\max(b_{kl}) \geq \epsilon$ run the algorithm again.</p>
--	--

Figure 1: Jacobi rotation algorithm for tridiagonal matrix