

Outlook

From	andile.mthembu@collections.ops.keystonebank.co.za
To	analytics.retail@keystonebank.co.za
Cc	credit.risk.retail@keystonebank.co.za, customer.experience@keystonebank.co.za
Sent	Tue, 09 Feb 2021 08:13:00 -0000

Separate self-cure from successful collection

Morning,

I need help with something that sounds simple but probably is not.

Collections wants to reduce unnecessary contact by identifying customers who usually restore the account without intervention. Several teams are reporting more failed or retried activity than they expected, but they do not agree on the cause. The first extracts did not reconcile, which means source precedence and exclusions must be part of the answer.

The questions I would ask in the room are:

1. What would we expect to see if repeat NSF customers need a different treatment?
2. What evidence would instead support that balance recovery and payment timing identify self-cure?
3. How do we rule out that past contact creates the apparent cure?

This has returned because the previous answer was useful but not strong enough to become a standing rule. Use February 2021 as the new evidence point rather than recycling the earlier conclusion. Please work towards which accounts can receive a light-touch reminder and which need an agent. A useful output would be a working Excel file with the exception population and clear columns.

Use collections cases, promises to pay and recovery payments; transaction-level timestamps, channels, amounts, statuses and error context; customer contacts, complaints and suggestions; debit-order mandates, collection dates, suspensions and cancellations. One caution: Observed cure after contact is not proof that the contact caused it.

Regards