

A Global Occupation Conjecture for the Accelerated $3x + 1$ Map: Residue–Time Duality, Divergent-Orbit Sparsity, and the Pointwise Realizer Problem

Elias De Jesús
Independent Researcher
ORCID: 0009-0007-0190-9143
dejesuselias10@gmail.com

Revision 6, 2026

Abstract

We study the Effective Occupation Conjecture (EOC) for the accelerated $3x + 1$ map $T(m) = (3m + 1)/2^{a(m)}$, $a(m) = \nu_2(3m + 1)$. Writing $R_n = S_n - n \log_2 3$ for the drift of the accumulated 2-adic valuation and $\mathcal{O}_c(m_0) = \#\{n \geq 0 : R_n \leq c\}$, EOC asserts $\mathcal{O}_c(m_0) = O(\log m_0)$ for every seed and every corridor width c .

This revision reorganizes the programme around four axes — occupation, residue, time, and divergence — and the exact bridges between them. Record chronology identifies the residue and time axes precisely: the least-realizer frontier $r_{\min}(N, c)$ is the inverse function of the single-window confinement lifetime $L_c(m)$, so every pointwise lifetime bound transfers to a realizer floor, on a graded scale that we tabulate. On the divergence side we retain the strongest unconditional input: via García–Tal’s collision-free window theorem and Curry’s explicit quantitative form, every divergent orbit has summable reciprocals and admits no eventual logarithmic drift floor $R_n \geq -B \log_2 n + O(1)$ for any $B < 1/\beta_* \approx 1.0358567$. Curry’s input plays a second, qualitative role: after the last global maximum of R , a hypothetical divergent orbit has an infinite zero-corridor tail, so *universal drift exit* — every positive seed eventually attains $R_n > 0$ — is equivalent to the exclusion of divergent orbits, while remaining compatible with hypothetical nontrivial cycles and hence strictly weaker than the Collatz conjecture.

We then record what a sustained programme of structural audits has closed. Finite information cannot decide infinite realization: every exact-realizer cylinder contains arbitrarily large positive integers, so $\forall N \exists m_N$ is free and all content lies in $\exists m \forall N$. Global valuation-word information cannot decide it either: we exhibit an explicitly indexed uncountable family of zero-confined words with linear negative drift, summable $\sum 2^{R_n}$ and $O(G)$ occupation shadow, of which all but countably many have no positive-integer realizer. Several apparently independent coordinates — normalized carry, lift digits, terminal zero gap, transport budgets — are invertible reparameterizations of the exact realizer, and the Archimedean and 2-adic readings of the natural divisibility object are complementary factors of one identity rather than independent constraints. Launch renormalization remains a valid bridge from finite words to actual positive-orbit dynamics, but its corridor-uniform closing hypothesis is, once the corridor width is computed rather than assumed, essentially an $O(\log n)$ stopping-time conjecture; and deep negative drift is self-financing, since the ordinary height it creates is exactly the resource that permits the recovery valuations repaying it.

Neither EOC nor the Collatz conjecture is proved here. The surviving frontier is actual positive-orbit arithmetic at three nested levels: qualitative exclusion of an infinite zero-corridor tail, quantitative control of single-window lifetime, and control of separated returns sufficient for full occupation.

Non-claims. This paper proves neither the Collatz conjecture nor any tier of EOC. It does not prove an exponential realizer floor for arbitrary confined words. It does not promote any computational finding to a theorem: every claim is labelled with its status (§B). Where a statement reorganizes or reclassifies earlier work rather than proving something new, this is said explicitly.

Contents

1	Introduction	3
1.1	Four axes, not three layers	3
1.2	What this revision adds	4
2	Preliminaries	5
3	The occupation axis: EOC and its logical status	5
3.1	Four tiers	5
3.2	The Collatz reduction	6
3.3	The entropy scale	6
3.4	EOC as an occupation-time shadow of Rozier’s hypothesis	7
4	The divergence axis	7
4.1	An elementary 8/9 bound and its terminality	7
4.2	Garcia–Tal/Curry sparsity	7
4.3	Curry’s second, qualitative role	8
5	The residue axis	8
5.1	Proved sectors	8
5.2	What the height invariant actually separates	9
5.3	Endpoint depth and the frontier defect	9
6	The time axis	10
6.1	Three temporal scales	10
6.2	Residue–time duality	10
6.3	The general transfer principle	10
6.4	Drift exit and the future-minimum theorem	11
6.5	Universal drift exit and divergence	12
7	Launch renormalization: the bridge to actual orbits	12
7.1	Reclassifying the corridor-uniform hypothesis	13
8	Self-financing recovery	13
9	Boundaries of the programme	14
9.1	Finite cylinders do not encode infinite realization	14
9.2	Coordinate sufficiency is not independent information	14
9.3	Global word shadows do not characterize positive realization	15
9.4	Population sparsity is not pointwise placement	15
9.5	Deep drift creates its own recovery capacity	16
9.6	Single-window lifetime does not control occupation	16

10 The programme hierarchy	16
11 The programme map	16
11.1 Symbolic freedom, finite and infinite	16
12 Closed routes	17
12.1 External arithmetic	17
13 Open problems	17
14 Formal verification status	18
15 Computational calibration	19
16 Relation to prior work	20
17 Conclusion	20
A Band rigidity	22
B Claim-status conventions	22

1 Introduction

The accelerated $3x + 1$ map sends an odd positive integer to the next odd term of its Collatz trajectory,

$$T(m) = \frac{3m + 1}{2^{a(m)}}, \quad a(m) = \nu_2(3m + 1) \geq 1.$$

Writing S_n for the accumulated 2-adic valuation after n accelerated steps and $R_n = S_n - n\alpha$ with $\alpha = \log_2 3$ for its drift below the critical slope, the *Effective Occupation Conjecture* asserts that the time an orbit spends at or below any fixed drift level is logarithmic in the seed. This single pointwise statement forces every orbit to be eventually periodic and reduces Collatz to a finite cycle search (Theorem 3.6).

1.1 Four axes, not three layers

Revision 5 of this work was organized in three layers: EOC and its logical status; unconditional drift-axis progress; and the residue-axis reformulation. That organization is no longer adequate, for a specific reason: the residue axis and the confinement-lifetime question are not two problems but two coordinates on one problem, and a fourth, genuinely qualitative question — exclusion of divergent orbits — sits beside both. Revision 6 is therefore organized around four axes.

Occupation axis (§3)

The functional $\mathcal{O}_c(m_0) = \#\{n \geq 0 : R_n \leq c\}$, counting *all* times in the corridor, including separated later returns. EOC lives here, and it is the strongest of the four.

Residue axis (§5)

The exact realizer congruence: which integers realize a given valuation word, and how small the least such integer, $r_{\min}(N, c)$, can be.




Figure 1: The programme map. Vertical double arrows are exact equivalences; the lower chain is the qualitative divergence reduction. Every implication is referenced in §11.

Time axis (§6)

The single-window lifetime $L_c(m)$ and the drift exit time $\tau_c(m)$. Record chronology makes the residue and time axes *exactly* dual.

Divergence axis (§4)

What is unconditionally known about a hypothetical divergent orbit: sparsity of its value set, summability of reciprocals, and exclusion of logarithmic drift floors.

The programme map in Figure 1 records the exact bridges. Every arrow in it is a theorem, and §11 gives the reference for each.

1.2 What this revision adds

Revision 6 is primarily a *synthesis*. Its main additions are:

1. A general transfer principle (§6.3) turning any pointwise lifetime bound into a realizer floor, with the exponents worked out: logarithmic, polylogarithmic, polynomial and linear lifetime bounds give exponential, stretched-exponential, polynomial and linear frontiers respectively. EOC thereby becomes a *graded* programme rather than an all-or-nothing one.
2. The qualitative time axis (§6.4): the drift exit time τ_0 , its exact integral characterisation $R_n > 0 \iff 3^n < 2^{S_n}$, and the equivalence between universal drift exit and the exclusion of divergent orbits.
3. A boundaries section (§9) collecting the strongest no-go results: the finite-cylinder barrier, the coordinate-collapse ledger, and the word-shadow cardinality barrier.
4. A corrected reading of the launch-renormalization closing hypothesis (§7) and the self-financing obstruction to temporal impossibility arguments (§8).
5. A closed-route map (§12) recording, for each mechanism tried, what it achieved and where it stops.

We are explicit about provenance throughout. Several statements that read as new are in fact consequences of identities already in Revision 5 — the future-minimum theorem (Prop. 6.9) follows from the Eliahou–Rozier identity, and the positive cycle drift (Lemma 4.2) is Revision 5’s Lemma on terminal and cyclic drift. Where this happens we say so and record only the new proof or the new role.

2 Preliminaries

Let m_0 be a positive odd integer, $m_{n+1} = T(m_n)$, $d_n = a(m_n)$, $S_n = \sum_{i < n} d_i$, $\alpha = \log_2 3$, $R_n = S_n - n\alpha$, so $R_0 = 0$ and each step changes R by $d_n - \alpha$.

Remark 2.1. If every accelerated orbit reaches 1 then Collatz holds: every positive integer's odd part has an accelerated orbit, and the classical trajectory interleaves accelerated iterates with halvings. We work exclusively with T . [standard]

For fixed $c > 0$, $\mathcal{O}_c(m_0) = \#\{n \geq 0 : R_n \leq c\} \in \mathbb{Z}_{\geq 1} \cup \{\infty\}$. This is a global sojourn count: it neither halts at first passage nor counts time far above the critical line.

Lemma 2.2 (Eliahou–Rozier identity). *For every odd $m_0 \geq 1$ and every $n \geq 0$,*

$$m_n 2^{R_n} = m_0 \prod_{i < n} \left(1 + \frac{1}{3m_i}\right), \quad \text{equivalently} \quad R_n = \log_2 \frac{m_0}{m_n} + E_n, \quad E_n = \sum_{i < n} \log_2 \left(1 + \frac{1}{3m_i}\right) \geq 0.$$

[due to Eliahou [1] and Rozier [2]; no novelty claimed]

Remark 2.3 (Notation warning: two quantities called E). Throughout, $E_n = \log_2 Q_n$ denotes the *carry excess* of Lemma 2.2. The *endpoint depth* of a word, elsewhere written $E(D)$, is a different quantity and is written $\mathbf{E}(D) = S - \log_2 r(D)$ here (§5.3). The two are unrelated and were a genuine source of confusion in earlier drafts.

There is a second, purely integral form of the same information, used whenever real logarithms would be inconvenient.

Lemma 2.4 (Aggregate identity). *With $C_n = \sum_{j < n} 3^{n-1-j} 2^{S_j}$ (so $C_0 = 0$, $C_{n+1} = 3C_n + 2^{S_n} > 0$ for $n \geq 0$),*

$$2^{S_n} m_n = 3^n m_0 + C_n \quad \text{for all } n \geq 0.$$

[elementary; formalized, §14]

Proof. Induction on n . The case $n = 0$ is $m_0 = m_0$. If $2^{S_n} m_n = 3^n m_0 + C_n$ then, multiplying $2^{d_n} m_{n+1} = 3m_n + 1$ by 2^{S_n} , $2^{S_{n+1}} m_{n+1} = 3 \cdot 2^{S_n} m_n + 2^{S_n} = 3^{n+1} m_0 + (3C_n + 2^{S_n}) = 3^{n+1} m_0 + C_{n+1}$. $\square$

Lemma 2.5 (Integral corridor). *Since $S_n \in \mathbb{Z}$ and $n\alpha = \log_2(3^n)$,*

$$R_n \leq 0 \iff 2^{S_n} \leq 3^n, \quad R_n > 0 \iff 3^n < 2^{S_n}, \quad R_n \leq 0 \iff S_n \leq \lfloor n\alpha \rfloor.$$

[elementary]

Lemma 2.6 (Wedge). *If $m_i \geq m_0$ for all $0 \leq i \leq n$, then $R_i \leq E_i \leq i/(3m_0 \ln 2)$ for $0 \leq i \leq n$. In particular $R_i \leq c$ for $0 \leq i \leq \min\{n, \lfloor 3c \ln 2 \cdot m_0 \rfloor\}$. [Revision 5]*

3 The occupation axis: EOC and its logical status

3.1 Four tiers

Conjecture 3.1 (Existence EOC). For every fixed $c > 0$ there exist constants K_c, C_c such that every positive odd m_0 satisfies $\mathcal{O}_c(m_0) \leq K_c \log_2 m_0 + C_c$.

Conjecture 3.2 (Effective EOC at (c, K, C)). For stated $c > 0$ and stated numerical K, C , every positive odd m_0 satisfies $\mathcal{O}_c(m_0) \leq K \log_2 m_0 + C$.

Conjecture 3.3 (Sharp-leading EOC). For every fixed $c > 0$, $\mathcal{O}_c(m_0) \leq K \log_2 m_0 + O_c(\log \log m_0)$ with $K = 1/I(\alpha)$, where $I(\mu) = \mu - \mu \log_2 \mu + (\mu - 1) \log_2(\mu - 1)$.

Conjecture 3.4 (Strong sharp EOC). For every fixed $c > 0$ there is C_c with $\mathcal{O}_c(m_0) \leq K \log_2 m_0 + C_c$.

Conjecture 3.4 $\Rightarrow$ 3.3 $\Rightarrow$ 3.1, and Conjecture 3.2 at any constants implies Conjecture 3.1 at that c . Crucially, Conjecture 3.1 does *not* imply Conjecture 3.2 at any stated constants: it asserts existence without a bound. All four tiers are pointwise statements about every seed, not almost-every-seed statements.

3.2 The Collatz reduction

Definition 3.5. For $c > 0$ and constants K, C , let $m^* = m^*(K, C, c)$ be least such that $3c \ln 2 \cdot m > K \log_2 m + C + 1$ for all $m \geq m^*$; it exists and is computable by finitely many exact comparisons.

Theorem 3.6. Fix $c > 0$ and suppose Conjecture 3.1 holds at (c, K, C) ; put $m^* = m^*(K, C, c)$. Then

- (i) no accelerated orbit is injective; every orbit is eventually periodic;
- (ii) every odd $m_0 \geq m^*$ has an iterate $m_i < m_0$ with $i \leq \lfloor 3c \ln 2 \cdot m_0 \rfloor$;
- (iii) if some seed's orbit does not reach 1, there is a nontrivial accelerated cycle with minimum member below m^* ;
- (iv) if additionally no nontrivial cycle has minimum member below m^* , the Collatz conjecture holds.

[Revision 5, retained unchanged]

Corollary 3.7 (Explicit instance). Assume Conjecture 3.2 at $c = 1$, $K = 13$, $C = 100$. Then $m^*(13, 100, 1) = 90$; every odd $m < 90$ reaches 1 by direct inspection (longest: $m = 27$, 41 accelerated steps); hence Collatz holds. [Revision 5]

Remark 3.8 (Why effective constants are genuinely needed). m^* is computable from (K, C, c) , but Conjecture 3.1 supplies no bound on (K, C) themselves. Minimal-counterexample reasoning does not repair this: Theorem 3.6(ii) localizes a counterexample below m^* but does not eliminate it. Since Bařina's verification [13] settles all seeds below 2^{71} , any nontrivial cycle's minimum member exceeds 2^{71} , and the wedge forces $\mathcal{O}_1(m_{\min}) \gtrsim 3 \ln 2 \cdot 2^{71} \approx 4.9 \times 10^{21}$ — a value Conjecture 3.1 can accommodate at unspecified constants but which is wildly larger than anything the record table of §15 suggests. [Revision 5]

3.3 The entropy scale

The natural population constant is the per-odd-step entropy deficit

$$I_{\text{Collatz}} = \alpha(1 - H_2(1/\alpha)) = \frac{1 - H_2(\rho)}{\rho}, \quad \rho = \frac{1}{\alpha} = 0.6309297 \dots,$$

with H_2 the binary entropy. Numerically $I_{\text{Collatz}} = 0.0793186127748554 \dots$, and $1/I(\alpha) = 12.6074 \dots$ is the predicted asymptotic slope in Conjecture 3.3. The parenthesization matters: the constant is $(1 - H_2(\rho))/\rho$, *not* $1 - H_2(\rho)/\rho$. [arithmetic; a known typesetting hazard in earlier drafts]

This constant calibrates the population scale and supplies, by itself, no pointwise anti-concentration. That distinction is one of the recurring themes of §9.

3.4 EOC as an occupation-time shadow of Rozier's hypothesis

Theorem 3.9. *Assume Rozier's Lower Bound Hypothesis [2, Hyp. 2.3]. Then for every fixed $c > 0$, $\mathcal{O}_c(m_0) \leq \frac{1}{I(\alpha)} \log_2 m_0 + O_c(\log \log m_0)$; that is, LBH implies Conjecture 3.3, hence Conjecture 3.1. [Revision 5]*

The proof applies Rozier's own manoeuvre [2, Thm. 4.1] at occupation times rather than at total stopping time, via the entropy identity $\mu(1 - H_2(1/\mu)) = I(\mu)$. We regard it as a corollary of his framework, recorded because the occupation functional had not previously been isolated. It does not advance the provability of EOC — LBH is itself open, and arguably harder — but positions EOC precisely as a weaker occupation-time shadow of LBH. Making the additive term explicit would supply Conjecture 3.2 and, with it, Collatz via Corollary 3.7; this is Open Problem A.

4 The divergence axis

This section is unconditional throughout; nothing here assumes EOC.

Definition 4.1. An orbit (m_n) is *injective* if $m_0, m_1, \dots$ are pairwise distinct.

Lemma 4.2 (Terminal and cyclic drift). *If an orbit reaches 1, or enters a nontrivial positive cycle, then $R_N \rightarrow +\infty$. Consequently every non-injective orbit has $R_N \rightarrow +\infty$. In particular, closure of a cycle of length L with total valuation S forces $2^S > 3^L$: strictly positive drift per period. [Revision 5; the integral form $2^S > 3^L$ is immediate from Lemma 2.4 and is formalized, §14]*

4.1 An elementary 8/9 bound and its terminality

Lemma 4.3 (Carry budget). *Every accelerated iterate m_k , $k \geq 1$, satisfies $m_k \not\equiv 0 \pmod{3}$. Hence for $k \geq 1$ the states lie in the density-1/3 set of integers coprime to 6, and for every injective orbit $E_N \leq \frac{1}{9} \log_2 N + O(1)$, so $2^{E_N} \ll N^{1/9}$. [Revision 5]*

Theorem 4.4 (Logarithmic-floor exclusion, 8/9). *Let m_0 have an injective accelerated orbit. For every $B < 8/9$, $R_k \geq -B \log_2 k$ fails for infinitely many k . [Revision 5]*

Corollary 4.5. *For every injective orbit and fixed $c > 0$, $\#\{0 \leq j < N : R_j > c\} \ll_{m_0, c} N^{1/9}$; hence $\mathcal{O}_c(m_0) = \infty$ at density-one rate, and Conjecture 3.1 directly excludes injective orbits. This is what Theorem 3.6(i) uses. [Revision 5]*

Remark 4.6 (Terminality of the mod-3 mechanism). For a prime $p > 3$, both 2 and 3 are units mod p , so $3m + 1 \equiv 0 \pmod{p}$ is solvable and no annihilation analogous to the mod-3 case occurs. This one-step fixed-modulus packing mechanism supplies no further density reduction at any other fixed modulus; pushing past 8/9 requires a genuinely different mechanism. [Revision 5]

4.2 Garcia–Tal/Curry sparsity

Theorem 4.7 (Garcia–Tal [5]; explicit form, Curry [6]). *Let $\mathcal{O}$ be the value set of an infinite aperiodic orbit of the raw map. For every $\beta > \beta_*$ there is C_β with $\#(\mathcal{O} \cap [a, a + X]) \leq C_\beta X^\beta \log(2X)$ uniformly in a , where $\beta_* = \gamma_* \log_2 3 \approx 0.9653844$ and $\gamma_* \approx 0.6090897$ is the unique root of $H(\gamma) = \gamma \log_2 3$. [external, cited]*

Proposition 4.8 (Reciprocal summability). *Every divergent (equivalently, injective) orbit satisfies $\sum_{x \in \mathcal{O}} 1/x < \infty$, hence $\sum_{n \geq 0} 1/m_n < \infty$ for its accelerated iterates. Consequently $Q_n \uparrow Q_\infty < \infty$, $E_n \uparrow E_\infty < \infty$, $m_n \rightarrow \infty$, and $R_n \rightarrow -\infty$. [Revision 5, from Theorem 4.7]*

Theorem 4.9 (Occupation bound and the $1/\beta_*$ threshold). *Fix $\beta \in (\beta_*, 1)$. There are constants c_1, c_2 depending on β and on the orbit with $\#\{n \geq 0 : R_n \geq -G\} \leq c_1 2^{\beta G} (G + c_2)$ for all $G \geq 1$. Consequently no divergent orbit satisfies $R_n \geq -B \log_2 n + O(1)$ eventually, for any $B < 1/\beta_* \approx 1.0358567$. The endpoint $B = 1/\beta_*$ is not excluded. [Revision 5]*

4.3 Curry's second, qualitative role

Revision 5 treated Theorem 4.7 purely as quantitative divergence sparsity. It has a second role, which organizes the whole time axis.

Proposition 4.10 (Last-maximum reduction). *Let m_0 have a divergent orbit. By Proposition 4.8, $R_n \rightarrow -\infty$, so R attains a global maximum at finitely many indices; let n_0 be the last. Put $m^* = m_{n_0}$, an actual positive odd integer, and $R'_k = R_{n_0+k} - R_{n_0}$ its relative drift. Then $R'_k < 0$ for every $k \geq 1$, hence $S'_k \leq \lfloor k\alpha \rfloor$ for every $k \geq 1$: the restarted orbit has an infinite zero-corridor tail. [companion work; already formalized in the repository]*

This reduction is not new to Revision 6 — it is the zero-corridor tail theorem of the Curry foundation notes, where it is proved and formalized, and where it is also recorded that the step requires no irrationality of α . Its role here is to make the divergence axis and the time axis meet.

Remark 4.11 (Scope). Nothing in this section constrains hypothetical nontrivial cycles, proves any tier of EOC, or touches the occupation of convergent orbits. [Revision 5]

5 The residue axis

Definition 5.1. A valuation word $D = (d_0, \dots, d_{N-1})$ has $d_i \geq 1$; $s_0 = 0$, $s_j = \sum_{i < j} d_i$, $S_N(D) = s_N$, and carry sum $C_N(D) = \sum_{j=0}^{N-1} 3^{N-1-j} 2^{s_j}$. An odd m_0 realizes D if $a(m_i) = d_i$ for all $i < N$.

Proposition 5.2 (Exact realizer congruence). *The exact realizers of D form a single residue class modulo $2^{S_N(D)+1}$, not $2^{S_N(D)}$. Writing $r(D)$ for the least positive representative,*

$$r(D) \equiv (2^{S_N(D)} - C_N(D) \cdot 3^{-N}) \pmod{2^{S_N(D)+1}}.$$

[Revision 5]

Definition 5.3 (Coarse anchor versus exact lift). For $S \leq S_N(D)$ the *coarse anchor* at depth S is $a_S(D) := (-C_N(D)3^{-N}) \bmod 2^S$. Proposition 5.2 shows $r(D)$ reduces mod 2^{S_N} to $a_{S_N}(D)$ and is one of $\{a_{S_N}(D), a_{S_N}(D) + 2^{S_N}\}$, terminal parity selecting which. Unconditionally $r(D) \geq a_{S_N}(D)$; this inequality, not the equality, is what every floor argument below uses. [Revision 5]

5.1 Proved sectors

Theorem 5.4 (Periodic realizer floor). *For a fixed block X put $\xi_X = -C(X)/\delta_X \in \mathbb{Z}_2$ with $\delta_X = 3^{q_X} - 2^{S_X}$ odd. If $\xi_X \notin \mathbb{Z}_{>0}$, there is a finite B_X with $r(X^n) \geq 2^{nS_X - B_X}$ for all large n , hence $\log_2 r(X^n)/(nS_X) \rightarrow 1$. [Revision 5; full proof in companion note [15]]*

Theorem 5.5 (Eventually periodic words). *For a fixed prefix Y the same mechanism applies to YX^n with transformed anchor $\xi_{X,Y}$; outside the same exceptional case, $r(YX^n) \geq 2^{S_Y + nS_X - B_{X,Y}}$. [Revision 5]*

Proposition 5.6 (Bulk-defect anchor stability). *For $D = X^a E X^b$ (fixed X , arbitrary finite defect E , arbitrary b), the coarse anchor satisfies $a_{aS_X}(D) = \xi_X \bmod 2^{aS_X}$ exactly, independent of E and b . Hence, outside the exceptional case, $r(D) \geq 2^{aS_X - B_X}$. [Revision 5]*

Remark 5.7 (Arbitrary irregular words: open). For a general confined word the coarse anchor $\xi_N(D) = -C_N(D)3^{-N}$ changes with N : there is no single fixed rational whose periodicity can be invoked. This is the moving-anchor sector, and it is where actual Collatz orbits live. [Revision 5]

5.2 What the height invariant actually separates

Revision 5 framed the split as periodic versus irregular. Later audits sharpened it. Every realizer condition can be written

$$2^{S_N} \mid A + \chi\Lambda$$

for integers A, Λ *generated by the word*: for a general word $(A, \Lambda) = (C_N, 3^N)$, while for a purely periodic word of period p the carry collapses to $C_{Np} \cdot \Delta = c_0(3^{Np} - 2^{N\sigma})$ with c_0, Δ independent of N . The only general tool converting divisibility into a size statement is that a nonzero multiple of 2^m has absolute value at least 2^m , which gives a *dichotomy*: either $A + \chi\Lambda = 0$, pinning χ to the single rational $-A/\Lambda$, or

$$|\chi| \geq 2^{S_N}/H - 1, \quad H = \max(|A|, |\Lambda|).$$

Writing $\rho_N = \log_2 H/S_N$, the second branch is useful exactly when ρ_N is bounded away from 1.

Observation 5.8 (Height amortization, not complexity). Periodic, eventually periodic and one-defect words have $\rho_N \rightarrow 0$, at rate $1/(\text{number of repetitions})$: a fixed height is amortized over unboundedly many periods. Arbitrary confined words supply only $(C_N, 3^N)$, with $\log_2 H = \alpha N \geq S_N$, so $\rho_N \geq 1$ and the second branch is vacuous. The separating invariant is therefore height amortization, not symbolic complexity: the Beatty/Sturmian zero-corridor word, of minimal aperiodic subword complexity $p(n) = n + 1$, already has $\rho_N > 1$ at every depth examined, while its periodic continued-fraction approximants keep $\rho_N \approx 0.26$. [structural interpretation, supported by exact identities and by computation; not a theorem about all words]

5.3 Endpoint depth and the frontier defect

Write $E(D) = S - \log_2 r(D)$ for the endpoint depth (Remark 2.3) and $g(D) = S - \lfloor \log_2 r(D) \rfloor$ for the discrete terminal gap. Then

$$E(D) \leq g(D) < E(D) + 1, \quad \text{i.e. } g = \lceil E \rceil,$$

the two agreeing only when $r(D)$ is a power of 2. [proved; formalized, §14]

Remark 5.9 (Correction to Revision 5's Open Problem E). Revision 5 asked whether the frontier-defect target $E(D) \leq H_2(\rho_c)S + \Phi(N)$, $\Phi = o(N)$, is genuinely easier than an exponential floor, and recorded that question as open. It is not easier in any useful sense: by the bracket above the statement is equivalent, up to an additive 1, to $g(D) \leq H_2(\rho_c)S + \Phi(N)$, and under confinement $S \leq \lfloor \alpha N \rfloor$ this is an exponential floor with $\varepsilon = I_{\text{Collatz}} + o(1)$. Open Problem E of Revision 5 is therefore the sharp-leading form of Open Problem C, not an independent weaker route, and is restated as such in §13. [correction to Revision 5]

6 The time axis

6.1 Three temporal scales

Definition 6.1. Fix $c \geq 0$ and an odd seed m .

$$\begin{aligned} L_c(m) &= \max\{N \geq 0 : R_j(m) \leq c \text{ for all } 0 \leq j \leq N\} && \text{(single-window lifetime),} \\ \mathcal{O}_c(m) &= \#\{n \geq 0 : R_n(m) \leq c\} && \text{(total occupation),} \\ \tau_c(m) &= \inf\{n \geq 1 : R_n(m) > c\} && \text{(drift exit time).} \end{aligned}$$

These answer different questions. L_c measures the initial confined episode; $\mathcal{O}_c$ sums all episodes, including separated later returns; τ_c asks only whether the corridor is ever left. Always $L_c(m) \leq \mathcal{O}_c(m)$ and $\tau_c(m) = L_c(m) + 1$ when finite.

Remark 6.2 (Single window does not control occupation). The inequality $\mathcal{O}_c \geq L_c$ is strict in general, and dramatically so: $m_0 = 285175$ has $L_1 = 14$ but $\mathcal{O}_1 = 97$. Consequently a complete single-window theory — even an optimal one — does not yield EOC. [Revision 5, computational example retained]

6.2 Residue–time duality

Definition 6.3. Fix $c > 0$. A word D of length N is *c-confined* if $R_j(D) \leq c$ for $0 \leq j \leq N$. For the c -confined family at depth N , $r_{\min}(N, c) := \min_D r(D)$.

Proposition 6.4 (Record-chronology inversion). $r_{\min}(N, c) = \min\{m \text{ odd} : L_c(m) \geq N\}$. [Revision 5, Proposition 5.6]

Proposition 6.5 (Single-window equivalence). Fix $c > 0$ and $\varepsilon \in (0, 2]$. The following are equivalent:

- (i) $r(D) \geq 2^{\varepsilon N}$ for every c -confined D of length N with $r(D) \geq 3$;
- (ii) $L_c(m_0) \leq \varepsilon^{-1} \log_2 m_0$ for every odd $m_0 \geq 3$.

[Revision 5, Proposition 5.7]

Revision 5 presented Propositions 6.4 and 6.5 as technical observations inside the residue section. They deserve a more prominent role: together they say that *the residue and time axes are the same axis in two coordinates*. Every statement about how small a realizer can be is a statement about how long an orbit can stay confined, and conversely. Revision 6 takes this as an organizing principle rather than a lemma.

6.3 The general transfer principle

Proposition 6.5 matches one specific pair of scales. The inversion is general.

Proposition 6.6 (Lifetime–frontier transfer). Let $F : \mathbb{Z}_{>0} \rightarrow \mathbb{Z}_{>0}$ be nondecreasing and suppose $L_c(m) \leq F(m)$ for every odd m . Then

$$r_{\min}(N, c) \geq \min\{m : F(m) \geq N\} = F^{-1}(N).$$

[proved here; the finite form is formalized, §14]

Proof. By Proposition 6.4, $r_{\min}(N, c)$ is the least odd m with $L_c(m) \geq N$. If $m < F^{-1}(N)$ then $F(m) < N$ by monotonicity, so $L_c(m) \leq F(m) < N$; hence no such m qualifies. $\square$

Writing the hypothesis in terms of $x = \log_2 m$ gives the scales of interest.

Lifetime hypothesis	Frontier obtained	Character
$L_c(m) \leq K \log_2 m$	$\log_2 r_{\min}(N, c) \geq N/K$	exponential (Open Problem C)
$L_c(m) \leq K(\log_2 m)^p, p > 1$	$\log_2 r_{\min}(N, c) \geq (N/K)^{1/p}$	stretched exponential
$L_c(m) \leq Km^\theta, \theta > 0$	$r_{\min}(N, c) \geq (N/K)^{1/\theta}$	polynomial
$L_c(m) \leq Km$	$r_{\min}(N, c) \geq N/K$	linear

Remark 6.7 (A graded programme). This table is the most constructive consequence of the recent audits. EOC's residue axis has usually been posed all-or-nothing — prove an exponential floor or prove nothing. Transfer shows otherwise: *any* unconditional pointwise lifetime bound, however weak, yields a corresponding unconditional frontier bound. A bound as weak as $L_c(m) \leq Km$ would already give a linear frontier, and none is currently known. We record this as Open Problem D. [proved; the assessment of what is currently known is in §6.8]

Remark 6.8 (No unconditional lifetime bound can be easy). No unconditional bound $L_0(m) \leq F(m)$ is currently known, and none can be proved without excluding divergent orbits: applied to the restarted seed m^* of Proposition 4.10, which has $L_0(m^*) = \infty$, any such bound is immediately contradicted. The transfer table therefore grades a programme whose every entry is open. [proved]

6.4 Drift exit and the future-minimum theorem

Proposition 6.9 (Future minimum). *If $R_n \leq 0$ then $m_n > m_0$. Hence an orbit with $R_n \leq 0$ for all $n \geq 1$ has its seed as a strict minimum of its entire future. Contrapositively, $m_n \leq m_0$ already forces $R_n > 0$.*

Proof. Two proofs. Analytically, Lemma 2.2 gives $R_n = \log_2(m_0/m_n) + E_n$ with $E_n > 0$ for $n \geq 1$, so $m_n \leq m_0$ implies $R_n \geq E_n > 0$. Integrally, Lemma 2.4 and $C_n > 0$ give $2^{S_n}m_n = 3^n m_0 + C_n > 3^n m_0 \geq 2^{S_n}m_0$ when $2^{S_n} \leq 3^n$, so $m_n > m_0$. $\square$

Remark 6.10 (Provenance). This is *not* a new theorem. The analytic proof is an immediate consequence of the Eliahou–Rozier identity already in Revision 5 (Lemma 2.2), and is close in content to Revision 5's wedge (Lemma 2.6). What is new is only the second proof, which uses no logarithms and is therefore convenient to formalize, and the role the statement plays below. [reclassification]

Proposition 6.11 (Drift exit is weaker than ordinary descent). *Let $\sigma(m) = \inf\{n \geq 1 : m_n < m\}$ be the (Terras) stopping time. Then $m_n < m_0 \iff R_n > E_n$, whence $\tau_0(m) \leq \sigma(m)$, and $\sigma(m) < \infty \Rightarrow \tau_0(m) < \infty$. The inequality is strict at cycle minima: for $m = 1$, $R_n = E_n = (2 - \alpha)n$ exactly, so $\tau_0(1) = 1$ while $\sigma(1) = \infty$. [proved]*

Remark 6.12 (How much weaker, quantitatively). Off cycle minima the gap is small. At $n = \tau_0$, strictness needs $R_n \leq E_n$; under confinement for $j < n$, Proposition 6.9 gives $m_j > m_0$, so $E_n \leq (3 \ln 2)^{-1} \sum_{j < n} 1/m_j \leq 0.4809 n/m_0$, while integrality of S_n with $R_n > 0$ forces $R_n \geq 1 - \{\alpha n\}$. Strictness at depth n therefore requires

$$r_{\min}(n, 0) \leq m_0 \leq 0.4809 n / (1 - \{\alpha n\}),$$

whose right side is polynomial in n because α has finite irrationality measure. So *any* superpolynomial realizer floor forces $\tau_0 = \sigma$ away from cycle minima. An exhaustive search of every odd m below this size bound at every depth $n < 200$ where the bound is not already violated found no strict example. [the inequality is proved; the search is a computational diagnostic]

6.5 Universal drift exit and divergence

Definition 6.13 (Universal drift exit).

$$(DE) \quad \text{for every odd } m \geq 1 \text{ there is } n \geq 1 \text{ with } 3^n < 2^{S_n(m)}.$$

Equivalently, by Lemma 2.5, $\tau_0(m) < \infty$ for every m . The phrase “drift exit time” is terminology used here; we found no standard name for this quantity in the literature (§16).

Theorem 6.14 (Divergence equivalence). *(DE) holds if and only if there is no divergent positive accelerated orbit.*

Proof. ($\Leftarrow$, contrapositive.) If a divergent orbit exists, Proposition 4.10 produces an actual positive odd integer m^* with $R'_k < 0$ for every $k \geq 1$, so $\tau_0(m^*) = \infty$ and (DE) fails.

($\Rightarrow$, contrapositive.) Suppose (DE) fails at m , so $2^{S_n} \leq 3^n$ for all $n \geq 1$. The orbit cannot repeat a state: a repeat makes it eventually periodic, and Lemma 4.2 then gives $R_N \rightarrow +\infty$, contradicting $R_n \leq 0$. So the orbit is injective, hence divergent. $\square$

Corollary 6.15 (Cycle compatibility). *(DE) is compatible with hypothetical nontrivial positive cycles: by Lemma 4.2 every point of a cycle, and every transient point entering one, has finite τ_0 . Hence (DE) is strictly weaker than the Collatz conjecture, which also excludes cycles.*

Remark 6.16 (Provenance, emphatically). Neither direction is new mathematics. The forward direction is the zero-corridor tail theorem of the companion Curry foundation, already proved and formalized there; the reverse direction is Revision 5’s Lemma 4.2. Moreover the *statement* (DE) is, in negated form, exactly the frontier already posed in that companion work: “determine whether an infinite zero-confined valuation word can be realized by a positive ordinary integer.” Revision 6 contributes the reclassification — recognising this as the qualitative endpoint of the *time* axis rather than an isolated realizability question — together with the integral form and its exact position in the hierarchy of §10. We do not coin a new acronym for it; in particular it is distinct from the repository’s ZCRE statement, which is an equivalence about bounded prefix realizers. [reclassification, not a new theorem]

7 Launch renormalization: the bridge to actual orbits

Finite word and residue analysis becomes powerful exactly when it produces an *actual* positive-orbit segment. Launch renormalization does this. We summarize only what the programme map needs; the full treatment is the companion note [17].

Theorem 7.1 (Launch-state collapse, [17]). *Let D be a word and j any index with $2^{S_j} > r(D)$. Then $m_j^* := (3^j r(D) + C_j(D))/2^{S_j}$ is a positive integer, in fact the genuine j -th accelerated iterate of $r(D)$, and the letters $(d_{j+1}, \dots, d_N)$ are exactly the accelerated valuation word of m_j^* . [companion work]*

Call D of length N ε -dangerous if it is c -confined with $\log_2 r(D) < \varepsilon N$. For such a word the pinning index t_1 and the launch data satisfy

$$t_1 \leq \varepsilon N + O(1), \quad L \geq (1 - \varepsilon)N - O(1), \quad \log_2 m_{t_1}^* \leq \alpha \varepsilon N + O(\log N),$$

with inherited *relative* corridor $A \leq c + (\alpha - 1)\varepsilon N + O(1)$. The leading coefficient in the launch-size bound is $\alpha \varepsilon$, not ε . The residual irregular class is renormalization-stable: the forced suffix of a dangerous word in that class is again a small integer with a long, near-critically confined injective orbit segment, so finite structural enumeration cannot close the programme [17].

7.1 Reclassifying the corridor-uniform hypothesis

The companion note closes the route conditionally on a corridor-uniform lifetime hypothesis: for some $K < \infty$, every injective initial segment of m with $R_i \leq A$ for $0 \leq i \leq L$ has $L \leq K(\log_2 m + A)$. This reads as a mild statement about corridors. It is not.

Proposition 7.2 (The corridor is automatic). *For every positive orbit and every n , $2^{S_n} \leq 3^n m_0 + C_n$, i.e. $R_n \leq \log_2 m_0 + E_n$; and equality holds as soon as the orbit reaches 1. Hence the least valid corridor width for a full orbit is exactly $\log_2 m_0 + E$, not merely bounded by it. [proved; formalized, §14]*

Proof. Immediate from Lemma 2.4 and $m_n \geq 1$, with equality when $m_n = 1$. $\square$

Corollary 7.3 (Reclassification). *Instantiating the corridor-uniform hypothesis at its own automatic corridor gives $L \leq K(2\log_2 m + E)$. For an injective segment the iterates are distinct odd integers, so $E_L \leq (3\ln 2)^{-1} \sum_{j < L} 1/m_j \leq 0.2404 \ln L + O(1)$, which grows slower than L and therefore bounds L . Consequently the hypothesis implies that no accelerated orbit is injective forever — i.e. it implies the absence of divergent orbits — and, with that, an $O(\log m)$ bound on accelerated total stopping time. Conversely a bound $L \leq K' \log_2 m$ returns the hypothesis for every $A \geq 0$. [proved]*

So the corridor-uniform hypothesis is *noncircular* with respect to Open Problem C — it is not a repackaging of the realizer floor — but it is substantially *stronger* than the Collatz conjecture. It should not be presented as a plausibly easier closing hypothesis. The companion note’s weakened variant, which exempts segments majority-covered by a bounded-period terminal repetition, does not change this: over all odd $m < 60001$ the exemption applies to none of the 29 999 orbits tested. [computational diagnostic only]

8 Self-financing recovery

A natural strategy for the time axis is to argue that a deep negative drift excursion cannot be repaid, because the recovery would demand an impossibly large valuation. This fails at its source.

Proposition 8.1 (Self-financing). *A single step of valuation d from state m_n achieves drift exit as soon as $2^d m_0 > 3m_n$. The available valuation is capped by $2^{d_n} \leq 3m_n + 1$, since $2^{d_n} \mid 3m_n + 1$. [proved; formalized, §14]*

Proof. For the cap, $2^{d_n} m_{n+1} = 3m_n + 1$ with $m_{n+1} \geq 1$. For the sufficient condition, multiply the target $3^{n+1} < 2^{S_{n+1}}$ by m_n : by Lemma 2.4, $2^{S_{n+1}} m_n = 2^d 2^{S_n} m_n \geq 2^d 3^n m_0 > 3 \cdot 3^n m_n = 3^{n+1} m_n$ whenever $2^d m_0 > 3m_n$. $\square$

The cap therefore exceeds the sufficient threshold $3m_n/m_0$ by a factor of about m_0 . The underlying reason is the identity $\log_2 m_n = \log_2 m_0 - R_n + E_n$ of Lemma 2.2: sustained negative drift makes the ordinary iterate *larger*, and the ordinary height so created is exactly the resource that permits the large valuations needed to repay the deficit.

Remark 8.2. What this rules out is a class of arguments, not a theorem: any argument of the form “a sufficiently deep drift excursion cannot recover, because the required valuation exceeds what the current iterate can supply” is blocked before it starts. Over all odd $m < 200001$, the minimum over all steps of $\log_2(3m_n + 1) - (\alpha - R_n)$ — available valuation minus what a one-step return to $R = 0$ requires — is $+1.83$, never negative. [the inequality is proved; the margin is a computational diagnostic]

9 Boundaries of the programme

This section collects the strongest no-go results. Their scope is deliberately narrow: each says that a specific class of information cannot distinguish positive-integer realization. None says that Collatz or EOC is unprovable.

9.1 Finite cylinders do not encode infinite realization

Proposition 9.1. *For $k \geq 1$, every odd residue class mod 2^k contains arbitrarily large positive odd integers. Consequently every exact-realizer cylinder is nonempty and unbounded, and no property of integers determined by finitely many 2-adic digits can be equivalent to positivity. [proved; formalized, §14]*

Proof. $r + 2^k(B + 1)$ lies in the class and exceeds B ; oddness is preserved since $2 \mid 2^k$. For the second claim, 1 and $1 - 2^k$ share a class mod 2^k and differ in sign. $\square$

The consequence is a quantifier distinction that should be displayed:

$\forall N \exists m_N$ (free)	versus	$\exists m \forall N$ (the whole problem)
--------------------------------	--------	---

A valuation prefix of length N pins exactly one odd class mod 2^{S_N+1} , which contains unboundedly many positive integers; so finite-prefix realizability never obstructs. The content lies entirely in whether *one* integer works for all N , which is the bounded-prefix-realizer equivalence of the companion work.

Remark 9.2 (Scope). This does *not* say finite information is useless. Finite data can still prove quantitative lower bounds — Open Problem C is precisely such a statement at each finite N . The claim is only about *deciding* the infinite realization property. [scope statement]

9.2 Coordinate sufficiency is not independent information

Several coordinates introduced in the programme turn out to be invertible reparameterizations of the exact realizer. The ledger below records each, with the reference where the collapse is proved.

Coordinate	Status
Normalized carry Z_N	Same moving-anchor information as $-C_N 3^{-N}$; no independent obstruction.
Lift digits t_N	Exactly the binary digits of a fixed realizer, read in blocks cut at the valuation sums.
Eventually-zero lift tail	Equivalent to eventual constancy of the prefix realizers, hence to positive-integer realization.
Terminal zero gap $g(D)$	$\lceil E(D) \rceil$; a restatement of the target, not a mechanism (§5.3).
Transport depths F, M, H	Exact bookkeeping. On a genuine orbit the matching precision M_j is the length of the zero-run of the <i>seed</i> 's binary expansion beginning at bit S_j .
Cumulative excess regeneration	Bit intervals at distinct failures are disjoint, so the total is at most $\lfloor \log_2 m_0 \rfloor + 1$.
$B_N = C_N + 3^N \chi$	Equals $2^{S_N} m_N$ exactly for a realizing χ ; hence $v_2(B_N) = S_N$ exactly and $\log_2 B_N - v_2(B_N) = \log_2 m_N$.

The last row deserves emphasis. The Archimedean size and the 2-adic valuation of B_N are *complementary factors of one exact identity*, not two independent constraints; knowing the word

and either one determines the other. This is why attempts to intersect a p -adic condition with an Archimedean one at this object produced nothing: there was only ever one condition. It also explains why the product formula yields exactly the trivial bound $v_2(B) \leq \log_2 |B|$ here, and why the odd-prime support of B_N — being that of the arbitrary odd integer m_N — is unconstrained. [proved in companion audits; identities verified computationally without exception]

9.3 Global word shadows do not characterize positive realization

Theorem 9.3 (Word-shadow cardinality barrier). *For $b \subseteq \mathbb{N}$ define the valuation word (digits indexed from 0)*

$$w_b(i) = \begin{cases} 2 & \text{if } i \equiv 2 \pmod{3} \text{ and } \lfloor i/3 \rfloor \in b, \\ 1 & \text{otherwise.} \end{cases}$$

Then every w_b is an admissible valuation word satisfying

- (a) $\sum_{i < n} w_b(i) \leq n + \lfloor n/3 \rfloor$, hence $2^{S_n} \leq 3^n$ for every n : zero confinement, in the integral form of Lemma 2.5;
- (b) $R_n \leq (4/3 - \alpha)n = -0.2516 \dots n \rightarrow -\infty$ linearly;
- (c) $\sum_n 2^{R_n} < \infty$;
- (d) $\#\{n : R_n \geq -G\} = O(G)$.

The map $b \mapsto w_b$ is injective, so the family has cardinality $2^{\aleph_0}$. Since one positive integer realizes at most one infinite valuation word, all but countably many members have no positive-integer realizer. [proved; formalized, §14]

Proof. (a) At most one digit in three is a 2, so $S_n \leq n + \lfloor n/3 \rfloor$; and $2^{n+\lfloor n/3 \rfloor} \leq 3^n$ follows by cubing both sides, since $3(n + \lfloor n/3 \rfloor) \leq 4n$ and $16 \leq 27$. (b) $4/3 < \alpha$. (c) The bound in (b) makes 2^{R_n} dominated by a geometric series. (d) $R_n \geq -G$ forces $n \leq G/0.2516$. Injectivity: evaluate w_b at $i = 3k + 2$. The last sentence is Cantor's theorem: a realizing assignment would give an injection $\mathcal{P}(\mathbb{N}) \rightarrow \mathbb{N}$. $\square$

Remark 9.4 (What this does and does not establish). Every global condition expressible from the valuation word alone — zero confinement, linear negative drift, summability of $\sum 2^{R_n}$, Curry-scale occupation shadows, aperiodicity, subword complexity — is satisfied by all members of this family, while all but countably many are unrealizable. Hence *no collection of word-level global properties can characterize positive-integer realization*. This is a statement about characterization, not about quantitative bounds: it does not preclude word-level information yielding realizer lower bounds. [scope statement]

Remark 9.5 (Explicit small witnesses). The barrier is not an artefact of uncountability. The all-ones word (realizer -1) and the word $1, 2, 1, 2, \dots$ (realizer -5) are both zero-confined with $R_n \rightarrow -\infty$ linearly, and neither has a positive-integer realizer. [computational verification of an exact statement]

9.4 Population sparsity is not pointwise placement

Theorem 4.7 and the entropy constant of §3.3 are population statements: they count, respectively, orbit values in a window and words in a shell. Open Problem C is a statement about the *minimum* over a family. A density-zero exceptional set may contain every least-realizer champion, and no argument currently rules this out. The same gap defeats density-one results in the literature (§16) as applied to launch integers, which by construction form an exceptional set. [structural interpretation]

9.5 Deep drift creates its own recovery capacity

§8. [proved]

9.6 Single-window lifetime does not control occupation

Remark 6.2. [Revision 5, computational example]

Remark 9.6 (Net effect). These boundaries narrow the remaining search to actual positive-orbit arithmetic and to genuinely pointwise or cross-scale mechanisms. They do not constitute an argument that the programme cannot succeed.

10 The programme hierarchy

Level	Statement	Status
0	Finite word realizability	Solved exactly (Prop. 5.2); every cylinder is nonempty and unbounded (Prop. 9.1).
1	Universal drift exit (DE)	Equivalent to excluding divergent orbits (Thm. 6.14); compatible with nontrivial cycles; <i>open</i> .
2	Quantitative single-window: bounds on $L_c(m)$	Transfers to frontier bounds (Prop. 6.6); no unconditional bound known (Rem. 6.8).
3	Exponential single-window floor	Open Problem C; equivalent to $L_c(m) = O(\log m)$ (Prop. 6.5).
4	Total occupation $\mathcal{O}_c(m) = O(\log m)$	Full EOC (Conjecture 3.1); strictly stronger than Level 3 (Rem. 6.2).
5	Effective EOC + finite cycle verification	Collatz (Thm. 3.6, Cor. 3.7).

Levels $5 \Rightarrow 4 \Rightarrow 3 \Rightarrow 2$ and $3 \Rightarrow 1$ hold; $2 \Rightarrow 1$ holds by Remark 6.8. No converse holds in general. Level 1 is strictly weaker than Collatz (Cor. 6.15), and Level 4 is strictly stronger than Level 3.

11 The programme map

Axis	Object	Exact bridge	Strongest proved	Open target	Closed routes
Occupation	$\mathcal{O}_c(m)$	Thm. 3.6 to Collatz	Reduction to a finite cycle search, given effective constants	Separated-return control	—
Time	$L_c(m), \tau_0(m)$	Record chronology (Prop. 6.4); transfer (Prop. 6.6)	(DE) $\iff$ no divergence (Thm. 6.14)	Any unconditional bound on L_c	Self-financing recovery blocks excursion arguments (§8)
Residue	$r_{\min}(N, c), r(D)$	Exact congruence (Prop. 5.2) + inversion	Fixed-anchor sectors (Thms. 5.4–5.5, Prop. 5.6)	Arbitrary irregular floor	Coordinate collapse (§9.2)
Divergence	R_n, Q_n , value set	Curry sparsity; last-maximum reduction (Prop. 4.10)	Reciprocal summability; $B < 1/\beta_*$ (Thm. 4.9)	Exclude divergence	Word shadows insufficient (Thm. 9.3)
Symbolic	Valuation words; return labels	Realizer correspondence	Finite and infinite symbolic freedom	Positive realization	Cardinality barrier (Thm. 9.3)

11.1 Symbolic freedom, finite and infinite

Revision 5 established that no *finite* forbidden-pattern mechanism can settle pointwise balance for a natural mod-32 return observable: every finite binary history is realizable by a genuine orbit,

with prescribed events consecutive, and this was formally verified. The symbolic conclusion is now stronger. A machine-checked construction produces an infinite zero-confined valuation sequence carrying arbitrary prescribed binary labels at designated event positions: a *full shift* at the word level, inside the zero corridor.

The distinction that matters is therefore sharper than “no finite forbidden patterns”: even substantial *infinite* symbolic freedom coexists with zero confinement. Positive-integer realization of such words remains open and, by Theorem 9.3, cannot be decided by word-level information at all. The missing restriction is not local symbolic dynamics. [finite universality: Revision 5, formalized; infinite full shift: companion formalization]

12 Closed routes

For each mechanism, what it achieved and where it stops. Status codes: T formal result, C computational diagnostic, I interpretive closure.

	Mechanism	Where it stops	
1	Fixed-modulus drift packing	Terminal at 8/9 (Rem. 4.6); surpassed by a different mechanism	T
2	Bulk entropy / I_{Collatz}	Population scale only; no pointwise anti-concentration	I
3	Shell injectivity	Counts, does not place	I
4	Moving-anchor collision law	Symbolic prefix geometry only	I
5	Chang finite/local exclusions	Finite <i>and</i> infinite symbolic freedom (§11.1)	T
6	Lift digits	Binary blocks of the realizer (§9.2)	T
7	Terminal zero gap	[E]: target, not mechanism (§5.3)	T
8	Normalized carry	Same anchor information	T
9	Transport survival budget	Exact bookkeeping; on genuine orbits matching precision collapses to seed bits, cumulative excess $O(\log m_0)$	T
10	External p -adic arithmetic	Wrong complexity scale (§12.1)	I
11	Global word shadows	Cardinality barrier (Thm. 9.3)	T
12	Corridor-uniform lifetime hypothesis	Equivalent in strength to $O(\log n)$ stopping (Cor. 7.3)	T
13	Deep-drift recovery impossibility	Self-financing (§8)	T

12.1 External arithmetic

We record only the durable conclusion. The natural cleared-denominator object attached to a confined word is $B_N(D, \chi) = C_N + 3^N \chi$, an $(N+1)$ -term sum of $\{2, 3\}$ -units whose 2-adic valuation one would like to bound. Three families of deep results were examined against it. p -adic lower bounds for linear forms in logarithms bound $\alpha_1^{b_1} \cdots \alpha_n^{b_n} - 1$, a *product* minus one, never a sum of three or more terms; their constants are at best geometric in the number of logarithms; and at $p = 2$ with 2 among the bases the hypotheses fail and the quantity is identically zero. Quantitative results on S -unit equations bound the *number* of non-degenerate solutions, not their size, and are ineffective beyond two variables; moreover explicit lower bounds for that count forbid any bound uniform in the number of variables. The p -adic Subspace Theorem does apply formally and yields a bound of the right shape, but it is ineffective, its subspace count and height threshold depend on the dimension, and here the dimension is $N+1$. In every case the obstruction is the same: the number of additive terms grows with N , while the available theorems are quantified “for each fixed dimension” with constants that provably blow up. [applicability audit; external theorems cited but not reproved]

13 Open problems

Open Problem A (Effective EOC constants). Establish Conjecture 3.2 at any explicit (c, K, C) , or any explicit upper bound on admissible constants in Conjecture 3.1. This closes the Collatz

implication outright (Corollary 3.7).

Open Problem B (Cycle exclusion below threshold). Show no nontrivial accelerated cycle has minimum member below $m^*(K, C, c)$. Settled for $m^* \leq 2^{71}$ by [13]; open beyond.

Open Problem C (Exponential realizer floor / logarithmic lifetime). Find $\varepsilon > 0$ with, equivalently (Proposition 6.5),

$$(\text{residue}) \ r_{\min}(N, c) \geq 2^{\varepsilon N - O(1)} \iff (\text{time}) \ L_c(m) \leq \varepsilon^{-1} \log_2 m + O(1),$$

for all relevant c -confined words. This is the residue–time core of EOC. Revision 5’s separate Open Problems C and D are the same question at two levels of precision and are merged here; Revision 5’s Open Problem E (frontier defect) is its sharp-leading form, not an independent weaker target (Remark 5.9).

Open Problem D (Intermediate time-axis bounds). Prove *any* unconditional pointwise upper bound on $L_c(m)$. By Proposition 6.6 even $L_c(m) \leq Km$ yields a linear frontier, and $L_c(m) \leq K(\log_2 m)^2$ yields a stretched-exponential one. Nothing of this kind is known (Remark 6.8), and any such bound would be new.

Open Problem E (Universal drift exit). Show that for every positive odd m there is $n \geq 1$ with $3^n < 2^{S_n(m)}$. By Theorem 6.14 this is equivalent to excluding divergent orbits and is strictly weaker than Collatz. This is the previously posed zero-corridor realizability frontier, reclassified as the qualitative endpoint of the time axis (Remark 6.16); it is not newly invented here.

Open Problem F (Separated returns). Control the number of separated confined episodes. Single-window theory does not capture this (Remark 6.2); full EOC requires both episode length and episode count.

Open Problem G (Density-one EOC). Can a density-one form of EOC be proved by the methods of Tao [9]? This remains meaningful: divergent-orbit sparsity says nothing about the occupation of orbits that do converge.

Open Problem H (Pointwise return balance). For an infinite aperiodic orbit, does the natural mod-32 return label equidistribute along the orbit? By §11.1 no finite — and no word-level infinite — symbolic restriction settles this. Demoted here from a top-level EOC problem to an auxiliary one.

14 Formal verification status

The accompanying Lean 4 development (Mathlib) formalizes the elementary bridges of this paper. The following are machine-checked, with no `sorry`, `admit`, added axiom, or `opaque` escape; each depends only on the standard axioms `propext`, `Classical.choice`, `Quot.sound` (several on a strict subset).

Module	Paper reference	Content
CurryFoundation	Prop. 4.10	Curry identities; last-global-maximum existence; zero-corridor tail; deficit recurrence.
ChangHistory, ChangFullShift	§11.1	Finite return-history universality; infinite zero-corridor full shift.
ZCRERealizerGrowth	§9.1	Bounded prefix realizers $\iff$ positive-integer realizability.
RealizerLiftDigit, TerminalZeroGap	§9.2, §5.3	Lift digits are binary blocks; discrete gap and its bracket $g = \lceil E \rceil$.
TransportRegeneration, SuperBudgetCancellation	§9.2	Transport bookkeeping; forced/excess split; disjoint-run bound.
ConstraintHeight	§5.2	Height inequality and the separation criterion.
GlobalSeparation	Prop. 9.1, Thm. 9.3	Cylinder positivity; finite-precision sign impossibility; the family w_b , its confinement and injectivity; the countability barrier.
OrbitLifetime	Lem. 2.4, Prop. 7.2	Aggregate identity; growth envelope; automatic corridor.
DriftExit	Props. 6.9, 8.1, 6.6	Future-minimum theorem; cycle drift; self-financing inequality; frontier inversion.

Statements in this paper that are *not* formalized include Theorem 4.7 and Proposition 4.8 (external input), the proved-sector floors of §5 (companion note), and every conjecture.

15 Computational calibration

We retain only computations that materially support interpretation, and label each as a diagnostic.

- *Frontier records.* A direct seed scan, justified by Proposition 6.4, produces the least realizers of confined words at each depth. The maximum observed ratio of occupation to $\log_2$ seed remains under 10 in the scanned range.
- *Identity verification.* The identities of Lemmas 2.2, 2.4 and 2.5, the future-minimum theorem, the descent criterion $m_n < m_0 \iff R_n > E_n$, and the factorization $B_N = 2^{S_N} m_N$ were each verified without exception over the ranges scanned.
- *Self-financing margin.* §8, Remark 8.2.
- *Drift-exit diagnostics.* $\tau_0 = \sigma$ held for every odd seed tested away from cycle minima, consistent with Remark 6.12.

Remark 15.1 (Discipline). We make no empirical saturation claims. In particular we do not assert that any normalized stopping-time or lifetime ratio is bounded: a comparison of a limited scan against published record tables corrected an earlier such claim in the underlying audits. Where a ratio is quoted, the scanned range is stated and no trend is inferred beyond it. Units also require care: the classical total stopping time σ_∞ counts *even* steps — equivalently the valuation depth S_L — whereas L counts *odd* accelerated steps; the two must not be compared without conversion. [methodological]

16 Relation to prior work

The occupation functional and its four tiers are, to our knowledge, not isolated elsewhere; the Collatz reduction of Theorem 3.6 is ours. Lemma 2.2 is due to Eliahou [1] and Rozier [2]. Theorem 4.7 is Garcia–Tal [5] with Curry’s explicit form [6]. The 2-adic conjugacy underlying the realizer correspondence is Bernstein [19] and Bernstein–Lagarias [20]; cycle-structure bounds are Steiner [10], Simons–de Weger [11] and Hercher [12]; verification is Bařina [13]; stochastic models are Lagarias–Weiss [7] and Kontorovich–Lagarias [8]. Sturmian words are treated in Lothaire [14]. The return observable and its structural reduction are Chang’s [21, 22].

On the time axis specifically: the condition “ $\exists n : S_n > \alpha n$ ” does not appear in the surveyed literature under a standard name. It is not Terras’s stopping time [3], which is the condition $m_n < m_0$, i.e. $R_n > E_n$ (Proposition 6.11). The classical results on stopping time — Terras [3], Everett [4], and in a different framework Tao [9] — are density statements and carry no pointwise content; none bounds the number of steps an orbit may spend at or above its starting value in terms of that value, and such a bound would prove Collatz by strong induction. [literature audit]

17 Conclusion

EOC is best viewed as a programme with an exact residue–time duality and a strictly stronger global occupation layer. The residue axis is solved for finite realizability and for several fixed-anchor sectors, and remains open at the arbitrary-word frontier. The time axis separates into a qualitative drift-exit problem — equivalent, under the current divergence theory, to excluding divergent orbits, and strictly weaker than Collatz — and quantitative lifetime bounds that invert directly to realizer floors on a graded scale. Garcia–Tal/Curry supplies strong unconditional divergence-side sparsity and the last-maximum reduction, but not drift exit.

A sustained programme of structural audits has narrowed what can supply the missing information. Finite cylinders cannot encode infinite realization; several apparently independent coordinates are invertible reparameterizations of the exact realizer, and the Archimedean and 2-adic readings of the natural divisibility object are complementary factors of one identity; global valuation-word shadows cannot characterize positive realization, by an explicit uncountable family; launch renormalization validly reaches actual positive-orbit dynamics, but its original corridor-uniform closure is essentially an $O(\log n)$ stopping-time conjecture; and deep negative drift is self-financing with respect to the recovery valuations it requires.

The surviving frontier is therefore actual positive-orbit arithmetic, at three nested levels: qualitative exclusion of an infinite zero-corridor tail; quantitative control of single-window lifetime, at any scale; and, beyond that, recurrence control sufficient for total occupation. Neither EOC nor the Collatz conjecture is proved here.

References

- [1] S. Eliahou, *The $3x + 1$ problem: new lower bounds on nontrivial cycle lengths*, Discrete Mathematics **118** (1993), 45–56.
- [2] O. Rozier, *The $3x + 1$ problem: a lower bound hypothesis*, Functiones et Approximatio Commentarii Mathematici **56** (2017), no. 1, 7–23.
- [3] R. Terras, *A stopping time problem on the positive integers*, Acta Arithmetica **30** (1976), 241–252.

- [4] C. J. Everett, *Iteration of the number-theoretic function $f(2n) = n$, $f(2n+1) = 3n+2$* , Advances in Mathematics **25** (1977), 42–45.
- [5] M. V. P. Garcia and F. A. Tal, *A note on the generalized $3n+1$ problem*, Acta Arithmetica **90** (1999), no. 3, 245–250.
- [6] M. J. Curry, *An Explicit Windowed Sparsity Bound for Divergent $3x+1$ Orbits, with Logarithmic-Floor Exclusion beyond the Harmonic Barrier*, Zenodo, 2026.
- [7] J. C. Lagarias and A. Weiss, *The $3x+1$ problem: two stochastic models*, Annals of Applied Probability **2** (1992), 229–261.
- [8] A. V. Kontorovich and J. C. Lagarias, *Stochastic models for the $3x+1$ and $5x+1$ problems and related problems*, in: The Ultimate Challenge: The $3x+1$ Problem, Amer. Math. Soc., 2010, 131–188.
- [9] T. Tao, *Almost all orbits of the Collatz map attain almost bounded values*, Forum of Mathematics, Pi **10** (2022), e12.
- [10] R. P. Steiner, *A theorem on the Syracuse problem*, Proc. Seventh Manitoba Conf. Numer. Math. and Computing, 1977, 553–559.
- [11] J. Simons and B. de Weger, *Theoretical and computational bounds for m -cycles of the $3n+1$ problem*, Acta Arithmetica **117** (2005), 51–70.
- [12] C. Hercher, *There are no Collatz m -cycles with $m \leq 91$* , Journal of Integer Sequences **26** (2023), Article 23.3.5.
- [13] D. Bařina, *Improved verification limit for the convergence of the Collatz conjecture*, The Journal of Supercomputing **77** (2021), 2681–2688; and **81** (2025), 810.
- [14] M. Lothaire, *Algebraic Combinatorics on Words*, Cambridge University Press, 2002. (Sturmian words: Chapter 2, by J. Berstel and P. S  ebold.)
- [15] E. De Jes  s, *Periodic and Eventually Periodic Realizer Floors in the Accelerated Collatz Word Model*, companion note.
- [16] E. De Jes  s, *The Entropy-Deficit Identity in the Accelerated Collatz Word Model*, Zenodo, 2026.
- [17] E. De Jes  s, *Conditional Obstructions and Launch Renormalization in Accelerated Collatz Dynamics*, companion note, 2026.
- [18] E. De Jes  s, *Transport Deficits and 2-Adic Survival Budgets in Accelerated Collatz Words*, companion note, 2026.
- [19] D. J. Bernstein, *A non-iterative 2-adic statement of the $3N+1$ conjecture*, Proc. Amer. Math. Soc. **121** (1994), 405–408.
- [20] D. J. Bernstein and J. C. Lagarias, *The $3x+1$ conjugacy map*, Canadian Journal of Mathematics **48** (1996), 1154–1169.
- [21] E. Y. Chang, *A Structural Reduction of the Collatz Conjecture to One-Bit Orbit Mixing*, preprint.
- [22] E. Y. Chang, *Exploring Collatz Dynamics with Human–LLM Collaboration*, preprint.

A Band rigidity

We retain the following structural observation from Revision 5, which constrains how a drift path may sit inside a unit band.

Theorem A.1 (Band rigidity). *Let $\theta = \alpha - 1$. If a word's drift path satisfies $\beta \leq R_j < \beta + 1$ for all j in a window, then the valuation letters in that window are determined up to the Beatty structure of θ : the admissible letters are confined to $\{1, 2\}$ and their placement follows the Beatty gap sequence of α . [Revision 5, retained]*

The relevance to this revision is limited but worth recording: band rigidity says that a *narrow* corridor forces near-Sturmian structure, which is exactly the regime where Observation 5.8 shows height amortization fails. Narrow corridors are therefore not automatically easier.

B Claim-status conventions

Every substantive statement carries a bracketed status: [Revision 5] retained unchanged from the previous revision; [proved] proved in this paper; [companion work] proved in a cited companion note; [external, cited] a cited theorem of others, used but not reproved; [formalized] machine-checked, see §14; [computational diagnostic] supported by computation only, never promoted to a theorem; [structural interpretation] an organizing reading of proved facts; [reclassification] a previously known statement placed in a new role; [correction to Revision 5] a superseded claim, corrected here.