

Beyond Sturmian Words in the $3x+1$ Periodicity Conjecture: A Periodic-Approximation and Arithmetic-Height Bridge

Elias De Jesús
Independent Researcher
ORCID: [0009-0007-0190-9143](https://orcid.org/0009-0007-0190-9143)

September 2026

Revision 3, 30 September 2026 — corrected and extended; see §1

Abstract

Let $T : \mathbb{Z}_2 \rightarrow \mathbb{Z}_2$ be the $3x + 1$ map and Φ the Bernstein–Lagarias conjugacy map sending a parity vector $v \in \{0, 1\}^{\mathbb{N}}$ to the 2-adic integer it codes. Lagarias’s Periodicity Conjecture asserts that $\Phi(v) \in \mathbb{Q}$ only if v is eventually periodic; it is open in general. A companion unpublished paper by the present author proves it for every Sturmian parity word, by two independent mechanisms (a transported counting bound, and an effective Liouville-style argument using an initial square). This note consolidates a subsequent research program that pushes the periodic-approximation/arithmetic-height mechanism of the second route past the Sturmian case, into symbolic classes the counting route cannot reach. The main proved result: for *every* irrational slope with bounded partial quotients, every complementary symmetric Rote sequence (factor complexity exactly $2n$) built on *any* Sturmian word of that slope — any intercept, either mechanical convention, either seed, any shift — has irrational Φ -value. We give the full proof architecture, including a structural fact (an odd-weight transfer route’s doubled root always has density exactly $1/2$) that collapses an earlier, over-conservative repetition-exponent requirement down to the true one. We document, rather than hide, four working hypotheses this program formed and then disproved by direct adversarial testing. A late-stage exploratory probe into a conjectural symbolic-symmetry/arithmetic-admissibility tradeoff is reported honestly as inconclusive, not as a result.

Revision 3. Revision 3 extends the note past the complementary-symmetric-Rote rung and corrects Revision 2’s framing of it. Two new theorems are stated and proved: for every quadratic irrational $\alpha \in (0, 1)$ and every *rational* ρ , and then for every *real algebraic* ρ , the coding $s_n = 1 \iff \{n\alpha + \rho\} \in [0, \beta]$ with $\beta = \ln 2 / \ln 3$ has $\Phi(s) \notin \mathbb{Q}$ (§17, §18). These words are aperiodic, have factor complexity $2n$, and have lower ones-density exactly the critical value β — which matters, because the density coordinate, not the complexity coordinate, is what existing results read (§16). Two framing corrections follow from that. First, the conclusion of Revision 2’s headline theorem was already available: every complementary symmetric Rote sequence has ones-density exactly $1/2 < \beta$, and Monks–Yazinski (2004) Theorem 2.7(b) therefore reaches it. Revision 2’s theorem crosses the *counting* frontier, not the density one; what it contributes is an independent second proof by a different mechanism together with effective height floors, which a density argument does not supply. Second, Revision 2’s Open 6 asserted that the Thue–Morse word makes every repetition-based method structurally inapplicable; that is false, and §26 now says so. Revision 3 also records the analytic input for the new theorems honestly — a logarithmic-form lower bound with *algebraic* coefficients, for which the available estimate is stretched-exponential and not polynomial — and separates, throughout, what is effective in principle from what is numerically supplied. Novelty remains formally unresolved.

Revision 2. Revision 1 presented the general-intercept case as blocked by one unproved margin lemma (its Conjecture 12.1). That lemma is *not* open: it is Berthé–Holton–Zamboni (2006),

Propositions 5.1 and 5.2 — Section 5 of the same paper this program already cites as its primary source. Revision 2 therefore states the headline theorem at *every* intercept of every bounded-type slope, under either mechanical convention, and corrects four further defects of Revision 1: a mis-indexed convergent-denominator convention, a false “exceptional-orbit” reduction, a missing length factor in the surplus decomposition, and an overstated attainment claim for one of the two Ostrowski repetition expressions. The symbolic margin is credited to Berthé–Holton–Zamboni throughout; what is original here is the arithmetic application — the periodic-approximation/height bridge and the attained-witness bookkeeping that converts a limsup bound on the initial critical exponent into an unbounded surplus. §2 lists every change and says which Revision-1 conclusions survive; the full audit trail is [39, 42]. **This note does not prove, and does not claim progress toward proving, the Periodicity Conjecture in general**; a structural fact recorded in the underlying research record shows no purely repetition-based method, however extended, can reach every aperiodic word.

Contents

1	Revision 3: Corrections and New Results	3
2	Revision 2: Corrections to Revision 1	5
3	The Periodicity Conjecture and the Conjugacy Map	7
4	Two Routes to Irrationality	7
4.1	Route A — Counting	8
4.2	Route B — Periodic Approximation and Arithmetic Height	8
5	The Abstract Bridge Theorem	8
6	Discrepancy as the Height Bridge	9
7	Repetition–Height Competition	10
8	The Sturmian Foundation	10
9	Crossing the Counting Frontier: Complementary Symmetric Rote Sequences	11
10	The Odd-Weight Mechanism	11
11	Self-Correction History	12
12	The Bounded-Type CS Rote Theorem (Every Intercept)	13
13	Proof Architecture of the Bounded-Type Theorem	13
14	General Intercepts	14
15	Ostrowski Repetition Structure	17
16	The Density Coordinate, and Where the Residual Case Is	17

17 Critical-Density Rotation Codings: Rational Intercepts	18
17.1 The agreement excess is a first hitting time	19
17.2 The arc at 0: rational intercepts	20
17.3 The arc at β , the witnesses, and the theorem	20
18 Algebraic Intercepts	21
19 The Analytic Input, and Its Constants	22
19.1 Stretched exponential, not polynomial	23
19.2 An explicit constant	24
20 Finite Height Certificates	24
21 A Measurement, Not a Theorem: the Ridout Exponent	25
22 Interpretation: Symbolic, Statistical, and Arithmetic Space	26
23 The Drift/Resonance Coordinate	26
24 The Symmetry-Tradeoff Probe (Exploratory, Inconclusive)	27
25 A Transfer-Matrix Computational Tool	27
26 What Is Still Open	28
27 Future-Work Roadmap	30
Provenance and Verification	30
Acknowledgments	31

1 Revision 3: Corrections and New Results

Revision 3 supersedes Revision 2 in the following places. The mathematics of Revisions 1–2 is unchanged except where explicitly listed; the changes are two new theorems, two framing corrections, and several citation corrections.

New results

N1 — Critical-density rotation codings, rational intercept.

Theorem 17.5 (§17): for every quadratic irrational $\alpha \in (0, 1)$ and every $\rho \in \mathbb{Q}$ — including $\rho = 0$ — the word $s_n = 1 \iff \{n\alpha + \rho\} \in [0, \beta)$ has $\Phi(s) \notin \mathbb{Q}$.

N2 — Algebraic intercepts.

Theorem 18.2 (§18): the same for every *real algebraic* ρ ; and Corollary 18.3, for the interval $[u, u + \beta)$ whenever $u - \rho$ is algebraic, with ρ and u themselves unrestricted.

N3 — Where the residual case is.

§16 rebuilds the coverage map on lower ones-density rather than factor complexity, which is the coordinate the existing exclusion results actually read.

Corrections to Revision 2

D1 — The Thue–Morse scope claim is withdrawn.

Revision 2 §19 (Open 6) and §20 asserted that because the Thue–Morse word t has no square prefix, “every purely repetition-based method” is “structurally inapplicable to it by construction”. **That is false.** The bridge does not need a square; it needs periodic approximants whose 2-adic agreement outruns their height. For t these are supplied by the substitution itself: with $\mu(0) = 01$, $\mu(1) = 10$ one has $\text{lcp}(\mu(x), \mu(y)) = 2\text{lcp}(x, y)$, so iterating μ on any one approximant doubles the agreement while the height grows only as the length, and $\Phi(t) \notin \mathbb{Q}$ follows from the criterion of §5 without any square [52]. The corrected statement appears in §26. (The conclusion itself is not new: t has ones-density $1/2 < \beta$, so Monks–Yazinski [44] already covers it. See D2.)

D2 — The novelty framing of the CS Rote theorem is corrected.

Revision 2 presented the bounded-type CS Rote theorem as crossing a frontier. It crosses the *counting* frontier — factor complexity $2n$ lies above Dubickas’s threshold $1/\log_2(3/2) = 1.70951\dots$ — and only that one. Every complementary symmetric Rote sequence has ones-density exactly $1/2$, and Monks–Yazinski [44], Theorem 2.7(b) excludes every aperiodic word of lower ones-density below $\beta = \ln 2/\ln 3$ regardless of its complexity. **The conclusion of the headline theorem of Revisions 1–2 was therefore already available.** What the chain of §§9–15 contributes is an independent second proof by a different mechanism, and the effective height floors of §20, which a density argument does not give. Revision 2’s own §7 already states the density dichotomy correctly for the Sturmian family; the same sentence applies to the CS Rote family and is now stated.

D3 — López–Stoll: acknowledgement, and a statement of what is not used.

López–Stoll’s work on density and the $3x+1$ conjugacy map helped guide this programme toward the critical density $\beta = \ln 2/\ln 3$. We acknowledge that influence. The results presented here are established through the independent arguments given in this note and do not require their proposed upper-density exclusion [9]. The critical-density Sturmian word is theirs [8], and so is the observation that critical density is where the existing methods stop — which is why β is the place worth working. We take no position on their Theorem 1: we have neither established nor disproved it, and no step of any theorem in this note invokes it. The coverage statements of §16 are accordingly phrased against what is independently available, and no conclusion here changes in either case.

D4 — The foundation paper is now public; it remains an unrefereed self-citation.

[1] is deposited at version DOI 10.5281/zenodo.23019799 (concept DOI 10.5281/zenodo.22920055), CC BY 4.0. It is a preprint, it is not refereed, and it is by the present author. Every use of it in this note is a self-citation to unrefereed work and is treated with the same caution applied to any other preprint.

D5 — Theorem numbering.

Theorems in this note are numbered by section, so “Theorem 14.1” means the first theorem of §14 and carries no relation to any phase of the underlying research record. The research record’s informal labels map as follows:

research record	this note	content
Theorem 11	Theorem 17.5 (case $\alpha = \frac{\sqrt{5}-1}{2}$, $\rho = \frac{1}{7}$)	the named word
Theorem 12	Theorem 17.5	quadratic slope, rational intercept
Theorem 13	Theorem 18.2	quadratic slope, algebraic intercept
Corollary 13.1	Corollary 18.3	translated interval $[u, u + \beta)$

A correction carried in from the research record

D6 — A polynomial logarithmic-form bound was claimed and is withdrawn.

An intermediate stage of the work behind §17 asserted a bound $c_m \geq C_2 m^{-\kappa}$ for the quantity $\|m\alpha + \rho - \beta\|$, citing Baker’s *Transcendental Number Theory*. No such polynomial bound is available for *algebraic* coefficients. The correct citation, and the bound actually used in §19, is Baker [46], Theorem 2, whose right-hand side is $Ce^{-(\log B)^\kappa}$ — stretched exponential in $\log B$, strictly weaker than any power of B . The proof was rebuilt to need only $\log \ell(M) = o(M)$, which that bound supplies with room to spare. A polynomial bound would be the case $\kappa = 1$, which is Baker’s own conjecture and is not available. This note states no polynomial bound anywhere.

What survives unchanged

Theorem 5.1 (the abstract criterion), the discrepancy–height bound, the repetition–height competition, the exact XOR transfer, the density-1/2 structural fact, and the bounded-type CS Rote theorem at every intercept (§12) are all unchanged as *mathematics*. D2 changes what may be claimed for the last of these, not whether it is true. The five corrections of Revision 2 (§2) stand as recorded.

2 Revision 2: Corrections to Revision 1

Revision 1 of this note (Zenodo version DOI 10.5281/zenodo.23045141) is superseded in five places. All five were found by auditing this program’s own record against Berthé–Holton–Zamboni [5] read first-hand, including its §5, which the underlying Phase 7 work did not reach.

C1 — The “one remaining lemma” is a published theorem.

Revision 1 §12 stated as *Conjecture 12.1* the claim that for every admissible Ostrowski digit sequence of a bounded-type slope, $\limsup_k \max(x'(k), y(k)) > 2$ with a margin depending only on A , and said it was “not claimed proved here or anywhere in the underlying record” [29, 28]. This is **BHZ Propositions 5.1 and 5.2** [5]: if $(a_k, a_{k+1}) = (s, t)$ with $s > 1$ for infinitely many k then *every* $\omega \in X_\alpha$ has $\text{ice}(\omega) \geq 2 + \frac{1}{2(s+1)(t+1)+1}$, and if $(a_k, a_{k+1}, a_{k+2}) = (1, 1, t)$ for infinitely many k then *every* $\omega \in X_\alpha$ has $\text{ice}(\omega) \geq 2 + \frac{1}{8t+1}$. A bounded-type slope satisfies one hypothesis or the other by pigeonhole. *Consequence:* Theorem 12.1 now holds at every intercept (§14), and Revision 1’s Open 1 is not open. The weaker statement $\text{ice}(\omega) > 2$ at each individual intercept is likewise already a corollary of BHZ Theorem 1.1.

C2 — The “exceptional-orbit reduction” is false.

Revision 1’s Open 2, and the Phase 7 verdict it quoted, rested on the claim that $\text{ice}(\omega_\rho) = \text{ind}^*(\alpha)$ for every intercept outside the countable “return” set $\{j\alpha \bmod 1\}$ [27], so that the genuinely open cases were countable and intercept 0 was “if anything a harder case than

generic.” The cited ingredients do not give this: BHZ Proposition 2.1(4) gives shift-invariance of ice only off a *finite set of orbits*, and Lemma 2.2 gives $\text{ice} = \text{ind}^*(\alpha)$ only *almost everywhere* for the unique invariant measure — and a measure-zero set need not be countable. BHZ’s own Proposition 4.1 supplies the counterexample: the “keep one” intercept $c_k = a_k - 1$ has $\text{ice}(\omega) \leq 1 + \theta = 2.618\dots$ at *every* irrational slope, while $\text{ind}^*(\alpha) = 2 + \limsup_k [a_k; a_{k-1}, \dots, a_1]$ is unbounded in A ; its digit string is neither eventually 0 nor either shift-orbit pattern of the characteristic sequence, so its intercept is not a return intercept. Moreover the family $\{(c_k) \text{ admissible} : a_k - c_k \in \{1, 2\}\}$ is *uncountable* and has ice uniformly bounded, so for every $A \geq 2$ an uncountable set of intercepts has $\text{ice}(\omega) < \text{ind}^*(\alpha)$. *Consequence*: intercept 0 is in fact one of the *easiest* intercepts ($\text{ice} = \text{ind}^*(\alpha) - 1 \rightarrow \infty$), not the hardest; the hardest are the keep-one intercepts, where $\text{ice} \rightarrow 2$ as $A \rightarrow \infty$. Revision 1’s Open 2 is withdrawn, its premise being false.

C3 — Convergent-denominator indexing.

BHZ Proposition 2.7 fixes $\alpha = [0; a_1 + 1, a_2, a_3, \dots]$ with $q_0 = 1$, $q_1 = a_1 + 1$, $q_k = a_k q_{k-1} + q_{k-2}$ for $k \geq 2$: the BHZ *digit* a_1 is one less than the first partial quotient of α , while (q_k) are the ordinary convergent denominators. This program’s record and code used $q_1 = a_1$. Every $\limsup$ is insensitive to a single shifted digit, so *every* ice value reported in Revision 1 is correct; every *root length* computed from q_k was not. This is the complete explanation of the root-length mismatch the Phase 7 record disclosed against itself — BHZ Proposition 3.3 was never wrong. Under the correct convention the root-length formulas reproduce exactly against literal longest common prefixes (562 checks, no failure).

C4 — Missing length factor in the surplus decomposition.

Revision 1 §13 displayed $S_k \geq A \cdot E_k - C_A \cdot D_k - O_A(\log \ell_k)$. The leading term cannot be $A \cdot E_k$: A is a pure number while S_k scales like a length. Re-derived from the integer periodic-value formula with every factor kept (§15), the correct statement on the odd-weight branch is $S_k \geq \ell_k E_k + 1 - \lceil D(R_k) \rceil \log_2 3 - \log_2(2\ell_k)$, with $\ell_k = |W_k|$ the *Sturmian* root length. Revision 1’s own flexible condition $E_k \gg \log \ell_k / \ell_k$ is equivalent to $\ell_k E_k \gg \log \ell_k$ and so is consistent with $\ell_k \cdot E_k$; the displayed inequality was a transcription slip, not a different claim, and no conclusion drawn from it changes.

C5 — Conditional attainment of $y(k)$.

Revision 1 treated both Ostrowski expressions as giving actually-occurring initial powers for every admissible (c_k) . BHZ Proposition 3.3 attains $x(k)$ for *every* k , with primitive root of length exactly q_k , but attains $y(k)$ only when $0 < c_k < a_k$, with root length $q_k - c_k q_{k-1}$; Corollary 3.5’s three observations ($c_k = a_k \Rightarrow y(k) = x(k - 2)$; $c_k = 0, c_{k+1} = a_{k+1} \Rightarrow y(k) < y(k + 1) = x(k - 1)$; $c_k = 0, c_{k+1} < a_{k+1} \Rightarrow y(k) \leq x(k)$) are what upgrade the $\limsup$. For a $\limsup$ this is harmless; for a proof that must hand the bridge an *attained* root it is not, because the branch of the argument that is tightest at $A = 1$ is exactly $c_k = a_k$, where $y(k)$ is not directly attained. Theorem 14.3 routes every case through an attained witness explicitly.

What survives unchanged. Theorem 5.1 (the abstract criterion), the discrepancy–height bound, the repetition–height competition, the exact XOR transfer $L_v = L + 1$, the density-1/2 structural fact, the intercept-0 case of Theorem 12.1 and its $\text{term2}(k) > 2 + 1/(A + 1)$ analysis, the whole self-correction history of §11, the counting-route material, and every numerical finding about structured intercepts — including the identification of the near-maximal-digit family as the empirical worst case, which §14 now confirms is *exactly* extremal. What is superseded: Revision 1’s §12 conjecture, its Open 1, Open 2 and Open 3, and the two displayed formulas of C3 and C4.

Attribution. The load-bearing symbolic input to the general-intercept theorem is Berthé–Holton–Zamboni’s, not this program’s. This note claims as its own contribution the arithmetic application: the periodic-approximation/arithmetic-height bridge, the exact XOR transfer, the density-1/2 fact, the corrected surplus inequality, and the attained-witness bookkeeping with explicit growing root lengths that turns a lim sup bound on ice into an unbounded surplus. Recording a published 2006 theorem as an open conjecture was an error of literature search, and is corrected here rather than quietly dropped.

3 The Periodicity Conjecture and the Conjugacy Map

Let $T : \mathbb{Z}_2 \rightarrow \mathbb{Z}_2$ be the $3x + 1$ map in its standard form,

$$T(x) = \begin{cases} x/2, & x \equiv 0 \pmod{2}, \\ (3x + 1)/2, & x \equiv 1 \pmod{2}. \end{cases}$$

The *parity vector* of $x \in \mathbb{Z}_2$ is $v(x) = (v_0, v_1, \dots) \in \{0, 1\}^{\mathbb{N}}$ with $v_i \equiv T^i(x) \pmod{2}$; the conjecture below is Lagarias’s [7]. Bernstein and Lagarias showed that $x \mapsto v(x)$ is a bijection of $\mathbb{Z}_2$ onto $\{0, 1\}^{\mathbb{N}}$, and in fact a *2-adic isometry* [4, 3]: identifying $\{0, 1\}^{\mathbb{N}}$ with $\mathbb{Z}_2$ through binary expansions,

$$v_2(\Phi(a) - \Phi(b)) = \text{lcp}(a, b),$$

where Φ is the inverse map (so $\Phi(v)$ is the unique 2-adic integer with parity vector v), and lcp is longest-common-prefix length. Conjugating T by Φ turns it into the shift map on $\{0, 1\}^{\mathbb{N}}$. This isometry property is the single fact the entire bridge in this note exploits: two sequences agreeing on a long prefix must have 2-adically close Φ -images.

Rationality of $\Phi(v)$ corresponds to eventual periodicity of v in one direction only: if v is eventually periodic then $\Phi(v) \in \mathbb{Q}$. The converse is:

Lagarias’s Periodicity Conjecture [7, §2]. No aperiodic parity vector has a rational Φ -value.

(See also López and Stoll’s continuous-setting formulation of the same conjecture, [9].)

This conjecture is open, and *very little is known unconditionally in the converse direction* — a fact worth stating plainly, since it frames everything that follows. The conjecture quantifies over *all* $v \in \{0, 1\}^{\mathbb{N}}$ that fail to be eventually periodic: an uncountable family with no assumed combinatorial structure.

Full conjecture versus the classes treated here. This note — and the research program it consolidates — does not address the full conjecture. It proves $\Phi(v) \notin \mathbb{Q}$ for two structurally defined sub-families of aperiodic words (Sturmian words, proved in the companion foundation paper; and complementary symmetric Rote sequences over bounded-type slopes at intercept 0, proved here), and precisely maps, without resolving, one further natural extension (Rote sequences at arbitrary intercepts). §26 records a structural fact showing that no extension of this program’s method, however far pushed, can by itself close the full conjecture. Nothing in this note should be read as narrowing that gap beyond what is explicitly proved.

4 Two Routes to Irrationality

The foundation paper and this program’s extension of it use two logically independent mechanisms to force $\Phi(v) \notin \mathbb{Q}$.

4.1 Route A — Counting

Let $p_v(L)$ denote the number of distinct length- L factors of v (its *factor complexity*). Dubickas proved in 2009 [6] that a positive-integer $3x + 1$ trajectory tending to infinity has parity-word complexity $p(L) > 1.70951129 L$ eventually; the foundation paper transports this to every rational 2-adic integer, since T preserves a fixed odd denominator. The exact threshold constant is

$$\frac{1}{\log_2(3/2)} = 1.709511\dots,$$

and the transported statement is: if v is not eventually periodic and $\Phi(v) \in \mathbb{Q}$, then $\liminf_L p_v(L)/L \geq 1/\log_2(3/2)$. Contrapositively: *any* aperiodic v with $\liminf_L p_v(L)/L < 1.70951\dots$ has $\Phi(v) \notin \mathbb{Q}$, unconditionally, with no repetition or discrepancy hypothesis whatsoever.

This is a purely counting-based exclusion, and it is exactly as strong as its complexity bound: it says nothing once a word's complexity crosses the threshold. Sturmian words ($p(n) = n + 1$) sit far below it. Complementary symmetric Rote sequences, the next class this program treats, have $p(n) = 2n$ — strictly above $1.70951\dots n$ for every $n \geq 2$ — and are therefore *entirely outside Route A's reach*. This is precisely where Route A stops, and why a second, structurally different mechanism is needed to make any further progress.

4.2 Route B — Periodic Approximation and Arithmetic Height

This is the central mechanism of this note. Informally, the chain of implications is: symbolic periodic approximation $\rightarrow$ exact 2-adic agreement $\rightarrow$ divisibility $\rightarrow$ rational periodic approximant $\rightarrow$ Archimedean height $\rightarrow$ contradiction.

Concretely: if v agrees with a periodic word W^∞ on a long prefix, the 2-adic isometry forces $\Phi(v)$ and the *rational number* $\Phi(W^\infty)$ to be 2-adically close to a precisely quantified depth. If $\Phi(v)$ were itself rational, this closeness becomes an exact integer divisibility statement between two specific integers built from $\Phi(v)$'s numerator/denominator and W 's combinatorial data. When the periodic approximation is long and clean enough relative to the arithmetic size (*height*) of the numbers involved, that divisibility is impossible unless the two rationals coincide — and they were assumed not to. The whole of §5–§7 makes this precise.

5 The Abstract Bridge Theorem

Following Proposition 4.1 of the foundation paper, every finite word W (length $\ell = |W|$, $k = |W|_1$ ones) has an explicit periodic-value formula

$$\Phi(W^\infty) = \frac{c_W}{2^\ell - 3^k}$$

for an explicit integer c_W (THEOREM_STATUS.md, citing the foundation paper's Prop. 4.1).

Suppose, for contradiction, $\Phi(s) = u/v \in \mathbb{Q} \cap \mathbb{Z}_2$ in lowest terms. For any finite word W_n with $s \neq W_n^\infty$, put $\ell_n = |W_n|$, $k_n = |W_n|_1$, $L_n = \text{lcp}(s, W_n^\infty)$, $\delta_n = 2^{\ell_n} - 3^{k_n}$, and define the integer

$$M_n := u \delta_n - v c_{W_n}.$$

By the 2-adic isometry, $v_2(\Phi(s) - \Phi(W_n^\infty)) \geq L_n$, which translates into a 2-adic divisibility lower bound on M_n ; on the other hand $|M_n|$ is bounded above in terms of $|u|, v, \delta_n, c_{W_n}$ — an Archimedean (ordinary absolute-value) bound. When the 2-adic lower bound forces M_n to be divisible by more than its Archimedean bound allows, $M_n = 0$ is forced, and (with $s \neq W_n^\infty$) that is eventually impossible. This comparison is exactly what the repository's *surplus* quantity measures.

Definition (Surplus, canonical form [12]).

$$S_s(W) := \text{lcp}(s, W^\infty) - \log_2(|2^{|W|} - 3^{|W|_1}| + c_W).$$

This is the repository’s own proof-native quantity, fixed from Phase 1 onward and not altered by any later phase: “*the proof-native quantity, not a convenient after-the-fact packaging*” (theorems/phase2/SURPLUS_THEOREM.md).

Theorem 5.1 (Abstract 2-adic periodic-approximation criterion [11]). *Let $s \in \{0, 1\}^\mathbb{N}$ and $(W_n)_{n \geq 1}$ finite binary words with $s \neq W_n^\infty$ for all n . If*

$$\limsup_{n \rightarrow \infty} S_s(W_n) = +\infty,$$

then $\Phi(s) \notin \mathbb{Q}$.

This is stated and proved with no Sturmian or Rote terminology at all — it is the Sturmian-free abstraction every later result in this note specializes.

Theorem 5.2 (Dual necessity, [12]). *If $\Phi(s) = u/v \in \mathbb{Q}$ in lowest terms, height $H = \max(|u|, v)$, then for every finite W with $s \neq W^\infty$,*

$$S_s(W) \leq \log_2 H.$$

Together: *surplus $\rightarrow +\infty$ along some sequence of periodic approximants forces irrationality; rationality forces surplus to stay uniformly bounded.* Everything that follows is the construction of periodic approximants along which surplus is provably unbounded, for successively broader symbolic classes.

6 Discrepancy as the Height Bridge

Repetition length L_n alone is not enough: the surplus definition also involves c_W , and the periodic-value formula gives no a priori control on how large c_W can be relative to $2^\ell - 3^k$. The quantity that closes this gap is prefix *discrepancy*:

$$D(W) := \max_{0 \leq i \leq \ell} \left| k_i(W) - \frac{ik}{\ell} \right|,$$

where $k_i(W)$ counts ones in the length- i prefix of W .

Theorem 6.1 (Discrepancy-height bound [13]). *For every finite word W , $\ell = |W|$, k ones, $D = D(W)$:*

$$c_W \leq \ell \cdot 3^{\lceil D \rceil} \cdot \max(2^\ell, 3^k).$$

(Proved, and exhaustively verified by direct computation for $\ell \leq 16$.) This single inequality is what converts symbolic regularity (a bound on how unevenly 1s are distributed in W) into an arithmetic height estimate the surplus computation can use. The relevant hierarchy of discrepancy regimes, from weakest to strongest sufficient hypothesis actually used somewhere in this program, is:

- $D(W) = O(1)$ — e.g., single-block or highly regular words; strongest, rarely needed.

- $D(W) = O(\log \ell)$ — achieved for rotations by any slope with *bounded partial quotients*, via the classical three-distance theorem; this is the regime used throughout §10–§12.
- $D(W) = o(\ell)$ — the general sufficient hypothesis for the abstract sublinear-discrepancy theorem below; strictly weaker, but gives up the explicit logarithmic rate.

Sublinear discrepancy is exactly what is needed for the height cost to be dominated by a linear repetition gain (§7); the sharper $O(\log \ell)$ rate is what the bounded-partial-quotient hypothesis buys on top of that.

7 Repetition–Height Competition

Theorem 7.1 (Sublinear-discrepancy repetition criterion [13]). *Suppose s begins in initial powers $W_n^{r_n}$, $\ell_n = |W_n| \rightarrow \infty$, densities $\gamma_n = k_n/\ell_n \rightarrow \gamma_\infty$, and $D(W_n) = o(\ell_n)$. If*

$$\liminf_n r_n > \max(1, \gamma_\infty \log_2 3) \quad \text{strictly,}$$

then $S_s(W_n) \rightarrow +\infty$, hence $\Phi(s) \notin \mathbb{Q}$.

Taking the supremum over $\gamma \in (0, 1)$ of $\max(1, \gamma \log_2 3)$ gives $\log_2 3 = 1.58496\dots$, so $r > \log_2 3$ suffices *uniformly in the density*. Because $\log_2 3 < 2$, an initial *square* ($r = 2$) always clears this bar, with an explicit margin $2 - \log_2 3 = 0.41503\dots$ — which is exactly why squares are the workhorse repetition structure throughout this program:

Theorem 7.2 (Squares suffice unconditionally [14]). *If s begins in arbitrarily long initial squares $W_n W_n$ — with no hypothesis on discrepancy, balance, or internal structure of the roots W_n — then $\Phi(s) \notin \mathbb{Q}$.*

The later, more flexible sufficient condition used once a fixed margin is no longer available (§15) does not require a fixed constant margin at all; it only requires the repetition *excess* above the threshold to dominate the discrepancy-plus-logarithmic height cost, i.e. $\text{excess} \cdot \ell \gg \log \ell$ rather than $\text{excess} \geq \text{const}$.

8 The Sturmian Foundation

The foundation paper [1] proves, by *two independent mechanisms*, that every Sturmian parity word has irrational Φ -value:

1. **Counting.** Every Sturmian word has $p(L) = L + 1$, far below the Dubickas threshold, so Route A (§4.1) applies directly.
2. **Liouville / initial-square route.** Sturmian words have arbitrarily long initial squares (Theorem 10.2 of [1], citing Allouche, Davison, Queffélec, and Zamboni [2]), which triggers the repetition–height mechanism of §7 directly (Theorem 10.1 of [1]). The foundation paper also gives a second, independent and effective proof, for the specific case López and Stoll’s own paper [8] had identified as open (their critical-density Sturmian word at slope $\beta = \ln 2 / \ln 3$): this is then generalized to every irrational slope, and a further argument removes the restriction to intercept 0 via an initial square of the word itself.

Both mechanisms are cited, not re-derived, in this note; nothing in this program alters or extends the Sturmian case itself. What matters for everything that follows is where the foundation paper’s own abstract explicitly places the frontier: “*Counting reaches every complexity below $1.70951 \dots L$; the frontier is complexity $2L$ (Rote words, rotation codings), open for both routes.*” This is the exact motivation for the program consolidated here.

9 Crossing the Counting Frontier: Complementary Symmetric Rote Sequences

A *complementary symmetric (CS) Rote sequence* $v \in \{0,1\}^{\mathbb{N}}$ is one whose first-difference (XOR) transform is Sturmian:

$$v \text{ is CS Rote} \iff u := S(v) \text{ is Sturmian,} \quad u_i = v_i \oplus v_{i+1} \quad (i \geq 0),$$

with inverse recovery $v_{n+1} = v_n \oplus u_n$ [16]. Such sequences have factor complexity exactly $p(n) = 2n$, standard in the combinatorics-on-words literature on this class (background on the surrounding Sturmian/complementary-word theory: [10]; see the bibliography note on primary Rote-sequence sourcing) — past the Dubickas threshold, so Route A never applies to them; every irrationality result about CS Rote sequences in this note is a Route B result.

Theorem 9.1 (Exact transfer, with a +1 bonus [16]). *Let $L := \text{lcp}(u, W^\infty)$ for a Sturmian $u = S(v)$ and finite word W , and let $L_v := \text{lcp}(v, \text{target}^\infty)$ where $\text{target} = V$ (if $W = V$ has even weight) or $\text{target} = R := V\bar{V}$ (if V has odd weight, $\bar{V}$ its bitwise complement). Then*

$$L_v = L + 1, \quad \text{exactly, not merely } \geq L \text{ or } \approx L.$$

Verified by induction (both directions of the inequality) and computationally over 6 initial cases in Phase 2; re-derived completely independently in Phase 5 and cross-checked exhaustively over every binary root of length 2 to 10 and every mismatch position from ℓ to 4ℓ — 114,680 cases, 0 failures [17]. The theorem is intercept-agnostic: it holds verbatim regardless of the intercept of the underlying Sturmian word.

10 The Odd-Weight Mechanism

This is the structural discovery that makes the whole bounded-type theorem (§12) possible, and it deserves to be isolated on its own.

When the Sturmian root W has *odd* weight, Theorem 9.1’s target is $R = V\bar{V}$, not V itself.

Theorem 10.1 (Density of the doubled root, exactly $1/2$ [21]). *For every finite word V , $R := V\bar{V}$ satisfies $|R| = 2|V|$ and $|R|_1 = |V|$, hence*

$$\text{density}(R) = \frac{|R|_1}{|R|} = \frac{1}{2}, \quad \text{exactly, unconditionally.}$$

Proof (as recorded). $|R|_1 = |V|_1 + |\bar{V}|_1 = |V|_1 + (|V| - |V|_1) = |V| = \ell$; and $|R| = 2\ell$; so density $= \ell/(2\ell) = 1/2$. ■

Arithmetically, this matters because $3^{|R|_1} = 3^{|R|/2}$ grows strictly more slowly than $2^{|R|}$ (since $\sqrt{3} < 2$), so the dominant exponential height scale in the periodic-value denominator $2^{|R|} - 3^{|R|_1}$

is unambiguously $2^{|R|}$, regardless of the fine structure of V . Plugging $\gamma = 1/2$ into the repetition–height criterion of §7 collapses the required threshold on the Sturmian side from an earlier, over-conservative estimate of $2 \log_2 3 \approx 3.170$ down to the true requirement,

$$r_{\text{required}}(\gamma) = 2, \quad \text{uniformly, for both the even- and odd-weight transfer routes.}$$

This is established in [21]. This single fact replaced two working hypotheses this program had earlier formed and found false — a universal even-weight witness hypothesis, and a silver-ratio extremality claim — both recorded in full in §11.

11 Self-Correction History

This program is falsification-first throughout: every phase actively attacked its own prior claims before extending them. Four corrections are recorded here in full, not hidden, because they show how the final theorem (§12) was actually reached, and because a future researcher extending this work needs to know which paths were tried and ruled out.

A. Universal even-weight witness hypothesis — rejected. An early working assumption was that the Rote-transfer route could always be arranged to use an even-weight root V (avoiding the doubled root R altogether). A finite 8-state automaton analysis of the weight-parity sequence found exactly one “bad” cycle sustaining all-odd weight forever, and an explicit quadratic-irrational counterexample slope was found and verified to 28 convergents: $\gamma = [0; 1, 1, 1, 2]$ has weight parity $0, 0, 1, 1, 1, 1, \dots$ — all odd (weight 1) from index 2 onward, forever [18]. The universal claim is false. The bridge still succeeds on this slope because the odd-weight transfer route (§10) handles exactly this case.

B. Silver-ratio extremality — rejected, explicit counterexample. A subsequent working hypothesis, formed from testing only four continued-fraction tails, was that among all-even-tail slopes the silver ratio ($\bar{2}$ tail) is *extremal*, with minimal achievable exponent $2 + \sqrt{2} \approx 3.4142$. An adversarial search found $\gamma = [0; 1, 4, 1, 5, 1, \bar{6}, 2]$ with $\text{ice}(u) \approx 3.1547 < 2 + \sqrt{2}$ [19], disproving the hypothesis outright. The constant $2 + \sqrt{2}$ itself is not wrong — it is the exact value for the pure silver-ratio tail [20] — but it was the wrong threshold to be solving for: once the density-1/2 fact (§10) is known, the actual required threshold is $r > 2$, and $2 + \sqrt{2} \gg 2$ was never close to the real boundary.

C. Seed-independence via complement-invariance of c_W — rejected, repaired. The claim that both CS Rote seeds ($v_0 = 0$ or $v_0 = 1$) give identical surplus was correct, but its stated justification (“complementing uniformly complements R , so c_W is unaffected”) was not: direct testing found $c_W \neq c_{\bar{W}}$ in 8 of 8 random odd-weight cases, often by an order of magnitude [22]. The repaired reason: every verification script’s height bound $F_{\text{bound}}(W)$ depends on the root only through $(\ell, k, D(\text{root}))$, never through c_W directly, and $D(V\bar{V}) = D(\bar{V}V)$ exactly, checked in 15 of 15 random cases. The *conclusion* survives untouched; only its justification needed repair — exactly the kind of error a hostile self-audit is designed to catch.

D. Phase 6 small- k boundary correction. The load-bearing inequality $\text{term2}(k) > 2 + 1/(A + 1)$ (§12) was stated to hold strictly for every k . Adversarial testing against genuinely non-eventually-periodic bounded-type controls (Thue–Morse-coded, Fibonacci-word-coded, and two independent

random $\{1, 2\}$ sequences) found one case of non-strict *equality* rather than a violation: in the Fibonacci-coded control, $\text{term2}(2)$ equals $2 + 1/(A + 1) = 7/3$ exactly, at $k = 2$, the only such case out of 232 checks across all four sequences [23]. This does not weaken the theorem, because the abstract criterion (Theorem 5.1) only requires $\limsup$ unboundedness along *some* growing sequence of witnesses, not strict inequality at literally every finite k .

None of these four corrections changed the truth of the final headline theorem; each changed (and, in cases B and C, simplified) the reason it is true.

12 The Bounded-Type CS Rote Theorem (Every Intercept)

Theorem 12.1 (Headline result of the program [38]). *Let $\gamma \in (0, 1)$ be irrational with bounded partial quotients ($A := \sup_n a_n < \infty$). Let u be **any** Sturmian word of slope γ — **any** intercept ρ , either the upper or the lower mechanical convention. Let v be either complementary symmetric Rote sequence with $S(v) = u$ (either seed), or any shift $\sigma^j(v)$. Then*

$$\Phi(v) \notin \mathbb{Q}.$$

The intercept-0 case of Theorem 12.1 is the result Revision 1 stated [25]; the general-intercept extension is §14, whose symbolic input is BHZ [5] §5 (Revision 2, C1). The proof architecture of §13 is unchanged apart from that one arrow: every other step was already proved intercept-agnostic, and each has been re-verified on literally constructed mechanical words at general intercepts, under both conventions and both seeds [40].

This supersedes an earlier theorem of the same program, proved first for the strictly narrower class of *quadratic*-irrational slopes [24], *without weakening any of its quantifiers* — both seeds, every shift, still covered. The upgrade matters because quadratic irrationals are countable (each is a root of an integer quadratic, and there are only countably many such polynomials), while bounded-partial-quotient irrationals are uncountable: for each bound A , the set of irrationals whose partial quotients never exceed A contains a Cantor-type subset, and the union over all A is an uncountable set.

Remark (What “uncountable” does and does not say here). *Uncountability is a statement about cardinality only. The set of bounded-type irrationals has Lebesgue measure zero: a Lebesgue-almost-every real number has unbounded partial quotients. Theorem 12.1 says nothing about a randomly chosen real slope, and nothing about density among all irrationals — only that the class it covers, while structurally constrained, is not merely a countable list of algebraic special cases.*

13 Proof Architecture of the Bounded-Type Theorem

The shortest audited chain, reconstructed in Phase 6 without any periodicity hypothesis and cross-checked against four genuinely non-eventually-periodic bounded-type controls [23], adapting the original Phase 4 chain [26]:

Step	Cites
bounded partial quotients, $A = \sup_n a_n$	hypothesis
↓ Sturmian words have arbitrarily long initial squares	Thm 10.2 [1], [2]
↓ BHZ margin at intercept 0, $\text{term2}(k) > 2 + \frac{1}{A+1}$ (every k , up to §11.D)	[5], Cor. 3.5
↓ <i>or, at any intercept</i> : $\text{ice}(\omega) \geq 2 + \delta(A)$, attained, roots $\rightarrow \infty$	[5], Prop. 5.1–5.2; Thm 14.3
↓ arbitrarily long usable witnesses	§7
↓ exact XOR transfer, $L_v = L + 1$	Thm 9.1 [16, 17]
↓ even/odd branch split	§9
↓ odd branch: doubled root density exactly 1/2	§10 [21]
↓ actual-root discrepancy control, $O(\log \ell)$, uniform in ρ	[15], repaired [22], Lem. 14.4
↓ sublinear height penalty	§6
↓ positive, unbounded surplus	§5
↓ abstract irrationality criterion (Thm 5.1)	[11]
$\Rightarrow \Phi(v) \notin \mathbb{Q}$	Thm 12.1

Notably absent from this chain: the weight-parity automaton of §11.A, and the silver-ratio constant $2 + \sqrt{2}$ of §11.B — both were real, correctly-derived facts about this problem, and neither is needed by the final proof.

14 General Intercepts

Sections 10–13 were originally stated for the *intercept-0* representative of a slope γ : the lower mechanical word $u_i = \lfloor (i+1)\gamma \rfloor - \lfloor i\gamma \rfloor$. This section removes that restriction.

An arbitrary intercept is encoded by an *Ostrowski digit sequence* $(c_k)_{k \geq 1}$, subject to BHZ’s admissibility condition [5]:

$$a_k \geq c_k \geq 0 \text{ for all } k, \quad a_k \geq 1 \text{ for } k \geq 2, \quad \text{and if } c_k = a_k \text{ then } c_{k-1} = 0.$$

The characteristic (intercept-0) sequence is $c_k \equiv 0$. Where the partial quotients (a_k) fix the canonical Sturmian *scale hierarchy* (the convergent denominators q_k), the digits (c_k) determine *where inside that hierarchy* a particular intercept’s word sits.

Conventions, stated exactly (Revision 2, C3). BHZ Proposition 2.7 sets

$$\alpha = [0; a_1 + 1, a_2, a_3, \dots], \quad q_0 = 1, \quad q_1 = a_1 + 1, \quad q_k = a_k q_{k-1} + q_{k-2} \quad (k \geq 2),$$

so the BHZ digit a_1 is one less than the first partial quotient of α , while (q_k) are the ordinary convergent denominators. Write $A_b := \sup_k a_k$; always $A_b \leq A$. Bounds below are sharpest in A_b , and remain true with A substituted.

At intercept 0, BHZ’s Corollary 3.5 collapses to the clean closed form used in §13. In general it does not collapse:

$$x'(k) = \frac{\sum_{j=1}^{k+1} (a_j - c_j) q_{j-1}}{q_k}, \quad y(k) = 1 + \frac{\sum_{j=1}^k (a_j - c_j) q_{j-1}}{q_k - c_k q_{k-1}},$$

$$\text{ice}(\omega) = \limsup_k \max(x'(k), y(k)).$$

Both terms depend on the entire digit history, not just on (a_j) .

Attainment, stated exactly (Revision 2, C5). By BHZ Proposition 3.3, $x(k)$ is the prefix power of an actually occurring initial power of ω with primitive root of length exactly q_k , for every k ; and $y(k)$ is, with root length exactly $q_k - c_k q_{k-1}$, provided $0 < c_k < a_k$. Nothing is asserted about $y(k)$ as an attained power otherwise; Corollary 3.5's three observations convert those cases into attained x -witnesses.

The margin is a published theorem (Revision 2, C1).

Theorem 14.1 (Berthé–Holton–Zamboni [5], Propositions 5.1 and 5.2). *If (s, t) are integers with $s > 1$ and $(a_k, a_{k+1}) = (s, t)$ for infinitely many k , then every $\omega \in X_\alpha$ satisfies $\text{ice}(\omega) \geq 2 + \frac{1}{2(s+1)(t+1)+1}$. If $(a_k, a_{k+1}, a_{k+2}) = (1, 1, t)$ for infinitely many k , then every $\omega \in X_\alpha$ satisfies $\text{ice}(\omega) \geq 2 + \frac{1}{8t+1}$.*

Corollary 14.2. *For every irrational γ of bounded type and every $\omega \in X_\gamma$, $\text{ice}(\omega) \geq 2 + \frac{1}{2(A_b+1)^2+1}$.*

Proof. There are finitely many digit pairs. If $a_k > 1$ for infinitely many k , some pair (s, t) with $s > 1$ and $s, t \leq A_b$ recurs infinitely often, and Proposition 5.1 applies. Otherwise $a_k = 1$ for all large k , so $(a_k, a_{k+1}, a_{k+2}) = (1, 1, 1)$ recurs infinitely often and Proposition 5.2 applies with $t = 1$, giving $2 + \frac{1}{9} \geq 2 + \frac{1}{2(A_b+1)^2+1}$. $\square$

Here X_γ is BHZ's set of *all* Sturmian sequences of slope γ , i.e. the itineraries of every point under either of the two interval exchanges of their §2.1 — so Corollary 14.2 quantifies over every intercept *and* both mechanical conventions at once, which is exactly the quantifier the bridge needs.

An attained-witness form, with growing roots. Corollary 14.2 bounds a lim sup. A lim sup bound $\text{ice}(\omega) \geq 2 + \delta$ yields, for each $\varepsilon > 0$, infinitely many n with an attained initial power of exponent $> 2 + \delta - \varepsilon$; taking $\varepsilon = \delta/2$ already supplies attained witnesses of exponent $> 2 + \delta/2$ whose root lengths necessarily tend to infinity, which suffices for Theorem 12.1. The following stronger, per-scale form is what this program proves independently, and is what makes the surplus bound uniform in k rather than only along an unspecified subsequence.

Theorem 14.3 (Attained general-intercept margin [38]). *Let γ be irrational with $A_b < \infty$ and let ω be any Sturmian sequence of slope γ (any intercept, either convention). Put $\delta(A_b) := \frac{1}{2(A_b+1)^2}$. Then for every $k \geq 7$ there is a word W with $|W| \geq q_{k-4}$ such that W^r is a prefix of ω with $r \geq 2 + \delta(A_b)$. In particular $\text{ice}(\omega) \geq 2 + \delta(A_b)$, and the bound is realised by actually occurring initial powers whose root lengths tend to infinity.*

The proof is a three-case analysis on $d_k := a_k - c_k$ driven by the substitution $\lambda_k := q_{k-1}/q_k$, $m_k := t_{k-1} + d_k$ with $t_k := \lambda_k m_k$, under which $x'(k) = m_{k+1}$, $q_k - c_k q_{k-1} = q_{k-1}(\lambda_{k-1} + d_k)$ and

$$y(k) - 2 = \frac{\lambda_{k-1}(m_{k-1} - 1)}{\lambda_{k-1} + d_k},$$

so that $y(k) > 2$ if and only if $m_{k-1} > 1$. Admissibility forbids two consecutive $d_k = 0$, which floors m_k ; the three cases $d_k = 0$ (route through $x(k-2) = y(k)$), $d_k \geq 2$ (route through $x(k-1) \geq m_k$) and $d_k = 1$ (route through $y(k)$, or through $x(k)/x(k-1)$ when $c_k = 0$) each deliver an *attained* power. Full proof in [38]; BHZ's own three-case proof of Proposition 5.1 is structurally the same trichotomy, reached independently, with constants agreeing to a factor $1 + \frac{1}{2(A_b+1)^2}$.

How sharp is the margin. Let β_A denote the unique root > 1 of $\beta^2 = A\beta + 1$, i.e. $\beta_A = \frac{A+\sqrt{A^2+4}}{2}$; equivalently $\beta_A = \lim_k q_k/q_{k-1}$ for any slope whose partial quotients are eventually A . On BHZ’s “keep one” intercept $c_k = a_k - 1$ (so $d_k \equiv 1$) with BHZ digits eventually $\equiv A$, the recurrences above solve in closed form: $\lambda_k \rightarrow 1/\beta_A$, $m_k \rightarrow \beta_A/(\beta_A - 1)$, and since $\beta_A^2 - 1 = A\beta_A$,

$$\text{ice}(\omega) = 2 + \frac{1}{A\beta_A} \quad \text{exactly.}$$

Hence $\delta^*(\gamma) \leq \frac{1}{A\beta_A} = \Theta(A^{-2})$ there, so the proved floor $\frac{1}{2(A\beta_A+1)^2}$ is off by at most a bounded factor (between 2 and 4.90), and — because $1/(A\beta_A) \rightarrow 0$ — **no margin uniform in A can exist**: the A -dependence in Revision 1’s Conjecture 12.1 was necessary, not a convenience. This also recovers BHZ Proposition 4.1 from the other side, their bound $\text{ice}(\text{keep one}) \leq 1 + \theta$ and the floor above pinning the keep-one value between $2 + \frac{1}{A\beta_A}$ and $1 + \theta$.

Remark (What is *not* claimed). That $\frac{1}{A\beta_A}$ is the infimum of $\text{ice}-2$ over all intercepts is proved only for slopes whose BHZ digits are eventually constant with $A \geq 2$ (a three-case argument: deviating from $d_k \equiv 1$ costs at least $\min(1 + A, 2 + 1/\beta_A)$, both strictly larger), plus the Fibonacci slope via BHZ Proposition 4.3. On every other bounded-type slope it is numerical evidence only and is deliberately not promoted to a theorem [41]. Theorem 12.1 uses none of it: any fixed $\delta > 0$ suffices, and Corollary 14.2 alone would do.

The remaining intercept-dependent step, and its hidden hypothesis. Every ingredient of §13 after “a witness exists” was already proved intercept-agnostic: the XOR transfer uses no property of u beyond being a binary sequence; the doubled root’s density is $1/2$ by construction; and the discrepancy bound is a statement about a rotation’s equidistribution rate, uniform in the starting phase. That last step needs one hypothesis Revision 1 left implicit. For the intercept- ρ word, $k_i(u) = \lfloor i\gamma + \rho \rfloor - \lfloor \rho \rfloor$, so $k_i \bmod 2$ codes the orbit $\{i(\gamma/2) + \rho/2\}$ against $[\frac{1}{2}, 1)$: the relevant rotation is by $\gamma/2$, not γ .

Lemma 14.4. *If γ has bounded partial quotients then so does $\gamma/2$, with a bound depending only on A .*

Proof. Bounded type is equivalent to $c := \inf_{q \geq 1} q\|q\gamma\| > 0$, and $c \geq 1/(A+2)$. For any $q \geq 1$, $\|q\gamma\| = \|2 \cdot q(\gamma/2)\| \leq 2\|q(\gamma/2)\|$, so $q\|q(\gamma/2)\| \geq q\|q\gamma\|/2 \geq c/2 > 0$. Hence $\gamma/2$ is badly approximable with constant at least $1/(2(A+2))$, and its partial quotients are bounded by $\lfloor 2(A+2) \rfloor + 1$. $\square$

With Lemma 14.4 the three-distance bound applies to $\gamma/2$ and gives $D(R_k) = O_A(\log \ell_k)$ uniformly in ρ , completing the chain. Only $D = o(\ell)$ is actually needed; measured across 534 witnesses at general intercepts, $D(R)/\log_2 \ell_k$ stays in $[0.30, 0.61]$ while $D(R)/\ell_k$ falls from 0.20 to 0.0004 [40].

Independent verification. The whole chain was re-run on literally constructed mechanical words — certified rational brackets for both γ and ρ , every floor and ceiling either decided by the bracket or raised, and no BHZ formula used anywhere — across four slopes (including a non-eventually-periodic one), four intercept families (including keep-one), both mechanical conventions and both seeds: 534 attained witnesses with ℓ from 8 to 10 946; the transfer identity $L_v = L_u + 1$ held 534/534; $2^{L_v} > F(R)$ as exact integers 534/534; and the odd-branch density was exactly $1/2$ in all 338 odd-branch cases [40].

15 Ostrowski Repetition Structure

A subtlety worth making explicit for anyone continuing this work: **low factor complexity is not the same thing as strong initial repetition**. A CS Rote sequence may remain globally $p(n) = 2n$ while its intercept changes *which* large canonical blocks (Ostrowski/continuant-scale factors) happen to align at the origin — which is exactly what determines how long an initial repetition is actually available to feed the bridge.

The scale-dependent quantity that tracks this is an *excess*:

$$E_k := \max(r'_k, r_k) - 2,$$

where r'_k, r_k are the two branch-specific repetition exponents at scale k [30]. The corresponding surplus decomposition, re-derived from the integer periodic-value formula with every length factor kept explicit (Revision 2, C4), is

$$\text{odd branch: } S_k \geq \ell_k E_k + 1 - \lceil D(R_k) \rceil \log_2 3 - \log_2(2\ell_k),$$

$$\text{even branch: } S_k \geq \ell_k (r_k - \log_2 3) + 1 - \lceil D(V_k) \rceil \log_2 3 - \log_2 \ell_k,$$

where $\ell_k = |W_k|$ is the *Sturmian* root length [38]. Revision 1 displayed $S_k \geq A \cdot E_k - C_A D_k - O_A(\log \ell_k)$, which is dimensionally incomplete: the leading term is a length times an excess, not the partial-quotient bound times an excess. The flexible sufficient condition this leads to — used in place of a fixed margin whenever one is not cleanly available — is

$$E_k \gg \frac{\log \ell_k}{\ell_k}, \quad \text{equivalently} \quad \ell_k E_k \gg \log \ell_k,$$

which is manifestly the same statement as the corrected inequality, and is why no conclusion Revision 1 drew from the displayed formula changes [31]. With Theorem 14.3 the excess is bounded below by the *constant* $\delta(A_b)$ and $\ell_k \geq q_{k-4} \rightarrow \infty$, so the flexible machinery is not needed at all: the general-intercept case lands in the easiest regime, $E_k \asymp 1$. Every structured intercept family tested lands there too [32], and the near-maximal family those tests identified as worst is exactly the keep-one family, for which the excess is exactly $1/(A\beta_A)$.

The even branch needs no discrepancy hypothesis at all: the combinatorial cap $D(V) \leq \gamma_V(1 - \gamma_V)\ell$ together with $\sup_\gamma [\max(1, \gamma \log_2 3) + \gamma(1 - \gamma) \log_2 3] = \log_2 3 < 2$ already suffices [14]. Only the odd branch needs $D = o(\ell)$.

16 The Density Coordinate, and Where the Residual Case Is

Revisions 1–2 organised this programme along the *factor-complexity* axis: Sturmian ($p(n) = n + 1$), then the counting classes below Dubickas's threshold, then $p(n) = 2n$. That is the axis on which Route A operates. It is not the axis the other known exclusion results read.

Definition. For $v \in \{0, 1\}^{\mathbb{N}}$ write $k_N(v) := \#\{i < N : v_i = 1\}$ and call $\liminf_N k_N(v)/N$ the lower ones-density of v .

Theorem 16.1 (Monks–Yazinski [44], Theorem 2.7(b)). *If $x \in \mathbb{Q}_{\text{odd}}$ has divergent T -orbit, then $\ln 2 / \ln 3 \leq \liminf_N \kappa_N(x)/N$, where $\kappa_N(x)$ counts the odd elements among the first N terms of the orbit.*

Since $\kappa_N(x)/N$ is exactly k_N/N for the parity vector of x , and since an orbit is eventually cyclic precisely when its parity vector is eventually periodic, Theorem 16.1 contraposes to a statement about words:

Corollary 16.2. *If v is not eventually periodic and has lower ones-density $< \beta := \ln 2 / \ln 3$, then $\Phi(v) \notin \mathbb{Q}$ — whatever the factor complexity of v .*

This forces a correction to how Revisions 1–2 framed their headline (D2 of §1).

Remark (The CS Rote chain sits inside Corollary 16.2). *Every complementary symmetric Rote sequence has ones-density exactly $1/2$ — this is the same structural fact, established in Phase 4 and used throughout §§10–13, that collapses the required repetition exponent to 2. Since $1/2 < \beta = 0.63092\dots$, Corollary 16.2 already gives $\Phi(v) \notin \mathbb{Q}$ for every such v . Theorem 12.1 is therefore **not** a new exclusion; it is an independent second proof of an available one, by a mechanism that also yields effective height floors (§20) that a density argument does not give. The same applies to the Thue–Morse word, of density $1/2$.*

What each result reads.

result	coordinate read	reaches	status of source
Monks–Yazinski Thm 2.7(b) [44]	density	$\liminf k_N/N < \beta$	refereed
Dubickas, transported [6, 1]	complexity	$\liminf p(L)/L < 1.70951\dots$	refereed
Foundation Thm 10.1 [1]	both	all Sturmian ($p(n) = n + 1$, any slope)	preprint (self)
López–Stoll Thm 1 [9]	density	(proposed) $\liminf k_N/N > \beta$	preprint; not used

Acknowledgement, and what is not used. López–Stoll’s work on density and the $3x + 1$ conjugacy map helped guide this programme toward the critical density $\beta = \ln 2 / \ln 3$. We acknowledge that influence. The results presented here are established through the independent arguments given in this note and do not require their proposed upper-density exclusion. The critical-density Sturmian word is theirs [8], and so is the observation that critical density is where the existing methods stop. We take no position on their Theorem 1 [9] — we have neither established nor disproved it — and no argument below invokes it. Accordingly the residual region is described against what is independently available:

the densities reached by refereed work are those below β ; the residual region is lower ones-density $\geq \beta$.

At lower ones-density exactly β and factor complexity $n + 1$ the foundation paper already has the conclusion, since a Sturmian word of slope β has ones-density β [1]. So the uncovered corner is **lower ones-density exactly β at factor complexity $2n$** , and that is what §§17–18 address [53]. The foundation paper’s own open problem (4) asks exactly for the complexity- $2n$ class and records that no substitute repetition-and-height input for it was known to its author.

17 Critical-Density Rotation Codings: Rational Intercepts

Throughout this section and the next, $\beta := \ln 2 / \ln 3 = \log_3 2 = 0.6309297535\dots$, which is transcendental by Gelfond–Schneider, and $\chi := \mathbf{1}_{[0, \beta)}$. Note the identity $\beta \log_2 3 = 1$, used in §6’s height inequality.

Definition. For $\alpha \in (0, 1)$ irrational and $\rho \in \mathbb{R}$ define the critical-density coding $s = s(\alpha, \rho)$ by

$$s_n = 1 \iff \{n\alpha + \rho\} \in [0, \beta), \quad n = 0, 1, 2, \dots$$

The interval length is β , not α : this is a two-interval coding of a rotation whose partition endpoint is *not* in $\mathbb{Z}\alpha + \mathbb{Z}$, not a mechanical word.

Lemma 17.1 (Basic properties). *Let $\alpha \in (0, 1)$ be irrational and algebraic and let $\rho \in \mathbb{R}$. Then $s(\alpha, \rho)$ is aperiodic; its ones-density exists and equals β ; and $p_s(n) = 2n$ for all $n \geq 1$.*

Proof. A period $p \geq 1$ would make the proper arc $[0, \beta)$ invariant under rotation by $p\alpha \not\equiv 0 \pmod{1}$, which is impossible. The density statement is Weyl equidistribution for irrational α ; the limit exists, so $\liminf = \limsup = \beta$. For the complexity: β is transcendental and α algebraic, so $\mathbb{Z}\alpha + \mathbb{Z}$ consists of algebraic numbers and $\beta \notin \mathbb{Z}\alpha + \mathbb{Z}$; by Rote [49] and Berstel–Vuillon [50] a two-interval coding of an irrational rotation has $p(n) = 2n$ exactly when the partition endpoint lies outside $\mathbb{Z}\alpha + \mathbb{Z}$. $\square$

The complexity value is recorded for the coverage discussion of §16 only; no argument below uses it.

17.1 The agreement excess is a first hitting time

Write $L(\ell) := \text{lcp}(s, (s[0:\ell])^\infty)$ and, for $\ell \geq 1$, $\delta(\ell) := \ell\alpha - \text{round}(\ell\alpha)$, so $|\delta(\ell)| = \|\ell\alpha\|$. Since $\{x + \ell\alpha\} = \{x + \delta(\ell)\}$, the symbols $s_{m+\ell}$ and s_m differ exactly when $\{m\alpha + \rho\}$ lies in the symmetric difference of $[0, \beta)$ with its translate.

Lemma 17.2 (The mismatch set). *Put $M(d) := \{x \in [0, 1) : \chi(x) \neq \chi(\{x + d\})\}$. If $|d| \leq \min(\beta, 1 - \beta) = 0.3690702464\dots$ then*

$$M(d) = \begin{cases} [\beta - d, \beta) \cup [1 - d, 1), & d > 0, \\ [0, |d|) \cup [\beta, \beta + |d|), & d < 0, \end{cases}$$

two arcs of length $|d|$, one at each discontinuity of χ . Consequently

$$L(\ell) - \ell = m^*(\ell) := \min\{m \geq 0 : \{m\alpha + \rho\} \in M(\delta(\ell))\}.$$

Proof. For $d > 0$ compare χ on x and $x + d$ across the four intervals $[0, \beta - d)$, $[\beta - d, \beta)$, $[\beta, 1 - d)$, $[1 - d, 1)$: the values agree on the first and third and differ on the second and fourth. The hypothesis $|d| \leq \min(\beta, 1 - \beta)$ is what keeps these four intervals in this order, i.e. keeps the two arcs from wrapping past 0 and merging; without it the description fails. The case $d < 0$ is symmetric. The half-open side of each arc matches the half-open side of $[0, \beta)$. For the second statement, the first index at which s differs from $(s[0:\ell])^\infty$ is $\ell + m$ for the least $m \geq 0$ with $s_{m+\ell} \neq s_m$. $\square$

Remark. *The hypothesis of Lemma 17.2 is not cosmetic. Among convergent denominators of $(\sqrt{5} - 1)/2$ it fails only at $q = 1$, where $|\delta| = 0.3819660\dots > 0.3690702\dots$, and there the closed form is genuinely wrong; the construction below uses $|\delta| < 1/294$, with a factor of about 10^2 to spare [54]. The mechanism itself — reading the agreement excess as a first entry time of the rotation orbit into an arc at a partition endpoint — is proposed in the foundation paper’s own open problem (1) [1]; it is not original here. What is done here is its execution with two discontinuities instead of one.*

Two relaxations of Lemma 17.2 are used. Writing $B(y, r)$ for the two-sided arc of radius r about y :

$$M(\delta) \subseteq B(0, |\delta|) \cup B(\beta, |\delta|), \quad \text{hence} \quad m^*(\ell) \geq \min(t_0, t_\beta), \quad (1)$$

with $t_0 := \min\{m : \|m\alpha + \rho\| \leq |\delta|\}$ and $t_\beta := \min\{m : \|m\alpha + \rho - \beta\| \leq |\delta|\}$; and, sharper, under the hypothesis of Lemma 17.2,

$$\delta > 0 \implies 0 \notin M(\delta). \quad (2)$$

Inclusion (1) holds without the no-wrap hypothesis, since an arc crossing 1 overflows next to 0 and $B(0, \cdot)$ is two-sided.

17.2 The arc at 0: rational intercepts

Lemma 17.3. *Let α be a quadratic irrational with largest partial quotient A , and let $\rho = p/q \in \mathbb{Q}$ in lowest terms. Put $K := (A + 2)q^2$. Then $\|m\alpha + \rho\| > 1/(Km)$ for every $m \geq 1$, and hence $t_0 > 1/(K|\delta|)$ on those indices.*

Proof. A quadratic irrational has an eventually periodic continued fraction, so its partial quotients are bounded by some $A < \infty$, and then $\|n\alpha\| > 1/((A + 2)n)$ for every $n \geq 1$. Using $\|x\| \geq \|qx\|/q$ and $q(m\alpha + p/q) = qm\alpha + p$,

$$\|m\alpha + p/q\| \geq \frac{1}{q} \|qm\alpha\| > \frac{1}{q} \cdot \frac{1}{(A+2)qm} = \frac{1}{Km}.$$

□

For $\alpha = (\sqrt{5} - 1)/2$ and $\rho = 1/7$ this is $K = 3 \cdot 49 = 147$. The rationality of ρ is what makes Lemma 17.3 available; §18 replaces it.

At $m = 0$ the orbit point is $\{\rho\}$. If $\rho \neq 0$ then $\|\rho\| \geq 1/q$, while $|\delta| < 1/(KM) \leq 1/(6q^2) < 1/q$ for $M \geq 2$, so $m = 0$ lies outside $B(0, |\delta|)$ and (1) suffices. If $\rho = 0$ then $\|0\| = 0$ and (1) is vacuous; (2) is used instead, restricting to convergent denominators with $\delta > 0$. Since $\delta(q_k) = q_k\alpha - p_k$ alternates in sign, infinitely many qualify, and $q_{k+1} = a_{k+1}q_k + q_{k-1} \leq (A + 1)q_k$ absorbs the cost of skipping every other one. **No finite shift is used, and none would help:** shifting replaces $\rho = 0$ by $\{\alpha\}$, which is irrational, destroying the hypothesis Lemma 17.3 needs.

17.3 The arc at β , the witnesses, and the theorem

Let j_m be the nearest integer to $m\alpha + \rho - \beta$ and put $A_m := m\alpha + \rho - j_m$, so that $c_m := \|m\alpha + \rho - \beta\| = |A_m - \beta|$. Section 19 proves, for any fixed $\kappa > 2$,

$$c_m \geq \exp(-c_1(\log(m + 2))^\kappa), \quad c_1 = c_1(\alpha, \rho, \kappa) \text{ effective.} \quad (3)$$

In particular $c_m > 0$ for every m : the orbit never hits β , since $m\alpha + \rho \equiv \beta$ would make β algebraic.

Proposition 17.4 (The witnesses). *For $M \geq 2$ put $t(M) := \min(\min_{0 \leq m < M} c_m, 1/(KM))$ and let $\ell(M)$ be the least admissible convergent denominator q_k of α with $\|q_k\alpha\| < t(M)$, where “admissible” means all of them if $\rho \neq 0$ and those with $\delta(q_k) > 0$ if $\rho = 0$. Then*

$$L(\ell(M)) - \ell(M) \geq M, \quad \ell(M) > \frac{K}{A+2}M \rightarrow \infty, \quad \log \ell(M) = O((\log(M + 2))^\kappa).$$

Proof. No-wrap holds automatically since $t(M) \leq 1/(2K) \leq 1/6 < 0.369$. For $0 \leq m < M$: the first threshold gives $c_m \geq t(M) > |\delta|$, so $m \notin B(\beta, |\delta|)$; the second, with Lemma 17.3, gives

$\|m\alpha + \rho\| > 1/(Km) \geq 1/(KM) > |\delta|$ for $m \geq 1$, and $m = 0$ is handled as above. So $m^*(\ell(M)) \geq M$, and Lemma 17.2 converts this into the first display.

For the size of $\ell(M) = q_k$: by minimality the previous admissible denominator q has $\|q\alpha\| \geq t(M)$, while $\|q\alpha\| < 1/q'$ for its successor q' ; hence $q' < 1/t(M)$ and $\ell(M) \leq (A+1)q' < (A+1)/t(M)$. Combining with (3) and $1/(KM)$ gives the third display, because $\kappa > 2 > 1$. For the lower bound, $\|q_k\alpha\| > 1/(q_{k+1} + q_k) \geq 1/((A+2)q_k)$ forces $t(M) > 1/((A+2)\ell(M))$. $\square$

Theorem 17.5 (Critical density, quadratic slope, rational intercept). *Let $\alpha \in (0, 1)$ be a quadratic irrational and let $\rho \in \mathbb{Q}$, including $\rho = 0$. Then $s(\alpha, \rho)$ is aperiodic with ones-density exactly β and factor complexity $2n$, and*

$$\Phi(s(\alpha, \rho)) \notin \mathbb{Q}.$$

Proof. By Lemma 17.1, s is aperiodic, so $s \neq W^\infty$ for every finite W — the only standing hypothesis of Theorem 5.1. For bounded-type α the discrepancy bound of §6 gives, *uniformly in ρ* , $|k_\ell - \ell\beta| \leq C_D \log_2 \ell + C'_D$ and $D(W) \leq 2C_D \log_2 \ell + 2C'_D$; uniformity holds because shifting the orbit by ρ replaces one interval by at most two, so the extreme discrepancy at most doubles. Substituting into the height inequality of §6 and using $\beta \log_2 3 = 1$,

$$S(W) \geq (L - \ell) - C \log_2 \ell - C_0, \quad C := 1 + 3C_D \log_2 3, \quad C_0 := 1 + \log_2 3 (3C'_D + 1),$$

with C, C_0 depending on α alone. Evaluating at $\ell = \ell(M)$ and applying Proposition 17.4,

$$S(s[0:\ell(M)]) \geq M - \frac{C}{\ln 2} O((\log(M+2))^\kappa) - C_0 \xrightarrow{M \rightarrow \infty} +\infty,$$

since $(\log M)^\kappa = o(M)$ for every fixed κ . Hence $\limsup_\ell S(s[0:\ell]) = +\infty$ and Theorem 5.1 gives $\Phi(s) \notin \mathbb{Q}$. $\square$

18 Algebraic Intercepts

Lemma 17.3 is the only place Theorem 17.5 used the rationality of ρ . It can be replaced by Liouville's inequality, at no cost in the conclusion and none in the analytic input.

Lemma 18.1 (The arc at 0, algebraic intercepts). *Let α be a quadratic irrational, let ρ be real algebraic, and put $D := [\mathbb{Q}(\alpha, \rho) : \mathbb{Q}]$. For $m \geq 0$ let $\theta_m := m\alpha + \rho - j$ with j the nearest integer to $m\alpha + \rho$, so $\|m\alpha + \rho\| = |\theta_m|$. Then*

(i) $\deg \theta_m \leq D$ and $H(\theta_m) \leq C_h m^D$ for a constant $C_h = C_h(\alpha, \rho)$, where H is the classical height;

(ii) $\theta_m = 0$ for some m if and only if $\rho \in \mathbb{Z}\alpha + \mathbb{Z}$, and then that index is unique;

(iii) if $\theta_m \neq 0$ then $|\theta_m| \geq (1 + H(\theta_m))^{-D}$, hence $\|m\alpha + \rho\| \geq c_L m^{-D^2}$ with $c_L = c_L(\alpha, \rho) > 0$.

Proof. (i) $\theta_m \in \mathbb{Q}(\alpha, \rho)$, so its degree divides D . Each conjugate is $m\alpha^{(i)} + \rho^{(i)} - j$ with $|j| \leq m+1$, hence $O(m)$; the coefficients of the characteristic polynomial of multiplication by θ_m are the elementary symmetric functions of these D conjugates, hence $O(m^D)$, and the denominators of α, ρ clear once and for all.

(ii) $\theta_m = 0$ means $m\alpha + \rho \in \mathbb{Z}$, i.e. $\rho \in \mathbb{Z}\alpha + \mathbb{Z}$. Two indices $m \neq m'$ would give $\alpha = (j - j')/(m' - m) \in \mathbb{Q}$.

(iii) Let $P(X) = \sum_{i \leq d} a_i X^i$ be the primitive integer minimal polynomial of $\theta := \theta_m \neq 0$, so $a_0 \neq 0$ and $|a_0| \geq 1$. Then $\prod |\text{roots}| = |a_0/a_d| \geq 1/H$ and every root has modulus $\leq 1 + H/|a_d| \leq 1 + H$, whence $|\theta| \geq (1 + H)^{-d} \geq (1 + H)^{-D}$. Combining with (i) gives the stated power of m . $\square$

No transcendence theory enters Lemma 18.1: the bound is Liouville's, and it is *polynomial* in m , which is more than the construction needs. The exceptional case of (ii) is handled by (2) exactly as $\rho = 0$ was in §17, and this is insensitive to the value of the exceptional index — which, unlike the rational case, need not be 0: for $\alpha = (\sqrt{5} - 1)/2$ and $\rho = 3 - 5\alpha$ it is $m = 5$.

Theorem 18.2 (Critical density, quadratic slope, algebraic intercept). *Let $\alpha \in (0, 1)$ be a quadratic irrational and let ρ be any real algebraic number (interpreted modulo 1). Then $s(\alpha, \rho)$ is aperiodic with ones-density exactly β and factor complexity $2n$, and*

$$\Phi(s(\alpha, \rho)) \notin \mathbb{Q}.$$

Proof. As for Theorem 17.5, with Lemma 18.1(iii) in place of Lemma 17.3 and $t(M) := \min(\min_{0 \leq m < M} c_m, c_L M^{-D^2})$. Then $t(M) \geq \exp(-O((\log(M+2))^\kappa))$, since a power of M dominates $\exp(-c_1(\log M)^\kappa)$ from above, and Proposition 17.4 goes through verbatim: the upper bound on $\ell(M)$ becomes $\log \ell(M) \leq \max(c_1(\log(M+2))^\kappa, D^2 \log M + O(1)) + O(1)$, still $O((\log(M+2))^\kappa)$, and $\ell(M) > 1/((A+2)t(M)) \rightarrow \infty$. The β -arc bound (3) holds for algebraic ρ with D in place of 2 (§19), and the discrepancy bound never sees ρ at all. $\square$

Remark (The exceptional intercepts reduce, too). *If $\rho \in \mathbb{Z}\alpha + \mathbb{Z}$, say $\rho = j_0 - m_0\alpha$, then $\{n\alpha + \rho\} = \{(n - m_0)\alpha\}$ for all n , so $s(\alpha, \rho)$ is the $\rho = 0$ coding read from index $-m_0$. Since $\Phi \circ S = T \circ \Phi$ and both branches $y \mapsto 2y$, $y \mapsto (2y - 1)/3$ of T^{-1} preserve rationality, rationality of Φ is invariant under the shift in both directions, so these cases also follow from the $\rho = 0$ case of Theorem 17.5. Both routes were verified [56].*

Corollary 18.3 (Translated intervals). *Let $\alpha \in (0, 1)$ be a quadratic irrational and let $u, \rho \in \mathbb{R}$ with $\rho - u$ algebraic. If $s_n = 1 \iff \{n\alpha + \rho\} \in [u, u + \beta) \pmod{1}$, then $\Phi(s) \notin \mathbb{Q}$.*

Proof. $\mathbf{1}_{[u, u+\beta) \pmod{1}}(x) = \mathbf{1}_{[0, \beta)}(\{x - u\})$ — both sides are the indicator of the same arc of length β , and the identity holds also when $u + \beta > 1$ and the arc wraps. So s is the coding $s(\alpha, \rho - u)$, and Theorem 18.2 applies. $\square$

Corollary 18.3 is strictly stronger than Theorem 18.2: what must be algebraic is the *offset* between the intercept and the interval's left endpoint, not the intercept. Both may be transcendental — $\rho = \pi$, $u = \pi - 1/7$, say. Accordingly the remaining obstruction on this axis is a transcendental offset, where both boundary bounds fail at once: Lemma 18.1 needs algebraicity, and $\|m\alpha + \rho - \beta\|$ stops being a logarithmic form in algebraic data.

Remark (Two widenings that are not available). *The proof needs α algebraic (for §19) and of bounded type (for Lemma 17.3/18.1 and Proposition 17.4). It therefore holds verbatim for every algebraic α with bounded partial quotients; whether any of degree ≥ 3 exists is a well-known open problem, so that generality is presently vacuous, which is why the theorems are stated for quadratics. For transcendental α the method fails outright: A_m is then transcendental and no logarithmic-form estimate applies.*

19 The Analytic Input, and Its Constants

The one analytic ingredient is a lower bound for $|A_m \log 3 - \log 2|$ where $A_m = m\alpha + \rho - j_m$ is algebraic. We use Baker's algebraic-coefficient theorem; to avoid a clash with the density β we write the coefficients as b_i .

Theorem 19.1 (Baker [46], Theorem 2). *Let $\alpha_1, \dots, \alpha_n$ and $b_1, \dots, b_n$ be non-zero algebraic numbers. Suppose that either $\log \alpha_1, \dots, \log \alpha_n$ or $b_1, \dots, b_n$ are linearly independent over the rationals. Suppose further that $\kappa > n$, and let d and B denote respectively the maximum of the degrees and heights of $b_1, \dots, b_n$. Then*

$$|b_1 \log \alpha_1 + \dots + b_n \log \alpha_n| > C e^{-(\log B)^\kappa}$$

for some effectively computable $C = C(n, \alpha_1, \dots, \alpha_n, \kappa, d) > 0$.

Lemma 19.2 (Hypotheses, verified). *Take $n = 2$, $\alpha_1 = 3$, $\alpha_2 = 2$, $b_1 = A_m$, $b_2 = -1$. Then:*

- (i) $A_m > \beta - \frac{1}{2} = 0.1309297536 \dots > 0$ for every $m \geq 0$, every α and every ρ ;
- (ii) $\Lambda_m := A_m \log 3 - \log 2 \neq 0$;
- (iii) $\log 3$ and $\log 2$ are linearly independent over $\mathbb{Q}$;
- (iv) $\deg A_m \leq D$ and $H(A_m) \leq C'_h m^D$, with $D = [\mathbb{Q}(\alpha, \rho) : \mathbb{Q}]$.

Proof. (i) j_m is the nearest integer to $m\alpha + \rho - \beta$, so $|A_m - \beta| \leq \frac{1}{2}$; and $\beta > \frac{1}{2}$. (ii) $\Lambda_m = 0$ would make $\log_3 2 = A_m$ algebraic, contradicting Gelfond–Schneider. (iii) A relation $a \log 2 + b \log 3 = 0$ with $(a, b) \neq 0$ gives $2^a = 3^{-b}$, contradicting unique factorisation. (iv) As in Lemma 18.1(i). $\square$

Item (i) deserves emphasis: it makes Baker’s non-vanishing hypothesis hold with *no* case analysis — in particular at $m = 0$, and at an exceptional index of Lemma 18.1(ii). This matters because the *second* branch of the disjunction in Theorem 19.1 — linear independence of b_1, b_2 , i.e. $A_m \notin \mathbb{Q}$ — genuinely fails at $m = 0$ when $\rho \in \mathbb{Q}$; it is the first branch, item (iii), that is load-bearing.

Since C depends on d and not on B , Lemma 19.2 gives (3) with a single constant for all m , for any fixed $\kappa > 2$. When $D > 2$ the degree $\deg A_m$ may vary with m , and it is then cleaner to invoke Baker’s Part I [45], Theorem 1.1, which is stated for all algebraic $b_1, \dots, b_n$, not all zero, of degree at most d , and so supplies one constant for every degree $\leq D$; the price is $\kappa > n + 1$ and the extra hypothesis that $\log \alpha_1, \dots, \log \alpha_n$ and $2\pi i$ be $\mathbb{Q}$ -independent, which holds here by taking imaginary parts. The index $m = 0$ needs no theorem at all: $c_0 = \|\rho - \beta\|$ is one explicit positive number.

19.1 Stretched exponential, not polynomial

Remark (No polynomial bound is available, and none is used). *Theorem 19.1 gives $e^{-(\log B)^\kappa}$ with $\kappa > n$, not $B^{-\kappa}$. A polynomial bound is the case $\kappa = 1$, which is Baker’s own conjecture and is unproved. It is also unnecessary: Proposition 17.4 needs only $\log \ell(M) = o(M)$, and $(\log M)^\kappa = o(M)$ for every fixed κ . D6 of §1 records that an intermediate stage of this work claimed the polynomial bound; that claim is withdrawn.*

Remark (No reduction to integer coefficients). *It is tempting to clear denominators: for $\rho = p/q$,*

$$q\Lambda_m = (qm)(\alpha \log 3) + (p - qj_m) \log 3 - q \log 2,$$

which has rational integer coefficients. But the first “logarithm” is $\alpha \log 3 = \log(3^\alpha)$, and 3^α is transcendental by Gelfond–Schneider, α being algebraic irrational. Linear-forms-in-logarithms theorems require algebraic arguments, so the sharp integer-coefficient estimates — Baker–Wüstholz, Laurent–Mignotte–Nesterenko — do not apply, with or without clearing denominators. The algebraic-coefficient theory is the only route.

19.2 An explicit constant

Waldschmidt’s algebraic-coefficient measure supplies a numerical constant, and a better exponent.

Theorem 19.3 (Waldschmidt [47], Theorem 10.1). *Let $\ell_1, \dots, \ell_m$ be logarithms of algebraic numbers, $\alpha_i = \exp(\ell_i)$, let $b_1, \dots, b_m$ be algebraic, let D be the degree of $\mathbb{Q}(\alpha_1, \dots, \alpha_m, b_1, \dots, b_m)$, and let $A_1, \dots, A_m, B \geq e$ satisfy $\log A_i \geq h(\alpha_i)$, $D \log A_i \geq e|\ell_i|$, $\log B \geq h(1 : b_1 : \dots : b_m)$ and $B \geq (50mD \log A)^{3m}$ with $A = \max A_i$. If $\Lambda = b_1 \ell_1 + \dots + b_m \ell_m \neq 0$ then*

$$|\Lambda| > \exp\{-(50m)^{3m} D^{m+2} (\log B)^2 \log A_1 \cdots \log A_m\}.$$

The hypotheses hold for our form by Lemma 19.2; note that Theorem 19.3 imposes *no* independence condition on the coefficients, which removes the one place where $m = 0$ fell outside a branch of Theorem 19.1. With $m = 2$, $h(3) = \log 3$, $h(2) = \log 2$, $\log A_1 = \max(1, \log 3, e \log 3/D)$ and $\log A_2 = \max(1, \log 2, e \log 2/D)$:

D	$\log A_1$	$c_1 = (50 \cdot 2)^6 D^4 \log A_1 \log A_2$	$\log B_{\min}$
2	1.49317	2.389×10^{13}	34.20
3	1.09861	8.899×10^{13}	34.79
4	1.09861	2.812×10^{14}	36.51
6	1.09861	1.424×10^{15}	38.95

So $c_m > \exp(-c_1(\log B_m)^2)/\log 3$ with $\log B_m = \max(\log B_{\min}, \log H(A_m) + 2)$ and every constant numerically supplied. The exponent is exactly 2, sharper than Theorem 19.1’s $\kappa > 2$.

Remark (What the explicit constant costs). *Propagated through Proposition 17.4, and taking the discrepancy constant optimistically as $C = 1$, the surplus first becomes positive near $M \approx 2.7 \times 10^{17}$, at a witness length $\ell(M)$ of order $10^{8 \times 10^{16}}$. The explicit route therefore buys rigour, not numbers: it replaces “effectively computable in principle” by a written-down bound. The numerically useful height floors remain those of §20, which need no analytic input at all.*

Remark (Effective in principle versus numerically supplied). *These are different, and this note keeps them apart.*

constant	role	numerically supplied?
the floors of §20	height bounds at computed witnesses	yes , exactly
$K = (A + 2)q^2$	arc at 0, rational ρ	yes
c_L, D	arc at 0, algebraic ρ	yes (Liouville, per α, ρ)
c_1	arc at β	yes , via Theorem 19.3
C, C_0	drift and discrepancy	no — requires the Kuipers–Niederreiter constants

One constant therefore stands between Theorem 17.5 and a fully numerical $H \geq 2^{f(M)}$ at every M . Making it explicit is bookkeeping in a classical estimate [48], not new mathematics [58].

20 Finite Height Certificates

Independently of every analytic input, the bridge yields explicit lower bounds on the height of any rational that could equal a given conjugacy value. If $\Phi(s) = u/v$ in lowest terms and $H =$

$\max(|u|, v)$, then $2^L \leq H F(W)$ with $F(W) = |2^\ell - 3^k| + c_W$, so $H \geq 2^{L - \log_2 F(W)}$. Each row below is an exact integer comparison $2^L > F(W) \cdot 2^S$; no floating-point value and no analytic estimate enters.

ℓ	k	L	$H \geq$
377	238	846	2^{462}
610	385	1456	2^{838}
1597	1008	2478	2^{872}
2584	1630	5062	2^{2469}
6765	4268	22773	2^{15998} (a 4816-digit bound)

These are for $\alpha = (\sqrt{5} - 1)/2$, $\rho = 1/7$. Analogous certificates were computed across the families of Theorems 17.5 and 18.2 — ten (α, ρ) pairs with $\rho \in \mathbb{Q}$, giving floors from 2^{140} to 2^{8069} , and seven with ρ algebraic of field degree 2, 4 and 6, giving 2^{67} to 2^{598} [55, 56].

Remark (These are not new as a kind of result). *Producing effective height floors by this bridge is the foundation paper’s own claimed contribution [1]; the certificates above are new instances of an existing method, not a new method. They are also not reached by the density results of §16, which are contradiction arguments and bound nothing.*

21 A Measurement, Not a Theorem: the Ridout Exponent

One observation is recorded because it is the most promising open direction the audit surfaced, and it is recorded as an observation only.

Theorem 21.1 (Ridout; as stated by Bugeaud–Kekeç [51], Theorem 1.3). *Let ξ be a p -adic number and $\varepsilon > 0$. Suppose there are rationals x_n/y_n with $\gcd(x_n, y_n) = 1$, heights $2 \leq |x_1, y_1| < |x_2, y_2| < \dots$, and $0 < |\xi - x_n/y_n|_p < |x_n, y_n|^{-2-\varepsilon}$ for all n . Then ξ is transcendental.*

Our approximants have $|\Phi(s) - c_W/\delta_W|_2 = 2^{-L}$ and reduced height $H_{\text{red}} \leq F(W) = 2^{\ell + O(\log \ell)}$, so the relevant exponent is $E(\ell) := L(\ell)/\log_2 H_{\text{red}}(\ell)$, and Theorem 21.1 asks for $E > 2 + \varepsilon$ infinitely often. Measured at every convergent denominator $3 \leq q \leq 28657$ of $(\sqrt{5} - 1)/2$ for the word $\alpha = (\sqrt{5} - 1)/2$, $\rho = 1/7$ [57]:

ℓ	L	$\log_2 H_{\text{red}}$	E	$E > 2$
8	24	9.12	2.631	yes
55	147	59.46	2.472	yes
377	846	383.66	2.205	yes
610	1456	617.26	2.359	yes
987	2443	995.13	2.455	yes
4181	9243	4190.94	2.205	yes
6765	22773	6774.98	3.361	yes

Seven of the twenty convergents tested exceed the threshold; the maximum observed is 3.361.

Remark (What is and is not claimed). *Nothing is claimed. Theorem 21.1 requires infinitely many such n with a fixed ε , which by Lemma 17.2 is the statement that $m^*(\ell)/\ell > 1 + \varepsilon$ infinitely often — an open question about hitting times. Seven of twenty, with no visible pattern, is not evidence*

of infinitude, and the witness family of Proposition 17.4 deliberately drives $E \rightarrow 1$, so it is no evidence either way. If the statement did hold, the conclusion of Theorem 17.5 would strengthen from irrationality to transcendence, answering the foundation paper’s open problem (3) for this word. Controlling $m^*(\ell)$ from below along a subsequence needs the opposite of what Proposition 17.4 arranges, so this would be a different argument, not a refinement of the present one.

22 Interpretation: Symbolic, Statistical, and Arithmetic Space

This section is conceptual, not a theorem statement, and should be read as orientation for a reader extending this work rather than as a claim.

It is useful to separate three levels at which a parity word can be described:

Symbolic space. The actual word: its literal order, block structure, and factor-by-factor arrangement.

Statistical projection. Aggregated summaries of the symbolic object — density $\gamma = k/\ell$, discrepancy $D(W)$, block frequencies, factor complexity. Many distinct symbolic objects can share the same statistical projection.

Arithmetic admissibility. Which symbolic objects, specifically, correspond to a rational Φ -value. This is a pointwise, not a statistical, property.

The relationship is a *fiber*: symbolic object $\rightarrow$ statistical state is generally many-to-one. A crucial consequence, easy to state and easy to get wrong in either direction: **statistical sparsity of a class (low complexity, small discrepancy, controlled density) does not by itself imply arithmetic impossibility of rational Φ -values within it**, and conversely, statistical richness does not by itself force irrationality. The bridge proved in this note works precisely because it does *not* stop at a statistical argument: it lifts symbolic and statistical information (repetition length, discrepancy) back into a pointwise arithmetic contradiction (Theorem 5.1’s exact integer divisibility argument) for the *specific* word s under study, not for a statistical ensemble containing it. This is why the method scales from Sturmian words to CS Rote sequences without a change in kind — and also why it stalls exactly where the statistical control (discrepancy, excess) it depends on stalls (§14).

23 The Drift/Resonance Coordinate

For a finite word W ($\ell = |W|$, $k = |W|_1$), define

$$R(W) := \ell - k \log_2 3.$$

Then $2^\ell - 3^k = 2^\ell (1 - 2^{-R(W)})$, and near $R = 0$ (i.e. near the “resonance” $\ell \approx k \log_2 3$), $|2^\ell - 3^k| \approx 2^\ell \ln 2 |R(W)|$.

This coordinate is explanatory, not structural: it was investigated directly in Phase 7 as a candidate new invariant, and the record is explicit that it is not one. “No new information is encoded by R that γ did not already carry” [33]; “Resonance did not materially help or hurt in any tested case” [34]. $R(W)$ is a reparameterization of the density margin $\gamma \log_2 3$ already used throughout §7–§10, useful for visualizing how close a given witness sits to the arithmetic boundary, but supplying no additional proof-theoretic leverage found anywhere in this program.

24 The Symmetry-Tradeoff Probe (Exploratory, Inconclusive)

This section reports an exploratory investigation. Nothing in it is a theorem, and nothing in it should be read as evidence for or against any conjecture stated elsewhere in this note. It is included so a future researcher does not repeat the same probe without knowing what was already tried and why it did not resolve anything.

The idea investigated: does *approach to symmetry* in the Ostrowski digit sequence (roughly, how often the maximal admissible continuation is available) trade off against arithmetic *resonance* (how close $R(W)$, §23, sits to 0)? Two experiments were run.

Experiment 1 correlated a normalized arithmetic-slack quantity $A(W) := |1 - 2^{-R(W)}|$ against the exact surplus $S(W)$ and against the achieved repetition ratio L/ℓ , using 74 genuine witnesses from the general-intercept search of §14. Result: $A(W)$ vs. $S(W)$, $r \approx 0.208$; $A(W)$ vs. L/ℓ , $r \approx -0.125$. Both fall below the probe’s own 0.3 threshold for “worth a theorem-seeking follow-up,” and $n = 74$ is too small for $r = 0.208$ to clear the standard significance threshold for that sample size. Separately, the first correlation is only weak evidence in any case: the surplus machinery already leans on the same $\ell - k \log_2 3$ resonance quantity that defines $A(W)$, so a correlation there may be partly definitional rather than an independent discovery.

Experiment 2, testing a genuinely independent quantity (a windowed frequency of admissible-digit freedom, N_{sym} , against $A(W)$), did not run to completion in its original form: it materialized an explicit substitution-image word whose length grows exponentially in the search depth k (for slope $A = 6$, a projected $\sim 9.9 \times 10^{14}$ symbols by $k = 19$), exhausting memory. This was diagnosed and repaired: the word’s length and weight can be obtained exactly, without ever materializing it, via a 2×2 integer transfer-matrix representation of the same substitutions (τ_0, τ_1 ; see §25), cross-checked against brute-force construction in 150 of 150 small cases with exact agreement. With this fix, Experiment 2 completed on 724 points: $A(W)$ vs. N_{sym} , $r \approx -0.0058$; on a log-log scale, $r \approx -0.0319$. **These numbers must not be read at face value:** 596 of the 724 points (82%) had $A(W)$ pinned to exactly 1.0 by the probe’s own floating-point underflow clamp ($|R(W)| > 60$), an artifact of the diagnostic’s float arithmetic, not a genuine measurement in the near-resonance regime where any real effect would have to live.

Verdict: INCONCLUSIVE. Neither experiment produces a correlation free of confounding structure (definitional entanglement in Experiment 1; a numerical clamping artifact dominating Experiment 2). This is consistent with, and independently corroborates, the Phase 7 drift-coordinate finding of §23 that resonance carries no separately exploitable signal in every case tested so far — but it establishes nothing new, and the *symbolic-symmetry/arithmetic-freedom tradeoff hypothesis* is neither supported nor contradicted by this probe.

A methodologically sound re-run would need $A(W)$ computed at sufficient precision to avoid the clamp, restricted to (or stratified by) genuine near-resonance witnesses, before any correlation from it could be informative.

25 A Transfer-Matrix Computational Tool

This section records a computational/structural tool, not a new irrationality theorem. It is preserved here because it is broadly useful and easy to lose track of.

The Sturmian substitutions used throughout this program (e.g., in constructing explicit BHZ witness words, §14) are

$$\tau_0 : 0 \mapsto 0, 1 \mapsto 01; \quad \tau_1 : 0 \mapsto 10, 1 \mapsto 1.$$

Applying either substitution repeatedly can make the resulting word’s length grow exponentially in the number of applications — and that length is all that is needed for most downstream computations (arithmetic slack $A(W)$, weight κ , density). It need never be materialized. Tracking only (n_0, n_1) , the counts of 0s and 1s, the two substitutions act linearly:

$$\tau_0 : (n_0, n_1) \mapsto (n_0 + n_1, n_1), \quad \tau_1 : (n_0, n_1) \mapsto (n_0, n_0 + n_1),$$

i.e. by the integer matrices $M_0 = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$, $M_1 = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$ (both determinant 1; their products are exactly the continuant recurrence generating the convergent denominators q_k). Applying τ_i exactly e times in a row is M_i^e ; composing across an entire substitution schedule of length k is k integer matrix–vector multiplications, giving the word’s exact length $\ell = n_0 + n_1$ and weight $\kappa = n_1$ in $O(k)$ exact-integer steps regardless of how astronomically large ℓ becomes — verified against brute-force construction with 150/150 exact agreement wherever brute force was still feasible to run.

A related, separate observation: counting the number of admissible Ostrowski digit sequences (c_k) (rather than computing one fixed sequence’s word) is itself governed by a minimal two-state automaton, not by any larger state. The admissibility rule (§14) depends only on whether the immediately preceding digit was exactly 0: from that state, the next digit has $a_k + 1$ admissible choices; from any nonzero-previous state, only a_k . No memory of the actual previous digit value, nor of anything beyond the current index into (a_k) , is needed. This directly mirrors the admissible-sequence generator already used throughout the repository’s scripts, and is recorded here as a structural fact about that generator, not as a separately theorem-proving result.

26 What Is Still Open

Open 1 — General intercept margin. CLOSED (Revision 2, C1).

Revision 1’s Conjecture 12.1 is BHZ Propositions 5.1 and 5.2 (Theorem 14.1), with an attained-witness refinement in Theorem 14.3. Theorem 12.1 now holds at every intercept. What remains here is only a sharpness question, not an obstruction: is $\frac{1}{A\beta_A}$ the infimum of $\text{ice} - 2$ on slopes whose digit tail is *not* eventually constant? Proved for eventually-constant tails with $A \geq 2$; numerical elsewhere [41]. Nothing depends on the answer.

Open 2 — Return/exceptional intercept set. WITHDRAWN (Revision 2, C2).

The premise was false: the set of intercepts with $\text{ice}(\omega) < \text{ind}^*(\gamma)$ is uncountable, not a countable return set, and intercept 0 sits among the *easiest* intercepts rather than the hardest. There is no distinguished countable family left to classify.

Open 3 — Intrinsic-square bypass. SUPERSEDED (reading confirmed).

The Revision 1 analysis was correct — bare squares ($r = 2$) give a fixed margin $2 - \log_2 3$ on the even-weight branch and *exactly zero* on the odd-weight branch [35] — and Theorem 14.3 makes the bypass unnecessary by supplying $r > 2$ strictly at every intercept. The route retains one distinct virtue, recorded in Open 4: Allouche–Davison–Queffélec–Zamboni’s square theorem has *no* hypothesis on γ , so it is the only part of the chain that survives unbounded partial quotients.

Open 4 — Unbounded partial quotients. THE LIVE FRONTIER.

Remove the bounded-partial-quotient hypothesis from Theorem 12.1. This is no longer merely “flagged as hard”: BHZ Theorem 1.1 *characterises* the slopes carrying an intercept with $\text{ice}(\omega) = 2$ exactly — for each pair (s, t) , $s > 1$, only finitely many k with $(a_k, a_{k+1}) = (s, t)$ or $(a_k, a_{k+1}, a_{k+2}) = (1, 1, t)$ — and every such slope has unbounded partial quotients. At such an intercept the odd-weight branch has margin exactly 0, so the present sufficient criterion cannot apply, independently of any discrepancy estimate. Three distinct questions, in increasing order of ambition: (i) can the *even* branch be forced infinitely often, where bare squares already give $2 - \log_2 3$ and the discrepancy hypothesis can be dropped entirely? (ii) is the $O(\log \ell)$ discrepancy bound replaceable by an exact height estimate that tolerates $E_k = 0$? (iii) does a CS Rote sequence carry intrinsic initial repetition of its own, beyond what transfers from u ? See `theorems/phase8/PHASE8_NEXT_TARGET.md`.

Open 5 — Higher complexity.

Only once the $p(n) = 2n$ rung is fully closed (Open 1–4): which symbolic classes beyond Rote sequences — higher, structured complexity classes; general two-interval rotation codings, for which the repetition-existence half of the argument is itself still open [36]; general morphic words — can this bridge reach next?

Open 6 — The full Periodicity Conjecture.

What mechanism, if any, could cover aperiodic words that admit no useful repetition structure at all? **Revision 2 answered this badly and the answer is withdrawn** (D1 of §1). It asserted that because the Thue–Morse word t has no square prefix below half-length 2048 [37], every purely repetition-based method is structurally inapplicable to it. That does not follow, and it is false: the bridge needs periodic approximants whose 2-adic agreement outruns their height, not a square. For t the substitution supplies them — $\mu(0) = 01$, $\mu(1) = 10$ satisfy $\text{lcp}(\mu(x), \mu(y)) = 2\text{lcp}(x, y)$, so iterating μ doubles the agreement while the height grows only with the length — and $\Phi(t) \notin \mathbb{Q}$ follows from Theorem 5.1 with no square anywhere [52]. (The conclusion is in any case available from Corollary 16.2, t having density $1/2 < \beta$.) What remains genuinely open is the conjecture itself, which quantifies over an uncountable family of which every class treated here is a measure-zero subset; and the honest statement is that no mechanism in this note is known to extend to that family, not that a structural obstruction has been identified.

Open 7 — Beyond an algebraic offset.

Corollary 18.3 covers the coding of a quadratic rotation by an interval of length β whenever the offset between intercept and left endpoint is algebraic. When that offset is transcendental both boundary bounds fail at once: Lemma 18.1 needs algebraicity, and $\|m\alpha + \rho - \beta\|$ is no longer a logarithmic form in algebraic data. Is there any route there, or is a transcendental offset a real barrier?

Open 8 — Explicit constants.

Evaluate the Kuipers–Niederreiter constants [48] for the interval $[0, \beta)$ and bounded-type α , obtaining C and C_0 of Remark 19.2 numerically. With c_1 already explicit (Theorem 19.3) this would give a written-down $H \geq 2^{f(M)}$ at every M — the first fully numerical statement in this programme beyond the finite certificates.

Open 9 — Transcendence.

Does $m^*(\ell)/\ell > 1 + \varepsilon$ hold for infinitely many convergent denominators, for a fixed $\varepsilon > 0$?

If so, §21 upgrades Theorem 17.5 from irrationality to transcendence. Measured data are suggestive and prove nothing.

27 Future-Work Roadmap

The roadmap below is organized directly around the six open problems of §26 and the repository's own stated priorities.

Near term.

Open 1–3 are settled or withdrawn, so the near-term work is not mathematical but editorial and hygienic: propagate the $q_1 = a_1 + 1$ convention through the underlying record's code and re-derive every root length reported there (no ice value changes), and fold Lemma 14.4 into the discrepancy audit where it was previously implicit.

Medium term.

Open 4(i): a general-intercept weight-parity criterion. The even-weight branch survives $\text{ice}(\omega) = 2$ with margin $2 - \log_2 3 \approx 0.415$ and needs no discrepancy hypothesis, so it suffices to force even weight infinitely often. At intercept 0 this was the Phase 3 automaton; at general intercepts the prefix-count parity is the classical Ostrowski-digit sum $k_i(u) = \lfloor i\gamma + \rho \rfloor - \lfloor \rho \rfloor \pmod 2$, so a literature search on Ostrowski-digit parity should precede any new derivation. This is the single highest-value target identified by Revision 2.

Longer term.

Open 4(ii)–(iii): an exact height estimate tolerating zero repetition excess, or intrinsic Rote repetition not inherited from u . Both would be genuinely new mechanisms rather than extensions, and both should be attempted only after 4(i) is decided.

Structural.

Open 6 is not, as Revision 2 claimed, blocked by a structural obstruction; D1 of §1 withdraws that claim. What is true is weaker and still discouraging: no mechanism in this note is known to reach an arbitrary aperiodic word, and the classes reached so far are parametrised families of measure zero. The nearest concrete targets are Open 7–9.

Process.

Continue the falsification-first discipline of §11: every extension of this program should include an adversarial audit pass against genuinely non-periodic, non-quadratic controls before any theorem upgrade is accepted, exactly as Phases 5 and 6 did for the bounded-type theorem.

Provenance and Verification

Revision 3 was prepared from the `periodicity-conjecture-bridge` repository on an isolated branch, from the Phase 9–13 deliverables (`theorems/phase9b/`, `theorems/phase10/`, `theorems/phase11/`, `theorems/phase11b/`, `theorems/phase12/`, `theorems/phase13/` with the matching `scripts/` and `data/` directories), which was not pushed during this preparation and from which nothing was published. No Phase 1–8 research file was rewritten; the Phase 7, 10, 11, 11b, 12 and 13 records are preserved verbatim and carry conspicuous correction and clarification banners. Every statement in §§16–21 was extracted from the corresponding file under `theorems/phase{9b,10,11,11b,12,13}/` and re-checked against the verification scripts named in

the bibliography; none was reconstructed from memory. Every pass/fail decision in those scripts is an exact integer or exact algebraic-number comparison, and comparisons against β use a certified 300-digit bracket that either decides or raises. Theorem 17.5 was additionally rebuilt from the definitions in independent code that imports none of the earlier phase scripts [55]. The deposited Revision 2 PDF and its erratum were not rebuilt or modified in the preparation of this revision.

Revision 2 was prepared from the `periodicity-conjecture-bridge` repository on the isolated branch `phase8-general-intercept`, whose Phase 8 deliverables (`theorems/phase8/`, `scripts/phase8/`, `data/phase8/`) carry every corrected statement, and which was not pushed during this preparation. No Phase 1–7 research file was rewritten: the Phase 7 record is preserved verbatim and carries conspicuous correction banners pointing to `theorems/phase7/PHASE7_CORRECTIONS.md` [43]. Revision 1 was consolidated from the same repository at commit `a4ac46aebf17fb2ce14cea50f5ce60a58704da15` (“Phase 7: map general-intercept CS Rote obstruction”). Every theorem statement, definition, and quoted result above was extracted directly from the corresponding file under `theorems/phase{1..7}/` and cross-checked against `THEOREM_STATUS.md` and `LADDER.md`; none was reconstructed from memory. SHA-256 checksums of every file under `theorems/`, `scripts/`, `data/`, and `reference/` at the time of writing are recorded in `technical-note/PRESERVATION_RECORD.md`. A full statement-by-statement classification (known/cited, proved in foundation paper, proved in this program, computationally verified, counterexample, rejected hypothesis, open, or exploratory only) is recorded in `technical-note/SOURCE_OF_TRUTH.md`. No Phase 1–7 research file was modified in the preparation of this note.

Revision 1 is deposited as Zenodo version DOI 10.5281/zenodo.23045141 and Revision 2 as a new version of the same record. Revision 3 is likewise intended as a *new version*, leaving both earlier revisions retrievable, not as an edit of either. At the time of writing Revision 3 is a draft and has not been deposited.

Acknowledgments

I thank Josefina López and Peter Stoll, whose work on density and the $3x+1$ conjugacy map guided this programme toward the critical density $\beta = \ln 2 / \ln 3$; the results here are established by the independent arguments given above and do not require their proposed upper-density exclusion. I also record that a technical question about one step of [9] was raised publicly by a third party; I have not verified it in either direction, and no argument in this note depends on it.

The author is an independent researcher whose professional experience is outside mathematics. **This note has not been refereed**, and neither has the foundation paper [1], which is the present author’s own preprint. Specialist review is invited: see `docs/INVITATION_TO_DOMAIN_EXPERTS.md` in the repository for the review and extension targets most likely to repay attention, and `docs/DEPENDENCE_ON_PRIOR_RESULTS.md` for exactly which external results the conclusions rest on.

AI assistance (Anthropic’s Claude) was used for repository archaeology, adversarial diagnostic work (including diagnosing and repairing the Experiment 2 performance bug in §24), literature cross-checking against primary sources, exact-arithmetic verification scripting, and drafting of this note, consistent with the acknowledgment practice of the foundation paper [1]. All mathematical claims, their framing, and any errors are the author’s.

References

- [1] Elias De Jesús, *The $3x+1$ Conjugacy Map Sends Every Sturmian Word to an Irrational 2-Adic Integer*, Zenodo, 28 September 2026, version DOI [10.5281/zenodo.23019799](https://doi.org/10.5281/zenodo.23019799) (concept DOI [10.5281/zenodo.22920055](https://doi.org/10.5281/zenodo.22920055)), CC BY 4.0. **Preprint; not refereed; by the present author — a self-citation.** (Foundation paper for this research program.)
- [2] Jean-Paul Allouche, J. L. Davison, Michel Queffélec, and Luca Q. Zamboni, *Transcendence of Sturmian or morphic continued fractions*, Journal of Number Theory **91** (2001), no. 1, 39–66.
- [3] Daniel J. Bernstein, *A noniterative 2-adic statement of the $3N + 1$ conjecture*, Proceedings of the American Mathematical Society **121** (1994), no. 2, 405–408.
- [4] Daniel J. Bernstein and Jeffrey C. Lagarias, *The $3x + 1$ conjugacy map*, Canadian Journal of Mathematics **48** (1996), no. 6, 1154–1169.
- [5] Valérie Berthé, Charles Holton, and Luca Q. Zamboni, *Initial powers of Sturmian sequences*, Acta Arithmetica **122** (2006), no. 4, 315–347.
- [6] Artūras Dubickas, *On integer sequences generated by linear maps*, Glasgow Mathematical Journal **51** (2009), no. 2, 243–252.
- [7] Jeffrey C. Lagarias, *The $3x + 1$ problem and its generalizations*, The American Mathematical Monthly **92** (1985), no. 1, 3–23.
- [8] Josefina López and Peter Stoll, *The $3x + 1$ conjugacy map over a Sturmian word*, Integers **9** (2009), no. A13, 141–162.
- [9] Josefina López and Peter Stoll, *The $3x + 1$ periodicity conjecture in $\mathbb{R}$* , arXiv:2101.12747, 2021. **Preprint.** Cited for context; its Theorem 1 is not used in this note.
- [10] M. Lothaire, *Algebraic Combinatorics on Words*, Encyclopedia of Mathematics and its Applications, vol. 90, Cambridge University Press, 2002, Chapter 2 (Sturmian words, by J. Berstel and P. Séébold). Cited here as the standard reference for the combinatorics-on-words background of complementary symmetric Rote sequences; the primary source defining this class (G. Rote, 1994) should be verified directly against primary literature before formal submission, as this repository’s record does not independently confirm its exact publication details.
- [11] `theorems/phase2/ABSTRACT_IRRATIONALITY_CRITERION.md`, periodicity-conjecture-bridge repository, commit `a4ac46a`.
- [12] `theorems/phase2/SURPLUS_THEOREM.md` and `theorems/phase1/SURPLUS_INVARIANT.md`, *ibid.*
- [13] `theorems/phase1/DISCREPANCY_HEIGHT_THEOREM.md`, *ibid.*
- [14] `theorems/phase1/REPETITION_HEIGHT_CRITERION.md`, *ibid.*
- [15] `theorems/phase2/ROTE_ROOT_DISCREPANCY.md`, *ibid.*
- [16] `theorems/phase2/ROTE_TRANSFER_THEOREM.md`, *ibid.*
- [17] `theorems/phase5/TRANSFER_REDERIVATION.md`, *ibid.*

- [18] `theorems/phase3/WEIGHT_PARITY_CLASSIFICATION.md`, *ibid.*
- [19] `theorems/phase4/SILVER_EXTREMALITY.md`, *ibid.*
- [20] `theorems/phase4/SILVER_CONSTANT_AUDIT.md`, *ibid.*
- [21] `theorems/phase4/REQUIRED_EXPONENT.md`, *ibid.*
- [22] `theorems/phase5/HOSTILE_REFeree_REPORT.md`, *ibid.*
- [23] `theorems/phase6/ARROW_BY_ARROW_RECONSTRUCTION.md`, *ibid.*
- [24] `theorems/phase4/QUADRATIC_CS_ROTATE_THEOREM.md`, *ibid.*
- [25] `theorems/phase6/BOUNDED_TYPE_THEOREM.md`, *ibid.*
- [26] `theorems/phase4/FINAL_PROOF_CHAIN.md`, *ibid.*
- [27] `theorems/phase7/BHZ_GENERAL_INTERCEPT_MAP.md`, *ibid.*
- [28] `theorems/phase7/PHASE7_VERDICT.md`, *ibid.*
- [29] `theorems/phase7/INTERCEPT_SUFFICIENT_CONDITION.md`, *ibid.*
- [30] `theorems/phase7/SURPLUS_DECOMPOSITION.md` and `theorems/phase7/INTERCEPT_EXCESS.md`, *ibid.*
- [31] `theorems/phase7/FLEXIBLE_MARGIN_BOUNDARY.md`, *ibid.*
- [32] `theorems/phase7/STRUCTURED_INTERCEPTS.md`, *ibid.*
- [33] `theorems/phase7/DRIFT_COORDINATES.md`, *ibid.*
- [34] `theorems/phase7/DRIFT_VS_DIGITS.md`, *ibid.*
- [35] `theorems/phase7/INTRINSIC_SQUARE_ROUTE.md`, *ibid.*
- [36] `theorems/phase2/ROTATION_EXAMPLE_THEOREM.md`, *ibid.*
- [37] `theorems/phase1/FULL_PERIODICITY_GAP.md`, *ibid.*
- [38] `theorems/phase8/PHASE8_PROOF_OR_OBSTRUCTION.md`, *periodicity-conjecture-bridge repository*, branch `phase8-general-intercept`.
- [39] `theorems/phase8/PHASE8_TARGET_AUDIT.md`, *ibid.*
- [40] `theorems/phase8/PHASE8_CHAIN_AUDIT.md`, *ibid.*
- [41] `theorems/phase8/PHASE8_MARGIN_STATUS.md`, *ibid.*
- [42] `theorems/phase8/PHASE8_VERDICT.md`, *ibid.*
- [43] `theorems/phase7/PHASE7_CORRECTIONS.md`, *ibid.*
- [44] Kenneth G. Monks and Jonathan Yazinski, *The autoconjugacy of the $3x + 1$ function*, *Discrete Mathematics* **275** (2004), no. 1–3, 219–236.

- [45] Alan Baker, *Linear forms in the logarithms of algebraic numbers I*, Mathematika **13** (1966), 204–216.
- [46] Alan Baker, *Linear forms in the logarithms of algebraic numbers III*, Mathematika **14** (1967), 220–228. (Theorem 2 is the homogeneous algebraic-coefficient case; reviewed in MR0220680.)
- [47] Michel Waldschmidt, *Diophantine Approximation on Linear Algebraic Groups: Transcendence Properties of the Exponential Function in Several Variables*, Grundlehren der mathematischen Wissenschaften **326**, Springer, 2000; and the companion survey *Linear independence of logarithms of algebraic numbers*, Theorem 10.1.
- [48] Lauwerens Kuipers and Harald Niederreiter, *Uniform Distribution of Sequences*, Wiley, 1974, Chapter 2, Theorem 3.4.
- [49] Gerhard Rote, *Sequences with subword complexity $2n$* , Journal of Number Theory **46** (1994), no. 2, 196–213.
- [50] Jean Berstel and Laurent Vuillon, *Coding rotations on intervals*, arXiv:math/0106217, 2001.
- [51] Yann Bugeaud and Gülcan Kekeç, *On Mahler’s classification of p -adic numbers*. (Theorem 1.3 there is Ridout’s 1958 p -adic analogue of Roth’s theorem, quoted in §21.)
- [52] `theorems/phase9b/THUE_MORSE_INDEPENDENT_PROOF.md` and `theorems/phase9b/PHASE9B_VERDICT.md` (repository record; script `scripts/phase9b/tm_independent_proof.py`).
- [53] `theorems/phase10/DENSITY_COMPLEXITY_LADDER.md` and `theorems/phase10/PRIOR_RESULT_COVERAGE.md` (repository record).
- [54] `theorems/phase11b/THEOREM11_DEPENDENCY_AUDIT.md`, `theorems/phase11b/BAKER_SOURCE_AND_APPLICATIONS.md` and `theorems/phase11b/REPAIRED_WITNESS_GROWTH.md` (repository record).
- [55] `theorems/phase12/QUADRATIC_RATIONAL_INTERCEPT_THEOREM.md` and `theorems/phase12/THEOREM11_INDEPENDENT_AUDIT.md` (repository record; scripts `scripts/phase12/independent_audit.py`, `scripts/phase12/general_quadratic.py`).
- [56] `theorems/phase13/ALGEBRAIC_INTERCEPT_THEOREM.md` and `theorems/phase13/CONSOLIDATED_THEOREM_AND_APPLICATIONS.md` (repository record; scripts `scripts/phase13/numfield.py`, `scripts/phase13/algebraic_intercept.py`).
- [57] `theorems/phase13/FAMILY_THEOREM_PRIOR_COVERAGE.md` §5 (repository record; script `scripts/phase13/ridout_exponent.py`).
- [58] `theorems/phase13/EXPLICIT_CONSTANTS_STATUS.md` (repository record; script `scripts/phase13/explicit_constants.py`).