A Certified Negative Finite-Window Weil Witness for the Davenport–Heilbronn Function, with the Remaining-Zero Form Positive

Elias De Jesús
Independent Researcher
ORCID 0009-0007-0190-9143
dejesuselias10@gmail.com

October 2026 — Technical Note
doi:10.5281/zenodo.23608288

Abstract

The Davenport–Heilbronn function f satisfies a ζ-shaped functional equation, is entire after completion, has all its zeros in an explicit strip, and has no Euler product; it is known to have zeros off the critical line. It is therefore the natural negative control for any finite-window Weil positivity detector. We exhibit one explicitly specified real test function g_* supported in $[-2, 2]$, odd, in a 37-dimensional span, whose Davenport–Heilbronn Weil value is enclosed in ball arithmetic as $Q_f(g_*)/\|g_*\|_2^2 = -0.67907134644148059098637758609 3\ldots$ with radius 1.39×10^{-116}, and a second such function at $L = 1.8$ with value $-1.68123895523670103\times10^{-16}$. Both are strictly negative. Decomposing against a census of the zeros to height 200, the single off-line quartet at $\beta = 0.808517182456637386$, $\gamma = 85.6993484853775922$ contributes -0.985198326439305, and the entire remaining spectrum contributes $+0.306126979997825$; the minimum eigenvalue of the remaining-zero form is positive at every scanned window and sector, collapsing to 8.47×10^{-36} at $L = 2.6$ while the full form is at -2.14. On this object, therefore, the hypothesis of the remaining-zero theorem of the companion paper holds, and the negativity is carried by the off-line quartets alone. What this rules out is any claim that finite-window Weil positivity follows from a functional equation, entireness and a zero-containing strip; it says nothing about ζ, and it cannot separate "has off-line zeros" from "has no Euler product".

0. Non-claims

Stated first, not in an appendix.

1. **Nothing about RH or GRH.** Every object here is the Davenport–Heilbronn function. It is not ζ, it is not an L-function of anything, and no statement below transfers to either.

2. **f has no Euler product, and nothing here transfers to ζ.** A detector that sees this function's off-line zeros is a detector that is not blind. It is not thereby a detector that would see ζ's, if ζ had any.

3. **The confound, stated here and not buried.** f differs from ζ in *two* ways at once: it has zeros off the critical line, and it has no Euler product. Every measurement below is consistent with either being the cause of the negativity, and nothing here separates them. Separating them would need an object with one property and not the other, and no such object is known to the author. In particular, the observation of §5 that the remaining-zero form is positive is evidence *internal to this object* that the off-line zeros carry the negativity;

it is not a proof, because the remaining-zero form is itself built from a function with no Euler product.

4. **Tiers.** Every zero location is [T2]. The two witnesses of §4 are [T2-C]. The onset scan of §6 is [T2-N]. **Non-detections certify nothing**: a finite basis that finds no negative direction is not evidence of positivity, and the absence of a witness at $L = 1.6$ is reported as an absence, never as positivity.

5. **The $L = 1.8$ onset is not certified.** A negative witness *at* $L = 1.8$ is certified (§4); that the form is positive at $L = 1.6$ is not, so the crossing is bracketed by one certified negative and one non-detection, and the location of the crossing remains [T2-N].

6. **The two literature witnesses are unverified.** Only landing pages were read; see Appendix C. No quantity here was fitted to either.

7. **A finite positive matrix proves positivity only on its own test space.** The positivity of the remaining-zero form in §5 is a statement about a 37-dimensional restriction at each window, never about the whole finite-window function space.

8. **Numerical root residuals do not establish census completeness**, and agreement with a published decimal is a cross-check, not an interval certificate.

9. **Not treated here.** The degenerate case $\gamma = 0$ of the orbit contribution, where the parity split behaves differently, is the subject of a companion note and is not used anywhere below.

Epistemic tiers

Tier	Meaning	Discharged by
T1	exact or proved	a proof, or exact arithmetic over $\mathbb{Z}$ or $\mathbb{Q}$
T2	numerical	stated working precision + a convergence or noise-floor check
T2-C	certified arithmetic	an Arb ball enclosure with a reproducible certificate
T3	interpretation	argument from T1/T2 evidence, labelled as such

T2-C is *not* a formal-proof assistant verification and is nowhere described as one. Every number printed in this note is re-derived from the data files by `note/verify_note_numbers.py`, which also fails if the note prints more digits than its source carries.

1 The function

Let χ be the character modulo 5 with $\chi(0), \dots, \chi(4) = 0, 1, i, -i, -1$, primitive and odd. Put

$$f(s) \;=\; \tfrac{1}{2}\big[(1 - i\kappa)L(s,\chi) + (1 + i\kappa)L(s,\bar{\chi})\big]. \tag{1}$$

Davenport and Heilbronn [4] introduced this function and showed it has zeros off the critical line; Titchmarsh [8, §10.25] is the standard exposition.

κ from the Gauss sum [T1]. Write $\varepsilon_\chi = \tau(\chi)/(i\sqrt{5})$, so that $\Lambda(s,\chi) = \varepsilon_\chi \Lambda(1-s,\bar{\chi})$ with $\Lambda(s,\chi) = (5/\pi)^{(s+1)/2}\Gamma(\frac{s+1}{2})L(s,\chi)$. Substituting (1) into $\Lambda_f(s) = (5/\pi)^{(s+1)/2}\Gamma(\frac{s+1}{2})f(s)$ and demanding $\Lambda_f(s) = \Lambda_f(1-s)$, the coefficient of $\Lambda(s,\chi)$ matches precisely when

$$(1 - i\kappa)\,\varepsilon_\chi \;=\; 1 + i\kappa, \qquad \text{i.e.} \qquad \kappa = \tan\big(\tfrac{1}{2}\arg\varepsilon_\chi\big), \tag{2}$$

and the coefficient of $\Lambda(s,\bar\chi)$ then matches automatically because $\varepsilon_{\bar\chi}=\overline{\varepsilon_\chi}$. Note that (1) is self-reciprocal while the equation for $L(s,\chi)$ alone exchanges χ and $\bar\chi$; that exchange is exactly what (1) is built to undo. Evaluating, $\tan\arg\varepsilon_\chi=\sin 36^\circ/\sin 72^\circ=1/\varphi$ with φ the golden ratio, whence the closed form

$$\kappa \;=\; \frac{\sqrt{10-2\sqrt5}-2}{\sqrt5-1} \;=\; 0.28407904384041229603\ldots$$

Our κ is computed as $\tan(\frac12\arg\varepsilon_\chi)$ from the Gauss sum, not imported; it agrees with the closed form to 3.9×10^{-62}, and the residual of (2) is exactly 0 at the working precision. The Dirichlet coefficients are $(a_1,\ldots,a_5)=(1,\kappa,-\kappa,-1,0)$, real and periodic but *not* multiplicative: $a_2a_3=-\kappa^2$ while $a_6=a_1=1$.

Functional equation, measured [T2]. At twenty reproducibly chosen points avoiding the critical line, the real axis and the gamma poles, the scale-aware relative residual of $\Lambda_f(s)=\Lambda_f(1-s)$ is at worst 6.951×10^{-58} at 60 working digits and 8.092×10^{-99} at 100; it tracks the precision, which is the check that it is a precision floor and not a defect. The relative form is necessary: one of the twenty points has $|\Lambda_f|\approx2\times10^{-31}$, where an absolute residual would be meaningless.

Strip [T1]. For $\sigma\ge2$, $|f(s)-1|\le\sum_{n\ge2}|a_n|n^{-\sigma}\le\zeta(2)-1=0.644934066848\ldots<1$. Sharpening the same elementary estimate with the actual $|a_n|$ gives the abscissa σ_0 with $\sum_{n\ge2}|a_n|n^{-\sigma_0}=1$, namely $\sigma_0=1.395136158235$, so every zero of Λ_f lies in $1-\sigma_0<\operatorname{Re}s<\sigma_0$. The strip is wide — this function has zeros with $\operatorname{Re}s>1$ — and it is what makes the census contour of §3 meaningful without an arbitrary cutoff.

Coefficients of $-f'/f$ [T1] and [T2]. In $\operatorname{Re}s>2$ set $-f'(s)/f(s)=\sum_{n\ge2}b_nn^{-s}$, so that $b_n=a_n\log n-\sum_{d|n,\,2\le d<n}b_da_{n/d}$ with $b_1=0$. Exactly, $b_2=\kappa\log2$, $b_3=-\kappa\log3$, and

$$b_6 \;=\; (1+\kappa^2)\log6 \;=\; 1.9363560766210387863,$$

nonzero at a non-prime-power. For ζ the set of $n<50$ that are not prime powers and have $b_n\neq0$ is empty; here it has 15 members. The missing Euler product is therefore visible in the coefficients the detector consumes, not only in the zeros. A second computation of b_n through the Dirichlet inverse of (a_n) agrees with the recursion to 1.245×10^{-60} for $n\le200$.

2 The window form

Let g be real and supported in $[-L,L]$, and set

$$G(z)=\int g(u)e^{zu}\,du,\qquad h(v)=\int g(u+v)g(u)\,du,\qquad H(z)=G(z)G(-z),$$

so h is supported in $[-2L,2L]$ and $H(it)=|G(it)|^2$. The spectral form runs over the zeros of the *entire completed* function, $Q_f(g)=\sum_\rho\operatorname{mult}(\rho)H(\rho-\frac12)$, and all reported values are normalised as $Q_f(g)/\|g\|_2^2$. Moving the contour as in the companion paper's §4–5, with the conductor 5 and the odd archimedean factor $\psi(\frac34+\frac{it}{2})$ in place of the ζ data, gives

$$Q_f(g) \;=\; h(0)\log\frac{5}{\pi} \;+\; \frac{1}{2\pi}\int_{\mathbb R}|G(it)|^2\operatorname{Re}\psi\Big(\tfrac34+\tfrac{it}{2}\Big)\,dt \;-\; 2\sum_{2\le n\le e^{2L}}\frac{b_n\,h(\log n)}{\sqrt n}. \tag{3}$$

There is **no pole term**: Λ_f is entire, the trivial zeros of f at $s=-1,-3,\ldots$ cancelling the poles of the gamma factor. The arithmetic sum is **finite** because h vanishes beyond $2L$; at $L=2.6$

the cutoff is $n \le 181$ with 122 nonzero terms, and the bound is decided in integer arithmetic and then re-tested as $\log n < 2L$ at full precision, never by a binary float. The formula is of the Weil–Yoshida–Bombieri type [9, 2]; what is specific to f is that b_n is not supported on prime powers.

Zero contributions. An on-line conjugate pair $\{\frac{1}{2} \pm i\gamma\}$ contributes $2H(i\gamma) = 2|G(i\gamma)|^2 \ge 0$. The complete orbit of a non-real off-line zero, $O = \{\beta \pm i\gamma,\, 1-\beta \pm i\gamma\}$, contributes, with $\delta = \beta - \frac{1}{2}$,

$$Q_O(g) \;=\; 4\,\mathrm{Re}(A\overline{B}) \;=\; 4(|P|^2 - |S|^2), \qquad A = G(\delta + i\gamma),\; B = G(-\delta + i\gamma), \tag{4}$$

where $P = \int g\cosh(\delta u)e^{i\gamma u}du$ and $S = \int g\sinh(\delta u)e^{i\gamma u}du$. This is a *signed* rank-two form, with a positive cosh channel and a negative sinh channel, and it is not $|G(\delta+i\gamma)|^2$. As $\delta \to 0$ it tends to $4H(i\gamma)$, the contribution of *two double* on-line zeros, so multiplicity is preserved: four zeros in, four out.

Gate A [T2-N]. The arithmetic side (3) is compared with two independent evaluations and three exact-algebra controls. The contour route writes $Q_f(g) = (2/\pi)\,\mathrm{Re}\int_0^\infty (\Lambda_f'/\Lambda_f)(\sigma_c + it)H(\sigma_c - \frac{1}{2} + it)\,dt$ at $\sigma_c = 1.42$, to the right of every zero, with Λ_f'/Λ_f built from f and f' directly and *never* from the b_n; the zero route sums over the census of §3.

Check	Discrepancy	Declared budget	Tier
arithmetic vs. logarithmic-derivative contour	1.73×10^{-13}	1.316×10^{-6}	T2-N
arithmetic vs. zero-side sum over the census	5.65×10^{-11}	7.13×10^{-5}	T2-N
control: negate the selected quartet (sign flip)	1.83×10^{-62}	exact algebra	T1
control: omit b_6 at a window containing $n = 6$	1.94×10^{-61}	exact algebra	T1
control: flip each arithmetic term in turn	3.11×10^{-61}	exact algebra	T1

The quartet control negates the selected orbit's contribution on the zero side alone and checks that the discrepancy moves by exactly twice it; with $Q_{O_1}(g) = 8.720121523693614 \times 10^{-11}$ on that low-frequency test function, the predicted and observed changes agree to 1.83×10^{-62}. The b_6 control predicts a change of $+2b_6 h(\log 6)/\sqrt{6} = -0.379255712825982$ from $b_6 = 1.93635607662104$ and $h(\log 6) = -0.239879170384883$, and observes it to 1.94×10^{-61}. The term-level sign flip negates each of the archimedean, conductor and arithmetic terms in turn; the conductor row is the control that would catch a wrong conductor, and it reproduces $-2\log 5 = -3.21887582486820$ for a unit-normalised test function in every sector and window. **Classification: PASS-N** — converged numerical agreement, explicitly uncertified, because the quadrature in the contour route and the zero density in the zero-side tail are a library estimate and a measured quantity, not rigorous bounds. The three controls, which are what would catch a wrong factor, a wrong conductor or a dropped term, are exact.

3 The zero census to $T = 200$

Counting by the argument principle on the strip $-0.4 < \mathrm{Re}\, s < 1.4$, whose vertical edges are provably zero-free by §1, gives 130 zeros with $0 < \mathrm{Im}\, s < 200$ against a Riemann–von Mangoldt main term of 129.548; the worst deviation of a winding number from an integer over the whole ladder is 1.32×10^{-23}. Sign changes of the Hardy-type Z give 122 critical-line ordinates, and four off-line zeros with $\mathrm{Re}\, s > \frac{1}{2}$ are located by box subdivision and Newton, each carrying local winding number exactly 1 on a circle of radius 0.02, hence simple. The reconciliation

$$122 \;+\; 2 \cdot 4 \;=\; 130$$

is exact, and it is also what closes the band $\frac{1}{2} < \operatorname{Re} s < 0.55$ that the located search deliberately avoids: an unlocated zero there would break it.

	β	γ	$\delta = \beta - \frac{1}{2}$	$\lvert f(\rho)\rvert$	winding
O_1	0.808517182456637386	85.6993484853775922	0.308517182456637386	1.75×10^{-60}	1
O_2	0.650830080609737082	114.163342730756981	0.150830080609737082	3.21×10^{-60}	1
O_3	0.574356050450805991	166.479305913168156	0.0743560504508059907	5.99×10^{-60}	1
O_4	0.724257694626809780	176.702461242855825	0.224257694626809780	2.99×10^{-60}	1

The residuals are at the working precision of the census run (60 digits); at 140 digits the refined O_1 has $|f(\rho_1)| = 8.2 \times 10^{-141}$ with Newton step 6.5×10^{-141} and is stable to 3.17×10^{-106} between 100 and 140 digits. These are roots to the precision floor, not approximate ones.

Which are cross-checked [T2]. One of the four, and only against a secondary source. O_1 was found from a decimal seed supplied with the task; Newton moved that seed by 7.8×10^{-15}, so the seed's digits are correct to about fourteen places. The provenance of that decimal is a secondary report of the literature computations of Spira [7] and Balanzario and Sánchez-Ortiz [1]; **neither paper was read** — the AMS PDF of the latter returned HTTP 403 from the machine used — and no numerical value from either appears in this note. O_2, O_3 and O_4 are **not** cross-checked against anything: they are this computation's own. Bombieri and Ghosh [3] is the standard modern reference for the analytic theory of f; it was not read either, and no result or constant of theirs is used — in particular the counting function above is the computed argument-principle count, with the main term quoted only as a diagnostic.

4 The certified witnesses

[T2-C] Two test functions are frozen, with every coefficient stored as an exact 30-digit decimal string. Both are odd, both are in the span of a low block together with a block resonant at γ_1, and the Gram matrix of the normalised basis is the identity exactly (deviation 4.83×10^{-60} and 7.4×10^{-60}), so the generalised eigenproblem is the ordinary one and no direction was regularised or discarded. The coefficients of g_* are in Appendix A.

	g_*	$g_\sharp$
L	2.0	1.8
sector	odd	odd
N	37	57
arithmetic cutoff	$n \le 56$, 39 nonzero terms	$n \le 38$, 26 nonzero terms
$Q_f/\lVert g\rVert_2^2$	-0.679071346441480590986	$-1.68123895523670103 \times 10^{-16}$
ball radius	1.39×10^{-116}	8.15×10^{-177}
strictly negative	yes	yes

Why the certificate is complete. The enclosure is of the *whole* quantity, not of an approximation to it, and there is nothing outside the ball:

- **The arithmetic sum is finite and complete.** h vanishes beyond $2L$, so (3) has 39 terms at $L = 2$ and no tail is dropped.
- **No quadrature.** The archimedean term is the closed form in digamma, the Hurwitz zeta and the Lerch transcendent; Arb encloses all three.

- **No truncation and no spectral input.** $Q_f(g_*)$ is computed from the arithmetic side alone. It does not know where any zero is, so the census plays no part in it and the sign does not depend on the census being right.
- **The test function carries no error.** Its coefficients are exact finite decimals; the ball encloses Q_f for *this* g_*, not for a rounding of some other one.
- **Independent implementation.** The certifying code re-implements the cross-correlations, κ, the b_n and the archimedean closed form against Arb, sharing no code with the `mpmath` assembly; the two agree to all 44 digits `mpmath` carries, so the certificate is simultaneously a cross-check of both.

The only inputs are Arb's enclosures and the derivation of (3). κ enters through square roots only. This is [T2-C] and not a formal verification.

5 The decomposition

[T2-C] Both terms below are finite sums and both are enclosed. At g_*:

$$\underbrace{-0.679071346441480590986}_{Q_f(g_*)} = \underbrace{-0.985198326439305481794}_{Q_{O_1}(g_*)} + \underbrace{+0.306126979997824890807}_{Q_{\mathrm{rest}}(g_*)}, \tag{5}$$

with ball radii 1.39×10^{-116}, 4.06×10^{-117} and 1.63×10^{-116}, and additive reconciliation residual 2.99×10^{-116}. Here $Q_{\mathrm{rest}} = Q_f - Q_{O_1}$ is evaluated by subtracting the exact rank-two orbit form (4) from the arithmetic side, so *every* remaining zero — including every off-line zero above the census height and every zero never located — is included implicitly. Nothing is replaced by a finite zero list.

The selected quartet therefore carries more than the whole of the negativity, and the rest of the spectrum returns 31.07% of it. At $g_\sharp$ the same decomposition reads $-6.45917960933013 \times 10^{-14} + 6.44236721977777 \times 10^{-14}$: near the onset the cancellation is 99.7397%, which is why the witness there is of order 10^{-16} rather than 10^{-14}.

The remaining-zero form is positive on every space searched [T2-N]. $\lambda_{\min}$ of the restriction of Q_{rest} to the level-2 space, alongside $\lambda_{\min}(Q_f)$:

L	$\lambda_{\min}(Q_f)$, even	$\lambda_{\min}(Q_{\mathrm{rest}})$, even	$\lambda_{\min}(Q_f)$, odd	$\lambda_{\min}(Q_{\mathrm{rest}})$, odd
1.0	$4.034840454024 \times 10^{-5}$	$4.03625203581 \times 10^{-5}$	0.01356156739650	0.0132281100537
1.2	$8.780065301754 \times 10^{-9}$	$8.46979844459 \times 10^{-9}$	$9.702616267333 \times 10^{-6}$	$9.54583799747 \times 10^{-6}$
1.4	$5.087177960475 \times 10^{-14}$	$4.64125898422 \times 10^{-14}$	$6.538829304189 \times 10^{-11}$	$6.63381728826 \times 10^{-11}$
1.6	$2.549827775205 \times 10^{-19}$	$2.25086996550 \times 10^{-19}$	$2.153851664770 \times 10^{-16}$	$2.21332806419 \times 10^{-16}$
1.8	$7.580609464352 \times 10^{-24}$	$7.01854463036 \times 10^{-24}$	$1.007541951495 \times 10^{-20}$	$1.08891994498 \times 10^{-20}$
2.0	-0.4896520626161	$2.02946591361 \times 10^{-27}$	-0.6790713464415	$1.20623694960 \times 10^{-24}$
2.2	-1.219875374653	$1.22467880540 \times 10^{-30}$	-1.178435305399	$4.87721021342 \times 10^{-28}$
2.4	-1.503014486764	$1.81134931584 \times 10^{-33}$	-1.590268954821	$4.26137491134 \times 10^{-31}$
2.6	-2.137655567743	$8.47226605297 \times 10^{-36}$	-2.005507257484	$8.13613127375 \times 10^{-34}$

All 18 rows are positive. Once Q_f is macroscopically negative the remaining-zero form does not merely stay positive: it collapses towards zero, reaching 8.47×10^{-36} at $L = 2.6$ while Q_f is at -2.14. Removing O_1 from the frozen witness's space and re-optimising gives $\lambda_{\min} = +1.20623694960 \times 10^{-24}$; no other negative direction is found there, which is a statement about that space and not about positivity.

The consequence, stated exactly. The remaining-zero theorem of the companion paper assumes a remaining-zero form that is nonnegative, and states that assumption as *sufficient*, not necessary, for its conclusion. On this object that hypothesis **holds on every space searched**, so the theorem's conclusion applies here; and because its proof uses the hypothesis only through the single inequality $Q_{\mathrm{rest}} \geq 0$, nothing else about where the other zeros lie is needed. The masking contribution is 31.07% in the witness direction and does *not* reverse the sign. What is *not* established is that a positive finite restriction establishes the whole-window hypothesis: controlling the infinite-dimensional complement and its coupling to the block would be needed, and a positive Galerkin matrix supplies neither.

And the confound, once more. Q_{rest} is built from the on-line zeros of an object with no Euler product, and it is positive; that is the strongest evidence *internal to this object* that the negativity is attributable to the off-line quartets, and it is not a proof.

6 Onset

[T2-N] The scan runs $L = 1.0, 1.2, \ldots, 2.6$ in both parity sectors at three declared basis levels; enrichment can only lower $\lambda_{\min}$, and the levels are reported as enrichments, never folded into one another. Over all 54 rows the Rayleigh quotient of the returned eigenvector agrees with the eigenvalue to 5.76×10^{-60}, so the eigensolver is confirmed by direct evaluation. At the deepest level, $\lambda_{\min}$ falls by a factor 1.459×10^{17} from $L = 1.0$ to $L = 1.6$ while remaining positive, and then changes sign:

L	even, lvl 1	even, lvl 2	even, lvl 3	odd, lvl 1	odd, lvl 2
1.6	$1.97261814306 \times 10^{-13}$	$2.54982777520 \times 10^{-19}$	$2.66141166363 \times 10^{-22}$	$2.91871108168 \times 10^{-11}$	2.15385166477×10
1.8	$2.91410423267 \times 10^{-15}$	$7.58060946435 \times 10^{-24}$	$-2.66180729007 \times 10^{-21}$	$1.19062366907 \times 10^{-13}$	1.00754195149×10
2.0	-0.188728484539	-0.489652062616	-0.547390086184	-0.658652328928	-0.679071346441

Twenty-six of the 54 rows are negative. The crossing is not gradual: the form sits on a knife edge through most of the scanned range and then falls off it, from -1.68×10^{-16} at $L = 1.8$ to -0.71 at $L = 2.0$.

Which sector detects first. The scan does not resolve this. Both sectors are positive at $L = 1.6$ and negative at $L = 1.8$ at level 3, so on this grid they cross together; the odd value is deeper by a factor 6.316×10^4 at $L = 1.8$ and by 0.16 at $L = 2.0$. Since $K_{\mathrm{even}}(2.0, \gamma_1)/K_{\mathrm{odd}}(2.0, \gamma_1) = 1.00634$ at this height, the two sectors have essentially the same isolated sensitivity, and **no structural explanation is offered** for the depth difference.

Two comparisons. Both are computed by the verifier from the Phase-A values of (δ_1, γ_1), not quoted from elsewhere.

(i) Bandwidth scale. With $L_{\mathrm{pred}} = \frac{1}{2}\log(5\gamma_1/2\pi) = 2.11120303$, and the measured onset bracketed below by the certified negative witness at $L = 1.8$ and above by the macroscopic one at $L = 2.0$,

$$\frac{L_{\mathrm{pred}}}{1.8} = 1.17289, \qquad \frac{L_{\mathrm{pred}}}{2.0} = 1.0556.$$

This is a reference scale, not a predicted onset: its provenance is a bandwidth heuristic that was not verified, and the bracket is one-sided in the sense of non-claim 5.

(ii) The δ^2 coefficient. In the odd sector the exact extremal constant is $K_{\mathrm{odd}}(L, \gamma) = \int_{-L}^{L} u^2 \cos^2(\gamma u)\, du = 2.6582355735990895$ at $L = 2$, $\gamma = \gamma_1$, and the companion paper's δ^2 coefficient gives

$$4K_{\mathrm{odd}}(2.0, \gamma_1)\, \delta_1^2 \;=\; 1.01207377, \qquad \text{against} \qquad |Q_{O_1}(g_*)| = 0.985198326439305,$$

a ratio of 1.02728. Measured instead against the actual drop from the collapsed configuration, $Q_{O_1}(0) - Q_{O_1}(\delta_1) = 0.986028353$ with $Q_{O_1}(0) = 4|\hat{f}(\gamma_1)|^2 = 0.000830026353$, the ratio is 1.02641.

Is that inside what the remainder allows? Yes, and *so comfortably that the comparison is not a sharp test of the bound.* At $L\delta_1 = 0.617034$ the companion paper's δ^4 allowance is $(\frac{2}{5} + \frac{16}{3}e^{2L\delta})L^5\delta^4 = 5.42744$, while the observed discrepancy $|4K\delta^2 - \text{drop}|$ is 0.02605 — smaller by a factor 208.4. The 2.6% agreement is therefore *better* than the bound requires, by more than two orders of magnitude, and should not be read as confirming the bound's constant; the bound is simply very loose at this displacement. $L\delta_1 = 0.617$ is far outside any small-displacement regime, and no small-δ law is used anywhere in §4 or §5.

7 What this rules out, and what it does not

Ruled out. Any claim that finite-window Weil positivity follows from a functional equation of ζ's shape, together with entireness of the completion and a zero-containing strip. The Davenport–Heilbronn function has all three — (1) is self-reciprocal, Λ_f is entire, and every zero lies in $|\operatorname{Re} s - \frac{1}{2}| < 0.895136$ — and its window form at $L = 2.0$ is negative, at -0.679071346441, now by a *certified example at a stated window* rather than by citation of the existence of off-line zeros. The same example rules out the weaker claim that a positive finite Galerkin restriction of a remaining-zero form establishes a whole-window hypothesis: here $\lambda_{\min}(Q_{\text{rest}}) > 0$ on every space searched while Q_f is macroscopically negative at five of the nine windows.

Not ruled out, and not claimed. Nothing about ζ: no statement here has a ζ analogue, and the whole point of the control is that it runs where positivity is known to fail. Nothing that separates the two properties of f, off-line zeros and the absent Euler product, which is the confound of non-claim 3. Nothing about other L-functions, other conductors, other bases, or other windows than the nine scanned. Nothing about positivity at $L \le 1.6$: those are non-detections. And nothing certified about the *onset*, only about the two witnesses — certifying the crossing would need a certified lower bound at $L = 1.6$ over a whole space, which is a different and harder object than a certified value at a point.

Open, in four sentences. Whether $\lambda_{\min}(Q_{\text{rest}})$, already at 8.47×10^{-36} at $L = 2.6$, stays positive as the basis grows, or whether its near-null direction is the leading edge of a genuine negative one. Whether the onset can be bracketed by certified bounds at both ends rather than by a certified negative and a non-detection. Whether the census can be made rigorously complete, the winding numbers being integers to 1.32×10^{-23} by adaptive sampling rather than by interval arithmetic. And whether an off-line quartet of small δ at larger height — O_3 has $\delta = 0.0743560504508059907$ at $\gamma = 166.479305913168156$ — is seen earlier or later than O_1 by a basis resonant at *its* height, which the present scan, resonant only at γ_1, cannot say.

A The frozen witness g_*

$L = 2.0$, odd sector, $N = 37$. The test function is $g_*(u) = \sum_m c_m \phi_{k_m}(u)$ with $\phi_k(u) = \sin((k+1)\pi u/L)/\sqrt{L}$ on $[-L, L]$ and zero outside, so that $\|g_*\|_2^2 = \sum_m c_m^2$. The index set is the low block $k = 0, \dots, 15$ together with the resonant block $k = 44, \dots, 64$, chosen so that $w_k = (k+1)\pi/L$ brackets γ_1. The coefficients are exact finite decimals, read back from the certificate as strings; the table below is generated mechanically from `certificates/witness.json` and is regenerated and compared by the verifier.

m	k_m	c_m	m	k_m	c_m
0	0	$-0.00260883678800775017628484317 74$	19	47	0.016079408349982117195326094 1976
1	1	0.005936729798368368678676143712306	20	48	0.004989867805626867671955611514 28
2	2	$-0.018553406278023102392126189 6105$	21	49	0.000027705392952165417781488258 6319
3	3	0.005448463051510521275799587393 66	22	50	$-0.022217926726101455164643676 4475$
4	4	$-0.015639601737394463596695710 9136$	23	51	0.039586845834277726764109564134
5	5	$-0.010978316465511239065578918 8025$	24	52	0.094230299910623825585872804 7973
6	6	$-0.012590775431328615862660174 7424$	25	53	0.804187411949596545617290494929
7	7	$-0.014462804894923709347672683 6875$	26	54	$-0.559992992803302061132173856 712$
8	8	$-0.014634391489449079888172622 8996$	27	55	0.137708858583258544659438566561
9	9	0.010389852439123943709897434 9874	28	56	0.022177672667469742476663478 5707
10	10	0.004486077899024563022124890579 99	29	57	0.015045495540320733808379466 7482
11	11	0.024197386383040060930974723 9095	30	58	$-0.030309802449998409598133768 0134$
12	12	$-0.014154558071251219994613019 8503$	31	59	0.027392761708372298186338956 7813
13	13	0.008382913355790691852662303 6363	32	60	0.001127864409163977755739920517 311
14	14	0.003064724536680788545001660 29552	33	61	$-0.022628980112189728180711178 8614$
15	15	0.031971898922306640455049590 7318	34	62	0.002118602697675533696146791448 69
16	44	$-0.031189330491393555429287951 3548$	35	63	0.019589441263546411704813632605
17	45	$-0.005540657344275701976102674 00465$	36	64	0.030112760581951388414172362309
18	46	$-0.030419110162133506994324226 6561$			

The mass is concentrated on $k = 53$ and $k = 54$, whose frequencies bracket $\gamma_1 = 85.699348485377592$; the witness has normalised overlap 0.9513529401 with $u\cos(\gamma_1 u)$, the extremiser of the isolated derivative-evaluation functional, while the ground state of Q_{rest} is orthogonal to it to 9.07×10^{-12}.

B Retraction log

Six corrections were made during this work. They are reproduced from `RETRACTIONS.md`, in order.

R1. Two-sided zero-count normalisation. The Riemann–von Mangoldt main term was written $(T/\pi)\log(5T/2\pi e)$, which counts $|\operatorname{Im}\rho| < T$, while the census counts $0 < \operatorname{Im}\rho < T$. Corrected to $(T/2\pi)\log(5T/2\pi e)$. The near-exact factor of two between the winding numbers and the "main term" is what exposed it. Diagnostic only: the counts come from the argument principle, and no reported number changed.

R2. Off-line search contour too close to the critical line. The first implementation subdivided from $\operatorname{Re} s \geq 0.505$, which passes within 0.005 of the on-line zeros; $\arg f$ then swings by nearly π between samples and the adaptive refinement recursed to its depth limit on every band. Corrected by searching at $\operatorname{Re} s \geq 0.55$ and closing the uncovered band by the reconciliation of §3, which is a stronger check and cheap. Method only; no reported number changed.

R3. Division by zero at a removable singularity. The closed form for $\int \cos(wu)e^{zu}du$ has a $0/0$ at $z = \pm iw$. A unit test evaluating there caught it; an explicit limit branch was added. The condition never arose in a reported computation — H is evaluated at zero ordinates, not at basis frequencies — and the branch exists so the test can cover the case.

R4. The first independent archimedean route was too weak to be a check. It evaluated the frequency-side integral by quadrature; the integrand oscillates with period π/L and the result was accurate only to about 10^{-6}, which cannot distinguish a correct assembly from one with a small error. Replaced by Gauss's integral for ψ, giving a one-dimensional non-oscillatory integral with an exact tail. The two routes now agree to 6.225×10^{-21} against an analytic bound of 1.66×10^{-20} on the piece deliberately omitted near the origin.

R5. A gate threshold that could not be met. Phase B required agreement below 10^{-40} while its own declared budget was 10^{-20}. A gate that cannot pass is not a gate; the threshold was tied to the route's own budget. No reported value changed.

R6. Decimal window lengths parsed at the ambient precision. This is the one that mattered. `mpmath`'s `mpf` rounds a decimal string to the *current* working precision and freezes it. A window length built as `mpf('1.8')` before the precision was raised carried only 15 digits, and the subsequent precision ladder at 60, 100 and 160 digits then evaluated the form at that same slightly-wrong L three times, agreed with itself to 10^{-85}, and reported a value wrong in the *thirteenth* significant digit. A ladder that agrees with itself to 85 digits is not evidence that the number is right. It was caught only because two scripts froze the same witness independently and disagreed in the 13th digit — the ladder itself could not have caught it, and would not have, at any precision. Every decimal constant that feeds the form now goes through a parser fixed at 220 digits, and a guard test fails if any of them is reconstructed at the ambient precision. Blast radius: the certified witness sits at $L = 2.0$, exactly representable in binary, and is unaffected; the window scan ran at a consistent precision throughout and is unaffected; the $L = 1.8$ witness was recomputed, and its sign and first twelve digits are unchanged.

C The two literature witnesses

Two contemporaneous documents report negative Davenport–Heilbronn window witnesses. **Only their landing pages were read**; neither PDF was retrieved, and nothing below is a reading of their arguments.

The first is T. Ziad, *Localized Constrained Weil Forms: Near-Degeneracy and a Davenport–Heilbronn First Kernel*, Zenodo, doi:10.5281/zenodo.23145455, 7 October 2026, which reports "an exactly constrained negative witness at $a = 9/5$" for odd, *constrained* test functions, and a first loss-of-positivity scale strictly between $\frac{1}{2}\log 2$ and $9/5$. The second is a Zenodo series on cross-energy of Dirichlet partial sums, from which the sentence "positive up to a window $L \approx 3.6$ … $\lambda_{\min} = -0.73$ at $L = 4$, with a minimizer concentrated on its off-line zero $0.8085 + 85.70i$" was surfaced.

The two window numbers differ by exactly a factor of two. In the convention used here, g is supported in $[-L, L]$ and its autocorrelation h in $[-2L, 2L]$, so a factor of two is what one would expect between a document measuring the support of g and one measuring the support of h. **That is a hypothesis, not a finding**: nothing here establishes which convention either document uses, and until it is settled neither agreement nor disagreement with them can be asserted. Our level-3 value at $L = 2.0$, -0.7101755122300, is in the same region as the quoted -0.73 under the hypothesis, and is an upper bound for the true minimum because it is computed on a finite subspace — but that observation is worth exactly as much as the convention hypothesis it rests on.

Nothing in this note was fitted to either value. The scanned window range was fixed before either document was located, no threshold or basis was tuned, and no novelty is claimed: at least two contemporaneous efforts address the same control, one of them with its own interval certificate.

D Reproduction

Code and data: `https://github.com/innerlightr-wq/weil-window-control`, directory `extensions/davenport`. Python 3.14.7 (CPython), `mpmath` 1.3.0, `python-flint` 0.9.0 (Arb). No `numpy`, no `scipy`, and no `float64` on any path reaching a reported value. All conventions, precisions, basis levels, windows and thresholds are in `src/config.py`; no randomness is used.

Result	Script	Data
§1 function, κ, functional equation	`src/phaseA.py`	`results/phaseA.json`
§1 b_n, §2 form	`src/phaseB.py`	`data/dh_coefficients.csv`
§2 orbit conventions	`tests/test_orbits.py`	—
§3 census	`src/census.py`	`data/dh_zeros.csv`
§2 Gate A	`src/phaseC.py`, `src/gate_signflip.py`	`data/gate_checks.csv`, `data/gate_signflip.`
§6 scan	`src/phaseD.py`	`data/window_scan.csv`
§4 freeze	`src/freeze.py`	`certificates/witness*.json`
§4 certificates	`src/certify.py`	`certificates/ball_certificate*.txt`
§5 remaining form	`src/phaseE.py`	`data/remaining_form.csv`
§5 masking	`src/phaseF.py`	`data/masking.csv`
§6 K, perturbation	`src/phaseG.py`	`data/sensitivity.csv`

The two certificates are reproduced by

```
.venv/bin/python src/certify.py 400
.venv/bin/python src/certify.py 600 certificates/witness_smallest_L.json \
                                    ball_certificate_L1.8.txt
```

`make verify`, in the note directory, runs `note/verify_note_numbers.py`: it rebuilds every quantity printed above from the data files, scans this document for numeric literals, fails if any literal is not derived or is printed to more significant digits than its source carries, regenerates Appendix A's table from `certificates/witness.json` and compares it, and cross-checks the Gate A values against `REPORT.md`. The full suite is `./run_suite.sh`.

Licence

As the parent repository: code MIT, this note and the data CC BY 4.0. CC BY 4.0 requires attribution; if you use the note, the certificates or the data, please cite this note by its DOI, 10.5281/zenodo.23608288, and the parent paper [5] separately.

Acknowledgments

AI assistance was used for algebra checking, numerical verification, adversarial review, literature organization, and manuscript drafting. All conceptual development, research direction, interpretation, and final responsibility for the content belong to the author.

References

[1] E. P. Balanzario and J. Sánchez-Ortiz, *Zeros of the Davenport–Heilbronn counterexample*, Math. Comp. **76** (2007), no. 260, 2045–2049.

[2] E. Bombieri, *Remarks on Weil's quadratic functional in the theory of prime numbers, I*, Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei (9) Mat. Appl. **11** (2000), no. 3, 183–233.

[3] E. Bombieri and A. Ghosh, *Around the Davenport–Heilbronn function*, Russian Math. Surveys **66** (2011), no. 2, 221–270; Uspekhi Mat. Nauk **66** (2011), no. 2, 15–66.

[4] H. Davenport and H. Heilbronn, *On the zeros of certain Dirichlet series*, J. London Math. Soc. **11** (1936), 181–185; **11** (1936), 307–312.

[5] E. De Jesús, *A δ^2–L^3 Sensitivity Law for Finite-Window Weil Positivity, with a Function-Field Control of the Connes–van Suijlekom Pipeline*, Zenodo, 2026. doi:10.5281/zenodo.23250932.

[6] F. Johansson, *Arb: Efficient arbitrary-precision midpoint-radius interval arithmetic*, IEEE Trans. Comput. **66** (2017), no. 8, 1281–1292.

[7] R. Spira, *Some zeros of the Titchmarsh counterexample*, Math. Comp. **63** (1994), no. 208, 747–748.

[8] E. C. Titchmarsh, *The Theory of the Riemann Zeta-Function*, 2nd ed., revised by D. R. Heath-Brown, Oxford University Press, 1986, §10.25.

[9] H. Yoshida, *On Hermitian forms attached to zeta functions*, in: N. Kurokawa and T. Sunada (eds.), *Zeta Functions in Geometry*, Adv. Stud. Pure Math. **21**, Math. Soc. Japan, Kinokuniya, Tokyo, 1992, 281–325.