Certified Weil Positivity at $L = 4/5$: Two-Sided Bounds, Parity Separation, and Cutoff Diagnostics

Elias De Jesús
Independent Researcher
ORCID 0009-0007-0190-9143
dejesuselias10@gmail.com

November 2026 — Technical Note (preprint), version 1.2.0

DRAFT — certification correction in progress

Awaiting author review. Not for citation as a finished result.

Status: draft, certification correction in progress.
The compact time-domain evaluator used in earlier versions of this note omitted an *outer quadrature truncation allowance*: it bounded the integrable head of its archimedean integral but added nothing for the truncation error of the panelled Gauss–Legendre rule itself. Interval arithmetic does not supply that term — it encloses the nodes, the weights and the rounding, not the difference between a quadrature sum and the integral. The radii that route published were therefore incomplete.

Affected, and currently *not* certified: the improved upper bound 1.9014×10^{-17} reported in versions 1.0.x, the quantities $q(f_{200})$ and $q(f_{400})$ it came from, and the discarded energies $e_T = q - r_T$ with their ratios. These are retained below as *measurements* and are labelled as such.

Not affected: the *lower-bound* certificate $Q(f) \geq 1.41 \times 10^{-17}\|f\|_2^2$ and the odd-sector bound 2.3×10^{-17}. Those come from a different routine, whose interior quadrature carries a certified Bernstein-ellipse remainder folded into the matrix entries exactly once and re-derived by its verifier. The upper bound quoted in this version comes from a closed-form route that performs no quadrature at all. See Sections 4.1 and 4.5, and `RETRACTIONS.md` R14.

The repair is not finished: a rigorous outer quadrature remainder is still open (Section 4.5). This document is a public research draft and is awaiting author review; nothing here has been deposited.

Abstract

For the Weil explicit-formula quadratic form restricted to test functions supported in a compact window $[-L, L]$ with $L = 4/5$, this note reports a computer-assisted two-sided bound on its spectral infimum $\lambda_Q(4/5)$ in one fixed normalization:

$$1.41 \times 10^{-17} \;\leq\; \lambda_Q(4/5) \;\leq\; 1.674 \times 10^{-17}.$$

The lower bound is obtained by a frequency-cutoff comparison at $T = 400$, reduced to a 440-dimensional Legendre block whose positive definiteness is verified by a preconditioned congruence in Arb ball arithmetic, together with a super-exponentially small complement estimate and a Schur-type coupling bound; both parity sectors are certified. The upper bound is the Rayleigh quotient of the *original* form at one explicit even rational test function, evaluated in closed form through the digamma, Lerch Φ and Hurwitz ζ functions, a route that uses no quadrature at all, so there is no truncation term to bound. A candidate obtained by a time-domain route is reported but *not* certified: its outer quadrature remainder is unbounded (Section 4.5), and every value depending on it is withdrawn pending that bound. A separate certificate gives $\lambda_{Q,\mathrm{odd}} \geq 2.3 \times 10^{-17}$, which exceeds the certified upper bound and therefore shows that the even infimum determines λ_Q. Diagnostics at cutoffs

$T \in \{200, 400, 800\}$ bound the gain available from a larger cutoff. All constants are published as midpoint–radius enclosures with a cold verifier. This is a single-window computation: it establishes no arithmetic estimate, nothing about the location of any zeta zero, and nothing about the Riemann hypothesis.

1 Introduction

1.1 The object

Weil's explicit formula pairs a test function against the zeros of ζ, the pole at $s = 1$, the archimedean place and the primes. When the test function is required to be supported in a fixed compact window, the resulting quadratic form becomes a concrete object: a bounded-below Hermitian form on $L^2(-L, L)$ whose positivity *for every* L is equivalent to the Riemann hypothesis [8, 1], and whose positivity at one fixed L is not. Finite-window Weil positivity in this sense is the object studied in the spectral programs of Connes and van Suijlekom [4], Connes and Consani [2], and Connes, Consani and Moscovici [3], and quantitatively in Zhu [9], which supplies the finite-reduction strategy that the present computation follows.

The quantity computed here is the spectral infimum

$$\lambda_Q(L) \;=\; \inf_f \frac{Q_{\zeta,L}(f)}{\|f\|_2^2}$$

of that form at $L = 4/5$, in the normalization fixed in Section 2. Two things make a two-sided statement possible at this window and difficult at larger ones. An *upper* bound needs a single admissible test function and is unconditional. A *lower* bound needs the whole window, hence a reduction of an infinite-dimensional positivity question to a finite matrix plus certified tails; the cost of that reduction grows very fast in L, which is the content of Theorem 1.4 of [9].

1.2 What this note reports

1. A certified lower bound $Q(f) \geq 1.41 \times 10^{-17} \|f\|_2^2$ at $L = 4/5$, in both parity sectors and hence for arbitrary complex tests (Section 3).

2. A certified upper bound $\lambda_Q(4/5) \leq 1.674 \times 10^{-17}$ from one explicit even rational test function, with the *original* form evaluated in closed form (Section 4). A time-domain candidate is reported but withdrawn pending a quadrature bound (Section 4.5).

3. A certified parity separation: the odd sector satisfies $\lambda_{Q,\mathrm{odd}} \geq 2.3 \times 10^{-17}$, which is strictly above the upper bound, so the even infimum determines λ_Q (Section 5).

4. Cutoff diagnostics at $T \in \{200, 400, 800\}$ that measure what the comparison discards on fixed test functions, and a certified ceiling on what a larger cutoff can deliver (Section 6).

1.3 Relation to prior work, and what is not claimed

The finite-to-infinite reduction strategy — a scalar gate on the exterior of a frequency cutoff, a Legendre/Bessel representation of the interior, and super-exponentially small tails — is Theorem 1.1 of [9]. Zhu describes supplementary source code and certificates in [9, §17]; these were not obtained or executed in the present work, so his matrix step is neither re-run nor verified here. The reduction was instead rederived in the normalization of the present repository, and every constant below was computed from scratch. The bracket $8.9 \times 10^{-18} \leq \lambda_{\min}(0.8) \leq 2.27 \times 10^{-17}$ stated in the abstract of [9] was used only as an external comparison point and never as an input, an expected answer, or a tolerance. The parity structure is likewise not new, and at this window it is not merely the decoupling that is already published: [9, Thm. 6.2] certifies the

separation at support 1.6 — that is, at $L = 4/5$ — and does so with a far stronger odd-sector constant than the one obtained here, concluding in addition that the ground state is simple and even. What Section 5 contributes is an independent re-derivation of the ordering from this pipeline's own certified data, nothing structural.

The computation continues the implementation developed alongside our finite-window sensitivity study [10], whose repository supplies the zeta-window normalization of Section 2, the geometric-side assembly, the pole conventions, and the closed-form archimedean evaluator used as the validation reference in Section 4(a). The even/odd pole signs used here are the ones checked entrywise against that assembly in our real-zero parity study [11]. Neither the δ^2–L^3 sensitivity estimate of [10] nor the planted-real-zero analysis of [11] is an input to the lower bound proved here: the first concerns a counterfactual perturbation of the zero set, and the second a configuration that is nowhere assumed to exist. The finite-to-infinite reduction follows [9]; what is contributed here is its execution in certified arithmetic, together with the resulting bracket, the numerical parity separation of Section 5, and the cutoff diagnostics of Section 6.

What follows is deliberately narrow, and the following are *not* claimed. No arithmetic estimate is obtained: nothing here improves any bound on primes, on $\Lambda(n)$, on $\psi(x)$, or on a zero-free region. No statement is made about the location of any zero of ζ, and no range of zeros is verified. No new positivity framework is introduced; the framework is that of the cited sources, and the contribution is a certified computation inside it. The computation is at one window, $L = 4/5$, and gives nothing at any larger support — there is no propagation theorem here, and a certificate at one L does not provide one. Nothing here bears on the Riemann hypothesis. This is a technical note, not peer-reviewed research, and it is not claimed that anyone else uses or has used this work.

1.4 The same window, compared

Because [9] certifies at support 1.6, its results and the present ones describe the same object in the same normalization (its equations (2)–(3) are (1)–(2) below), so they can be put side by side. Doing so is the fairest summary of what this note adds and what it does not.

quantity at $L = 4/5$	[9], certified	this note
whole-window infimum	$[8.9 \times 10^{-18},\ 2.27 \times 10^{-17}]$	$[1.41 \times 10^{-17},\ 1.674 \times 10^{-17}]$
odd-sector lower bound	8.206×10^{-15}	2.3×10^{-17}
second even min–max value	$\geq 2.085 \times 10^{-12}$	not certified here
simplicity of the ground state	[9, Thm. 6.2]	not independently established here

The one place this note improves on the published result is the width of the whole-window bracket, which falls from 1.38×10^{-17} to 2.64×10^{-18}: approximately 80.9% narrower, a reduction by a factor of 5.23. The present bracket is strictly nested inside the published one, both endpoints moving inward. In the other three rows [9] is stronger, and in the odd sector by more than two orders of magnitude.

As a consistency check — not as an endpoint, and not as a replay of his certificate — [9] also reports a numerical reference estimate $\lambda_{\min}(Q|_{\text{window}}) = 1.656 \times 10^{-17}$ for this window, from his own exact time-domain assembly. That value lies inside the bracket certified here. It is a non-certified reference value in [9] and is used here only in that capacity.

2 Setup, conventions, and the statement

2.1 Normalization

Fix $L > 0$. For $f \in L^2(-L, L)$, extended by zero, write

$$F(t) \;=\; \int_{-L}^{L} f(u)\, e^{itu}\, du, \qquad \text{so} \qquad \int_{\mathbb{R}} |F(t)|^2\, dt \;=\; 2\pi \|f\|_2^2.$$

The window Weil form used throughout is the Hermitian form

$$Q(f) \;=\; 2\,\mathrm{Re}\big(F(i/2)\,\overline{F(-i/2)}\big) \;+\; \frac{1}{2\pi}\int_{\mathbb{R}} |F(t)|^2\, \Psi_L(t)\, dt, \tag{1}$$

$$\Psi_L(t) \;=\; H(t) - P_L(t), \qquad H(t) \;=\; \mathrm{Re}\,\psi\Big(\tfrac{1}{4} + \tfrac{it}{2}\Big) - \log\pi, \qquad P_L(t) \;=\; \sum_{\log n < 2L} \frac{2\Lambda(n)}{\sqrt{n}}\, \cos(t \log n). \tag{2}$$

Here ψ is the digamma function and Λ is the von Mangoldt function. Writing $C = \int f \cosh(u/2)\, du$ and $S = \int f \sinh(u/2)\, du$, so that $F(\mp i/2) = C \pm S$, the pole term of (1) is

$$2\,\mathrm{Re}\big(F(i/2)\,\overline{F(-i/2)}\big) \;=\; 2\big(|C|^2 - |S|^2\big), \tag{3}$$

which is the form that must be used for complex f: it is additive over the real and imaginary parts, whereas $2F(i/2)^2$ is not — replacing f by if reverses its sign. The computation never needs (3) in that generality. It is carried out on *real* test functions of definite parity, where C survives on even f and S on odd f, and (3) specializes to

$$2s\, F(i/2)^2, \qquad s = +1 \text{ (even)}, \qquad s = -1 \text{ (odd)},$$

the two signs recording that the pole helps the even sector and hurts the odd one. Arbitrary complex f is then reached through the decomposition of Section 3.9, not through (3) directly.

The two sector signs are not asserted here: in the parity study [11, §3(c)] the $\delta = 1/2$ real-pair perturbation is compared entrywise with this repository's own pole term $2\,\mathrm{pole_sign}(\varepsilon) P_j P_k$ and agrees in all nine tested cells to 6.22×10^{-61}, with $\mathrm{pole_sign} = -1$ on the odd sector, while the opposite sign convention is wrong by $O(1)$ to $O(10)$; and [11, Remark 6] identifies that $\delta = 1/2$ case as the parity splitting of [9, Lemma 6.1]. The prime sum in (2) is *finite and exact*: the autocorrelation of a function supported in $[-L, L]$ is supported in $[-2L, 2L]$, so only n with $\log n < 2L$ contribute and there is no truncation. At $L = 4/5$ the active prime powers are $\{2, 3, 4\}$.

This is the normalization of `src/zeta_window.py` in the repository accompanying this note, documented in [10], where the same object is written $T_R = A - 2M$ with $A = \mathrm{Pole} + \mathrm{Arch} - (\log\pi) I$ and $2M = \mathrm{Prime}$. It is an imported convention, not derived here, and the grouping of $-\log\pi$ is part of it. It was cross-checked against the repository's own independent closed form for the constant test function (Section 4, item (a)).

2.2 The admissible class

Since H is even, continuous, and increasing on $t > 0$ (Lemma 5), and $|P_L| \le A_L := \sum_{\log n < 2L} 2\Lambda(n)/\sqrt{n} < \infty$, the function Ψ_L is bounded below on $\mathbb{R}$ and bounded on compacts. Hence for *every* $f \in L^2(-L, L)$ the negative part of the integral in (1) is finite and $Q(f)$ is well defined in $(-\infty, +\infty]$. We therefore take the admissible class to be all of $L^2(-L, L)$, with the convention that a lower bound $Q(f) \ge \mu \|f\|_2^2$ is satisfied trivially where $Q(f) = +\infty$. This is a choice of convention made in this note; it strengthens rather than weakens the statement, and it removes any need for a density or form-domain argument.

2.3 Three spectral quantities, kept apart

Confusing the following three numbers is the easiest way to overstate a result of this kind, so they are named separately and used separately throughout.

Definition 1. With $T > 0$ a frequency cutoff, β a constant with $\Psi_L(t) \geq \beta$ for $|t| \geq T$, and V_N a finite-dimensional subspace of $L^2(-L, L)$:

$$\begin{aligned} \lambda_Q &= \inf\{\, Q(f)/\|f\|_2^2 \,\} && \text{over the admissible class,} \\ \lambda_R(T) &= \inf\{\, R_{T,\beta}(f)/\|f\|_2^2 \,\} && \text{over the same class, } R_{T,\beta} \text{ as in (4),} \\ \lambda_A(T, N, \varepsilon) &= \lambda_{\min}(\text{compression of } R_{T,\beta} \text{ to } V_N) && \text{in the parity sector } \varepsilon. \end{aligned}$$

Because $Q \geq R_{T,\beta}$ pointwise, and because an infimum over a subspace is at least the infimum over the whole space,

$$\lambda_R(T) \;\leq\; \lambda_Q, \qquad \lambda_R(T) \;\leq\; \lambda_A(T, N, \varepsilon),$$

and there is **no** inequality in either direction between λ_A and λ_Q. In particular a numerical least eigenvalue of a finite block is not an upper bound for λ_Q, and only the evaluation of Q itself on an explicit test function is.

2.4 The theorem

Theorem 2. *Let $L = 4/5$ and let Q be the form (1)–(2) on $L^2(-L, L)$.*

(i) (Lower bound.) *For every $f \in L^2(-4/5, 4/5)$,*

$$Q(f) \;\geq\; \tfrac{141}{10^{19}}\,\|f\|_2^2 \;=\; 1.41 \times 10^{-17}\,\|f\|_2^2,$$

in both parity sectors separately and hence for arbitrary complex f.

(ii) (Upper bound.) *There is an explicit even test function f_{160}, a combination of the 160 even cosine modes $\cos(k\pi u/L)$, $k = 0, \ldots, 159$, with rational coefficients of denominator 10^{180}, such that*

$$\lambda_Q(4/5) \;\leq\; \frac{Q(f_{160})}{\|f_{160}\|_2^2} \;\leq\; U \;=\; 1.673125534070681868077268377 36\times 10^{-17} \;<\; 1.674\times 10^{-17}.$$

(iii) (Parity separation.) *For every odd $f \in L^2(-4/5, 4/5)$, $Q(f) \geq 2.3 \times 10^{-17}\|f\|_2^2$. Since f_{160} is even and $2.3 \times 10^{-17} - U > 6.26 \times 10^{-18} > 0$,*

$$\inf_{f \text{ even}} \frac{Q(f)}{\|f\|_2^2} \;\leq\; U \;<\; 2.3 \times 10^{-17} \;\leq\; \inf_{f \text{ odd}} \frac{Q(f)}{\|f\|_2^2},$$

so the even infimum equals $\lambda_Q(4/5)$.

Parts (i) and (iii) are computer-assisted: they rest on interval arithmetic in the Arb midpoint–radius model [6], including all analytic error terms, on exact rational input data, and on a cold verifier that recomputes every decisive implication. Part (ii) needs one test function and is unconditional given the normalization.

Remark 3. Theorem 2(iii) identifies which sector carries the infimum. It does *not* assert that the infimum is attained, that a minimizer is unique, or that the least eigenvalue is simple. None of those is established here.

Remark 4. The bracket width is $1.674 \times 10^{-17} - 1.41 \times 10^{-17} = 2.64 \times 10^{-18}$, about 15.8% of the upper endpoint and 18.7% of the lower one. Section 6 measures where that width comes from and bounds how much of it the present method can remove.

3 The certified lower bound

The lower bound is assembled in five steps. Steps 1–5 are proved here in the normalization of Section 2; the strategy is that of [9, Thm. 1.1].

3.1 Step 1: the exterior gate

Lemma 5. *$t \mapsto \operatorname{Re}\psi(\frac{1}{4}+\frac{it}{2})$ is strictly increasing on $t>0$.*

Proof. From the digamma series,

$$\operatorname{Re}\psi\Big(\tfrac{1}{4}+\tfrac{it}{2}\Big) = -\gamma + \sum_{n\ge 0}\left[\frac{1}{n+1} - \frac{n+1/4}{(n+1/4)^2+t^2/4}\right],$$

and each summand is strictly increasing in $t>0$. Termwise differentiation is justified by local uniform convergence. □

Consequently, with $A_L = \sum_{\log n<2L} 2\Lambda(n)/\sqrt{n}$ and $\beta := H(T) - A_L$, we get $\Psi_L(t) \ge \beta$ for all $|t| \ge T$, because $P_L \le A_L$ pointwise. At $L = 4/5$,

$$A_L = 2.941973525223620455503909 \pm 3.91 \times 10^{-25},$$

and the gate values at the three cutoffs used in this note are

T	$\beta = H(T) - A_L$	status
200	$0.51846573324384671998735 36 \pm 2.62 \times 10^{-26}$	certified
400	$1.21161369505806454947150 2 \pm 1.89 \times 10^{-26}$	certified
800	$1.90476107093077688862120 6 \pm 1.68 \times 10^{-25}$	certified

with the increments $\beta_{400} - \beta_{200} = 0.693147961814217829484148 4 \pm 7.24 \times 10^{-27}$, numerically close to $\log 2$ as the asymptotics of H suggest. All three are *positive*, so the set $\{|t| \ge T : \Psi_L(t) < \beta\}$ is empty and no occupation or measure estimate on a bad exterior set is needed anywhere below.

3.2 Step 2: the comparison form

Because $\int_{|t|\ge T}(\Psi_L - \beta)|F|^2 \ge 0$, equation (1) gives $Q(f) \ge R_{T,\beta}(f)$ with

$$R_{T,\beta}(f) \;=\; \beta\|f\|_2^2 \;+\; 2sF(i/2)^2 \;+\; \frac{1}{2\pi}\int_{-T}^{T}(\Psi_L(t)-\beta)|F(t)|^2\,dt. \tag{4}$$

The discarded quantity is exactly

$$e_T(f) \;:=\; \frac{Q(f) - R_{T,\beta}(f)}{\|f\|_2^2} \;=\; \frac{1}{2\pi\|f\|_2^2}\int_{|t|>T}(\Psi_L(t)-\beta)|F(t)|^2\,dt \;\ge\; 0, \tag{5}$$

which is measured in Section 6.

Proposition 6 (monotonicity in β)**.** *By Plancherel, $R_{T,\beta}(f) = 2sF(i/2)^2 + \frac{1}{2\pi}\int_{|t|\le T}\Psi_L|F|^2 + \frac{\beta}{2\pi}\int_{|t|>T}|F|^2$, so for $\beta_2 \ge \beta_1$,*

$$R_{T,\beta_2}(f) - R_{T,\beta_1}(f) = \frac{\beta_2-\beta_1}{2\pi}\int_{|t|>T}|F|^2 \;\ge\; 0.$$

On the finite compression this reads $A(\beta_2) - A(\beta_1) = (\beta_2-\beta_1)(I - K_T)$, where K_T is the compression of the Fourier energy-concentration operator and $0 \le K_T \le I$.

Proposition 6 is recorded because it rules out a tempting move: *lowering* β cannot raise the exact comparison form, its infimum, or the least eigenvalue of its compression to a fixed subspace. Inspecting only the factor $\Psi_L - \beta$ inside the integral misses the $\beta\|f\|_2^2$ term. The identity was checked numerically with maximum residual 5.23×10^{-143} and diagonal entries of K_T in $[0.9420888637695,\ 0.9980132600038]$ at $T = 200$, $N = 16$.

3.3 Step 3: the Legendre basis

Let $\varphi_n(u) = \sqrt{(2n+1)/(2L)}\,P_n(u/L)$ be the orthonormal Legendre basis of $L^2(-L,L)$. Then

$$F_n(t) \;=\; i^n\sqrt{2L(2n+1)}\;j_n(Lt), \qquad \sum_{n\ge 0}(2n+1)j_n(x)^2 = 1,$$

with j_n the spherical Bessel function. The form is assembled in the real coordinates $y_n = \sigma_n x_n$ with $\sigma_n = (-1)^{\lfloor n/2\rfloor}$, which makes $v_n(t) = \sqrt{2L(2n+1)}\,j_n(Lt)$ the real working transform. This basis is used, rather than the cosine basis of the repository's original assembly, because $j_n(x)$ is super-exponentially small for $n > ex/2$, which is what makes Step 4 effective; a cosine basis gives only $O(n^{-2})$ tail decay and cannot reach the required magnitude at an acceptable dimension.

3.4 Step 4: the complement

For $f \in V_N^{\perp}$ with $\|f\|_2 = 1$, Cauchy–Schwarz in n gives $|F(t)|^2 \le 2L\,S_N(tL)$ with $S_N(x) = \sum_{n\ge N}(2n+1)j_n(x)^2$. Since S_N is increasing on $0 \le x < N$,

$$g_N \;:=\; \sup_{f\in V_N^{\perp},\,\|f\|=1}\frac{1}{2\pi}\int_{-T}^{T}|F|^2 \;\le\; \frac{2LT}{\pi}\,S_N(TL).$$

This is a lossy but rigorous surrogate for the $(N+1)$-st Slepian eigenvalue [7], and it is super-exponentially small past the Shannon number $2TL/\pi$. At $T = 400$, $N = 440$:

$$S_N(320) = 9.260051461963608\times 10^{-63}, \qquad g_N = 1.886442193224756\times 10^{-60}.$$

3.5 Step 5: the Schur-type block bound, with strict tests

Split $f = f_N + f_\perp$. With a a certified lower bound for the least eigenvalue of the compression of $R_{T,\beta}$ to V_N, d a certified lower bound for $R_{T,\beta}$ on $V_N^{\perp}$, and κ a certified bound on the coupling, a 2×2 comparison yields positivity as soon as $ad > \kappa^2$. The threshold κ^2/d is a *break-even point*, not the resulting bound. To avoid the cancellation-prone explicit eigenvalue formula, the final implication is tested as three *strict* interval signs for the proposed constant μ:

$$a > \mu, \qquad d > \mu, \qquad (a-\mu)(d-\mu) > \kappa^2. \tag{6}$$

Equality anywhere in (6) would give only a nonnegative comparison, so strictness is required and is checked as a sign of an interval, never of a midpoint. The coupling constant is $\kappa = 2|p_N||p_\perp| + M\sqrt{g_N}$ with $M = \sup_{|t|\le T}|\Psi_L - \beta|$ and p the pole vector $p_n = \sqrt{2L(2n+1)}\,i_n(L/2)$ taken in the same y-coordinates; $|p|^2 = \int_{-L}^{L} e^{-u}\,du = 2\sinh L$ exactly. M is obtained from the monotonicity of H alone, with no covering argument.

3.6 Verified positive definiteness

A naive interval LDL^{T} factorization is unusable at this dimension: entry radii of order 10^{-38} amplify by roughly 6.5 per pivot and reach $O(1)$ by step 47. Instead, positive definiteness of $B = A - aI$ is established by a preconditioned congruence in the sense of Rump [5]. Let R be the upper triangular Cholesky factor of the *midpoint* of B, treated as exact input, and set $C = R^{-1}$ and

$$G \;=\; C^{*}B\,C \;=\; R^{-\mathsf{T}}\,(A - aI)\,R^{-1},$$

computed by two explicit forward substitutions in ball arithmetic. The verified inequality is

$$\|I - G\|_{\infty} \;<\; 1.$$

This proves $B \succ 0$: the bound places every eigenvalue of the Hermitian G in $(0,2)$, so $G \succ 0$; if C were singular then $G = C^{*}BC$ would be singular, so C is nonsingular; and then for $x \neq 0$, writing $x = Cy$ with $y \neq 0$, $x^{*}Bx = y^{*}Gy > 0$. It is a congruence, not a one-sided inverse residual — the latter would establish nonsingularity only.

3.7 Quadrature

The interior integral in (4) is evaluated by panelled Gauss–Legendre quadrature with certified nodes and weights (FLINT's `legendre_p_root`). The symbol, prime comb and the full basis-transform vector are evaluated once per node and reused across all matrix entries, so no entry is integrated adaptively. The quadrature remainder is bounded by a Bernstein-ellipse estimate on genuinely holomorphic extensions: the half-sum

$$H_{\mathrm{hol}}(z) \;=\; \tfrac{1}{2}\Big[\psi\Big(\tfrac{1}{4}+\tfrac{iz}{2}\Big)+\psi\Big(\tfrac{1}{4}-\tfrac{iz}{2}\Big)\Big]-\log\pi$$

agrees with H on the real axis and has its nearest poles at $z=\pm i/2$ (note that $\operatorname{Re}\psi(\frac{1}{4}+\frac{iz}{2})$ is *not* a holomorphic extension), and the basis product is extended as $v_n(z)v_m(-z)$ rather than by conjugation at a complex argument. On the ellipses, $|v_n(z)|$ is bounded uniformly in n by

$$\Big|\int_{-L}^{L}\varphi(u)e^{izu}\,du\Big| \;\le\; \Big(\int_{-L}^{L}e^{-2u\operatorname{Im}z}du\Big)^{1/2} \;\le\; \sqrt{2L}\,e^{L|\operatorname{Im}z|} \tag{7}$$

for any L^2-normalized φ supported in $[-L,L]$. Inequality (7) is immediate from Cauchy–Schwarz and is *not* claimed as new; it is recorded because it replaces a direct interval evaluation of j_n on wide complex balls, which is numerically useless here. The remainder is folded into the matrix entries as a widening exactly once.

3.8 The $T = 400$ certificate

parameter	value
window	$L = 4/5$
cutoff	$T = 400$
dimension	$N = 440$ (Legendre degrees 0..439; 220 modes per parity)
quadrature	800 panels of half-length 1/4, Gauss–Legendre order 64
Bernstein data	$b_e = 0.30$ (inside the strip $\lvert\operatorname{Im}z\rvert < 1/2$), $\rho = 2.762049935$
remainder factor	$3.551349410 \times 10^{-57}$
precision	900 bits (Bessel), 384 bits (accumulation), 1400 bits (verification)

$N = 440$ was chosen from the actual inequalities, not from the heuristic $eLT/2 = 434.9$. The certified constants are

$$\beta = 1.21161369505806454947150 2, \qquad M = 9.52577063950735058720 8369,$$
$$\kappa = 1.30834377128538825902 9018 \times 10^{-29}, \qquad \kappa^2/d = 1.41279636475158 8 \times 10^{-58},$$

with pole tail $|p_\perp| = 7.99337215214956 4 \times 10^{-1282}$.

sector	a (certified)	$\lVert I-G\rVert_\infty$	d	μ	margin in (6)
even	1.42×10^{-17}	$1.4973903 \times 10^{-14}$	1.211613695058065	1.41×10^{-17}	$1.21161369506 \times 10^{-19}$
odd	2.4×10^{-17}	$4.3535838 \times 10^{-22}$	1.211613695058065	2.3×10^{-17}	$1.21161369506 \times 10^{-18}$

The whole-window constant is the smaller of the two sector constants, so $\mu = 1.41 \times 10^{-17}$ and *the even sector binds*. Assembly took 3532 s, the full run 3581 s, with peak resident memory 393.4 MiB; the two Rump factorizations at $N = 440$ took under 50 s together.

3.9 Parity and complex tests

The form decouples by parity: the interior matrix has $C_{nm} = 0$ when $n + m$ is odd, because $v_n v_m$ is then odd on a symmetric interval, and the pole term $2(C^2 - S^2)$ separates into its even and odd parts. For complex $f = u + iv$ the Hermitian pole term (3) gives $2(|C|^2 - |S|^2) = 2(C_u^2 - S_u^2) + 2(C_v^2 - S_v^2)$, and the multiplier term splits because $|F|^2 = |F_u|^2 + |F_v|^2$ pointwise, so $Q(f) = Q(u) + Q(v)$; each real part then splits by parity. Hence the two sector certificates cover the full class, which is what makes Theorem 2(i) a statement about arbitrary complex f.

Remark 7. A positive-definiteness check cannot serve as an oracle for the pole sign. The direct convention checks of Section 4 are retained for that purpose, because an earlier version of this computation carried a pole-phase error that produced a plausibly negative pivot (Appendix A, item 1).

4 The original-form upper bound

4.1 The certified bound, from the closed-form route

A variational upper bound needs one admissible test function and an evaluation of Q on it that is certified end to end. The evaluation used for Theorem 2(ii) is the *closed-form* one: in the even cosine basis $f_k(u) = \cos(k\pi u/L)$ the prime sum is finite and exact, and the archimedean term is summed analytically through the digamma, Lerch Φ and Hurwitz ζ functions, each evaluated as an Arb ball. **This route performs no quadrature of any kind**, so it carries no quadrature truncation error to bound — which is exactly why it, and not the sharper route of Section 4.2, supplies the certified endpoint.

The procedure has two stages, which are kept apart. An exploratory floating-point eigensolver *locates* a candidate vector in this basis; it contributes no certified claim, and a poor candidate costs a weaker bound, never a wrong one. The candidate is then frozen to exact rationals and its Rayleigh quotient enclosed in Arb inside an owned precision scope. Because the second stage is exact, the first needs only enough precision to find a good trial function, which is why the mode count can be pushed without an expensive eigensolve.

Raising the mode count lowers the bound, and the sequence is the useful record:

N	certified $Q(f_N)/\|f_N\|_2^2 \le$	
28	$2.1129926117214908 \times 10^{-17}$	(the phase-1 witness)
48	$1.9204310132 \times 10^{-17}$	
64	$1.8541624482 \times 10^{-17}$	
96	$1.7476544722 \times 10^{-17}$	
128	$1.6972723502 \times 10^{-17}$	
160	$1.6731255341 \times 10^{-17}$	used in Theorem 2(ii)

The witness f_{160} is a frozen exact rational vector over the 160 even modes $k = 0, \dots, 159$ with denominator 10^{180}. Re-certified from those exact rationals alone — no eigensolver, no floating point — at 2920 bits,

$$\frac{Q(f_{160})}{\|f_{160}\|_2^2} \;=\; 1.67312553407068186807726837736 \times 10^{-17} \pm 4.41 \times 10^{-47},$$

whose outward upper endpoint is

$$U \;=\; 1.67312553407068186807726837736 \times 10^{-17} \;<\; 1.674 \times 10^{-17},$$

giving Theorem 2(ii). The radius is a consequence of the route: with no quadrature and no truncation, only ball rounding contributes. For comparison, [9] reports a numerical reference

estimate 1.656×10^{-17} for this infimum (Section 1.4), so the certified endpoint above sits about 1% over it.

Remark 8 (on quoting an endpoint). A serialized enclosure must be consumed through the correct endpoint — lower for a lower bound, upper for an upper bound — and never through its midpoint or its printed decimal. Two earlier drafts of this note violated that: one quoted an enclosure's midpoint as the bracket's upper endpoint, and one quoted an endpoint obtained by re-parsing a stored ball at the ambient 53 bits. Both are corrected (Appendix A, item 6), and the safe decimals 2.1130×10^{-17} here and 1.41×10^{-17} in part (i) are the forms in which the result should be quoted.

4.2 A sharper candidate: the time-domain route

A variational upper bound requires Q itself, not $R_{T,\beta}$. Evaluating (1) directly would mean certifying an oscillatory integral over all of $\mathbb{R}$. Instead Q is evaluated on the geometric side via the autocorrelation

$$g(x) \;=\; \int f(s+x)f(s)\,ds,$$

supported in $[-2L, 2L]$, together with the compact archimedean identity ([9, Lemma 2.5], transcribed into the present normalization)

$$\mathrm{Arch}(g) = -\big[\gamma_E + \log\pi + \log(1-e^{-4L})\big]g(0) + \int_0^{2L} \frac{2\big[e^{-2x}g(0) - e^{-x/2}g(x)\big]}{1-e^{-2x}}\,dx, \tag{8}$$

$$Q(f) = 2sF(i/2)^2 + \mathrm{Arch}(g) - \sum_{\log n < 2L} \frac{2\Lambda(n)}{\sqrt{n}}\,g(\log n). \tag{9}$$

This is the route by which [9, §6.2] evaluates its own variational upper bounds at this window, there with shift-correlation integrands of degree at most 239 integrated exactly by a 160-node Gauss rule. Two features make it cheap and rigorous. They are set out below because the present error accounting depends on them, not as new observations.

Exact inner quadrature. If f is a polynomial of degree d on $[-L, L]$ then g restricted to $[0, 2L]$ is a polynomial of degree at most $2d+1$ — not $2d$: for $f \equiv 1$ one has $g(x) = 2L - x$. What governs the *inner* rule is the degree of the integrand $f(s+x)f(s)$ in the integration variable s, which is at most $2d$, so an m-node Gauss–Legendre rule is exact once $2m-1 \geq 2d$, i.e. $m \geq d+1$. The implementation uses $m = d+2$; for $d = 438$ that is 440 nodes. *Exactness* here means no polynomial quadrature truncation error in exact arithmetic — node, weight, coefficient and rounding enclosures are still carried as ball radii, and the outer integral in (8) keeps its own certified Bernstein remainder.

The removable endpoint, by an identity. The integrand of (8) is removable at $x = 0$ and its limit involves $c = -g'(0^+)$. For even f,

$$g'(x) \;=\; -f(L)f(L-x) + \int_{-L}^{L-x} f'(s+x)f(s)\,ds,$$

and at $x = 0$ the integral is $\frac{1}{2}[f^2]_{-L}^{L} = 0$, so

$$c \;=\; -g'(0^+) \;=\; f(L)^2 \qquad \text{exactly,} \tag{10}$$

and the integrand limit is $c - \frac{3}{2}g(0)$. This closed form is an implementation convenience obtained by elementary differentiation and integration; [9, Lemma 2.5] leaves c as the one-sided limit and remarks only that it is nonzero in general. Its absence from the displayed formulas there is not evidence that it is new, and no novelty is claimed for it. The outer integrand is analytic in $|\operatorname{Im} x| < \pi$; panels are geometric near the origin so that no Bernstein ellipse encloses it, and the residual head $[0, 10^{-40}]$ is bounded by an explicit constant.

4.3 Validation of the representation

(a) On the constant $f = \varphi_0$, the compact route gives $Q(\varphi_0) = 0.080274054959269611214 \pm 3.09\times10^{-22}$, matching the repository's independent closed form $Q[0,0] = 0.080274054959269611214$ to all 21 printed digits (difference -3.08×10^{-22}). That closed form is the evaluator of [10], which sums the archimedean series analytically through the digamma, Lerch Φ and Hurwitz ζ functions; it is an independent route to Q, not an independent implementation of the present one.

(b) A degree-0 check cannot see a pole-phase error, since $\sigma_0 = +1$. Checks at φ_2 and at the signed mixed function $\varphi_0 - \varphi_2$ were therefore added, with the pole moment computed *directly* from the physical-space function rather than from the pole-vector helper, and with the finite prime shifts re-enumerated independently of the symbol helper:

f	$Q(f)$	pole moment (direct)
φ_0	0.080274054959269611214	1.29891290374844725
φ_2	0.389336213205013655883	0.030516224225381024
$\varphi_0 - \varphi_2$	0.117009437226589934780	1.26839667952306623

All agree with the helper values, the first to 10^{-25}.

(c) *Disclosure.* This is a second *representation* of the same form, not an independent implementation: it shares the Arb library, the symbol helper for the comb data, and the pole-vector helper. It is not described as an independent check.

4.4 The frozen test functions of the time-domain route

Two trial functions are used, each a frozen exact rational vector in the Legendre y-coordinates with denominator 10^{40}:

- f_{200}: 110 even modes, degrees 0..218, $\|f\|_2^2 = 2.27524444365096536097055 0$;
- f_{400}: 220 even modes, degrees 0..438.

Both were *located* by an exploratory floating-point eigensolver, then rounded to exact rationals and certified; the eigensolver contributes no certified claim, and in particular supplies no eigenvalue.

For f_{200}, equation (9) gives, at 94 outer panels and head cut 10^{-40},

$$\text{pole} = 3.2303213242823875400, \qquad \text{Arch} = -3.1402989129419417722,$$

$$\text{comb} = 0.090022411340445684793,$$

whose signed sum is $8.30143811294573721 \times 10^{-17}$: a cancellation of about sixteen digits, which is why the evaluation is done at high precision in ball arithmetic and why the compact representation — with its exact inner quadrature and the exact identity (10) — matters. Dividing by the norm,

$$q(f_{200}) = 3.64859175290407702 \times 10^{-17} \pm 4.15 \times 10^{-35}.$$

This is above the then-current upper bound and so does not improve it; the expected outcome, since f_{200} minimizes a comparison form, not Q.

For f_{400},

$$q(f_{400}) = 1.90138291114546440 \times 10^{-17} \pm 6.03 \times 10^{-35},$$

which would improve on U — were the enclosure complete. It is not; see Section 4.5. The radii printed above reflect ball rounding and the explicit head bound only.

4.5 Status: the outer quadrature remainder is not bounded

The outer integral in (8) is evaluated by panelled Gauss–Legendre quadrature — geometric panels from $\eta = 10^{-40}$ to 10^{-3}, then 32 uniform panels of order 32 to $2L$ — and the implementation adds the explicit head bound on $[0, \eta]$ but **no bound on the quadrature truncation error of those panels**. Ball arithmetic carries the node, weight and rounding enclosures; it does not capture the difference between a Gauss–Legendre sum and the integral it approximates, which is a property of the rule rather than of the arithmetic. The radii printed in Section 4.2 therefore do not include that term.

Accordingly, **every value produced by the time-domain route is withdrawn as a certificate** pending such a bound: $q(f_{200})$, $q(f_{400})$, the endpoint once quoted from the latter, and the discarded energies $e_T = q - r_T$ of Section 6 together with their ratios. They are retained below as *measurements*, labelled as such.

What is *not* affected, and why:

- the lower bound of Section 3, whose interior quadrature carries a certified Bernstein-ellipse remainder folded into the matrix entries exactly once, re-derived by its verifier;
- the odd-sector bound 2.3×10^{-17}, from the same matrix route;
- the certified upper bound of Section 4.1, which uses no quadrature at all;
- the scalar values $R_T(f)$ of Section 6, which are computed by a separate routine that *does* add a certified remainder and publishes it; in particular the ceiling $\lambda_R(800) \le r_{800}(f_{400})$ does not involve Q at any point.

A rigorous remainder for (8) requires a certified bound on the analytic continuation of the outer integrand on a Bernstein ellipse around each panel, with the denominator's zeros at $x \in i\pi\mathbb{Z}$ kept outside. The degree statement corrected above is a prerequisite for that bound and not a substitute for it: knowing $\deg g \le 2d + 1$ says nothing by itself about the size of g off the real axis. The obvious route — bound $|g|$ on the ellipse by $\max_{[0,2L]} |g| \cdot R^{\deg g}$ — is far too lossy at $\deg g = 877$, forcing panels of width $\sim 10^{-3}$. A coefficient-sensitive bound should work instead, because the frozen minimizers concentrate on *low* Legendre orders, so estimating $|f(w)|$ off the real axis directly from the frozen coefficients is tight exactly where $R^{\deg g}$ is hopeless. That derivation is left open; it is recorded with the diagnostic in `phase5/src/outer_convergence.py`, which measures the refinement difference at $\pm 2.71 \times 10^{-35}$ — small, and not a bound.

Since the certified endpoint of Section 4.1 now sits *below* the withdrawn time-domain value, closing this gap would no longer improve the bracket; it would restore a second, independent route to it.

5 The parity separation

The $T = 400$ certificate of Section 3 delivers $a = 2.4 \times 10^{-17}$ in the odd sector against $a = 1.42 \times 10^{-17}$ in the even sector. Re-evaluating the three strict signs (6) from the exact certified data — not from rounded report values — at $\mu_{\mathrm{odd}} = 2.3 \times 10^{-17}$ gives

$$a > \mu_{\mathrm{odd}}, \qquad d > \mu_{\mathrm{odd}}, \qquad (a - \mu_{\mathrm{odd}})(d - \mu_{\mathrm{odd}}) > \kappa^2,$$

with margin $1.21161369506 \times 10^{-18}$. So the whole odd sector satisfies $Q(f) \ge 2.3 \times 10^{-17} \|f\|_2^2$. Combining with the certified endpoint U of Section 4.1, and using that the witness f_{28} is even,

$$\mu_{\mathrm{odd}} - U \;=\; 6.268744659293181 \times 10^{-18} \;>\; 0,$$

so

$$\inf_{\mathrm{even}} \frac{Q(f)}{\|f\|_2^2} \;\le\; U \;<\; \mu_{\mathrm{odd}} \;\le\; \inf_{\mathrm{odd}} \frac{Q(f)}{\|f\|_2^2},$$

which with the orthogonal parity decomposition identifies the even infimum as $\lambda_Q(4/5)$. The global lower bound remains 1.41×10^{-17}, governed by the even sector.

The witness used here is the closed-form one. An earlier draft ran this argument on the time-domain value for f_{400}; that value is withdrawn by Section 4.5, and the separation is re-established above on a witness whose evaluation uses no quadrature. The conclusion is unchanged.

As noted in Section 1.3, none of this structure is new: at this very window [9, Thm. 6.2] certifies the same ordering with $\lambda_1^{\mathrm{odd}} \geq 8.206 \times 10^{-15}$, more than two orders of magnitude above the constant obtained here. What is recorded above is only that this pipeline's own certified numbers separate the two sectors. Nothing about simplicity, uniqueness or attainment follows from them — simplicity and evenness at this window were previously established by [9, Thm. 6.2].

6 Cutoff diagnostics

6.1 The cutoff comparison, proved

Proposition 9. *Let $\Psi_L \geq \beta_i$ on $|t| > T_i$ for $i = 1, 2$, with $T_2 \geq T_1$ and $\beta_2 \geq \beta_1$. Then*

$$R_{T_2,\beta_2}(f) - R_{T_1,\beta_1}(f) = \frac{1}{2\pi} \int_{T_1<|t|\leq T_2} (\Psi_L - \beta_1)|F|^2 + \frac{\beta_2 - \beta_1}{2\pi} \int_{|t|>T_2} |F|^2 \;\geq\; 0,$$

both terms being nonnegative. Hence $\lambda_R(T_2) \geq \lambda_R(T_1)$ for the exact forms.

Two cautions. This does not imply that a *certified numerical* bound improves, since error terms change with the parameters. And it does not order finite matrices of different sizes: changing N changes the compression, so only matrices in the same subspace are comparable by this identity.

6.2 The scalar table

Each R_T entry below is a certified enclosure on a *fixed, frozen* function; no reoptimization is involved, so the columns compare exact forms on the same vector. The Q row comes from the time-domain route and is therefore a **measurement, not a certificate** (Section 4.5); it is shown because the diagnostics below are read off it.

	f_{200}	f_{400}
R_{200}	$1.0310220302459896 \times 10^{-17}$	$1.24275215 \times 10^{-17}$
R_{400}	$2.3520530467310079 \times 10^{-17}$	$1.4330724047355850 \times 10^{-17}$
R_{800}	3.01603×10^{-17}	1.6681×10^{-17}
Q measured	$3.6485917529040770 \times 10^{-17}$	$1.9013829111454644 \times 10^{-17}$

The R_{800} entries carry wider enclosures ($\pm 5.30 \times 10^{-23}$ and $\pm 4.60 \times 10^{-22}$) because the quadrature remainder grows with the cutoff; $R_{200}(f_{400})$ was evaluated as a direct scalar integral on the full degree-438 function, with no truncation and no enlarged matrix. Each column is strictly increasing in T, as Proposition 9 requires.

6.3 What the comparison discards

From (5), with $e_T(f) = q(f) - r_T(f)$ by rigorous interval subtraction. Since q is withdrawn as a certificate (Section 4.5), **every entry in this table is a measurement**, and no statement below rests on one:

	f_{200}	f_{400}
e_{200}	$2.617569722658087 \times 10^{-17}$	6.586308×10^{-18}
e_{400}	$1.296538706173069 \times 10^{-17}$	$4.683105064098794 \times 10^{-18}$
e_{800}	6.3256×10^{-18}	2.333×10^{-18}
e_{400}/e_{200}	$0.49532156 \pm 4.69 \times 10^{-9}$	$0.7110365 \pm 1.61 \times 10^{-8}$
e_{800}/e_{400}	$0.48788 \pm 9.07 \times 10^{-6}$	$0.498 \pm 2.36 \times 10^{-4}$

There is no decay law here. On f_{400} the successive ratios are 0.711 and then 0.498, which is not a constant rate; and the two functions disagree at the *same* doubling $200 \to 400$, giving 0.495 and 0.711, so the ratio is not a function of T alone. That both functions give approximately 0.49 across $400 \to 800$ is two samples of one step, not a rate in T, and comparing differently optimized vectors would establish less still. No uniform operator estimate is inferred from a fixed-vector ratio, and none of these ratios is used in any certified statement.

6.4 What $q_f - r_f$ is, and is not

For f_{400}, $q_f - r_f = 4.6831050640988 \times 10^{-18}$ is the discarded-term identity (5) evaluated on that function, inside its enclosure — a measured quantity, since q is withdrawn (Section 4.5). It is *not* $\lambda_Q - \lambda_R(400)$: both $\lambda_R(400) \le r_f$ and $\lambda_Q \le q_f$ are upper bounds, and two upper bounds do not make their difference an equality between infima.

6.5 Certified ceilings, and the stopping decision

Because f_{400} and f_{200} are admissible, each R_T value in the table is a variational *upper* bound for $\lambda_R(T)$, and any certificate resting only on $Q \ge R_{T,\beta}$ is capped by it. Relative to the lower bound in force:

route	certified ceiling	reference lower bound	headroom
R_{200}	$\lambda_R(200) \le r_{200}(f_{200}) \le 1.0310221 \times 10^{-17}$	1×10^{-17}	3.1023%
R_{400}	$\lambda_R(400) \le r_{400}(f_{400}) \le 1.4330725 \times 10^{-17}$	1.41×10^{-17}	1.6364%
R_{800}	$\lambda_R(800) \le r_{800}(f_{400}) \le 1.6681461 \times 10^{-17}$	1.41×10^{-17}	18.309%

Each ceiling above is the *outward-rounded upper endpoint* of the corresponding enclosure, not its midpoint: from $r_{800}(f_{400}) = 1.6681 \times 10^{-17} \pm 4.60 \times 10^{-22}$ the endpoint is $1.6681460 \times 10^{-17}$, quoted outward as $1.6681461 \times 10^{-17}$; quoting the midpoint 1.6681×10^{-17} as a ceiling would be the same error as in Remark 8.

So the $T = 400$ route is within about 1.64% of exhausted, and the entire $T = 800$ route is capped at a factor 1.1831 over the present lower endpoint; since a reoptimized f_{800} can only sit at or below that ceiling, the realizable gain is at most that. Measured costs on the machine used:

stage	measurement
assembly, $T = 200$, $N = 220$, 800×64 nodes	1338 s, peak 205.5 MiB
assembly, $T = 400$, $N = 440$, same node count	3532 s, peak 393.4 MiB
Rump factorizations, both sectors, $N = 440$	under 50 s
compact $Q(f_{200})$	389 s
cross table (R_{400}, $Q(f_{400})$)	1544 s
scalar leg $R_{200}(f_{400})$, 12800 nodes	933 s
scalar leg $R_{800}(f_{400})$, 51200 nodes	8049 s
scalar leg $R_{800}(f_{200})$, 51200 nodes	3651 s

The assembly grew by a factor 2.64 when N doubled at fixed node count, and the $T = 800$ scalar legs overran the node-count scaling by more than a factor of two, which indicates that per-Bessel cost grows with the argument rather than node count alone. Extrapolating — as an estimate anchored on these measurements, not a guarantee — a full $T = 800$ matrix at $N \approx 880$ with roughly 1600×64 nodes would cost on the order of ten hours of assembly. Against a ceiling of $+18.31\%$, the recommendation recorded in the accompanying repository is to **consolidate at** $T = 400$. Neither the pilot nor any failed certificate here is an intrinsic barrier; this is a cost–benefit judgement from measured data, and a separately scoped $T = 800$ attempt would be a reasonable thing to run with that ceiling stated in advance.

7 Reproducibility and limitations

7.1 Artifacts and verification

Every decisive quantity in this note is published as a midpoint–radius enclosure in machine-readable form, together with the exact rational input vectors and the code that produced them. Printed decimals in this note are summaries; the enclosures are authoritative. Nine *verification scripts* recompute the chain from the frozen artifacts and exit nonzero on any failure; on the run accompanying this version all nine exit 0, with 187 checks passing, plus a re-certification of the upper bound from its stored rationals, in 128 seconds. They are not independent implementations: they share the assembly and constants modules with the computation they check, and no independence is claimed for them. A seventh script, shipped with this note, re-derives every decimal quoted above from the artifacts and additionally checks each quoted bound's *rounding direction* and its scope and parity label; it is what caught the mis-quoted upper endpoint of Appendix A, item 6.

Within those scripts two kinds of check must be distinguished. Most recompute a *certificate-producing inequality* — a Schur sign, a Cholesky residual, a tail bound — in ball arithmetic, and a failure there invalidates a claim. A few are *auxiliary*: they compare the Arb assembly against the repository's independent `mpmath` implementation, or assert that a diagnostic spectrum is finite. Those use floating point and settle nothing about the certificates; one of them formerly compared two matrices after converting both to ordinary doubles while asserting agreement below 10^{-30}, a threshold that no double-precision comparison can reach. It now compares at full precision and states the tolerance its inputs actually support (Appendix A, item 9).

The verifiers fail closed. Any NaN, infinite radius, indeterminate enclosure, or unresolved sign produces an explicit failure or a refinement request; no skipped cell, panel or pivot is interpreted as success. This matters because an earlier version of the computation silently skipped cells where a library digamma call returned NaN on wide balls and thereby produced a wrong constant that was *used* (Appendix A, item 2).

7.2 Environment

Linux, Python 3.14.7, `mpmath 1.3.0`, `python-flint 0.9.0` (FLINT/Arb). Floating-point `mpmath` is used to *locate* candidate vectors, which are then frozen as exact rationals and certified in Arb, and in the auxiliary cross-implementation checks described above. It enters no certificate-producing inequality; it does appear in a small number of auxiliary verifier assertions, which is a weaker statement than the one made in earlier versions of this note. Arb working precisions on sign-critical paths range from 256 to 2000 bits and are stated per stage in the artifacts.

7.3 Limitations

1. **An unbounded outer quadrature remainder (open).** The time-domain evaluator of Section 4.2 omits the truncation error of its outer panelled rule, so every value it produced

is withdrawn as a certificate (Section 4.5). This is the principal open item. The certified endpoint of Section 4.1 does not depend on it, and neither does the lower bound.

2. **One window.** Everything is at $L = 4/5$. There is no propagation theorem, and nothing follows at any larger support. The repository's companion measurement at $L = 81/100$ indicates roughly $75\times$ the assembly work for a target that is certifiably at most 6.565×10^{-18}, which is why that window was not attempted here.

3. **An imported normalization.** The explicit-formula pairing in the convention of Section 2 is imported from the accompanying repository, not rederived. It was cross-checked only through the independent closed form for $Q[0, 0]$ and the degree-2 convention checks of Section 4.

4. **An unverified external matrix step.** Zhu describes supplementary source code and certificates in [9, §17]; these were not obtained or executed in the present work, so the matrix step of [9, Thm. 1.1] is not reproduced here. The present certificate is independent of it.

5. **Shared arithmetic.** The compact time-domain evaluator shares library code with the matrix assembly, and is reported as a second representation, not an independent implementation.

6. **No arithmetic content.** The only analytic sharpening proved here is the monotone gate of Lemma 5, which moves a constant by under one percent and does not touch the doubly exponential threshold that governs the cost in L.

7. **Scope.** No zero-free region, no verified range of zeros, no statement about the Riemann hypothesis.

7.4 Revision history

The three stages are kept distinguishable on purpose: the versions that carried the *original omission*, the version that made the *interim correction*, and any version carrying a *verified repair*. No version is rewritten in place.

status All versions below are **drafts**. Nothing has been deposited, released or archived, and the manuscript stays marked *awaiting author review* until the author authorizes publication — which remains true after the repair passes its checks.

1.0.x carried the omission. The upper bound 1.9014×10^{-17} and everything derived from it rested on an incomplete error budget. Those versions should not be cited for the upper bound.

1.1.0 interim correction. The affected claims were withdrawn and the endpoint reverted to a quadrature-free route.

1.2.0 improved, repair still open. The quadrature-free endpoint was tightened past the withdrawn value. The outer quadrature remainder itself is *not* repaired; that would be a later version.

1.0.0 (2026-11-10) First version.

1.0.1 (2026-11-10) Citation and provenance revision only. Adds the two earlier technical notes whose implementation and parity conventions this computation inherits ([10] and [11]), the lineage paragraph in Section 1.3, and the corresponding citations in Sections 2 and 4. No numerical result, bound, artifact or limitation changed; the certified bracket and the parity statement are exactly those of version 1.0.0.

1.0.2 (2026-11-10) Corrections after reading [9] in full, which had previously been consulted only through its abstract and rendered HTML. Three attributions were too weak and are strengthened: [9, Thm. 6.2] certifies the parity *separation* at this very window, with a far stronger odd-sector constant, and establishes simplicity and evenness of the ground state (Sections 1.3 and 5); [9, §6.2] already evaluates its variational upper bounds by the time-domain route used in Section 4; and the supplementary materials of [9, §17] are described there, rather than absent. Section 1.4 is new and compares the two sets of bounds at this window. Again no numerical result, bound or artifact changed.

1.1.0 (2026-11-10) **The certified upper endpoint changes.** The time-domain evaluator was found to omit the truncation error of its outer quadrature (Section 4.5), so every value it produced is withdrawn as a certificate and the upper bound reverts to the closed-form witness of Section 4.1: 2.1130×10^{-17} in place of 1.9014×10^{-17}. The bracket widens from 4.914×10^{-18} to 7.03×10^{-18}, the parity separation is re-established on the closed-form witness with margin 1.87×10^{-18}, and Section 1.4 is updated. The lower bound, the odd-sector bound, the R_T values and the ceilings are unaffected. Four further corrections: the Hermitian pole term for complex f (Section 2), the degree of the autocorrelation, outward rounding of the ceilings, and the scope of the verification claims (Section 7).

1.2.0 (2026-11-10) **The certified upper endpoint improves past the withdrawn one.** Rather than repair the unbounded quadrature remainder, the endpoint is tightened along the route that has no quadrature to bound: the closed-form evaluation is pushed from 28 even cosine modes to 160, giving $U = 1.6731255341 \times 10^{-17}$ in place of 2.1130×10^{-17}, and in place of the withdrawn time-domain value 1.9014×10^{-17} as well. The bracket narrows to 2.64×10^{-18}, 80.9% inside the published one, and the parity margin grows to 6.27×10^{-18}. The lower bound, the odd-sector bound, the R_T values and the ceilings are unchanged. The outer quadrature remainder of Section 4.5 remains open; it is no longer on the critical path.

A Error history

Errors found during this work are recorded because several of them produced plausible wrong numbers rather than failures, and because the countermeasures are part of what the artifacts are for. Items 1–5 are library- or discipline-level defects; items 6–8 are corrections to claims. A longer ledger, including retracted statements $R1$–$R13$, is kept with the artifacts.

1. **Pole-coordinate phase.** The form is assembled in the coordinates $y_n = \sigma_n x_n$ with $\sigma_n = (-1)^{\lfloor n/2 \rfloor}$, so the pole vector requires the same phase. Omitting it is invisible at degree 0 and wrong from degree 2 onward. It produced a genuinely negative pivot (-0.3949) that would have been reported as indefiniteness. Caught by comparing $|F(t)|^2$ and $F(i/2)$ against direct integration; the degree-2 checks of Section 4 now cover it.

2. **Digamma on wide balls.** The library digamma returns NaN on balls whose real part has radius $\gtrsim 0.1$, even strictly inside the right half-plane. A comparison loop silently skipped those cells and returned $M \approx 1831$ instead of 8.83 — and the wrong value was used. Replaced by the explicit recurrence $\psi(z) = \psi(z+K) - \sum_{k<K} (z+k)^{-1}$ with adaptive K, and by a covering-free monotonicity bound for M.

3. **Bessel functions on wide balls.** Interval evaluation of j_n on the complex ellipses is useless here (width $\sim 10^{32}$ at $|w| \sim 80$, unchanged at 3000 bits). Replaced by the uniform Cauchy–Schwarz bound (7).

4. **Interval factorization.** Naive interval LDL^{T} is unusable at these dimensions (radii amplify about 6.5× per pivot, reaching $O(1)$ by step 47). Replaced by the Rump preconditioned congruence.

5. **Precision ownership.** The Arb context precision is a process global. Routines set it on entry and did not restore it, so a callee silently changed its caller's context; this produced four separate wrong constants in this project, each plausible-looking rather than an error (a complement constant at $[\pm 7.4 \times 10^{132}]$, an even-sector d at $[\pm 2.8 \times 10^{183}]$, and two coefficient sets built at 53 bits). The repair is an owned-scope wrapper that asserts restoration on normal return *and* on exception, caches keyed by precision so a low-precision value can never be served to a high-precision caller, and an explicit *unresolved* result for any wide or non-finite enclosure. The lesson, and the harder half, is that *an input constructed outside the scope carries the ambient precision's radius, and entering a higher-precision scope afterwards does not recover it*: own your inputs as well as your scope.

6. **The upper endpoint, twice mis-quoted.** Two distinct errors, both in the safe direction numerically but both wrong as stated. (a) Internal drafts quoted the enclosure *midpoint* $1.90138291114546440 \times 10^{-17}$ as the bracket's upper endpoint; a midpoint is not an upper bound for its own ball. (b) A separate record quoted $1.9013829111454646892 \times 10^{-17}$ as "the exact upper endpoint"; that value came from re-parsing the serialized ball at the ambient 53 bits, which rounds the midpoint and then inflates the endpoint by about 2.8×10^{-33} — the precision-ownership defect of item 5 recurring in the very routine that documents it. It recurred twice more while that was being repaired: once extracting an endpoint after an owned scope had exited, once parsing a stored ball string at the ambient precision. Four occurrences, which is why deserialisation and endpoint extraction now have their own entry points in the precision module instead of being written inline. See Remark 8.

7. **An omitted quadrature remainder.** The time-domain evaluator adds an explicit bound for the integrable head on $[0, \eta]$ but none for the truncation error of its outer panelled Gauss–Legendre rule, while its module docstring asserted that a Bernstein-ellipse remainder was used. Ball arithmetic does not supply that term: it encloses the nodes, weights and rounding, not the difference between the rule and the integral. The published radii were therefore incomplete, and the bound they supported is withdrawn (Section 4.5). The certified endpoint now comes from a route that performs no quadrature. This is the one item in this list that changed a headline number.

8. **A float comparison asserting an unreachable tolerance.** An auxiliary verifier check compared the Arb assembly with the repository's `mpmath` assembly by converting both to ordinary doubles and requiring agreement below 10^{-30}. Doubles carry about sixteen digits, so the comparison could only report exact equality or a difference near 10^{-16}; the threshold was unreachable and the check was weaker than it read. It now compares at full precision against the tolerance its inputs support.

9. **A retracted sparsity target.** An earlier proposed next step — a power-saving sparsity bound on the near-resonant set of the prime comb — is false, and is retracted in full: by continuous Kronecker–Weyl equidistribution that set has positive density, confirmed numerically here to four significant figures across four decades of T. The replacement is the empty-exterior gate of Section 3, which needs no such estimate.

10. **Overstated interpretations, corrected.** Four further claims made alongside earlier versions were withdrawn or corrected: that no bound on the measure of a bad set could control $\int_E |F|^2$ (false — standard finite-measure and local-occupation estimates do exactly that); that a Schur *threshold* was a lower bound (it is a break-even point, and the final

test needs strict inequalities, hence (6)); that lowering β would buy a larger a (refuted by Proposition 6 — the $\beta\|f\|_2^2$ term had been ignored); and an accusation that the library's adaptive integrator returned a silently wrong value (withdrawn — it could not be reproduced; the wrong numbers came from a run that still carried the pole-phase error of item 1, at a different cutoff, with output from two runs conflated).

References

[1] E. Bombieri, *Problems of the Millennium: the Riemann Hypothesis*, Clay Mathematics Institute, 2000.

[2] A. Connes and C. Consani, *Weil positivity and trace formula: the archimedean place*, Selecta Mathematica **27** (2021), no. 4, art. 77.

[3] A. Connes, C. Consani and H. Moscovici, *Zeta zeros and prolate wave operators*, preprint, arXiv:2203.02261.

[4] A. Connes and W. D. van Suijlekom, *Spectral truncations in noncommutative geometry and operator systems*, Communications in Mathematical Physics **383** (2021), 2021–2067.

[5] S. M. Rump, *Verification methods: rigorous results using floating-point arithmetic*, Acta Numerica **19** (2010), 287–449.

[6] F. Johansson, *Arb: efficient arbitrary-precision midpoint-radius interval arithmetic*, IEEE Transactions on Computers **66** (2017), no. 8, 1281–1292.

[7] D. Slepian and H. O. Pollak, *Prolate spheroidal wave functions, Fourier analysis and uncertainty I*, Bell System Technical Journal **40** (1961), 43–63.

[8] A. Weil, *Sur les "formules explicites" de la théorie des nombres premiers*, Comm. Sém. Math. Univ. Lund (1952), 252–265.

[9] X. Zhu, *Weil positivity in compact windows: a finite reduction, certified two-sided bounds, and a Landau–Widom decay law*, preprint, arXiv:2608.24827v2 (2026).

[10] E. De Jesús, *A δ^2–L^3 Sensitivity Law for Finite-Window Weil Positivity, with a Function-Field Control of the Connes–van Suijlekom Pipeline*, technical note, Zenodo, 2026. doi:10.5281/zenodo.23250932

[11] E. De Jesús, *Parity Blindness of Finite-Window Weil Detectors at a Real Zero: the Even Sector Cannot See a Siegel Configuration and the Odd Sector Is Maximally Sensitive to It*, technical note, version 2, Zenodo, 2026. doi:10.5281/zenodo.23624161

[12] E. De Jesús, *weil-window-control*, research repository, branch `positive-weil-certificate-2026-11-09`. `https://github.com/innerlightr-wq/weil-window-control`