




Recurrent Neural Networks



Agenda

1. Theory on Recurrent Neural Networks
2. RNNs in TensorFlow
3. Exercise Time Series Prediction
4. A Message from the President

Sequence Prediction with RNNs

Types of Sequences

- › Time-Series: Stock Prices, Revenue, Customer
- › Speech: Automatic Translation, Autocomplete, Speech-to-Text, Sentiment Analysis (written & spoken)
- › Creative Tasks



magenta






Figure 5. A selection of evaluation results, grouped by human rating.

- › Works on sequences of arbitrary lengths
- › Loops



Recurrent Neural Network


- › Loops
- › Works on sequences of arbitrary lengths




Sequence Prediction with RNNs

Types of Sequences


one to one




one to many




many to one



many to many



many to many



RNN and Memory


Types of Sequences

- › Recurrent neuron memory since it preserves some state across time steps (**memory cell**)
- › Weight Sharing across cells
- › Backbone of NLP
- › **But** plain RNNs are not very good for anticipating long-term dependencies


Long-Short-Term-Memory Cells

Capturing long-term dependencies


- › 1997 introduced by Hochreiter and Schmidhuber
- › Solve for long-term dependency problem




LSTM Mechanics (1)



LSTM Mechanics (1)



LSTM Mechanics (1)




$$f_t = \sigma(W_f \cdot [h_{t-1}, x_t] + b_f)$$

$$i_t = \sigma(W_i \cdot [h_{t-1}, x_t] + b_i)$$

$$\tilde{C}_t = \tanh(W_C \cdot [h_{t-1}, x_t] + b_C)$$

LSTM Mechanics (1)




$$f_t = \sigma(W_f \cdot [h_{t-1}, x_t] + b_f)$$

$$i_t = \sigma(W_i \cdot [h_{t-1}, x_t] + b_i)$$

$$\tilde{C}_t = \tanh(W_C \cdot [h_{t-1}, x_t] + b_C)$$

$$C_t = f_t * C_{t-1} + i_t * \tilde{C}_t$$

LSTM Mechanics (1)



$$f_t = \sigma(W_f \cdot [h_{t-1}, x_t] + b_f)$$

$$i_t = \sigma(W_i \cdot [h_{t-1}, x_t] + b_i)$$

$$\tilde{C}_t = \tanh(W_C \cdot [h_{t-1}, x_t] + b_C)$$

$$C_t = f_t * C_{t-1} + i_t * \tilde{C}_t$$

$$o_t = \sigma(W_o \cdot [h_{t-1}, x_t] + b_o)$$

$$h_t = o_t * \tanh(C_t)$$

Agenda

1. Theory on Recurrent Neural Networks
2. RNNs in TensorFlow
3. Exercise Time Series Prediction
4. A Message from the President

RNNs in TensorFlow

`tf.contrib.rnn.BasicRNNCell()` – most basic RNN cell

`tf.contrib.rnn.BasicLSTMCell()` – basic LSTM RNN cell

`tf.contrib.rnn.BasicGRUCell()` – Gated Recurrent Unit cell

1. Define Basic Cell with hidden size
2. Stack cells

Agenda

1. Theory on Recurrent Neural Networks
2. RNNs in TensorFlow
3. Exercise Time Series Prediction
4. A Message from the President

let's do some practical stuff...

04_1_rnn_time_series.ipynb

Agenda

1. Theory on Recurrent Neural Networks
2. RNNs in TensorFlow
3. Exercise Time Series Prediction
4. A Message from the President

<https://trumpnet.fosterelli.co/>

Which tweet is real?

Help us train a neural network!

Rubio is weak on illegal immigration, with the worst voting record in the U.S. Senate in many years. He will never MAKE AMERICA GREAT AGAIN!

Senator wins in West Virginia and name. Get Was for than my business. That's well under fighting Republicans!

For $\bigoplus_{n=1,\dots,m} \mathcal{L}_{m,n} = 0$, hence we can find a closed subset \mathcal{H} in \mathcal{H} and any sets \mathcal{F} on X , U is a closed immersion of S , then $U \rightarrow T$ is a separated algebraic space.

Proof. Proof of (1). It also start we get

$$S = \text{Spec}(R) = U \times_X U \times_X U$$

and the comparicoly in the fibre product covering we have to prove the lemma generated by $\coprod Z \times_U U \rightarrow V$. Consider the maps M along the set of points \mathcal{Sch}_{fppf} and $U \rightarrow U$ is the fibre category of S in U in Section ?? and the fact that any U affine, see Morphisms, Lemma ???. Hence we obtain a scheme S and any open subset $W \subset U$ in $\mathcal{Sh}(G)$ such that $\text{Spec}(R') \rightarrow S$ is smooth or an

$$U = \bigcup U_i \times_{S_i} U_i$$

which has a nonzero morphism we may assume that f_i is of finite presentation over S . We claim that $\mathcal{O}_{X,x}$ is a scheme where $x, x', x'' \in S'$ such that $\mathcal{O}_{X,x'} \rightarrow \mathcal{O}'_{X',x'}$ is separated. By Algebra, Lemma ?? we can define a map of complexes $\text{GL}_{S'}(x'/S'')$ and we win. \square

To prove study we see that $\mathcal{F}|_U$ is a covering of X' , and \mathcal{T}_i is an object of $\mathcal{F}_{X/S}$ for $i > 0$ and \mathcal{F}_p exists and let \mathcal{F}_i be a presheaf of \mathcal{O}_X -modules on \mathcal{C} as a \mathcal{F} -module. In particular $\mathcal{F} = U/\mathcal{F}$ we have to show that

$$\tilde{\mathcal{M}}^\bullet = \mathcal{I}^\bullet \otimes_{\text{Spec}(k)} \mathcal{O}_{S,s} - i_X^{-1} \mathcal{F}$$

is a unique morphism of algebraic stacks. Note that

$$\text{Arrows} = (\mathcal{Sch}/S)_{fppf}^{\text{opp}}, (\mathcal{Sch}/S)_{fppf}$$

and

$$V = \Gamma(S, \mathcal{O}) \longrightarrow (U, \text{Spec}(A))$$

is an open subset of X . Thus U is affine. This is a continuous map of X is the inverse, the groupoid scheme S .

Proof. See discussion of sheaves of sets. \square

The result for prove any open covering follows from the less of Example ???. It may replace S by $X_{\text{spaces},\text{étale}}$ which gives an open subspace of X and T equal to S_{Zar} , see Descent, Lemma ???. Namely, by Lemma ?? we see that R is geometrically regular over S .

Lemma 0.1. Assume (3) and (3) by the construction in the description.

Suppose $X = \lim |X|$ (by the formal open covering X and a single map $\underline{\text{Proj}}_X(\mathcal{A}) = \text{Spec}(B)$ over U compatible with the complex

$$\text{Set}(\mathcal{A}) = \Gamma(X, \mathcal{O}_{X,\mathcal{O}_X}).$$

When in this case of to show that $\mathcal{Q} \rightarrow \mathcal{C}_{Z/X}$ is stable under the following result in the second conditions of (1), and (3). This finishes the proof. By Definition ?? (without element is when the closed subschemes are catenary. If T is surjective we may assume that T is connected with residue fields of S . Moreover there exists a closed subspace $Z \subset X$ of X where U in X' is proper (some defining as a closed subset of the uniqueness it suffices to check the fact that the following theorem

(1) f is locally of finite type. Since $S = \text{Spec}(R)$ and $Y = \text{Spec}(R)$.

Proof. This is form all sheaves of sheaves on X . But given a scheme U and a surjective étale morphism $U \rightarrow X$. Let $U \cap U = \coprod_{i=1,\dots,n} U_i$ be the scheme X over S at the schemes $X_i \rightarrow X$ and $U = \lim_i X_i$. \square

The following lemma surjective restrocomposes of this implies that $\mathcal{F}_{x_0} = \mathcal{F}_{x_0} = \mathcal{F}_{X,\dots,0}$.

Lemma 0.2. Let X be a locally Noetherian scheme over S , $E = \mathcal{F}_{X/S}$. Set $\mathcal{I} = \mathcal{J}_1 \subset \mathcal{I}'_n$. Since $\mathcal{I}^n \subset \mathcal{I}'_n$ are nonzero over $i_0 \leq p$ is a subset of $\mathcal{J}_{n,0} \circ A_2$ works.

Lemma 0.3. In Situation ???. Hence we may assume $q' = 0$.

Proof. We will use the property we see that p is the next functor (??). On the other hand, by Lemma ?? we see that

$$D(\mathcal{O}_{X'}) = \mathcal{O}_X(D)$$

where K is an F -algebra where δ_{n+1} is a scheme over S . \square

<https://github.com/martin-gorner/tensorflow-rnn-shakespeare>