
EEG PEAK PREDICTION

Project

Author

Xinyu Qian and Satyarth Arora

March 5, 2025

Contents

1	Abstract	3
2	Plan	3
2.1	Peaks in an EEG signal	3
2.2	Experiment Biases	3
2.3	Labeling and Validation of Peaks	3
2.4	Strategies for Isolating Peaks	3
2.5	Model/Algorithm Proposal	3
3	Visual Evoked Potential Peaks	4
3.1	N35	4
3.2	P50	4
3.3	N75	4
3.4	P100	4
3.5	N170	4
3.6	P200	4
3.7	P3	5
4	Peaks for Auditory	5
4.1	N0	5
4.2	P1	5
4.3	N1	5
4.4	P2	5
4.5	N2	5

1 Abstract

This paper proposes a peak detection algorithm/model to recognize and label specific peaks found in an EEG signal. The model uses audiovisual interaction EEG data collected in Dr. Ozdamar's lab supplemented with EEG data found in online public databases.

2 Plan

2.1 Peaks in an EEG signal

Our first plan of action is to extrapolate and explain in the most simplest terms, the different peaks observed in an EEG signal. We also will expand on meta properties about the peaks which will feed in to our algorithm as features.

2.2 Experiment Biases

As compositions of EEG signals are dependent on

- Equipment Setup: Type and number of electrodes used to record the data.
- Experiment Setup: Which part of the human brain is targeted for the experiment.

We will be describing the biases and providing justification on how our algorithm is bias independent/dependent.

2.3 Labeling and Validation of Peaks

To verify whether the peaks we are trying to find are accurate, we will be validating our EEG peak database from a domain expert (Dr. Ozdamar).

2.4 Strategies for Isolating Peaks

Next step is researching the state of the art on Isolation and peak extraction using signal processing or ML approaches.

2.5 Model/Algorithm Proposal

Upon doing the above, we will propose an architecture describing the best process (in our opinion) to achieve peak extraction and labelling.

3 Visual Evoked Potential Peaks

3.1 N35

N35 are peaks observed at around 35 ms post stimulus. They are recorded over the occipital or parietal regions. Observations have shown that this peak originates from the primary visual cortex (V1) and possibly extrastriate areas, reflecting the initial cortical response. It is influenced by low-level visual features. (What Features?)

3.2 P50

P50 are peaks observed at 45-60 ms post stimulus, produced by retinal ganglion cells and other retinal cellular elements, V1 is also involved. It is sensitive to stimulus features, the amplitude can be modulated by attention and stimulus saliency (frontal lobe involved).

3.3 N75

N75 are early negative peaks observed at around 75 ms post stimulus, a hallmark component of the pattern-reversal VEP, most prominent over the occipital cortex, often followed by P100. It is associated with the initial cortical response to visual contrast and pattern detection. It is also linked to the parvocellular pathway.

3.4 P100

P100 is a large positive peak observed at 80-130 ms post stimulus. P100 is a prominent peak that is used for analysis, as it has relatively little variation between subjects, minimal within subject interocular difference, and minimal variation with repeated measurement over time. P100 peak time is affected by nonpathophysiologic parameters such as pattern size, pattern contrast, mean luminance, signal filtering, patient age, refractive error, poor fixation and miosis.

3.5 N170

N170 is a negative peak around 140-180 ms, typically at 170 ms post stimulus, most prominent over the occipitotemporal regions. N170 is strongly associated with face perception but also responds to other complex visual objects. Examples include words, cars in experts. Primarily generated in the fusiform gyrus (FG, part of the fusiform face area, FFA) and superior temporal sulcus (STS).

3.6 P200

P200 is a positive peak observed at 180-250 ms post stimulus. Most prominent over frontal, central and parietal electrodes, but can also appear in occipitotemporal regions, depending on stimulus type. It is associated with higher order visual processing, including feature detection, attention allocation and stimulus categorization. Modulated by top-down attention and expectation.

3.7 P3

This is a positive peak observed at 250-500 ms. This is an important VEP peak. It is most prominent over the parietal and central regions, often visible at Pz and Cz sites. It reflects higher-order cognitive processing, particularly stimulus evaluation, decision-making and memory updating. It is most commonly observed in oddball paradigms. P3 amplitude is influenced by task difficulty, stimulus salience and attention. Larger P3 amplitudes are typically seen for attended or relevant stimuli. The latency of P3 is often used as an index of cognitive processing speed and is reflected by stimulus complexity and cognitive load.

4 Peaks for Auditory

4.1 N0

It's an early negative peak (Latency needs to be found). It is observed primarily over the vertex (Cz) and frontocentral regions, with contributions from temporal electrodes. It is likely generated by thalamic and primary auditory cortex activity. Sensitive to stimulus features such as intensity and frequency.

4.2 P1

This is an early positive peak observed at 50-100 ms post stimulus in adults (Earlier in children). It is maximal over frontocentral regions (Cz, Fz) and sometimes detected at temporal sites. P1 reflects early cortical processing of auditory stimuli, including sound detection and encoding. They are generated mainly in the primary auditory cortex with contributions from secondary auditory areas.

4.3 N1

This is a negative peak observed at 100 ms post stimulus. N1 reflects cortical processing of auditory stimuli, likely generated in the primary auditory cortex (A1) and secondary auditory areas. The amplitude and latency of N1 are modulated by attention, sound saliency and stimulus relevance.

4.4 P2

This is a positive peak observed at 150-200 ms post stimulus. P2 reflects further cortical processing of auditory stimuli, especially in terms of sound discrimination, attention and categorization. It is generated primarily in the secondary auditory cortex, but can involve other regions such as the prefrontal cortex and parietal areas, depending on the attentional demands of the task. Amplitude of P2 is modulated by stimulus saliency and attentional focus.

4.5 N2

This is a negative peak observed at 170-250 ms post stimulus. It is most prominent over frontocentral and central regions (Cz, Fz, Pz). N2 reflects higher-order auditory processing,

involving cognitive and attentional mechanisms. It is particularly associated with the cognitive control of auditory attention and the detection of target stimuli in oddball paradigms. The amplitude of N2 is modulated by the saliency of the auditory stimulus.

References