

# Entanglement and thermodynamics after a quantum quench in integrable systems

Vincenzo Alba<sup>a,1</sup> and Pasquale Calabrese<sup>a</sup>

<sup>a</sup>International School for Advanced Studies, Istituto Nazionale di Fisica Nucleare, Sezione di Trieste, 34136 Trieste, Italy

Edited by Subir Sachdev, Harvard University, Cambridge, MA, and approved June 6, 2017 (received for review March 3, 2017)

Entanglement and entropy are key concepts standing at the foundations of quantum and statistical mechanics. Recently, the study of quantum quenches revealed that these concepts are intricately intertwined. Although the unitary time evolution ensuing from a pure state maintains the system at zero entropy, local properties at long times are captured by a statistical ensemble with nonzero thermodynamic entropy, which is the entanglement accumulated during the dynamics. Therefore, understanding the entanglement evolution unveils how thermodynamics emerges in isolated systems. Alas, an exact computation of the entanglement dynamics was available so far only for noninteracting systems, whereas it was deemed unfeasible for interacting ones. Here, we show that the standard quasiparticle picture of the entanglement evolution, complemented with integrability-based knowledge of the steady state and its excitations, leads to a complete understanding of the entanglement dynamics in the space–time scaling limit. We thoroughly check our result for the paradigmatic Heisenberg chain.

entanglement | quantum quench | integrability | thermodynamics

Since the early days of quantum mechanics, understanding how statistical ensembles arise from the unitary time evolution of an isolated quantum system has been a fascinating question (1–7). A widely accepted mechanism is that, although the entire system remains in a pure state, the reduced density matrix of an arbitrary finite compact subsystem attains a long time limit that can be described by a statistical ensemble (8). In the last decade, groundbreaking experiments with cold atoms (9–19) simulated with incredible precision the unitary time evolution of many-body quantum systems, reviving the interest in this topic. The simplest out-of-equilibrium protocol in which these ideas can be theoretically and experimentally tested is the quantum quench (20, 21). A system is prepared in an initial state  $|\Psi_0\rangle$ , typically the ground state of a local Hamiltonian  $H_0$ , and it evolves with a many-body Hamiltonian  $H$ . At asymptotically long times, physical observables relax to stationary values, which for generic systems, are described by the Gibbs (thermal) ensemble (3–7), whereas for integrable systems, a generalized Gibbs ensemble has to be used (8, 22–45).

Although these results suggest a spectacular compression of the amount of information needed to describe steady states, state of the art numerical methods, such as the time-dependent density matrix renormalization group (46–49) (tDMRG), can only access the short-time dynamics. Physically, the origin of this conundrum is the fast growth of the entanglement entropy  $S \equiv -\text{Tr} \rho_A \ln \rho_A$ , with  $\rho_A$  being the reduced density matrix of an interval  $A$  of length  $\ell$  embedded in an infinite system. It is well-understood that  $S$  grows linearly with the time after the quench (50). This linear behavior implies an exponentially increasing amount of information manipulated during typical tDMRG simulations. Remarkably, the entanglement dynamics has been successfully observed in a very recent cold-atom experiment (19).

In this work, using a standard quasiparticle picture (50), we show that, in integrable models, the steady state and its low-lying excitations encode sufficient information to reconstruct the entanglement dynamics up to short times. According to the quasiparticle picture (50), the prequench initial state acts as a

source of pairs of quasiparticle excitations. Let us first assume that there is only one type of quasiparticles identified by their quasimomentum  $\lambda$  and moving with velocity  $v(\lambda)$ . Although quasiparticles created far apart from each other are incoherent, those emitted at the same point in space are entangled. Because these propagate ballistically throughout the system, larger regions get entangled. At time  $t$ ,  $S(t)$  is proportional to the total number of quasiparticle pairs that, emitted at the same point in space, are shared between  $A$  and its complement (Fig. 1A). Specifically, one obtains

$$S(t) \propto 2t \int_{2|v|t < \ell} d\lambda v(\lambda) f(\lambda) + \ell \int_{2|v|t > \ell} d\lambda f(\lambda), \quad [1]$$

where  $f(\lambda)$  depends on the production rate of quasiparticles. Eq. 1 holds in the space–time scaling limit  $t, \ell \rightarrow \infty$  at  $t/\ell$  fixed. When a maximum quasiparticle velocity  $v_M$  exists [e.g., because of the Lieb–Robinson bound (51)], for  $t \leq \ell/(2v_M)$ ,  $S$  grows linearly in time, because the second term in Eq. 1 vanishes. In contrast, for  $t \gg \ell/(2v_M)$ , the entanglement is extensive (i.e.,  $S \propto \ell$ ). This light-cone spreading of entanglement has been confirmed analytically only in free models (52–57), numerically in several studies (58–60), in the holographic framework (61–68), and in a recent experiment (19). The validity of the quasiparticle picture Eq. 1 for the entanglement dynamics has been proven for free models in ref. 52. In the presence of interactions, few results are known. For instance, the validity of Eq. 1 has been proven for rational CFTs (Conformal Field Theories) in ref. 69. In interacting integrable models à la Yang–Baxter, the quasiparticle picture has been used to describe the out-of-equilibrium dynamics after

## Significance

Understanding how statistical ensembles arise from the out-of-equilibrium dynamics of isolated pure systems has been a fascinating question since the early days of quantum mechanics. Recently, it has been proposed that the thermodynamic entropy of the long-time statistical ensemble is the stationary entanglement of a large subsystem in an infinite system. Here, we combine this concept with the quasiparticle picture of the entanglement evolution and integrability-based knowledge of the steady state to obtain exact analytical predictions for the time evolution of the entanglement in arbitrary 1D integrable models. These results explicitly show the transformation between the entanglement and thermodynamic entropy during the time evolution. Thus, entanglement is the natural witness for the generalized microcanonical principle underlying relaxation in integrable models.


Author contributions: V.A. and P.C. designed research, performed research, analyzed data, and wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

<sup>1</sup>To whom correspondence should be addressed. Email: valba@sissa.it.

This article contains supporting information online at [www.pnas.org/lookup/suppl/doi:10.1073/pnas.1703516114/-DCSupplemental](http://www.pnas.org/lookup/suppl/doi:10.1073/pnas.1703516114/-DCSupplemental).



**Fig. 1.** Entanglement dynamics after a quantum quench: theoretical scheme. (A) Quasiparticle picture. Full lines denote quasiparticles with maximum velocity emitted in the initial state. Shaded cones show the halo of slower quasiparticles. (B) Main steps to calculate the entanglement dynamics using Bethe ansatz and the quasiparticle picture.

an inhomogeneous quench, and it is at the foundation of the integrable hydrodynamics approach for transport in integrable models (70, 71).

## Results


In a generic interacting integrable model, there are different species of stable quasiparticles corresponding to bound states of an arbitrary number of elementary excitations. Integrability implies that different types of quasiparticles must be treated independently. It is then natural to conjecture that

$$S(t) = \sum_n \left[ 2t \int_{2|v_n|t < \ell} d\lambda v_n(\lambda) s_n(\lambda) + \ell \int_{2|v_n|t > \ell} d\lambda s_n(\lambda) \right], \quad [2]$$

where the sum is over the types of particles  $n$ ,  $v_n(\lambda)$  is their velocity, and  $s_n(\lambda)$  is their entropy. To give predictive power to Eq. 2, in the following, we show how to determine  $v_n(\lambda)$  and  $s_n(\lambda)$  in the Bethe ansatz framework for integrable models. Eq. 2 is straightforwardly generalized to the mutual information between two intervals (see *SI Materials and Methods* and *Fig. S1*).

The eigenstates of Bethe ansatz solvable models are in correspondence with a set of pseudomomenta (rapidities)  $\lambda$ . In the thermodynamic limit, these rapidities form a continuum. One then introduces the particle densities  $\rho_{n,p}(\lambda)$ , the hole (i.e., unoccupied rapidities) densities  $\rho_{n,h}(\lambda)$ , and the total densities  $\rho_{n,t}(\lambda) = \rho_{n,p}(\lambda) + \rho_{n,h}(\lambda)$ . Every set of densities identifies a thermodynamic “macrostate.” This macrostate corresponds to an exponentially large number of microscopic eigenstates, any of which can be used as a “representative” for the macrostate. The total number of representative microstates is  $e^{S_{YY}}$ , with  $S_{YY}$  as the thermodynamic Yang–Yang entropy of the macrostate

$$\begin{aligned} S_{YY} &= s_{YY}L = L \sum_{n=1}^{\infty} \int d\lambda [\rho_{n,t}(\lambda) \ln \rho_{n,t}(\lambda) \\ &\quad - \rho_{n,p}(\lambda) \ln \rho_{n,p}(\lambda) - \rho_{n,h}(\lambda) \ln \rho_{n,h}(\lambda)] \\ &\equiv L \sum_{n=1}^{\infty} \int d\lambda s_{YY}^{(n)}[\rho_{n,p}, \rho_{n,h}](\lambda). \end{aligned} \quad [3]$$




**Fig. 2.** Analytical predictions for the XXZ chain. Entanglement entropy per site  $S/\ell$  vs.  $v_M t/\ell$ , with  $v_M$  as the maximum velocity. Different panels correspond to the different initial states: Néel (A), tilted Néel (B), Dimer (C), and tilted ferromagnet (D and E). Different lines correspond to different  $\Delta$ . For  $\Delta \rightarrow \infty$ ,  $S \rightarrow 0$  for the Néel quench, whereas it saturates in the other cases. Note in E the substantial entanglement increase for  $v_M t/\ell > 1$ . (F) The numerical derivative  $S'(\ell) \times 100$  for the quench in E.


Physically,  $S_{YY}$  corresponds to the total number of ways of assigning the quasimomentum label to the particles, similar to free fermion models.

In the Bethe ansatz treatment of quantum quenches (72–74), local properties of the postquench stationary state are described by a set of densities  $\rho_{n,p}^*(\lambda)$  and  $\rho_{n,h}^*(\lambda)$ . Calculating these densities is a challenging task that has been performed only in a few cases (75–87). From the densities, the thermodynamic entropy of the stationary ensemble Eq. 3 is  $s_{YY}[\rho_{n,p}^*, \rho_{n,h}^*](\lambda)$ . Physically,  $e^{L s_{YY}}$  is the number of microscopic eigenstates entering in a generalized microcanonical ensemble for quenches, in which all of the microstates corresponding to the macrostate have the same probability.

We now present our predictions for the entanglement dynamics (Fig. 1B gives a survey of our theoretical scheme). First, in the stationary state, the density of thermodynamic entropy coincides with that of the entanglement entropy in Eq. 2, as it has been shown analytically for free models (52, 88, 89). This identification implies that  $s_n(\lambda) = s_{YY}[\rho_{n,p}^*, \rho_{n,h}^*](\lambda)$ . Moreover, it is natural to identify the entangling quasiparticles in Eq. 2 with the low-lying excitations around the stationary state  $\rho^*$ . Their group



**Fig. 3.** Bound-state contributions to the entanglement dynamics. On the x axis,  $n$  is the bound-state size. (A and B) Quench from the tilted ferromagnet. (A) Bound-state contributions to steady-state entropy density (second term in Eq. 2). (B) Bound-state contributions to the slope of the entanglement growth for  $t < \ell/v_M$  (first term in Eq. 2). Different histograms denote different tilting angles  $\vartheta$ . All of the data are for  $\Delta = 2$ . (C and D) Same as in A and B for the quench from the tilted Néel state.



**Fig. 4.** Comparison with numerical simulations. Entanglement entropy dynamics after the quench from the Néel state in the XXZ chain. (A) tDMRG results for a chain with  $L=40$  sites and  $\Delta=2$ . Different curves correspond to different subsystem sizes  $\ell$  (accompanying numbers). (B) The entropy saturation values (tDMRG results at  $t \approx 8$ ) as a function of the block length  $\ell$  for several  $\Delta$ . The dashed-dotted lines are the conjectured volume-law behavior  $S \propto s_{\gamma\gamma}^* \ell$  (compare with Eq. 3). (C) The scaling limit:  $S/\ell$  plotted vs.  $v_M t/\ell$ . The continuous curves are the tDMRG results for  $\ell = 5 - 20$ . The diamonds are numerical extrapolations to the thermodynamic limit. The dashed-dotted line is the conjecture Eq. 2.

velocities  $v_n$  depend on the stationary state, because the interactions induce a state-dependent dressing of the excitations. These velocities  $v_n$  can be calculated by Bethe ansatz techniques (90) (*SI Materials and Methods* and Figs. S2 and S3).

To substantiate our idea, we focus on the spin-1/2 anisotropic Heisenberg (XXZ) chain, considering quenches from several low-entangled initial states, namely the tilted Néel state, the Majumdar-Ghosh (dimer) state, and the tilted ferromagnetic state (*Materials and Methods*). For these initial states, the densities  $\rho_{n(h),p}^*$  are known analytically (77–79).


Fig. 2 summarizes the expected entanglement dynamics in the space-time scaling limit, plotting  $S/\ell$  vs.  $v_M t/\ell$ . Interestingly,  $S/\ell$  is always smaller than  $\ln 2$  (i.e., the entropy of the maximally entangled state). For the Néel quench, because the Néel state becomes the ground state of Eq. 4 in the limit  $\Delta \rightarrow \infty$ ,  $S/\ell \approx \ln(\Delta)/\Delta^2$  vanishes, whereas it saturates for all of the other quenches. For the Majumdar-Ghosh state, one obtains  $S/\ell = -1/2 + \ln 2$  at  $\Delta \rightarrow \infty$ . For the tilted ferromagnet with  $\vartheta \rightarrow 0$  (Fig. 2E),  $S/\ell$  is small at any  $\Delta$ , reflecting that the ferromagnet is an eigenstate of the XXZ chain. Surprisingly, the linear growth seems to extend for  $v_M t/\ell > 1$ . However,  $dS/dt$  (Fig. 2F) is flat only for  $v_M t/\ell \leq 1$ , which signals true linear regime only for  $v_M t/\ell \leq 1$ . This peculiar behavior is caused by the large entanglement contribution of the slow quasiparticles. In Fig. 3, we report the bound-state resolved contributions to the entanglement dynamics. Fig. 3A and C focuses on the steady-state entropy (second term in Eq. 2), whereas Fig. 3B and D shows the bound-state contributions to the slope of the linear growth (first term in Eq. 2). The contribution of the bound states, although never dominant, is crucial to ensure accurate predictions.

Fig. 4A shows tDMRG results for  $S(t)$  for the quench from the symmetrized Néel state  $(|\uparrow\downarrow\uparrow \dots\rangle + |\downarrow\uparrow\downarrow \dots\rangle)/\sqrt{2}$ . The data are for the open XXZ chain and subsystems starting from the chain boundary. The qualitative agreement with Eq. 2 is apparent. Fig. 4B reports the steady-state entanglement entropy as a function  $\ell$  (data at  $t \approx 8$  in Fig. 4A). The volume law  $S \propto \ell$  is visible. The dashed-dotted lines in Fig. 4B are fits to  $S \propto s_{\gamma\gamma}^* \ell$ , supporting the equivalence between entanglement and thermodynamic entropy. Fig. 4C focuses on the full-time dependence, plotting  $S/\ell$  vs.  $v_M t/\ell$ . The dashed-dotted line in Fig. 4C is Eq. 2 with  $t \rightarrow t/2$  because of the open boundary conditions (58). Deviations from Eq. 2 because of the finite  $\ell$  are visible. The diamonds in Fig. 4C are numerical extrapolations to the thermodynamic limit. The agreement with Eq. 2 is perfect. Finally, we provide a more stringent check of Eq. 2, focusing on the linear entanglement growth. Fig. 5 shows infinite time-evolving block decimation (iTEBD) results in the thermodynamic limit for

$S'(v_M t)$ , with  $S'(x) \equiv dS(x)/dx$  taken from ref. 91. For all of the quenches, the agreement with Eq. 2 (horizontal lines in Fig. 5) is spectacular.

## Conclusions

The main result of this work is the analytical prediction in Eq. 2 for the time-dependent entanglement entropy after a generic quantum quench in an integrable model. We tested our conjecture for several quantum quenches in the XXZ spin chain, although we expect Eq. 2 to be more general. Additional checks of Eq. 2 (e.g., for the Lieb-Liniger gas) are desirable. It would be also interesting to generalize Eq. 2 to quenches from inhomogeneous initial states, exploiting the recent analytical results (70, 71, 92). Although we are not able yet to provide an *ab initio* derivation of Eq. 2, we find it remarkable that it is possible to characterize analytically the dynamics of the entanglement entropy, whereas its equilibrium behavior is still an open challenge. Finally, we believe that Eq. 2 represents a deep conceptual breakthrough, because it shows in a single compact formula the relation between entanglement and thermodynamic



**Fig. 5.** Comparison with numerical simulations: the short-time regime. Derivative  $S'(v_M t)$  as a function of time. Different panels correspond to different initial states: tilted Néel (A), tilted ferromagnet (B), and dimer state (C);  $\vartheta$  is the tilting angle. In each panel, the different curves are iTEBD results for different  $\Delta$ . The horizontal lines are the conjecture Eq. 2.



entropy for integrable models. An analogous description for nonintegrable systems, where quasiparticles have finite lifetime or do not exist at all, could lead to a deeper understanding of thermalization (19).

## Materials and Methods

The anisotropic spin-1/2 Heisenberg chain is defined by the Hamiltonian

$$H = \sum_{i=1}^L \left[ \frac{1}{2} (S_i^+ S_{i+1}^- + S_i^- S_{i+1}^+) + \Delta \left( S_i^z S_{i+1}^z - \frac{1}{4} \right) \right], \quad [4]$$

where  $S_i^\alpha$  are spin-1/2 operators, and  $\Delta$  is the anisotropy parameter. Here, we considered as prequench initial states the tilted Néel state  $|\vartheta, \nearrow \searrow \dots\rangle \equiv e^{i\vartheta \sum_j S_j^y} |\uparrow \downarrow \dots\rangle$ , the Majumdar-Ghosh (dimer) state  $|MG\rangle \equiv (|\uparrow \downarrow\rangle - |\downarrow \uparrow\rangle)/2^{\otimes L/2}$ , and the tilted ferromagnetic state  $|\vartheta, \nearrow \nearrow\rangle \equiv e^{i\vartheta \sum_j S_j^y} |\uparrow \uparrow \dots\rangle$ . The Heisenberg spin chain is the prototype of all integrable models. Moreover, for all of the initial states considered in this

work, the postquench steady state can be characterized analytically via the macrostate densities  $\rho_{p(h)}^*$ . Specifically, a set of recursive relations for these densities can be obtained (SI Materials and Methods). The group velocities of the low-lying excitations around the steady state (i.e., the entangling quasiparticles) are obtained by solving numerically an infinite set of second-type Fredholm integral equations (details are in SI Materials and Methods).

The numerical data for the postquench dynamics of the entanglement entropy presented in Fig. 4 were obtained using the standard tDMRG (46–49) in the framework of matrix product states. For the implementation, we used the ITENSOR library (itensor.org). The data presented in Fig. 5 are obtained using the iTEBD method (93) and they are a courtesy of Mario Collura.

**ACKNOWLEDGMENTS.** We thank Lorenzo Piroli and Eric Vernier for sharing their results before publication. The iTEBD data presented in Fig. 5 are a courtesy of Mario Collura. We acknowledge support from European Research Council Starting Grant 279391 EDEQS. This project has received funding from the European Union's Horizon 2020 research and innovation program under Marie Skłodowska-Curie Grant 702612 OEMBS.

- von Neumann J (1929) Beweis des ergodensatzes und des H-theorems. *Z Phys* 57: 30–70.
- Jensen RV, Shankar R (1985) Statistical behavior in deterministic quantum systems with few degrees of freedom. *Phys Rev Lett* 54:1879–1882.
- Deutsch JM (1991) Quantum statistical mechanics in a closed system. *Phys Rev A* 43:2046–2049.
- Srednicki M (1994) Chaos and quantum thermalization. *Phys Rev E* 50:888–901.
- Rigol M, Dunjko V, Olshanii M (2008) Thermalization and its mechanism for generic isolated quantum systems. *Nature* 452:854–858.
- Rigol M, Srednicki M (2012) Alternatives to eigenstate thermalization. *Phys Rev Lett* 108:110601.
- D'Alessio L, Kafri Y, Polkovnikov A, Rigol M (2016) From quantum chaos and eigenstate thermalization to statistical mechanics and thermodynamics. *Adv Phys* 65: 239–362.
- Essler FHL, Fagotti M (2016) Quench dynamics and relaxation in isolated integrable quantum spin chains. *J Stat Mech Theor Exp* 2016:064002.
- Kinoshita T, Wenger T, Weiss DS (2008) A quantum Newton's cradle. *Nature* 440: 900–903.
- Hofferberth S, Lesanovsky I, Fischer B, Schumm T, Schiedmayer J (2007) Non-equilibrium coherence dynamics in one-dimensional Bose gases. *Nature* 449:324–327.
- Trotzky S, et al. (2012) Probing the relaxation towards equilibrium in an isolated strongly correlated one-dimensional Bose gas. *Nat Phys* 8:325–330.
- Gring M, et al. (2012) Relaxation and prethermalization in an isolated quantum system. *Science* 337:1318–1322.
- Cheneau M, et al. (2012) Light-cone-like spreading of correlations in a quantum many-body system. *Nature* 481:484–487.
- Langen T, Geiger R, Kuhnert M, Rauer B, Schmiedmayer J (2013) Local emergence of thermal correlations in an isolated quantum many-body system. *Nat Phys* 9:640–643.
- Meinert F, et al. (2013) Quantum quench in an atomic one-dimensional Ising chain. *Phys Rev Lett* 111:053003.
- Fukuhara T, et al. (2013) Quantum dynamics of a mobile spin impurity. *Nat Phys* 9:235–241.
- Langen T, et al. (2015) Experimental observation of a generalized Gibbs ensemble. *Science* 348:207–211.
- Islam R, et al. (2015) Measuring entanglement entropy in a quantum many-body system. *Nature* 528:77–83.
- Kaufman AM, et al. (2016) Quantum thermalization through entanglement in an isolated many-body system. *Science* 353:794–800.
- Calabrese P, Cardy J (2006) Time dependence of correlation functions following a quantum quench. *Phys Rev Lett* 96:136801.
- Polkovnikov A, Sengupta K, Silva A, Vengalattore M (2011) Colloquium: Nonequilibrium dynamics of closed interacting quantum systems. *Rev Mod Phys* 83:863–883.
- Rigol M, Dunjko V, Yurovsky V, Olshanii M (2007) Relaxation in a completely integrable many-body quantum system: An *ab initio* study of the dynamics of the highly excited states of 1D lattice hard-core bosons. *Phys Rev Lett* 98:050405.
- Cazalilla MA (2006) Effect of suddenly turning on interactions in the Luttinger model. *Phys Rev Lett* 97:156403.
- Barthel T, Schollwöck U (2008) Dephasing and the steady state in quantum many-particle systems. *Phys Rev Lett* 100:100601.
- Cramer M, Dawson CM, Eisert J, Osborne TJ (2008) Exact relaxation in a class of nonequilibrium quantum lattice systems. *Phys Rev Lett* 100:030602.
- Cramer M, Eisert J (2010) A quantum central limit theorem for non-equilibrium systems: Exact local relaxation of correlated states. *New J Phys* 12:055020.
- Calabrese P, Essler FHL, Fagotti M (2011) Quantum quench in the transverse-field Ising chain. *Phys Rev Lett* 106:227203.
- Cazalilla MA, Lucchi A, Chung M-C (2012) Thermalization and quantum correlations in exactly solvable models. *Phys Rev E* 85:011133.
- Calabrese P, Essler FHL, Fagotti M (2012) Quantum quench in the transverse field Ising chain II: Stationary state properties. *J Stat Mech Theor Exp* 2012:P07022.
- Sotiriadis S, Fioretto D, Mussardo G (2012) Zamolodchikov-faddeev algebra and quantum quenches in integrable field theories. *J Stat Mech Theor Exp* 2012:P02017.
- Collura M, Sotiriadis S, Calabrese P (2013) Equilibration of a Tonks-Girardeau gas following a trap release. *Phys Rev Lett* 110:245301.
- Collura M, Sotiriadis S, Calabrese P (2013) Quench dynamics of a Tonks-Girardeau gas released from a harmonic trap. *J Stat Mech Theor Exp* 2013:P09025.
- Fagotti M, Essler FHL (2013) Stationary behaviour of observables after a quantum quench in the spin-1/2 Heisenberg XXZ chain. *J Stat Mech Theor Exp* 2013: P07012.
- Kormos M, Collura M, Calabrese P (2014) Analytic results for a quantum quench from free to hard-core one-dimensional bosons. *Phys Rev A* 89:013609.
- Sotiriadis S, Calabrese P (2014) Validity of the GGE for quantum quenches from interacting to noninteracting models. *J Stat Mech Theor Exp* 2014:P07024.
- Ilievski E, et al. (2015) Complete generalized Gibbs ensembles in an interacting theory. *Phys Rev Lett* 115:157201.
- Alba V (2015) Simulating the generalized Gibbs ensemble (GGE): A Hilbert space Monte Carlo approach. arXiv:1507.06994.
- Essler FHL, Mussardo G, Panfil M (2015) Generalized Gibbs ensembles for quantum field theories. *Phys Rev A* 91:051602.
- Cardy J (2016) Quantum quenches to a critical point in one dimension: Some further results. *J Stat Mech Theor Exp* 2016:023103.
- Sotiriadis S (2016) Memory-preserving equilibration after a quantum quench in a 1D critical model. *Phys Rev A* 94:031605.
- Bastianello A, Sotiriadis S (2016) Quasi locality of the GGE in interacting-to-free quenches in relativistic field theories. arXiv:1608.00924.
- Vernier E, Cubero AC (2016) Quasilocality charges and the complete GGE for field theories with non diagonal scattering. arXiv:1609.03220.
- Vidmar L, Rigol M (2016) Generalized Gibbs ensemble in integrable lattice models. *J Stat Mech Theor Exp* 2016:064007.
- Gogolin C, Eisert J (2016) Equilibration, thermalisation, and the emergence of statistical mechanics in closed quantum systems. *Rep Prog Phys* 79:056001.
- Calabrese P, Essler FHL, Mussardo G (2016) Introduction to “quantum integrability in out of equilibrium systems.” *J Stat Mech Theor Exp* 2016:064001.
- White SR, Feiguin AE (2004) Real-time evolution using the density matrix renormalization group. *Phys Rev Lett* 93:076401.
- Daley AJ, Kollath C, Schollwöck U, Vidal G (2004) Time-dependent density-matrix renormalization-group using adaptive effective Hilbert spaces. *J Stat Mech Theor Exp* 2004:P04005.
- Schollwöck U (2005) The density-matrix renormalization group. *Rev Mod Phys* 77:259.
- Schollwöck U (2011) The density-matrix renormalization group in the age of matrix product states. *Ann Phys* 326:96–192.
- Calabrese P, Cardy J (2005) Evolution of entanglement entropy in one-dimensional systems. *J Stat Mech Theor Exp* 2005:P04010.
- Lieb EH, Robinson DW (1972) The finite group velocity of quantum spin systems. *Commun Math Phys* 28:251–257.
- Fagotti M, Calabrese P (2008) Evolution of entanglement entropy following a quantum quench: Analytic results for the XY chain in a transverse magnetic field. *Phys Rev A* 78:010306.
- Eisler V, Peschel I (2008) Entanglement in a periodic quench. *Ann Phys* 17:410–423.
- Nezhadhighi MG, Rajabpour MA (2014) Entanglement dynamics in short and long-range harmonic oscillators. *Phys Rev B* 90:205438.
- Coser A, Tonni E, Calabrese P (2014) Entanglement negativity after a global quantum quench. *J Stat Mech Theor Exp* 2014:P12017.
- Cotler JS, Hertzberg MP, Mezei M, Mueller MT (2016) Entanglement growth after a quench in free scalar field theory. *J High Energy Phys* 11:166.
- Buschikh AS, Fagotti M, Schachenmayer J, Essler FHL, Daley AJ (2016) Entanglement growth and correlation spreading with variable-range interactions in spin and fermionic tunneling models. *Phys Rev A* 93:053620.
- De Chiara G, Montangero S, Calabrese P, Fazio R (2006) Entanglement entropy dynamics in Heisenberg chains. *J Stat Mech Theor Exp* 2006:P03001.
- Läuchli AM, Kollath C (2008) Spreading of correlations and entanglement after a quench in the one-dimensional Bose-Hubbard model. *J Stat Mech Theor Exp* 2008:P05018.

60. Kim H, Huse DA (2013) Ballistic spreading of entanglement in a diffusive nonintegrable system. *Phys Rev Lett* 111:127205.
61. Hubeny VE, Rangamani M, Takayanagi TA (2007) Covariant holographic entanglement entropy proposal. *J High Energy Phys* 2007:062.
62. Abajo-Arriastia J, Aparicio J, Lopez E (2010) Holographic evolution of entanglement entropy. *J High Energy Phys* 1011:149.
63. Albash T, Johnson CV (2011) Evolution of holographic entanglement entropy after thermal and electromagnetic quenches. *New J Phys* 13:045017.
64. Allais A, Tonni E (2012) Holographic evolution of the mutual information. *J High Energy Phys* 1201:102.
65. Callan R, He J-Y, Headrick M (2012) Strong subadditivity and the covariant holographic entanglement entropy formula. *J High Energy Phys* 1206:081.
66. Liu H, Suh SJ (2014) Entanglement tsunami: Universal scaling in holographic thermalization. *Phys Rev Lett* 112:011601.
67. Balasubramanian V, Bernamonti A, Copland N, Craps B, Galli F (2011) Thermalization of mutual and tripartite information in strongly coupled two dimensional conformal field theories. *Phys Rev D Part Fields* 84:105017.
68. Liu H, Suh SJ (2014) Entanglement growth during thermalization in holographic systems. *Phys Rev D Part Fields* 89:066012.
69. Asplund CT, Bernamonti A, Galli F, Hartmann T (2015) Entanglement scrambling in 2D conformal field theory. *J High Energy Phys* 9:110.
70. Castro-Alvaredo OA, Doyon B, Yoshimura T (2016) Emergent hydrodynamics in integrable quantum systems out of equilibrium. *Phys Rev X* 6:041065.
71. Bertini B, Collura M, De Nardis J, Fagotti M (2016) Transport in out-of-equilibrium XXZ chains: Exact profiles of charges and currents. *Phys Rev Lett* 117:207201.
72. Caux J-S, Essler FHL (2013) Time evolution of local observables after quenching to an integrable model. *Phys Rev Lett* 110:257203.
73. Caux J-S (2016) The quench action. *J Stat Mech Theor Exp* 2016:064006.
74. Ilievski E, Quinn E, De Nardis J, Brockmann M (2016) String-charge duality in integrable lattice models. *J Stat Mech Theor Exp* 2016:063101.
75. Bertini B, Schuricht D, Essler FHL (2014) Quantum quench in the sine-Gordon model. *J Stat Mech Theor Exp* 2014:P10035.
76. De Nardis J, Wouters B, Brockmann M, Caux J-S (2014) Solution for an interaction quench in the Lieb-Liniger Bose gas. *Phys Rev A* 89:033601.
77. Wouters B, et al. (2014) Quenching anisotropic Heisenberg chain: Exact solution generalized Gibbs ensemble predictions. *Phys Rev Lett* 113:117202.
78. Pozsgay B, et al. (2014) Correlations after quenching XXZ spin chain: Fail generalized Gibbs ensemble. *Phys Rev Lett* 113:117203.
79. Piroli L, Vernier E, Calabrese P (2016) Exact steady states for quantum quenches in integrable Heisenberg spin chains. *Phys Rev B* 94:054313.
80. Brockmann M, et al. (2014) Quench action approach for releasing the neel state into the spin-1/2 XXZ chain. *J Stat Mech Theor Exp* 2014:P12009.
81. Mestyán M, Pozsgay B, Takács G, Werner MA (2015) Quenching the XXZ spin chain: Quench action approach versus generalized Gibbs ensemble. *J Stat Mech Theor Exp* 2015:P04001.
82. Bucciantini L (2016) Stationary state after a quench to the Lieb-Liniger from rotating BECs. *J Stat Phys* 164:621–644.
83. Bertini B, Piroli L, Calabrese P (2016) Quantum quenches in the sinh-Gordon model: Steady state and one-point correlation functions. *J Stat Mech Theor Exp* 2016:063102.
84. Alba V, Calabrese P (2016) The quench action approach in finite integrable spin chains. *J Stat Mech Theor Exp* 2016:043105.
85. Piroli L, Calabrese P, Essler FHL (2016) Multiparticle bound-state formation following a quantum quench to the one-dimensional Bose gas with attractive interactions. *Phys Rev Lett* 116:070408.
86. Piroli L, Calabrese P, Essler FHL (2016) Quantum quenches to the attractive one-dimensional Bose gas: Exact results. *SciPost Phys* 1:001.
87. Piroli L, Vernier E, Calabrese P, Rigol M (2017) Correlations and diagonal entropy after quantum quenches in XXZ chains. *Phys Rev B* 95:054308.
88. Collura M, Kormos M, Calabrese P (2014) Stationary entanglement entropies following an interaction quench in 1D Bose gas. *J Stat Mech Theor Exp* 2014:P01009.
89. Kormos M, Bucciantini L, Calabrese P (2014) Stationary entropies after a quench from excited states in the Ising chain. *Europhys Lett* 107:40002.
90. Bonnes L, Essler FHL, Läuchli AM (2014) Light-cone dynamics after quantum quenches in spin chains. *Phys Rev Lett* 113:187203.
91. Fagotti M, Collura M, Essler FHL, Calabrese P (2014) Relaxation after quantum quenches in the spin-1/2 Heisenberg XXZ chain. *Phys Rev B* 89:125101.
92. Dubail J, Stéphan J-M, Viti J, Calabrese P (2017) Conformal field theory for inhomogeneous one-dimensional quantum systems: The example of non-interacting Fermi gases. *SciPost Phys* 2:002.
93. Vidal G (2007) Classical simulation of infinite-size quantum lattice systems in one spatial dimension. *Phys Rev Lett* 98:070201.
94. Takahashi M (1999) *Thermodynamics of One-Dimensional Solvable Models* (Cambridge Univ Press, Cambridge, UK).
95. Yang CN, Yang CP (1969) Thermodynamics of a one-dimensional system of bosons with repulsive delta function interaction. *J Math Phys* 10:1115–1122.
96. De Nardis J, Piroli L, Caux J-S (2015) Relaxation dynamics of local observables in integrable systems. *J Phys A* 48:43FT01.
97. Calabrese P, Cardy J (2007) Quantum quenches in extended systems. *J Stat Mech Theor Exp* 2007:P06008.
98. Gambassi A, Calabrese P (2011) Quantum quenches as classical critical films. *Europhys Lett* 95:66007.
99. Delfino G (2014) Quantum quenches with integrable pre-quench dynamics. *J Phys A* 47:402001.
100. Collura M, Calabrese P, Essler FHL (2015) Quantum quench within the gapless phase of the spin-1/2 Heisenberg XXZ spin-chain. *Phys Rev B* 92:125131.
101. Calabrese P, Cardy J (2016) Quantum quenches in 1+1 dimensional conformal field theories. *J Stat Mech Theor Exp* 2016:064003.
102. Barbiero L, Dell'Anna L (2016) Spreading of correlations in quenched repulsive and attractive one dimensional lattice system. arXiv:1609.07387.
103. Kormos M, Collura M, Takács G, Calabrese P (2016) Real time confinement following a quantum quench to a non-integrable model. *Nat Phys* 13:246–249.
104. Bertini B, Essler FHL, Groha S, Robinson N (2016) Thermalization and light cones in a model with weak integrability breaking. *Phys Rev B* 94:245117.
105. Chiocchetta A, Tavora M, Gambassi A, Mitra A (2016) Short-time universal scaling and light-cone dynamics after a quench in an isolated quantum system in d spatial dimensions. *Phys Rev B* 94:134311.
106. Asplund CT, Bernamonti A (2014) Mutual information after a local quench in conformal field theory. *Phys Rev D Part Fields* 89:066015.
107. Leichenauer S, Moosa M (2015) Entanglement tsunami in (1+1)-dimensions. *Phys Rev D Part Fields* 92:126004.
108. Fagotti M, Calabrese P (2010) Entanglement entropy of two disjoint blocks in XY chains. *J Stat Mech Theor Exp* 2010:P04016.