

Figure 1: a Cortical regions (Allen CCF, Beryl level) assigned to the default mode network by four rodent resting-state studies. Lu et al. (2012): rat, BOLD rsfMRI under dexmedetomidine/isoflurane, group ICA (30 components); rat areas mapped to mouse (PtP → VISa/VISrl, V2M → VISam/VISpm; dorsal CA1 tentative). Whitesell et al. (2021): mouse, BOLD rsfMRI under halothane, group ICA (5 components) registered to CCFv3; regions with >50% of voxels in the core DMN mask. Grandjean et al. (2020): 17 mouse rsfMRI datasets, ACA seed and group ICA; DMN components named prefrontal (PFC), cingulate/RSP and temporal association areas; † prefrontal subregions not specified, mapped here to PL, ILA and ORB. Mandino et al. (2022): mouse rsfMRI co-activation patterns and viral-tracer projection similarity to an ACA seed; cortical DMN labelled ACA and RSP (a medial visual extension is not labelled). Subcortical DMN elements (dorsal striatum, dorsal and lateral thalamus, zona incerta, periaqueductal grey, peri-hippocampal areas) are not shown. Bold: the conservative core used in this report (ACA, RSP, PL, ORBm). ACA and RSP are in all four maps; PL and ORB in three (Mandino et al. name only cingulate regions). ORBl and ORBvl have the same support as ORBm and could be added.