


Figure 1: **a** IBL passive protocol after the task (iblrig `passiveChoiceWorld`): frame2ttl spacers separate the spontaneous period, during which Bonsai clears the screen to grey with the sync square off and shows no stimulus (`SPONTANEOUS_ACTIVITY_SECONDS` = 600), receptive-field mapping (sparse noise) and the replay of task stimuli, valve clicks and tones. Task length is schematic. The interval comes from `_ibl_passivePeriods.intervalsTable`. **b** Grey-screen duration over the 396 BWM sessions (130 mice) that have it; 600 s in 86%. **c** Delay between the end of the last task trial and the start of the grey screen (values above 8 min clipped). **d** Fraction of the period that is still: 50 ms bins without wheel movement (>0.05 rad/s), licks or motion energy above the session's 60th percentile, with a one-bin margin (the first second is skipped); median 6.5 min of 10. In the 35 sessions without motion energy, stillness rests on wheel and licks only (median fraction 0.81 vs 0.67). **e** Example session with median stillness: wheel speed, left-camera motion energy (normalised to the session median; dashed, the stillness threshold; wheel in rad/s), the resulting still bins, and a raster of the 20 neurons with the highest firing rate in this window (of 236 good units on both probes; 0.5 s bins, each row scaled between its 1st and 99th percentile, black = high; labels give Beryl region, probe and mean rate). This session's probes are in cerebellum and brainstem, so these are not DMN neurons.