


Figure 2: **a** IBL passive protocol after the task (iblrig `passiveChoiceWorld`): frame2ttl spacers separate the spontaneous period, during which Bonsai clears the screen to grey with the sync square off and shows no stimulus (`SPONTANEOUS_ACTIVITY_SECONDS` = 600), receptive-field mapping (sparse noise) and the replay of task stimuli, valve clicks and tones. Task length is schematic. The interval comes from `_ibl_passivePeriods.intervalsTable`. **b** Grey-screen duration over the 396 BWM sessions (130 mice) that have it; 600 s in 86%. **c** Delay between the end of the last task trial and the start of the grey screen (values above 8 min clipped). **d** Fraction of the period that is still: 50 ms bins without wheel movement ($>$0.05 rad/s), licks or motion energy above the session's 60th percentile, with a one-bin margin (the first second is skipped); median 6.5 min of 10. In the 35 sessions without motion energy, stillness rests on wheel and licks only (median fraction 0.81 vs 0.67). **e** Example session with neurons in the conservative DMN core (CSH_ZAD_019, 55 of 141 good units in RSPagl, RSPd and RSPv; still fraction 0.70, close to the median): wheel speed, left-camera motion energy (normalised to the session median; dashed, the stillness threshold; wheel in rad/s) and the resulting still bins. Orange band: the 10 s shown in f. **f** Zoom into 10 s of e at the native 50 ms resolution: wheel speed (dashed, the 0.05 rad/s movement threshold), motion energy (dashed, stillness threshold), still bins and spike counts per 50 ms bin of the 20 most active core-DMN neurons of the session (each row scaled to its 99th percentile over the whole period, black = high; labels give region, probe and mean rate). The window was chosen automatically as the one after the first 30 s whose still fraction is closest to 0.5 and that contains wheel movement.