

Figure 8: **a** Grey-screen co-fluctuation of region signals (still bins only; wheel, lick and motion-energy regression), as residual from pairs at the same distance and probe relation. Difference DMN-DMN minus DMN-other cortical pairs for each region set, all pairs (circles) and pairs on different probes only (squares), 1 s (black) and 0.5 s (grey) bins; 95% mouse-bootstrap CI; number of mice with a DMN-DMN pair in brackets. Orange tick: the 1 s estimate after dropping the single most coupled DMN-DMN pair. The distance baseline uses cortical and CA1 pairs only. With about 20 such contrasts, CIs that just exclude 0 would not survive correction. The conservative core has only 4 cross-probe DMN-DMN pairs (4 mice), too few to estimate. **b** Every DMN edge (conservative core, Whitesell et al. or Lu et al.) recorded in $\geq$3 mice: mean over mice of the residual coupling, 95% bootstrap CI; brackets give the number of mice and the median distance between the two regions. Blue: both regions in the conservative core; orange: other literature DMN edges. **c** Cross-validated rest-preference network: cortical regions ranked by grey-screen rest preference (RI) in one half of the mice (terciles), coupling measured in the other half; mean and 2.5–97.5% range over 400 half-splits; rest–rest exceeds rest–task in 52% of splits. **d** Within-session contrast DMN-DMN minus DMN-other, mean over mice (number of mice above bars).