

Figure 1: Wheel-speed correlates are rare in the mesoscope but present: among thousands of neurons per session the best 10 track wheel speed in held-out trials as well as the best BWM units, but a random set of 30 neurons, as in a small spatial bin, contains far fewer of them than a BWM set of the same size. Sessions with $\geq$30 neurons and $\geq$50 trials. **a** Fraction of neurons per session whose correlation with wheel speed (over all wheel bins) exceeds 0.1 in both the even and the odd trials: median 3.5% (mesoscope, 151 sessions), 21.7% (BWM isocortex, 137), 18.2% (BWM units within 640 μm of the surface, 29). **b** Correlation with wheel speed in the odd trials of the mean z-scored activity of the 10 neurons that correlate best on the even trials, against the number of neurons in the session (one dot per session). **c** The same with all neurons of a session (median mesoscope 0.50, BWM 0.47, p = 0.05) and with sessions subsampled to 30 random neurons (mean over 20 draws; 0.15 vs 0.37, $p < 10^{-32}$). Boxes: quartiles; whiskers: 1.5 inter-quartile ranges. **d** Same against trial-specific speed (wheel speed minus its per-bin mean over the even trials): 0.21 vs 0.19 (all neurons), 0.08 vs 0.15 (30 neurons).