Reducing Green House Gas Emissions With Congestion Control

Michael Welzl, Ozgu Alay, Peyman Teymoori, Safiqul Islam October 31, 2022

1 Internet energy saving: Why is it important?

By enabling online meetings, replacing snail mail with email, etc., the Internet helps to reduce Green House Gas (GHG) emissions. However, at the same time, the Internet itself needs a significant amount of energy, which translates into GHG emissions, and hence an effect on global warming. Appendix A is an attempt to estimate the magnitude of the Internet's impact, on the basis of well-known prior studies; it arrives at an approximate range of 0.5% - 1.17% of worldwide GHG contributions, and an approximate global warming impact between 1/7th and 1/3rd of the aviation industry.


While these numbers are dwarved by, for example, road traffic or energy use in buildings, the absolute amount is still large, and reducing it is important. The Internet offers a particularly interesting opportunity here because, quite different from e.g. the aviation industry, it is possible to achieve an almost immediate worldwide impact by introducing standards and fostering their implementation. Operating System updates, for example, are commonly rolled out to millions of devices within a short time period.

2 Why congestion control?

It is extremely common to regard energy consumption as a trade-off, where good performance means high energy usage and vice versa. Such a trade-off is made explicit to the user by smart phones offering a "power saving" mode.

Congestion control is uniquely placed in that regard: when done efficiently, it reduces the Flow Completion Time (FCT), which is a very well accepted congestion control metric [DM06]. A shorter FCT means that transfers finish faster, which translates into *both* a performance gain and longer sleep times for devices, i.e. energy reduction — a win-win, attractive for everyone to permanently enable.

3 Is this real?






Figure 1: Data transfer energy consumption when transferring 10 (left) and 80 packets (right). Transfers using larger IW values finish faster, and they are more energy efficient.

Yes. Figure 1 shows a result of a test where we fed packet traces from data transfers with different values of TCP's Initial Window (IW) parameter into EnergyBox [VNTP14], a tool that provides an

estimation of an end hosts's Wi-Fi power consumption related to a certain traffic pattern. EnergyBox has been validated to achieve 95-99% accuracy, and it has been used by Spotify for quantifying their mobile application's energy consumption.

The focus on end hosts may seem surprising, but there are in fact good reasons to reduce the energy usage of "last mile" equipment in general; see Section A.1.3 for an elaboration.

4 What can we do?

Generally, methods to decrease the FCT can help (e.g., 0-RTT connection setup as in QUIC [IT21] or TCP Fast Open [CCRJ14], or increasing the IW when this is appropriate). It is probably useful to go beyond a simple general update of the IW constant, and apply dynamic methods—e.g., the authors of [NZL+19] describe a reinforcement learning based approach to tune the IW, successfully deployed over several years at Baidu. Other ways to decrease the FCT include the re-use of parameters from ongoing connections—e.g., by coupling congestion control [IWH+18] when appropriate, or multiplexing flows over a single connection as with QUIC's multi-streaming. Such mechanisms can work better when they are applied inside the network (e.g., in proxies) as the multiplexing potential increases.

It is also interesting to entertain the thought of energy usage as a *metric* for congestion control, in the same vein as, for example, throughput, delay, packet loss, and fairness. Energy usage is generally tied to the specific hard- and software of a system, but applying it as a metric would require designing a more abstract measure. Perhaps, from measuring multiple systems, we could find a common relationship between the energy use of sending vs. receiving vs. calculation overhead, and maybe such relationships could be represented by a new "energy usage" metric?

We refer to [CWT⁺22] for a broader discussion of energy saving possibilities in the Internet, including some that are related to congestion control. Further details about the test described in Section 3, including an explanation of why these results are still applicable for modern smartphones, are given in [Wel22], and a brief presentation is also planned for the meeting of the Internet Congestion Congestion Control Research Group (ICCRG) at IETF-115 in London on November 8, 2022.

A Green House Gas Emissions: the Internet vs. Aviation

In public discourse, air travel has commonly been associated with an activity that should urgently be reduced to help slow down global warming. The impact of the Internet on global warming is much less known, and deserves more attention. It is therefore useful to relate the Green House Gas (GHG) emissions that are caused by the Internet to the aviation industry. Hence, here, we offer some facts about this relationship.

A.1 Electricity and the Internet

Table 1 is used as the basis for our derivation of Internet GHG emissions in the following subsections. In this section, we use GHG and CO₂ interchangeably, as CO₂ makes up the vast majority of GHG emissions from the energy sector [EPA22].

Line	Fact	Year	Source
1	Power generation: 13 Gt CO2, i.e. 38% of energy CO2	2018	[IEA19]
2	Energy = 37.6 Gt CO2 , 75.6% of GHG emissions	2022	[GFV22]
3	Heat & Electricity = 15.8 Gt CO2, 31.8% of GHG emissions	2022	[GFV22]
4	USA GHG contribution of electricity: 25%	2019	[EPA22]
5	ICT CO2 = 2.7%	2020	[LCW19], citing "SMARTer 2030" report
6	ICT CO2: between 2.1% and 3.9%	2021	[FBLW ⁺ 21]
7	ICT sector = 6% of electricity, telecom = 1.8%		[Wel16], citing [LHV ⁺ 12]
8	Networks CO2 share of ICT: 24%	2020	2020 data in Fig. 3 of [FBLW ⁺ 21]
9	Internet = between 1.1% and 1.9% of electricity	2011	[RM11]

Table 1: Internet related GHG emissions source data.

A.1.1 How much does electricity contribute?

Lines 1 and 2: Energy contributes 75.6% of all GHG emissions. Electricity makes up 38% of the CO₂ contribution of energy. Hence, by combining these data from 2018 and 2022, we can derive that electricity contributes 28.7% of the total GHG emissions. Line 3: electricity contributes 31.8% of the total. Line 4: In the USA, the electricity contribution is 25% of the total. However, in the USA, transport is a much larger contributor to energy than the global average [RRR20], and hence these 25%, while probably too small, can serve as a reasonably reliable lower bound.

From multiple sources, we arrive at the following lower and upper bounds for the global contribution of electricity to GHG emissions:

$$MINGHG_{POWER} = 25\%, MAXGHG_{POWER} = 31.8\%$$

A.1.2 How much does ICT contribute?

Lines 5 and 6: The ICT CO₂ contribution is between 2.1% and 3.9%. From line 7:, ICT constituting 6% of electricity means between a CO₂ constribution between 1.5% (using $MINGHG_{POWER}$) and 1.9% (using $MAXGHG_{POWER}$). Both of these values are significantly below the numbers from lines 5 and 6, but the study underlying table line 7 [LHV⁺12] is also much older than the others (2012). We therefore resort to using this reference only for the relationship between telecommunication and ICT in the following discussion.

From lines 5 and 6 in Table 1, we get:

$$MINGHG_{ICT}=2.1\%, MAXGHG_{ICT}=3.9\%$$

A.1.3 How much does the Internet contribute?

At this point, things become extremely vague. Limiting the focus to the core and access network, data centers and undersea cables, and not considering embodied energy (the energy used for device production), the authors of [AMKF18] have shown that increasing energy efficiency has led to a decline in energy usage per byte. In two large ISPs, this has offset the exponential growth of Internet traffic, leading to roughly linear energy use in one case and even a slight decline in the other [KM21].

This matches a very recent study, which, using a similar focus, finds that the electricity consumption remained approximately constant from 2015 to 2018 in the networks of 15 telecom network operators with headquarters in Europe, representing operations in 21 European countries, and additionally including their operations in 27 countries overseas [LMBL22].

Importantly, both of these studies exclude enterprise networks and home networks with both Customer Premises Equipment (CPE) (e.g., WiFi access points) and User Equipment (UE) (e.g., laptops). When the focus is broadened, studies differ in various aspects, such as:

- What types of UE are considered? E.g., TVs cost much energy, but are not normally a part of such studies.
- Is CPE considered?
- Is embodied energy considered?
- Are data centers considered? They constitute a large part of the Internet's energy, yet due to a strong (monetary) incentive, large data centers now quickly move towards operation on sustainable energy. Microsoft, for example, promises to shift to 100 percent supply of renewable energy for all its data centers, buildings, and campuses by 2025.

Here, we consider three studies from Table 1. Line 8: 24% of $MINGHG_{ICT}$ and $MAXGHG_{ICT}$ means a 0.5% and 0.9% GHG contribution, respectively; this excludes UE and data centers, and it is not clear whether embodied (i.e., device production) energy is considered. Line 7: As we stated earlier, this line stands out for its age, and we therefore only use the telecom/ICT relationship from here (30%) instead of the absolute energy numbers. Using $MINGHG_{ICT}$ and $MAXGHG_{ICT}$ this yields a GHG contribution between 0.63% and 1.17%, focusing on use phase electricity consumption of telecom operator networks, office networks and customer premises equipment. Line 9: between 1.1% and 1.9% of $MINGHG_{POWER}$ becomes a GHG contribution range from 0.28% to 0.48%, and for $MAXGHG_{POWER}$, these values become 0.35% and 0.6%, respectively, i.e. the total range becomes 0.28% – 0.6%. We report about this study despite its age because it is specifically focused on the Internet, and attempts to consider all elements: embedded energy, the core network, CPE, and UE.

$_{ m GHG}$ %	Embodied	$\mathbf{U}\mathbf{E}$	Data centers	Year	Source
0.5 - 0.9	?	N	N	2020	2020 data in Fig. 3 of [FBLW ⁺ 21]
0.63 - 1.17	N	N	N	2012	[LHV ⁺ 12]
0.28 - 0.6	Y	Y	Y	2011	[RM11]

Table 2: Three studies on network energy use, with age and coverage. Due to the age of these studies, only the networks/ICT relationship was considered from [LHV⁺12], and we will not further work with the data from [RM11].

From Table 2, we arrive at a range for the Internet's GHG contribution:

$$MINGHG_{INTERNET} = 0.5\%, MAXGHG_{INTERNET} = 1.17\%$$

A.2 Aviation

Recent data on the global warming impact of aviation is available from [Rit20]. Here, the analysis is complicated by the fact that not only CO_2 plays a role (around 2.5% of global emissions). Warming effects are also caused by other gases that airplanes leave in the atmosphere, including water vapor. The entire impact, called "effective radiative forcing", accounts for roughly 3.5% of global warming.

A.3 The Internet vs. Aviation

Putting aviation data from the previous section in relation to $MINGH_{INTERNET}$ and $MAXGH_{INTERNET}$, we can arrive at the following conclusion:

Considering only CO_2 emissions from airplanes, the Internet's contribution to global warming is between 1/5th and approximately half of the contribution of the aviation industry.

¹ https://blogs.microsoft.com/blog/2020/01/16/microsoft-will-be-carbon-negative-by-2030/

Considering the entire airplane effect of "effective radiative forcing", the Internet's contribution to global warming is between 1/7th and 1/3rd of the contribution of the aviation industry.

A.4 What does this mean?

First, all of the reported numbers are very rough, and to be taken as such — they are ball-park figures that give a "feel" of the magnitude. While they are all small compared to the "bigger picture" (compared to, e.g., road traffic or energy use in buildings [RRR20]), they are nevertheless significant.

Different from large industries such as aviation, the Internet offers the possibility to have an immediate *global* impact via a change of standards. There are many examples of ideas that were developed by small research groups or even individuals, and were quite quickly deployed in the major Operating Systems, with a direct effect on millions of devices all over the world.

Due to its potential for impact at a global scale, energy-saving Internet research focused on standards can cause a very significant global reduction of GHG emissions.

As a side point, we stress that our analysis does *not* conclude that video conferencing would be almost as harmful as flying (or that, conversely, flying would equate 3-7 video conferences). Generally, the number of people flying is much smaller than the users of the Internet, and thus, the per-person CO_2 footprint of an individual flight is much higher [Rit20]. Regarding meetings, there is ample evidence that videoconferencing is better for the planet; e.g., the authors of [OMS14] find that videoconferencing takes at most 7% of the energy/carbon of an in-person meeting. Such per-person or per-meeting analyses are entirely different in scope.

A.5 Misinformation

Plenty of misinformation has been published regarding the Internet's energy use, often exaggerating it. This can be quite harmful: it reduces the credibility of proper analyses, and it makes efforts to reduce the Internet's energy seem questionable [KM21]. The present document is intended to place the real energy usage into a more realistic range, hopefully contributing to a better understanding of the issue and the value of improving things.

Some notable examples of misinformation are:

- "The energy used in our digital consumption collectively emits the equivalent amount of carbon as the entire airline industry." according to our analysis, this is almost certainly an exaggeration.
- "Perhaps unsurprisingly, the footprint of an email also varies dramatically, from 0.3g CO2e for a spam email to 4g (0.14oz) CO2e for a regular email and 50g (1.7oz) CO2e for one with a photo or hefty attachment, according to Mike Berners-Lee, a fellow at Lancaster University who researches carbon footprints. These figures, however, were crunched by Berners-Lee 10 years ago. Charlotte Freitag, a carbon footprint expert at Small World Consulting, the company founded by Berners-Lee, says the impact of emailing may have gone up." this statement appeard in a BBC article from March 2020.³ A later blog post⁴ states: "With new data in hand, Berners-Lee estimates that an individual email can be anywhere from from 0.03g CO2e to 26g CO2e.". Finally, another BBC article⁵ states: "Mike Berners-Lee, a respected professor on the topic whose research was used in the Ovo Energy work, told the Financial Times it was based on 'back-of-the-envelope' maths from 2010 and while useful to start conversations, there were bigger questions."
- "A typical office worker sends and receives around 140 emails per day, which, over the course of a year, creates as much CO2 as flying from London to Bruges or watching 955 movies." here, the numbers simply do not add up. An average email size may be about 75 KB⁷, which means that the amount of data over a year becomes 3.65 GB. A NetFlix movie, for example, can

²https://www.climateimpact.com/news-insights/insights/infographic-carbon-footprint-internet/

³https://www.bbc.com/future/article/20200305-why-your-internet-habits-are-not-as-clean-as-you-think

 $^{^{4} \}texttt{https://sedna.com/blog/the-carbon-footprint-of-carbon-copies-how-does-email-affect-the-environment/of-carbon-car$

⁵https://www.bbc.com/news/technology-55002423

 $^{^6}$ https://www.cwjobs.co.uk/insights/environmental-impact-of-emails/

https://www.lifewire.com/what-is-the-average-size-of-an-email-message-1171208

be between around 0.3 GB and 7 GB per hour in size⁸, depending on the chosen quality. Being "friendly" towards the calculation would mean to assume only 1.5 hours per movie and using the lowest quality in this range, which then means 8 movies instead of 955.

In the examples above, we have quoted online media both as a source of misinformation and to refute it; since the information is contradictory, either one of the sources contains misinformation. More examples are available in the slides⁹ from Jonathan Koomey's keynote speech at the IEEE iTherm 2021 conference.

References

- [AMKF18] Joshua Aslan, Kieren Mayers, Jonathan G. Koomey, and Chris France. Electricity intensity of internet data transmission: Untangling the estimates. *Journal of Industrial Ecology*, 22(4):785–798, 2018.
- [CCRJ14] Yuchung Cheng, Jerry Chu, Sivasankar Radhakrishnan, and Arvind Jain. TCP Fast Open. RFC 7413, December 2014.
- [CWT⁺22] Alexander Clemm, Cedric Westphal, Jeff Tantsura, Laurent Ciavaglia, and Marie-Paule Odini. Challenges and Opportunities in Green Networking. Internet-Draft draft-cx-greenps-01, Internet Engineering Task Force, October 2022. Work in Progress.
- [DM06] Nandita Dukkipati and Nick McKeown. Why flow-completion time is the right metric for congestion control. SIGCOMM Comput. Commun. Rev., 36(1):59–62, jan 2006.
- [EPA22] EPA. Sources of Greenhouse Gas Emissions. https://www.epa.gov/ghgemissions/sources-greenhouse-gas-emissions, 2022. [Online; accessed 13 October 2022].
- [FBLW⁺21] Charlotte Freitag, Mike Berners-Lee, Kelly Widdicks, Bran Knowles, Gordon S. Blair, and Adrian Friday. The real climate and transformative impact of ict: A critique of estimates, trends, and regulations. *Patterns*, 2(9):100340, 2021.
- [GFV22] Mengpin Ge, Johannes Friedrich, and Leandro Vigna. 4 charts explain green-house gas emissions by countries and sectors. https://www.wri.org/insights/4-charts-explain-greenhouse-gas-emissions-countries-and-sectors, 2022. [Online; accessed 14 October 2022].
- [IEA19] IEA. Global Energy & CO2 Status Report 2019. https://www.iea.org/reports/global-energy-co2-status-report-2019, 2019. [Online; accessed 14 October 2022].
- [IT21] Jana Iyengar and Martin Thomson. QUIC: A UDP-Based Multiplexed and Secure Transport. RFC 9000, May 2021.
- [IWH⁺18] Safiqul Islam, Michael Welzl, Kristian Hiorth, David Hayes, Grenville Armitage, and Stein Gjessing. ctrltcp: Reducing latency through coupled, heterogeneous multi-flow tcp congestion control. In *IEEE INFOCOM 2018 - IEEE Conference on Computer Commu*nications Workshops (INFOCOM WKSHPS), pages 214–219, 2018.
- [KM21] Jonathan Koomey and Eric Masanet. Does not compute: Avoiding pitfalls assessing the internet's energy and carbon impacts. *Joule*, 5(7):1625–1628, 2021.
- [LCW19] Josip Lorincz, Antonio Capone, and Jinsong Wu. Greener, energy-efficient and sustainable networks: State-of-the-art and new trends. *Sensors*, 19(22), 2019.
- [LHV⁺12] Sofie Lambert, Ward Van Heddeghem, Willem Vereecken, Bart Lannoo, Didier Colle, and Mario Pickavet. Worldwide electricity consumption of communication networks. Opt. Express, 20(26):B513–B524, Dec 2012.

 $^{{}^{8} \}texttt{https://www.buckeyebroadband.com/support/internet/how-much-data-does-streaming-netflix-use}$

⁹https://www.koomey.com/post/652805139647840256

- [LMBL22] Dag Lundén, Jens Malmodin, Pernilla Bergmark, and Nina Lövehagen. Electricity consumption and operational carbon emissions of european telecom network operators. Sustainability, 14(5), 2022.
- [NZL⁺19] Xiaohui Nie, Youjian Zhao, Zhihan Li, Guo Chen, Kaixin Sui, Jiyang Zhang, Zijie Ye, and Dan Pei. Dynamic tcp initial windows and congestion control schemes through reinforcement learning. *IEEE Journal on Selected Areas in Communications*, 37(6):1231–1247, 2019.
- [OMS14] Dennis Ong, Tim Moors, and Vijay Sivaraman. Comparison of the energy, carbon and time costs of videoconferencing and in-person meetings. *Computer Communications*, 50:86–94, 2014. Green Networking.
- [Rit20] Hannah Ritchie. Climate change and flying: what share of global co2 emissions come from aviation? https://ourworldindata.org/co2-emissions-from-aviation, 2020.
- [RM11] Barath Raghavan and Justin Ma. The energy and emergy of the internet. In *Proceedings* of the 10th ACM Workshop on Hot Topics in Networks, HotNets-X, New York, NY, USA, 2011. Association for Computing Machinery.
- [RRR20] Hannah Ritchie, Max Roser, and Pablo Rosado. Co2 and greenhouse gas emissions. https://ourworldindata.org/co2-and-other-greenhouse-gas-emissions, 2020.
- [VNTP14] Ekhiotz Jon Vergara, Simin Nadjm-Tehrani, and Mihails Prihodko. Energybox: Disclosing the wireless transmission energy cost for mobile devices. Sustainable Computing: Informatics and Systems, 4(2):118–135, 2014.
- [Wel16] Marcus K. Weldon. The Future X Network: A Bell Labs Perspective. CRC Press, Inc., USA, 2016.
- [Wel22] Michael Welzl. Not a trade-off: On the wi-fi energy efficiency of effective internet congestion control. In 2022 17th Wireless On-Demand Network Systems and Services Conference (WONS), pages 1–4, 2022.