

A Structural Horizon Transition in a Representative TIG Geometry: A Test Against Quadratic $f(R)$ Gravity

Kai Stefan Dietrich
Independent Research

Revised draft – wave 4, 2 October 2026

Abstract

We analyze the horizon structure of a specified static, spherically symmetric representative geometry within the Topological Integrity Gravity (TIG) research program. The chosen regular mass profile yields the exact dimensionless horizon equation $x^3 - x^2 + \beta^3 = 0$. We derive its discriminant, positive-root sectors, and critical point $x_c = 2/3$, $\beta_c = (4/27)^{1/3}$. The two positive roots coalesce in a nondegenerate local fold with square-root separation. The cubic is a consequence of the chosen mass profile, not a universal requirement of fold bifurcations or dimensional consistency. A direct trace test also excludes this nontrivial representative metric as an exact vacuum solution of constant-coefficient quadratic metric $f(R)$ gravity. This geometry is a reparameterized Hayward metric. The Schwarzschild limit and bounded-curvature core remain geometric properties. A prescribed reflecting model yields a conditional inverse-square-root round-trip delay on the horizonless side. Physical reflection, dynamical completion and observational forecasts remain separate tasks.

1 Introduction

General Relativity provides an accurate description of gravitational phenomena across a wide range of scales [1, 2]. Extensions such as $f(R)$ gravity introduce higher-curvature corrections and additional degrees of freedom [3, 4].

The number of positive horizon roots can depend on a control parameter and change at a critical coalescence. This parameter-dependent root structure is the subject of the present static analysis.

We analyze a specified representative metric and test its compatibility with quadratic metric $f(R)$ gravity. The algebraic horizon transition is established for that metric, while the direct vacuum test gives a negative result. Counting static Killing-horizon roots does not establish dynamical horizon formation.

2 Consistency Constraint and Model Choice

Structural consistency and the Schwarzschild limit constrain a model without uniquely specifying its mass profile or the degree of its horizon equation. A local fold equilibrium equation is quadratic, $y^2 - \lambda = 0$; a cubic potential can generate it by differentiation [5]. Indeed, $x^2 - x + \lambda = 0$ has a root $x = 1$ at $\lambda = 0$ and a fold at $(x, \lambda) = (1/2, 1/4)$. Lower-order equations therefore can capture fold behavior. The cubic below follows specifically from the selected regular mass profile, rather than from a universal bifurcation requirement.

3 Quadratic Gravity as a Consistency Test

We use the metric formalism, signature $(-, +, +, +)$, and geometrized units $G = c = 1$. To retain the convention $\mu = 16\pi\alpha$, the action is

$$S = \int d^4x \sqrt{-g} \left[\frac{R - 2\Lambda}{16\pi} + \alpha R^2 \right] + S_{\text{matter}} = \frac{1}{16\pi} \int d^4x \sqrt{-g} f(R) + S_{\text{matter}}, \quad (1)$$

where

$$f(R) = R - 2\Lambda + \mu R^2, \quad \mu = 16\pi\alpha. \quad (2)$$

Thus α here is the coefficient outside the Einstein prefactor. If the entire expression $R - 2\Lambda + \alpha_{\text{in}} R^2$ were placed inside that prefactor, its coefficient would instead satisfy $\mu = \alpha_{\text{in}}$. These are different parameter conventions, not simultaneous definitions of the same α . Here $[\mu] = [\alpha] = L^2$, $[R] = [\Lambda] = L^{-2}$.

The metric field equations are [3, 4]

$$f_R R_{\mu\nu} - \frac{1}{2} f g_{\mu\nu} + (g_{\mu\nu} \square - \nabla_\mu \nabla_\nu) f_R = 8\pi T_{\mu\nu}, \quad (3)$$

with $f_R = 1 + 2\mu R$. Their trace is

$$6\mu \square R - R + 4\Lambda = 8\pi T. \quad (4)$$

Linearization about a constant-curvature vacuum background yields $m_\phi^2 = 1/(6\mu)$ for the scalar curvature mode. Positive μ gives a positive squared mass; this alone is not a proof of full background stability. The scalar variable $\phi = f_R$ is not assumed to be a canonically normalized Einstein-frame scalar. No relation between μ and the core scale r_c follows from this action comparison.

4 Effective Geometry

We consider a static, spherically symmetric metric

$$ds^2 = -F(r)dt^2 + \frac{dr^2}{F(r)} + r^2 d\Omega^2. \quad (5)$$

For $M > 0$ and $r_c > 0$, we choose the representative effective mass profile

$$m(r) = \frac{Mr^3}{r^3 + r_c^3}, \quad (6)$$

leading to

$$F(r) = 1 - \frac{2m(r)}{r}. \quad (7)$$

The asymptotic Schwarzschild limit and the regular-core behavior do not select this profile uniquely: $\tilde{m}(r) = Mr^3/(r^2 + r_c^2)^{3/2}$ satisfies both requirements and gives a different horizon equation. The following calculations refer only to the specified profile.

For $r_c^3 = 2M\ell^2$ this is precisely Hayward's static metric [6]; its metric and horizon coalescence are not new geometric discoveries. A fixed- r_c scan is not the fixed- ℓ mass evolution of that dynamical example.

5 Vacuum Field-Equation Test

Set $a = r_c^3$. A direct curvature calculation for the stated metric gives

$$R(r) = \frac{12Ma(2a - r^3)}{(r^3 + a)^3}, \quad \square R = \frac{1}{r^2} \frac{d}{dr} \left(r^2 F \frac{dR}{dr} \right). \quad (8)$$

In vacuum, Eq. (4) must vanish identically. At infinity its left side tends to 4Λ , forcing $\Lambda = 0$. For that value,

$$\lim_{r \rightarrow \infty} r^6 (6\mu \square R - R) = 12Ma \neq 0, \quad \lim_{r \rightarrow 0^+} (6\mu \square R - R) = -\frac{24M}{a} \neq 0. \quad (9)$$

Therefore this nontrivial representative geometry is not an exact vacuum solution of constant-coefficient quadratic metric $f(R)$ gravity. This exclusion does not cover different metric ansatzes or specified matter models. With matter, its stress tensor and its own equations must be supplied; defining $T_{\mu\nu}$ from the metric residual alone does not demonstrate that completion. The Schwarzschild limit $r_c = 0$ at $r > 0$ is a separate vacuum case.

6 Critical Horizon Structure

Candidate static Killing horizons at positive radii satisfy

$$F(r_H) = 0. \quad (10)$$

This yields

$$r_H^3 - 2Mr_H^2 + r_c^3 = 0. \quad (11)$$

Introducing

$$x = \frac{r_H}{2M}, \quad \beta = \frac{r_c}{2M}, \quad (12)$$

the parameter β is the dimensionless ratio between the internal scale r_c and the gravitational radius $2M$. It measures the strength of regularization at the horizon scale.

We obtain

$$P(x, \beta) = x^3 - x^2 + \beta^3 = 0. \quad (13)$$

Here $F = P/(x^3 + \beta^3)$; the denominator is positive for $x > 0$, $\beta > 0$. At $\beta = 0$, the double zero $x = 0$ in the cleared polynomial is outside this equivalence domain and is not a Schwarzschild horizon. The cubic term β^3 originates in r_c^3 in the chosen mass profile; dimensionlessness alone does not require it.

The critical value is

$$\beta_c = \left(\frac{4}{27} \right)^{1/3}. \quad (14)$$

At criticality, $P = (x - 2/3)^2(x + 1/3)$. For $0 < \beta < \beta_c$ there are two positive roots and one negative root; at β_c one positive double root; above β_c the only real root is negative. The discriminant is $\Delta = \beta^3(4 - 27\beta^3)$. Figure 1 illustrates the approach of the outer branch to the analytically derived critical value. The positive branch count changes, while the branch locations remain continuous up to coalescence.




Figure 1: Positive horizon branches of the specified representative Hayward family, obtained from $x^3 - x^2 + \beta^3 = 0$. The positive branches merge at $x_c = 2/3$ and $\beta_c = (4/27)^{1/3}$. This static parameter diagram does not demonstrate dynamical horizon formation.

7 Local Horizon Scaling

The horizon equation is

$$f(x, \beta) = x^3 - x^2 + \beta^3 = 0. \quad (15)$$

The critical point is

$$x_c = \frac{2}{3}, \quad \beta_c = \left(\frac{4}{27}\right)^{1/3}. \quad (16)$$

Expanding around criticality yields

$$x_{\pm} - x_c = \pm\sqrt{3}\beta_c\epsilon^{1/2} + O(\epsilon), \quad \epsilon = \beta_c - \beta > 0. \quad (17)$$

The derivatives $P_{xx}(x_c, \beta_c) = 2$ and $P_{\beta}(x_c, \beta_c) = 3\beta_c^2$ are nonzero, so this is a nondegenerate local fold. The exact expansion is

$$P(x_c + \delta, \beta_c - \epsilon) = \delta^2 + \delta^3 - 3\beta_c^2\epsilon + 3\beta_c\epsilon^2 - \epsilon^3. \quad (18)$$

Thus the horizon-root displacement exponent in this parameterization is

$$\alpha_H = \frac{1}{2}. \quad (19)$$

8 Literature Context and Theorem Boundaries

Static horizon roots and dynamical formation are distinct. Hayward's time-dependent example supplies mass evolution and fluxes [6]; these are additional inputs, not consequences of our cubic. Trapping horizons need not be global event horizons.

The zeroth-law analysis of Ghosh and Sarkar [7] assumes a smooth stationary Killing horizon and appropriate field equations and matter conditions. Its Ricci-squared argument uses nonzero coupling, a compact connected boundaryless section and non-extremality. Pure $f(R)$ uses the mixed projection and requires $f_R \neq 0$ instead. Our critical horizon has $\kappa = 0$; spherical angular constancy does not prove a field-equation solution or horizon formation.

Bessa [8] studies scalar dust perturbations of flat FLRW in full quadratic gravity, including Weyl squared, with sub-horizon and quasi-static approximations. Its formal infrared continuation does not prove TIG perturbative stability or a controlled super-horizon limit.

Garay et al. [9] provide an indirect two-vertex LQG comparison, not ordinary quantum mechanics recovered from TIG. Finite-dimensional quantum reconstruction [10] requires its own operational assumptions. No TIG derivation of these assumptions or quantum recovery is supplied here.

9 Conditional Round-Trip Delay

In a connected static region with $F > 0$, radial null propagation gives $dt = \pm dr/F$. We additionally prescribe an inner reflector at $r_{\text{ref}} = 2Mx_{\text{ref}} > 0$ and a partially reflecting, transmitting outer scattering boundary at $r_b = 2Mx_b$. Neither boundary is derived from TIG. The geometric round-trip time in asymptotically normalized t is

$$T_{\text{rt}} = 2 \int_{r_{\text{ref}}}^{r_b} \frac{dr}{F} = 4M \int_{x_{\text{ref}}}^{x_b} \frac{x^3 + \beta^3}{x^3 - x^2 + \beta^3} dx. \quad (20)$$

Echo-like output additionally requires excitation and appropriate scattering; this formula supplies neither a waveform nor a gravitational perturbation equation. A minimally coupled massless test scalar would have

$$(-\partial_t^2 + \partial_{r_*}^2 - V_\ell)\psi_\ell = 0, \quad \frac{dr_*}{dr} = \frac{1}{F}, \quad V_\ell = F \left[\frac{\ell(\ell+1)}{r^2} + \frac{F'}{r} \right]. \quad (21)$$

An ideal Dirichlet inner boundary reflects that test field; its physical origin and the required outer scattering remain assumptions. The horizonless-object calculation of Cardoso et al. [11] is a methodological comparison with different geometry and boundary data. Its one-way delay is distinct from the round trip defined here.

For fixed M , set $\eta = \beta - \beta_c > 0$, $\delta = x - x_c$, $D_0 = 4/9$ and $s = \sqrt{3}\beta_c\sqrt{\eta}$. Above criticality $F > 0$ at all positive radii. The exact numerator is

$$P = \delta^2 + \delta^3 + 3\beta_c^2\eta + 3\beta_c\eta^2 + \eta^3. \quad (22)$$

Its bottleneck has $F \sim (\delta^2 + s^2)/D_0$. If the fixed endpoints satisfy $0 < x_{\text{ref}} < x_c < x_b$, the scaled integral in Eq. (20) contains $\int_{-\infty}^{\infty} dy/(1+y^2) = \pi$ and hence

$$T_{\text{rt}} \sim \frac{16\pi M}{9\sqrt{3}\beta_c} (\beta - \beta_c)^{-1/2}, \quad \beta \downarrow \beta_c, \quad \beta > \beta_c. \quad (23)$$

The ratio to this leading term tends to one. The local scaled integrand is dominated by a constant times $(1+y^2)^{-1}$; regions away from x_c have bounded travel time. One fixed endpoint at x_c halves the coefficient; two fixed endpoints strictly on the same side have a finite limit. A fixed exterior observer with a finite positive limiting lapse measures $\Delta\tau_{\text{obs}} = \sqrt{F(r_{\text{obs}})} T_{\text{rt}}$.

Below criticality, let $\epsilon = \beta_c - \beta > 0$ and $s = \sqrt{3}\beta_c\sqrt{\epsilon}$. Returning exterior paths must stay at $x > x_+$. Fixed $x_c < x_{\text{ref}} < x_b$ give a finite critical limit. Only the additional tracking prescription $x_{\text{ref}} = x_+ + (q-1)s$, with fixed $q > 1$, gives

$$T_{\text{rt}} \sim \frac{2MD_0}{\sqrt{3}\beta_c} \ln \frac{q+1}{q-1} \epsilon^{-1/2}. \quad (24)$$

This prescribes a coordinate gap, not a fixed proper gap. At fixed subcritical β , a mirror approaching the simple outer horizon instead gives $T_{\text{rt}} = -\kappa_+^{-1} \ln(h/L) + O(1)$, where $h = r_{\text{ref}} - r_+ \rightarrow 0^+$ and $L > 0$ is fixed. At exact criticality, $x_{\text{ref}} = x_c + d$ gives $T_{\text{rt}} \sim 16M/(9d)$ as $d \rightarrow 0^+$. Crossing a future outer horizon is not a returning exterior signal path. The former unrestricted $(\beta_c - \beta)^{-1/2}$ echo claim therefore does not hold for arbitrary boundaries. Equation (23) is a conditional geometric delay, not a demonstrated physical TIG echo.

10 Conclusion

The specified representative TIG geometry has an exact cubic horizon equation and a non-degenerate fold with horizon-root exponent $1/2$. Neither structural consistency nor a generic fold uniquely requires this cubic mass-profile law. The nontrivial metric fails the vacuum trace equation of the tested quadratic metric $f(R)$ action; no exact solution with specified matter has been demonstrated. The algebraic horizon result remains model-conditional, while dynamical completion, stability, and echo forecasts require separate evidence.

The static family is identified with Hayward’s earlier metric. With additionally prescribed reflection and fixed endpoints straddling the horizonless bottleneck, the round-trip delay has exponent $-1/2$ on the $\beta > \beta_c$ side. Subcritical delays depend on the mirror prescription. Neither this kinematic calculation nor the comparison literature establishes physical TIG echoes, dynamical horizon formation or quantum-mechanical recovery.

References

- [1] Albert Einstein. Die feldgleichungen der gravitation. *Sitzungsberichte der Preussischen Akademie der Wissenschaften zu Berlin*, pages 844–847, 1915.
- [2] Robert M. Wald. *General Relativity*. University of Chicago Press, Chicago, 1984.
- [3] Thomas P. Sotiriou and Valerio Faraoni. $f(R)$ theories of gravity. *Reviews of Modern Physics*, 82: 451–497, 2010. doi:10.1103/RevModPhys.82.451. URL <https://arxiv.org/abs/0805.1726v4>. arXiv:0805.1726v4, published-version metadata.
- [4] Antonio De Felice and Shinji Tsujikawa. $f(R)$ theories. *Living Reviews in Relativity*, 13:3, 2010. doi:10.12942/lrr-2010-3. URL <https://arxiv.org/abs/1002.4928v2>. arXiv:1002.4928v2, published version.
- [5] Jeffrey R. Chasnov. Applied linear algebra and differential equations, 2020. URL <https://www.math.hkust.edu.hk/~machas/applied-linear-algebra-and-differential-equations.pdf>. HKUST lecture notes, preface January 2020, Section 11.2.1.
- [6] Sean A. Hayward. Formation and evaporation of nonsingular black holes. *Physical Review Letters*, 96:031103, 2006. doi:10.1103/PhysRevLett.96.031103. URL <https://arxiv.org/abs/gr-qc/0506126v2>. arXiv:gr-qc/0506126v2.
- [7] Rajes Ghosh and Sudipta Sarkar. How Universal Is the Black Hole Zeroth Law?, 2026. URL <https://arxiv.org/abs/2609.24648v1>. arXiv:2609.24648v1; Preprint, arXiv v1 submitted 21 September 2026.
- [8] Pedro Bessa. Matter density perturbations in Quadratic Gravity, 2026. URL <https://arxiv.org/abs/2609.29046v1>. arXiv:2609.29046v1; Preprint, arXiv v1 submitted 24 September 2026; author reports submission to PRD; journal receipt date not verified.
- [9] Ñaki Garay, Luis J. Garay, Diego H. Gugliotta, Sergio Rodríguez-González, and Raül Vera. Global variables and dynamics of emergent cosmology in loop gravity, 2026. URL <https://arxiv.org/abs/2609.31123v1>. arXiv:2609.31123v1; Preprint, arXiv v1 submitted 25 September 2026.
- [10] Giulio Chiribella, Giacomo Mauro D’Ariano, and Paolo Perinotti. Informational derivation of quantum theory. *Physical Review A*, 84:012311, 2011. doi:10.1103/PhysRevA.84.012311. URL <https://arxiv.org/abs/1011.6451v3>. arXiv:1011.6451v3; Version 3, 15 July 2011; finite-dimensional reconstruction.
- [11] Vitor Cardoso, Edgardo Franzin, and Paolo Pani. Is the gravitational-wave ringdown a probe of the event horizon? *Physical Review Letters*, 116:171101, 2016. doi:10.1103/PhysRevLett.116.171101. URL <https://arxiv.org/abs/1602.07309v4>. arXiv:1602.07309v4, includes Erratum.