

Analytical Research Studies

Dr. Mustafa Haytham

Faculty of Medicine | Department of Community Medicine & Epidemiology

Interpeko
Research Club

Interpeko
Research

Learning Objectives

- To critically evaluate the theoretical foundations and philosophical underpinnings of analytical epidemiological study designs within the positivist and post-positivist paradigms.
- To synthesise the methodological complexities of analytic epidemiology, including counterfactual reasoning, potential outcomes frameworks, and directed acyclic graphs (DAGs) for confounding control.
- To critically appraise the design, conduct, and interpretation of case–control studies, with particular emphasis on selection mechanisms, exposure ascertainment, and the assumptions underlying the odds ratio as an approximation of relative risk.
- To evaluate the theoretical and practical dimensions of cohort study design, including immortal time bias, time-varying covariates, competing risks, and the Cox proportional hazards model.
- To apply advanced causal inference methods—including inverse probability weighting, marginal structural models, and Mendelian randomisation—to observational data.

Taxonomy of Epidemiological Study Designs

- The hierarchy of epidemiological evidence places randomised controlled trials (RCTs) at the apex, yet observational analytical studies constitute the predominant source of etiological knowledge for exposures that cannot be ethically or practically randomised.
- Observational designs are taxonomically divided into descriptive studies (generating hypotheses) and analytical studies (testing hypotheses), the latter forming the empirical basis of modern causal inference.
- Analytical studies encompass ecological designs (susceptible to the ecological fallacy), cross-sectional designs (prevalence studies with limited temporal resolution), case–control designs (retrospective sampling from outcome-defined populations), and cohort designs (prospective or retrospective follow-up of exposure-defined populations).
- Experimental designs—principally RCTs and quasi-experimental designs—establish causality through manipulation and control, yet their external validity is frequently constrained by selective eligibility and artificial settings.
- The Bradford Hill criteria (1965) and modern causal inference frameworks (Pearl's do-calculus, Rubin's potential outcomes) provide the epistemological scaffolding for interpreting analytical study findings.


Analytical Epidemiology: Theoretical Foundations

- Analytical epidemiology is conceptualised as the systematic investigation of determinants (risk factors, exposures, interventions) of health-related states or events in human populations, operationalised through the comparison of exposure-defined or outcome-defined groups.
- The discipline is rooted in the counterfactual framework: the causal effect of an exposure is defined as the contrast between the outcome observed under exposure and the outcome that would have been observed under non-exposure (the counterfactual) for the same individual.
- The fundamental analytical strategy comprises: (i) precise operationalisation of exposure and outcome variables; (ii) formulation of falsifiable hypotheses grounded in biological plausibility; and (iii) comparative analysis controlling for confounding, selection bias, and information bias.
- Measures of effect—risk difference (causal risk difference), risk ratio, rate ratio, and odds ratio—quantify the magnitude and direction of association, while measures of association strength (e.g., attributable fraction) inform public health significance.
- Statistical inference employs frequentist (P-values, confidence intervals) and Bayesian (credible intervals, posterior probabilities) approaches, with multivariable regression (logistic, Cox, Poisson) serving as the primary analytical engine for confounding adjustment.

Analytical Study Designs: A Comparative Taxonomy

- **Ecological Studies:** Aggregate-level analysis of exposure–outcome relationships; highly susceptible to the ecological fallacy (cross-level bias) and confounding by group-level variables; useful for generating hypotheses but inadequate for causal inference.
- **Cross-Sectional Studies:** Simultaneous measurement of exposure and outcome at a single time point; provide prevalence estimates and prevalence odds ratios; incapable of establishing temporal precedence; vulnerable to prevalence–incidence bias (Neyman bias).
- **Case–Control Studies:** Retrospective sampling from a source population conditional on outcome status; efficient for rare diseases and long-latency outcomes; the odds ratio approximates the incidence odds ratio under the rare-disease assumption (incidence < 10%).
- **Cohort Studies:** Follow-up of exposure-defined groups over time; the only observational design permitting direct incidence estimation and relative risk calculation; prospective cohorts minimise recall bias, while retrospective (historical) cohorts exploit pre-existing records.
- **Nested Case–Control and Case–Cohort Designs:** Hybrid designs that combine the efficiency of case–control sampling with the temporal structure of cohort studies, enabling cost-effective biomarker assessment while preserving incidence data.

Temporal Architecture of Analytical Study Designs



Methodological Pillars of Analytical Studies

- **Comparison Groups:** The fundamental analytical operation requires the construction of contrast groups (exposed vs. unexposed, or cases vs. controls) drawn from a common source population to ensure exchangeability and minimise confounding bias.
- **Hypothesis Testing:** Analytical studies test a priori causal hypotheses derived from biological plausibility, prior empirical evidence, and theoretical models; the null hypothesis of no association is evaluated against predefined significance thresholds ($\alpha = 0.05$).
- **Measures of Effect:** Risk difference (absolute effect), risk ratio (relative effect), rate ratio (incidence density ratio), and odds ratio (cross-product ratio) provide complementary information on association magnitude and public health significance.
- **Statistical Inference:** Frequentist confidence intervals (95% CI) quantify precision, while P-values assess compatibility with the null hypothesis; Bayesian methods incorporate prior knowledge through posterior probability distributions.
- **Bias Assessment:** Systematic evaluation of selection bias (non-random sampling), information bias (differential misclassification), and confounding (common causes of exposure and outcome) is mandatory for valid causal inference.

Applications in Doctoral and Public Health Research

- **Etiological Investigation:** Identifying modifiable and non-modifiable determinants of disease occurrence, progression, and prognosis across diverse populations and contexts.
- **Risk Factor Evaluation:** Quantifying the independent contribution of environmental (air pollution, occupational hazards), genetic (polygenic risk scores), and lifestyle (diet, physical activity) factors to disease burden.
- **Evidence Synthesis:** Informing systematic reviews, meta-analyses, and clinical practice guidelines through high-quality primary observational data that complement experimental evidence.
- **Intervention Assessment:** Evaluating the real-world effectiveness, safety, and equity of pharmaceuticals, vaccines, and public health programmes through pharmacoepidemiological and implementation science frameworks.
- **Policy Formulation:** Providing robust evidence for health policy, resource allocation, and strategic planning at national and international levels, including burden-of-disease estimation and health technology assessment.

Case–Control Studies: Theoretical and Methodological Architecture

- A case–control study is an observational analytical design in which sampling is conducted conditional on the outcome (disease) status: cases (individuals with the outcome) and controls (individuals without the outcome) are selected from a defined source population.
- The design is inherently retrospective, working backward from effect (disease at present) to cause (exposure in the past), thereby violating the temporal sequence required for unambiguous causal inference.
- The odds ratio (OR) is the principal measure of effect, calculated as the ratio of exposure odds among cases to exposure odds among controls; under the rare-disease assumption (cumulative incidence $< 10\%$), the OR closely approximates the risk ratio.
- Efficiency for rare diseases and long-latency outcomes is the primary advantage, as the design avoids the prolonged follow-up and massive sample sizes required by cohort designs for rare outcomes.
- The design is particularly suited for outbreak investigations, hypothesis generation for novel diseases, and settings where prospective follow-up is infeasible due to cost, time, or ethical constraints.

Methodological Protocol: Case–Control Studies

- Step 1 — Case Definition: Establish explicit, reproducible diagnostic criteria using standardised classification systems (ICD-11, DSM-5-TR, WHO case definitions); ensure diagnostic homogeneity to minimise disease misclassification.
- Step 2 — Control Selection: Select controls from the same source population as cases, using density sampling (incidence density sampling) or cumulative sampling; match on strong confounders (age, sex, ethnicity) or adjust analytically.
- Step 3 — Exposure Ascertainment: Collect exposure data using validated instruments (structured interviews, medical records, biological specimens); blind data collectors to case/control status to prevent differential information bias.
- Step 4 — Effect Estimation: Calculate the odds ratio and its 95% confidence interval; evaluate effect modification through stratified analysis and interaction terms; assess dose–response relationships if exposure is ordinal or continuous.
- Step 5 — Confounding Control: Apply multivariable logistic regression, propensity score methods (matching, stratification, weighting), or instrumental variable approaches to adjust for measured and unmeasured confounders.

Case–Control Studies: Critical Appraisal at Doctoral Level

Methodological Strengths

- Exceptional efficiency for rare diseases and outcomes with long latency periods.
- Rapid execution and relatively low cost compared to longitudinal cohort designs.
- Simultaneous evaluation of multiple risk factors and potential effect modifiers.
- No attrition or loss to follow-up, as the study is retrospective by design.
- Ethical feasibility for studying harmful exposures, as the outcome has already occurred.

Methodological Limitations

- Recall bias: differential accuracy of exposure memory between cases and controls.
- Selection bias: inappropriate control selection may violate the study base principle.
- Cannot estimate incidence or prevalence; relative risk is not directly calculable.
- Temporal ambiguity: exposure may have occurred after disease onset (reverse causation).
- High susceptibility to information bias and unmeasured confounding.

Seminal Case–Control Study: Reye's Syndrome and Aspirin Exposure

- **Historical Context:** During the 1970s–1980s, an alarming cluster of Reye's syndrome (acute encephalopathy with hepatic failure) among children prompted urgent etiological investigation.
- **Study Design:** Multiple case–control studies were conducted across the United States and United Kingdom, enrolling children with biopsy-confirmed or clinically diagnosed Reye's syndrome (cases) and age-matched hospital controls or community controls.
- **Exposure Assessment:** Structured parental interviews documented antecedent medication use during the prodromal viral illness; aspirin exposure was dichotomised and quantified by dose and duration.
- **Findings:** Consistently elevated odds ratios ($OR > 10$) were observed across studies, with strong dose–response gradients and biological plausibility (mitochondrial toxicity of salicylates).
- **Public Health Impact:** The US Surgeon General's advisory (1982) and subsequent FDA labelling requirements led to a $>90\%$ reduction in paediatric aspirin use and near-elimination of Reye's syndrome—a paradigmatic example of rapid translation from analytical epidemiology to policy action.

Cohort Studies: Theoretical Framework and Classification

- A cohort study is a longitudinal observational design in which a defined population (cohort) free of the outcome at baseline is classified by exposure status and followed over time to ascertain incident disease or mortality.
- The design preserves temporal sequence (exposure precedes outcome), satisfying a necessary condition for causal inference and enabling direct estimation of incidence rates and relative risks.
- Prospective (Concurrent) Cohorts: Exposure is measured at baseline (t_0) and subjects are followed forward into the future (t_1 to t_n); advantages include standardised data collection, minimal recall bias, and capacity to measure time-varying exposures.
- Retrospective (Historical) Cohorts: Exposure is ascertained from pre-existing records (employment archives, medical registries) and outcome is determined in the present; efficient when exposure data are already archived but susceptible to incomplete records.
- Ambidirectional Cohorts: Combine retrospective and prospective elements, using historical exposure data while continuing follow-up into the future; maximises efficiency for diseases with both short and long latency periods.

Methodological Protocol: Cohort Studies

- Step 1 — Cohort Assembly: Define explicit eligibility criteria (inclusion/exclusion) to ensure a homogeneous, at-risk population; consider sampling fractions and power calculations based on expected incidence, effect size, and follow-up duration.
- Step 2 — Baseline Exposure Assessment: Characterise exposure using validated instruments, biomarkers, or occupational records; quantify exposure intensity, duration, and latency; assess co-exposures and effect modifiers.
- Step 3 — Follow-up Protocol: Establish systematic follow-up procedures (annual examinations, registry linkage, telephone interviews); document losses to follow-up and reasons for dropout; minimise differential attrition.
- Step 4 — Outcome Ascertainment: Apply standardised diagnostic criteria and blinded outcome adjudication; use multiple sources (medical records, death certificates, cancer registries) to maximise sensitivity and specificity.
- Step 5 — Analytical Strategy: Calculate incidence rates, rate ratios, and hazard ratios; fit Cox proportional hazards models for time-to-event data; adjust for confounders; evaluate competing risks and time-varying covariates.

Cohort Studies: Critical Appraisal at Doctoral Level

Methodological Strengths

- Clear temporal sequence strengthens causal inference and minimises reverse causation.
- Direct incidence estimation and relative risk calculation without rare-disease assumption.
- Multiple outcomes can be studied from a single exposure, enhancing cost-efficiency.
- Assessment of dose–response relationships and time-varying exposures.
- Reduced recall and interviewer bias in prospective designs.

Methodological Limitations

- Large sample sizes and extended follow-up periods are resource-intensive.
- Loss to follow-up may introduce selection bias if attrition is differential by exposure.
- Inefficient for rare diseases unless cohorts are exceptionally large (e.g., >100,000).
- Exposure status may change during follow-up, requiring time-varying covariate models.
- Ethical concerns for harmful exposures; Hawthorne effect may alter participant behaviour.

Landmark Cohort Study: The Framingham Heart Study (1948–Present)

- **Historical Genesis:** Initiated in 1948 under the direction of the US Public Health Service, the Framingham Heart Study was conceived to investigate the emerging cardiovascular disease epidemic in an era dominated by infectious disease epidemiology.
- **Cohort Design:** 5,209 residents of Framingham, Massachusetts, aged 30–62 years and free of overt cardiovascular disease, were enrolled and subjected to comprehensive biennial examinations including physical examination, electrocardiography, and laboratory assessment.
- **Major Discoveries:** The study identified hypertension, hypercholesterolaemia, cigarette smoking, obesity, diabetes mellitus, and physical inactivity as major independent risk factors; it introduced the term 'risk factor' into the medical lexicon.
- **Methodological Legacy:** The offspring cohort (enrolled 1971) and third-generation cohort (enrolled 2002) enabled familial and genetic analyses; the Framingham Risk Score became the global standard for 10-year cardiovascular risk prediction.
- **Scientific Impact:** Over 3,000 peer-reviewed publications have emanated from this single cohort, demonstrating that cardiovascular disease is largely preventable through lifestyle modification and risk factor control.

Comparative Methodological Analysis: Case–Control vs. Cohort Studies

Methodological Feature	Case–Control Studies	Cohort Studies
Direction of inquiry	From disease to exposure (backward sampling)	From exposure to disease (forward follow-up)
Temporal orientation	Retrospective; exposure history reconstructed post-hoc	Prospective or retrospective; temporal sequence preserved
Sampling framework	Outcome-dependent sampling (cases and controls)	Exposure-dependent sampling (exposed and unexposed cohorts)
Sample size requirements	Smaller; efficient for rare diseases (incidence < 5%)	Larger; especially for rare outcomes (N > 10,000 often required)
Duration and cost	Short duration (months); relatively inexpensive	Long duration (years to decades); resource-intensive
Measures of effect	Odds ratio (OR); approximates RR under rare-disease assumption	Relative risk (RR), incidence rate ratio, hazard ratio (HR)
Key methodological strengths	Rare diseases; multiple exposures; rapid hypothesis testing; long-latency outcomes	Clear temporal sequence; direct incidence estimation; multiple outcomes; dose–response assessment
Key methodological limitations	Recall bias; selection bias; temporal ambiguity; cannot estimate incidence or prevalence	Loss to follow-up; inefficient for rare diseases; exposure change over time; Hawthorne effect

Doctoral Exercise 1: Study Design Identification

- **Scenario:** Investigators sought to examine the association between occupational exposure to asbestos and the occurrence of mesothelioma—a relatively rare malignancy with a long latency period (20–40 years).
- **Data Sources:** Employment records from several major manufacturing facilities documented worker asbestos exposure levels over the preceding four decades. Union health records, including death certificates and cancer registry linkages, documented mesothelioma occurrence.
- **Methodology:** Researchers classified workers employed in the 1940s into high-exposure (≥ 25 fibre-years) and moderate/low-exposure (< 25 fibre-years) groups, then compared cumulative mesothelioma incidence between these groups through 2020.
- **Question:** Identify the study design and justify your classification.
- **Answer:** Retrospective Cohort Study. Exposure was defined a priori from historical occupational records, and outcome incidence was compared between exposure-defined groups over an extended follow-up period. The design preserves temporal sequence and permits direct incidence estimation, distinguishing it from case–control designs.

Doctoral Exercise 2: Study Design Identification

- Scenario: Persons diagnosed with new-onset Lyme disease (*Borrelia burgdorferi* infection confirmed by serology) were recruited from a regional infectious disease clinic during the summer of 2024.
- Methodology: Cases were interviewed regarding their frequency of walking through wooded areas, use of insect repellent (DEET concentration), wearing of short sleeves and trousers, and residential proximity to tick habitats. Twice as many patients without Lyme disease from the same physician's practice were recruited as controls and asked identical questions using a standardised questionnaire.
- Analytical Approach: Conditional logistic regression was used to estimate odds ratios adjusted for age, sex, and outdoor occupation, with 95% confidence intervals.
- Question: Identify the study design, discuss its methodological strengths and limitations, and evaluate the appropriateness of the analytical approach.
- Answer: Case–Control Study. The design samples conditional on disease status and retrospectively assesses exposure. Strengths include efficiency for a relatively rare outcome; limitations include susceptibility to recall bias and the inability to estimate incidence. Conditional logistic regression is appropriate for matched designs.

Doctoral Exercise 3: Study Design Identification

- Scenario: A research team investigated the association between cigarette smoking and lung cancer incidence in a large industrial population.
- Methodology: They enrolled 7,000 current smokers and 3,000 never-smokers from a manufacturing cohort and followed both groups prospectively for five years (2020–2025). Smoking status was verified through cotinine assays at baseline. Annual chest radiography and cancer registry linkage were used for outcome ascertainment.
- Findings: During follow-up, 70 smokers developed lung cancer (incidence rate: 20 per 1,000 person-years), compared with only 3 cases among non-smokers (incidence rate: 0.2 per 1,000 person-years). The age-adjusted hazard ratio was 98.7 (95% CI: 30.6–318.4).
- Question: Identify the study design, calculate the appropriate measures of effect, and discuss the causal interpretation of findings.
- Answer: Prospective Cohort Study. The relative risk (cumulative incidence ratio) = $70/7000 \div 3/3000 = 10.0$; the rate ratio = $20/0.2 = 100$. The exceptionally large hazard ratio, combined with biological plausibility, temporal sequence, and dose–response gradients from prior studies, supports a causal interpretation.

Doctoral Conclusions and Future Directions

- Analytical epidemiological studies constitute the empirical foundation of modern public health science, bridging descriptive observations with causal inference and policy translation.
- Case–control and cohort studies are methodologically complementary: case–control designs excel in efficiency for rare diseases and hypothesis generation, while cohort designs provide the temporal structure and incidence data necessary for robust causal inference.
- Rigorous study design—including explicit case definitions, appropriate control selection, validated exposure assessment, and blinded outcome ascertainment—is essential to minimise systematic error and ensure internal validity.
- Advanced analytical methods—propensity scores, marginal structural models, instrumental variables, and Mendelian randomisation—have substantially enhanced the capacity of observational studies to approximate experimental causal inference.
- Future directions include the integration of multi-omics data (genomics, epigenomics, metabolomics), artificial intelligence for exposure prediction, and real-world evidence frameworks that leverage electronic health records for large-scale analytical epidemiology.

Interpeko[™] Research Club

THANK *You!*

To everyone who supports, participates,
and believes in our mission.



Together
we learn.



Together
we research.



Together
we create impact.

Your encouragement **drives us forward**
and helps us build a stronger research community.

We appreciate you!



Let's keep growing, learning
and making a difference—**together.**



| Interpeko Research Club

