

Hilbert Substrate Framework 1.3: A Finite-Hilbert Quantum Substrate with Gauge-Like Constraints, Defragmentation Potentials, and Emergent Exchange-Antisymmetric Ground States

Ben Bray^{*}
(Dated: November 21, 2025)

We present an updated formulation of the Hilbert Substrate Framework, a finite-dimensional quantum substrate designed to investigate how exchange antisymmetry, spatial exclusion, and nonclassical correlations may arise from local constraint structures and information-balance terms. The framework combines: (1) a two-excitation quantum substrate defined on a small periodic lattice; (2) a Gauss-like information-flux penalty; (3) a defragmentation potential that encourages spatial coherence; (4) a contact spin interaction analogous to the singlet–triplet structure of two-body quantum mechanics; and (5) a CHSH/Bell-sector subsystem used as a calibration of the quantum machinery.

In the two-excitation substrate, we find that the ground state is exactly exchange antisymmetric under swap, and that all spatial-overlap weight is concentrated in the spin-singlet channel. This provides a concrete demonstration that a constrained finite Hilbert system can prefer fermion-like symmetry sectors without explicitly postulating fermionic operators. The results are not claimed to reproduce Pauli exclusion or the Standard Model, but they illustrate that constraint-driven information balance can shape symmetry sectors in a controlled finite model. Throughout, we adopt the guiding principle: *if the data do not show it, we do not claim it.*

CONTENTS

I. Introduction	2
II. Configuration Space and the Substrate Hilbert Space	2
A. Configuration space in finite models	2
B. Local geometric structure: neighbors and periodicity	2
III. Hamiltonian Terms	2
A. Hopping	2
B. Mass term	3
C. Defrag potential	3
D. Gauss-like information-flux penalty	3
E. Contact spin interaction	3
IV. Ground State Results	3
A. Exchange antisymmetry	3
B. Spatial overlap and spin structure	4
C. Gauss expectation	4
D. Interpretation	4
V. CHSH Quantum Core	4
VI. Fermion Toy: Swap Operator Eigenstructure	5
VII. Caveats	5
VIII. Future Work	5
IX. Conclusion	5
References	5

^{*} (email omitted for draft)

I. INTRODUCTION

Hilbert Substrate 1.3 refines and formalizes the conceptual and computational investigations reported in previous versions of this framework. The central philosophical motivation is to study how *local constraints on information distribution* can shape the symmetry structure of low-energy states within a finite Hilbert space. The question is not whether this reproduces real fermions, gauge theories, or gravity, but whether constraint and routing principles can bias a Hilbert system toward familiar quantum-sector behavior.

To remain controlled and falsifiable, we restrict ourselves entirely to finite-dimensional Hilbert spaces, local Hamiltonians acting on them, and direct numerical diagonalization of the two-excitation sector. We then test the quantum engine with a CHSH subsystem whose results can be compared to the Tsirelson bound.

A pair of guiding principles shape the framework:

1. **Finite Hilbert substrate:** All degrees of freedom exist inside an explicitly-constructed finite Hilbert space. There are no continuum fields, no functional integrals, and no infinite sums.
2. **Empirical modesty:** We do not claim more structure than what emerges from the model. *If the data do not show it, we do not say it.*

In this version, the framework achieves a significant milestone: the ground state of the two-excitation substrate becomes *exactly exchange antisymmetric*, and the spin structure of same-site configurations becomes *purely singlet*. This is a clean, interpretable, finite-Hilbert analogue of “fermion-like” behavior arising from constraint rules.

II. CONFIGURATION SPACE AND THE SUBSTRATE HILBERT SPACE

A. Configuration space in finite models

For two distinguishable excitations on a lattice with N_s sites, each carrying a two-level internal degree of freedom (spin), the configuration space is

$$\mathcal{C} = \{(r_1, s_1; r_2, s_2) \mid r_i \in \{0, \dots, N_s - 1\}, s_i \in \{\uparrow, \downarrow\}\}. \quad (1)$$

The Hilbert space is

$$\mathcal{H} = \mathbb{C}^{(N_s \times 2) \times (N_s \times 2)} \cong \mathbb{C}^{4N_s^2}. \quad (2)$$

In the present work, $N_s = 4$ (a 2×2 lattice), so $\dim \mathcal{H} = 64$.

We emphasize: nothing in the construction presupposes bosonic or fermionic symmetry sectors. The swap operator

$$P_{\text{ex}} |r_1, s_1; r_2, s_2\rangle = |r_2, s_2; r_1, s_1\rangle \quad (3)$$

is available as an observable. The model is free to place weight in symmetric, antisymmetric, or mixed symmetry subspaces.

B. Local geometric structure: neighbors and periodicity

The lattice is periodic: each site has four neighbors. Hopping terms act only along these edges, so there is a clear notion of locality.

III. HAMILTONIAN TERMS

The substrate Hamiltonian is

$$H = H_{\text{hop}} + H_{\text{mass}} + H_{\text{defrag}} + H_{\text{Gauss}} + H_{\text{contact}}. \quad (4)$$

A. Hopping

Each excitation hops to its nearest neighbors with amplitude $-J_{\text{hop}}$. This produces delocalization and kinetic spread.

B. Mass term

Each excitation contributes $+m$ to the energy. This plays no structural role but keeps the model interpretable.

C. Defrag potential

A Gaussian potential centered on the lattice midpoint,

$$V_{\text{defrag}}(r) = -\exp\left(-\frac{\text{dist}(r, \text{center})^2}{2\sigma^2}\right), \quad (5)$$

encourages clumping and coherence. It introduces a “direction” toward which the substrate tends to compress or self-organize.

D. Gauss-like information-flux penalty

Let $\rho(r)$ be the occupancy of site r and $\rho_0 = 2/N_s$ the uniform occupancy. Define

$$H_{\text{Gauss}} = \frac{\lambda_G}{2} \sum_r (\rho(r) - \rho_0)^2. \quad (6)$$

This term enforces local “information balance”: states with uneven occupancy cost energy. In practice, this penalizes multi-occupation configurations and thus discourages spatial overlap.

E. Contact spin interaction

When $r_1 = r_2$, the excitations interact via a singlet–triplet structure:

$$H_{\text{contact}} = J_{\text{exch}} \vec{S}_1 \cdot \vec{S}_2 + \lambda_S \mathbb{1}_{\text{singlet}} + \lambda_T \mathbb{1}_{\text{triplet}, \parallel}. \quad (7)$$

Here $\vec{S} \cdot \vec{S}$ reproduces the usual $-3/4$ (singlet) and $+1/4$ (triplet) splitting, while λ_S and λ_T adjust the relative preference.

IV. GROUND STATE RESULTS

We diagonalized H on the full 64-dimensional Hilbert space. Parameters were:

$$\begin{aligned} L_x &= L_y = 2, \\ J_{\text{hop}} &= 1.0, \\ m &= 0.1, \\ g_{\text{defrag}} &= 1.0, \quad \sigma_{\text{defrag}} = 1.0, \\ \lambda_G &= 5.0, \\ \lambda_S &= -1.0, \quad \lambda_T = 0.0, \quad J_{\text{exch}} = 1.0. \end{aligned}$$

A. Exchange antisymmetry

Define the diagnostic:

$$A = 1 - \frac{\|\psi + P_{\text{ex}}\psi\|^2}{\|\psi\|^2}. \quad (8)$$

$A = 1$ means perfectly antisymmetric; $A = 0$ means no antisymmetric component.

Result:

$$A = 1.000000, \quad \text{antisymmetry violation} \approx 5 \times 10^{-28}.$$

Thus the ground state lies entirely in the antisymmetric sector.

B. Spatial overlap and spin structure

Probability that $r_1 = r_2$:

$$P_{\text{overlap}} = 0.151583.$$

Within this region:

$$P_{\text{singlet}} = 0.151583, \quad P_{\text{triplet}} = 0.$$

The overlap region is *purely* singlet.

C. Gauss expectation

$$\langle H_{\text{Gauss}} \rangle = 3.257917,$$

indicating the Gauss sector plays an active role.

D. Interpretation

This is a controlled, finite-dimensional demonstration that a combination of:

- local information-flux balancing,
- defragmentation geometry,
- contact spin structure,

can bias a substrate toward a fermion-like antisymmetric ground state.

No fermionic creation/annihilation operators were introduced. No antisymmetrization was imposed. The symmetry arises as an energetic preference.

We do *not* claim this explains real fermions.

V. CHSH QUANTUM CORE

As a calibration of the quantum engine, we include a two-qubit Heisenberg dimer. With the Bell state as initial condition, the CHSH value

$$S(t) = E(A, B) + E(A, B') + E(A', B) - E(A', B')$$

achieves

$$S_{\text{max}} \approx 2.828427,$$

the Tsirelson bound. This confirms that the quantum portion of the code behaves properly.

VI. FERMION TOY: SWAP OPERATOR EIGENSTRUCTURE

A two-qubit toy subsystem verifies the swap operator:

$$\langle \psi_{\text{sym}} | P_{\text{ex}} | \psi_{\text{sym}} \rangle = +1, \quad \langle \psi_{\text{asym}} | P_{\text{ex}} | \psi_{\text{asym}} \rangle = -1.$$

This validates the operator used in the substrate diagnostics.

VII. CAVEATS

We emphasize several limitations:

- The model is finite-dimensional and not relativistic.
- No claim is made about reproducing Pauli exclusion in the continuum.
- No Standard Model structure is invoked.
- The results are specific to parameter choices.

Everything shown is a property of the specific Hamiltonian used.

VIII. FUTURE WORK

Potential extensions include:

- Scaling to larger lattices (e.g. 3×3).
- Adding explicit gauge-field Hilbert spaces.
- Studying phase diagrams across $(\lambda_G, \lambda_S, J_{\text{exch}})$.
- Embedding CHSH correlations inside substrate degrees of freedom.

IX. CONCLUSION

Hilbert Substrate 1.3 provides a coherent finite-Hilbert environment in which information-flux constraints, defragmentation potentials, and spin-structured contact interactions combine to yield an emergent antisymmetric ground state. The framework is deliberately modest: we observe only what the model shows. Nevertheless, the result demonstrates that fermion-like structure can arise from local rules without being explicitly imposed.

-
- [1] J. S. Bell, “On the Einstein Podolsky Rosen paradox,” *Physics Physique Fizika* **1**, 195 (1964).
 [2] B. S. Tsirelson, “Quantum generalizations of Bell’s inequality,” *Lett. Math. Phys.* **4**, 93–100 (1980).