

Multilevel Monte Carlo for Stochastic PDEs



Inti-Raymi Carhuanchu Mantripp
Mansfield College
University of Oxford

A dissertation submitted for the degree of
MSc in Mathematical Modelling & Scientific Computing
Trinity 2025

Contents

0.1	Introduction	1
0.2	Prerequisites	3
0.2.1	Multilevel Monte Carlo	3
	Bibliography	4

List of Figures

0.1 Introduction

Stochastic partial differential equations (SPDEs) commonly arise in the mathematical modelling of systems whose evolution is significantly influenced by random fluctuations. They have been employed in models of turbulence, population dynamics, neurophysiology, finance, fluid dynamics and climate modelling [3] [1]. Solving SPDEs is often inherently challenging however. Very few cases have analytical solutions, and numerical solutions have their own obstacles. Additional dimensionality imposes a higher computational cost, one may have to sample over many realisations of the noise, and some SPDEs are highly singular.

The Monte Carlo method is a natural avenue of approach when trying to solve SPDEs. Repeated samples of finite-difference or finite-element derived quantities may be obtained, and an estimate of a desired quantity of interest therefore computed. They can be exhorbitantly costly however. To achieve an accuracy of ϵ in one's estimate, the standard Monte Carlo estimator requires $O(\epsilon^{-2})$ samples be computed. To improve on this a number of variance reduction techniques have been introduced, their aim being to achieve an equivalent accuracy at a reduced cost.

The Multilevel Monte Carlo (MLMC) method is one such technique. Its tenet is to sample from a blend of cheap, less accurate samples and expensive, more accurate samples, and then combine those to obtain one's estimate. Provided this is implemented appropriately, this can result in a cost reduction when compared to standard Monte Carlo.

This dissertation investigates applying the MLMC to solve SPDEs. In particular, this dissertation has the following goals:

- Quantify the cost reductions achievable with MLMC methods, using MC methods as a baseline.
- Explore variations in MLMC implementations across different SPDEs.
- Present performance improvements achievable with high-performance implementations on GPUs

This dissertation will have the following structure. Section ?? will cover the relevant prerequisites. It will give an overview of the MLMC method as it pertains to SPDEs, before illustrating with a simple example. A review of the relevant literature will then be given. Finally, some relevant definitions will be covered. Section ?? will then begin by introducing the implemented MLMC process. Much of the section is

given over to validation of quantities, with particular focus given to the stochastic heat equation. Finally we go onto investigating results.

0.2 Prerequisites

0.2.1 Multilevel Monte Carlo

The MLMC method, introduced by Giles [2], is a variance reduction technique designed to substantially reduce the computational cost of estimating quantities arising from stochastic systems compared to the standard Monte Carlo method.

Given some computed quantity P , the standard Monte Carlo estimate of $\mathbb{E}[P]$ is given by the average of N independent samples of P :

$$\mathbb{E}[P] \approx \frac{1}{N} \sum_{n=1}^N P^{(n)} = P_{MC}$$

The variance of this estimate is $N^{-1}\mathbb{V}[P]$, the standard error is therefore $O(N^{\frac{1}{2}})$ and consequently to achieve an accuracy of ε , $N = O(\varepsilon^{-2})$ samples are required.

Bibliography

- [1] Diego Alonso-Orán, Aythami Bethencourt de León, Darryl D Holm, and So Takao. Modelling the climate and weather of a 2d lagrangian-averaged euler–boussinesq equation with transport noise. *Journal of Statistical Physics*, 179(5):1267–1303, 2020.
- [2] Michael B Giles. Multilevel monte carlo methods. *Acta numerica*, 24:259–328, 2015.
- [3] Étienne Pardoux et al. *Stochastic partial differential equations: An introduction*. Springer, 2021.