

# MATH 301

## INTRODUCTION TO PROOFS

Sina Hazratpour

Johns Hopkins University

Fall 2021

- integers
- rational numbers
- real numbers

## Relevant sections of the textbook

- Section B.2. (incomplete!)

## Quotients by relations

Recall from problem 5 of homework #4 that for each a binary relation  $R$  on a set  $X$  we can construct a set  $X/R$  whose elements are  $R$ -classes

$$[x]_R = \{y \in X \mid R(x, y)\}$$

for all  $x \in X$ .

## Quotients by relations

Recall from problem 5 of homework #4 that for each a binary relation  $R$  on a set  $X$  we can construct a set  $X/R$  whose elements are  $R$ -classes

$$[x]_R = \{y \in X \mid R(x, y)\}$$

for all  $x \in X$ . Now collect all such  $R$ -classes into one set:

$$X/R =_{\text{def}} \{[x]_R \mid x \in X\}$$

## Quotients by relations

Recall from problem 5 of homework #4 that for each a binary relation  $R$  on a set  $X$  we can construct a set  $X/R$  whose elements are  $R$ -classes

$$[x]_R = \{y \in X \mid R(x, y)\}$$

for all  $x \in X$ . Now collect all such  $R$ -classes into one set:

$$X/R =_{\text{def}} \{[x]_R \mid x \in X\}$$

We call the set  $X/R$  the  $R$ -quotient of  $X$  by the relation  $R$ .

## Example

*Consider the set of natural numbers with the usual ordering  $\leq: \mathbb{N} \rightarrow \mathbb{N} \rightarrow \mathbf{2}$  defined for  $m \in \mathbb{N}$  recursively by*

*$m \leq 0$  if and only if  $m = 0$ , and*

*$m \leq \text{succ}(n)$  if and only if  $m = \text{succ}(n)$  or  $m \leq n$ .*

## Example

*Consider the set of natural numbers with the usual ordering  $\leq: \mathbb{N} \rightarrow \mathbb{N} \rightarrow \mathbf{2}$  defined for  $m \in \mathbb{N}$  recursively by*

*$m \leq 0$  if and only if  $m = 0$ , and*

*$m \leq \text{succ}(n)$  if and only if  $m = \text{succ}(n)$  or  $m \leq n$ .*

*We have classes  $[n]$  forming a chain in the subset relation ordering:*

*$[0] \supset [1] \supset [2] \supset \dots$*

## Example

Consider the set of natural numbers with the usual ordering  $\leq: \mathbb{N} \rightarrow \mathbb{N} \rightarrow \mathbf{2}$  defined for  $m \in \mathbb{N}$  recursively by

$m \leq 0$  if and only if  $m = 0$ , and

$m \leq \text{succ}(n)$  if and only if  $m = \text{succ}(n)$  or  $m \leq n$ .

We have classes  $[n]$  forming a chain in the subset relation ordering:  $[0] \supset [1] \supset [2] \supset \dots$ . Note that  $0 \leq 1$  but  $[0] \neq [1]$ .



So far  $R$  is just a general binary relation. In general we do not have that

## Proposition

*Prove that if  $R$  is reflexive then  $\forall x \in X, x \in [x]$ .*

So far  $R$  is just a general binary relation. In general we do not have that

### Proposition

*Prove that if  $R$  is reflexive then  $\forall x \in X, x \in [x]$ .*

### Proposition

*Prove that if  $R$  is transitive then  $\forall x, y \in X, R(x, y) \wedge R(y, x) \Rightarrow [x] = [y]$ .*

So far  $R$  is just a general binary relation. In general we do not have that

### Proposition

*Prove that if  $R$  is reflexive then  $\forall x \in X, x \in [x]$ .*

### Proposition

*Prove that if  $R$  is transitive then  $\forall x, y \in X, R(x, y) \wedge R(y, x) \Rightarrow [x] = [y]$ .*

### Proposition

*Prove that if  $R$  is symmetric and transitive then*

*$\forall x, y \in X, R(x, y) \Rightarrow [x] = [y]$ .*

### Proposition

*Prove that if  $R$  is reflexive, symmetric and transitive then*

*$\forall x, y \in X, R(x, y) \Leftrightarrow [x] = [y]$ .*

Suppose we have a graph  $G$ . Define a relation  $R$  on vertices of  $G$  by imposing that

$$R(a, b) \Leftrightarrow \text{there is an edge from } a \text{ to } b.$$

Suppose we have a graph  $G$ . Define a relation  $R$  on vertices of  $G$  by imposing that

$$R(a, b) \Leftrightarrow \text{there is an edge from } a \text{ to } b.$$

- What is a class  $[a]$  for a vertex  $a$ ?

Suppose we have a graph  $G$ . Define a relation  $R$  on vertices of  $G$  by imposing that

$$R(a, b) \Leftrightarrow \text{there is an edge from } a \text{ to } b.$$

- What is a class  $[a]$  for a vertex  $a$ ?
- Show that if  $R(a, b) \wedge R(b, a)$  then it is not necessarily true that  $[a] = [b]$ .

Suppose we have a graph  $G$ . Define a relation  $R$  on vertices of  $G$  by imposing that

$$R(a, b) \Leftrightarrow \text{there is a path from } a \text{ to } b.$$

Suppose we have a graph  $G$ . Define a relation  $R$  on vertices of  $G$  by imposing that

$$R(a, b) \Leftrightarrow \text{there is a path from } a \text{ to } b.$$

- What is a class  $[a]$  for a vertex  $a$ ?



Suppose we have a graph  $G$ . Define a relation  $R$  on vertices of  $G$  by imposing that

$$R(a, b) \Leftrightarrow \text{there is a path from } a \text{ to } b.$$

- What is a class  $[a]$  for a vertex  $a$ ?
- Show that if  $R(a, b) \wedge R(b, a)$  then  $[a] = [b]$ .

## Proposition

*Prove that the function  $q: X \rightarrow X/R$  assigning to each  $x$  in  $X$  the class  $[x]$  in  $X/R$  is a surjection.*

# The universal mapping property of quotient construction

## Proposition

*Let  $R$  be a symmetric and transitive relation on a set  $X$ . For any set  $Y$ , precomposing with  $q$  yields a bijection*

$$(X/R \rightarrow Y) \cong \{f: X \rightarrow Y \mid \forall x, y \in X, R(x, y) \Rightarrow f(x) = f(y)\}$$

Recall that a relation  $R$  on a set  $A$  is called an **equivalence** if it satisfies the following conditions:

- **reflexivity**:  $\forall a \in A, R(a, a)$ ,
- **symmetry**:  $\forall a, b \in A, R(a, b) \rightarrow R(b, a)$ , and
- **transitivity**:  $\forall a, b, c \in A, R(a, b) \Rightarrow R(b, c) \Rightarrow R(a, c)$ .

Recall that a relation  $R$  on a set  $A$  is called an **equivalence** if it satisfies the following conditions:

- **reflexivity**:  $\forall a \in A, R(a, a)$ ,
- **symmetry**:  $\forall a, b \in A, R(a, b) \rightarrow R(b, a)$ , and
- **transitivity**:  $\forall a, b, c \in A, R(a, b) \Rightarrow R(b, c) \Rightarrow R(a, c)$ .

We usually denote an equivalence relation by the symbol  $\sim$  (instead of  $R$ ).

## Quotients by equivalence relations

For each equivalence  $\sim$  on a set  $X$  we can construct a set  $X/\sim$  whose elements are **equivalence classes**

$$[x]_{\sim} = \{y \in X \mid x \sim y\}$$

for all  $x \in X$ .

## Quotients by equivalence relations

For each equivalence  $\sim$  on a set  $X$  we can construct a set  $X/\sim$  whose elements are **equivalence classes**

$$[x]_{\sim} = \{y \in X \mid x \sim y\}$$

for all  $x \in X$ . Now collect all such equivalence classes into one set:

$$X/\sim =_{\text{def}} \{[x]_{\sim} \mid x \in X\}$$

## Quotients by equivalence relations

For each equivalence  $\sim$  on a set  $X$  we can construct a set  $X/\sim$  whose elements are **equivalence classes**

$$[x]_{\sim} = \{y \in X \mid x \sim y\}$$

for all  $x \in X$ . Now collect all such equivalence classes into one set:

$$X/\sim =_{\text{def}} \{[x]_{\sim} \mid x \in X\}$$

We call the set  $X/\sim$  the **quotient of  $X$  by equivalence relation  $\sim$** .



For an equivalence relation  $\sim$ , the surjection  $q: X \rightarrow X/\sim$  has an extra nice property:

$$q(x) = q(y) \Leftrightarrow R(x, y)$$

## Example of quotient by an equivalence relation

Consider the relation  $\sim$  on  $\mathbb{N} \times \mathbb{N}$ . where

$$(m, n) \sim (m', n') \Leftrightarrow m + n' = n + m'.$$

Prove that this relation is an equivalence.

## Example of quotient by an equivalence relation

Consider the relation  $\sim$  on  $\mathbb{N} \times \mathbb{N}$ . where

$$(m, n) \sim (m', n') \Leftrightarrow m + n' = n + m'.$$

Prove that this relation is an equivalence.

The equivalence class  $[(0, 0)]$  is the set  $\{(0, 0), (1, 1), (2, 2), \dots\}$ .


## Example of quotient by an equivalence relation

Consider the relation  $\sim$  on  $\mathbb{N} \times \mathbb{N}$ . where

$$(m, n) \sim (m', n') \Leftrightarrow m + n' = n + m'.$$

Prove that this relation is an equivalence.

The equivalence class  $[(0, 0)]$  is the set  $\{(0, 0), (1, 1), (2, 2), \dots\}$ . What is the equivalence class  $[(0, 1)]$ ?



# Integers

We define the set  $\mathbb{Z}$  of integers as the quotient  $\mathbb{N} \times \mathbb{N} / \sim$ .

# Integers

We define the set  $\mathbb{Z}$  of integers as the quotient  $\mathbb{N} \times \mathbb{N} / \sim$ .

- In other words, a pair  $(m, n)$  represents the would-be integer  $m - n$ .

# Integers

We define the set  $\mathbb{Z}$  of integers as the quotient  $\mathbb{N} \times \mathbb{N} / \sim$ .

- In other words, a pair  $(m, n)$  represents the would-be integer  $m - n$ .
- In this case, there are *canonical representatives* of the equivalence classes: those of the form  $(n, 0)$  or  $(0, n)$ .



## Addition on integers

We can define the operation of addition on  $\mathbb{Z}$  by an assignment  $+_{\sim} : \mathbb{Z} \times \mathbb{Z} \rightarrow \mathbb{Z}$  which assigns to the pair  $([(m, n)], [(m', n')])$  the class  $[(m + m', n + n')]$ .

### Exercise

*Show that the assignment  $+_{\sim}$  is well-defined, i.e. it defines a function.*

## Addition on integers

We can define the operation of addition on  $\mathbb{Z}$  by an assignment  $+_{\sim} : \mathbb{Z} \times \mathbb{Z} \rightarrow \mathbb{Z}$  which assigns to the pair  $([(m, n)], [(m', n')])$  the class  $[(m + m', n + n')]$ .

### Exercise

*Show that the assignment  $+_{\sim}$  is well-defined, i.e. it defines a function.*

### Exercise

- *Show that for all integers  $a$  we have  $0 + a = a = a + 0$ .*
- *Show that for all integers  $a, b, c$  we have  $(a + b) + c = a + (b + c)$ .*
- *Show that for all integers  $a, b$  we have  $a + b = b + a$ .*

We can define the operation of multiplication on  $\mathbb{Z}$  by an assignment  $\cdot_{\sim} : \mathbb{Z} \times \mathbb{Z} \rightarrow \mathbb{Z}$  which assigns to the pair  $([(m, n)], [(m', n')])$  the class  $[(m \cdot m', n \cdot n')]$ .

### Exercise

*Show that the assignment  $\cdot_{\sim}$  is well-defined, i.e. it defines a function.*

We can define the operation of multiplication on  $\mathbb{Z}$  by an assignment  $\cdot_{\sim} : \mathbb{Z} \times \mathbb{Z} \rightarrow \mathbb{Z}$  which assigns to the pair  $([(m, n)], [(m', n')])$  the class  $[(m \cdot m', n \cdot n')]$ .

### Exercise

*Show that the assignment  $\cdot_{\sim}$  is well-defined, i.e. it defines a function.*

### Exercise

- *Show that for all integers  $a$  we have  $0 + a = a = a + 0$ .*
- *Show that for all integers  $a, b, c$  we have  $(a + b) + c = a + (b + c)$ .*
- *Show that for all integers  $a, b$  we have  $a + b = b + a$ .*

Every function  $f: A \rightarrow B$  gives rise to an equivalence relation  $\sim_f$  on  $A$  defined by

$$a \sim_f b \Leftrightarrow f(a) = f(b) .$$

Every function  $f: A \rightarrow B$  gives rise to an equivalence relation  $\sim_f$  on  $A$  defined by

$$a \sim_f b \Leftrightarrow f(a) = f(b) .$$

That is to say that  $a \sim_f b$  if  $a, b$  are in the same fibre of  $f$ .

Every function  $f: A \rightarrow B$  gives rise to an equivalence relation  $\sim_f$  on  $A$  defined by

$$a \sim_f b \Leftrightarrow f(a) = f(b) .$$

That is to say that  $a \sim_f b$  if  $a, b$  are in the same fibre of  $f$ .


### Exercise

*Show that the relation above is indeed an equivalence relation.*

# Image factorization

## Proposition

*Suppose  $f: A \rightarrow B$  is a function. We can factor  $f$  into a surjection followed by a bijection followed by an injection.*



*that is there are functions  $q, g, i$  such that  $f = i \circ g \circ q$ , where  $q$  is a surjection,  $g$  is a bijection, and  $i$  is an injection.*



## Proof.

We have to construct the sets  $C$ ,  $D$  and a surjection  $q$ , a bijection  $g$  and an injection  $i$ .

## Proof.

We have to construct the sets  $C$ ,  $D$  and a surjection  $q$ , a bijection  $g$  and an injection  $i$ . Now we define  $C$  to be  $A / \sim_f$ , and we define  $D$  to be the image  $f_*(A)$  of  $A$  under  $f$ .

## Proof.

We have to construct the sets  $C$ ,  $D$  and a surjection  $q$ , a bijection  $g$  and an injection  $i$ . Now we define  $C$  to be  $A / \sim_f$ , and we define  $D$  to be the image  $f_*(A)$  of  $A$  under  $f$ . We also define  $q$  to be the obvious quotient map and  $i$  to be the obvious inclusion. As shown before,  $q$  is surjective and  $i$  is injective.

## Proof.

We have to construct the sets  $C$ ,  $D$  and a surjection  $q$ , a bijection  $g$  and an injection  $i$ . Now we define  $C$  to be  $A / \sim_f$ , and we define  $D$  to be the image  $f_*(A)$  of  $A$  under  $f$ . We also define  $q$  to be the obvious quotient map and  $i$  to be the obvious inclusion. As shown before,  $q$  is surjective and  $i$  is injective. We define  $g$  to be the assignment which takes an equivalence class  $[x]$  to the element  $f(x) \in B$ .

## Proof.

We have to construct the sets  $C$ ,  $D$  and a surjection  $q$ , a bijection  $g$  and an injection  $i$ . Now we define  $C$  to be  $A / \sim_f$ , and we define  $D$  to be the image  $f_*(A)$  of  $A$  under  $f$ . We also define  $q$  to be the obvious quotient map and  $i$  to be the obvious inclusion. As shown before,  $q$  is surjective and  $i$  is injective. We define  $g$  to be the assignment which takes an equivalence class  $[x]$  to the element  $f(x) \in B$ . Note that  $g$  is well-defined, since if  $[x] = [y]$  then  $x \sim_f y$  and therefore, by the definition of  $\sim_f$ , we have  $f(x) = f(y)$ .

## Proof.

We have to construct the sets  $C$ ,  $D$  and a surjection  $q$ , a bijection  $g$  and an injection  $i$ . Now we define  $C$  to be  $A / \sim_f$ , and we define  $D$  to be the image  $f_*(A)$  of  $A$  under  $f$ . We also define  $q$  to be the obvious quotient map and  $i$  to be the obvious inclusion. As shown before,  $q$  is surjective and  $i$  is injective. We define  $g$  to be the assignment which takes an equivalence class  $[x]$  to the element  $f(x) \in B$ . Note that  $g$  is well-defined, since if  $[x] = [y]$  then  $x \sim_f y$  and therefore, by the definition of  $\sim_f$ , we have  $f(x) = f(y)$ . We now show that  $g$  is a bijection.

## Proof.

We have to construct the sets  $C$ ,  $D$  and a surjection  $q$ , a bijection  $g$  and an injection  $i$ . Now we define  $C$  to be  $A / \sim_f$ , and we define  $D$  to be the image  $f_*(A)$  of  $A$  under  $f$ . We also define  $q$  to be the obvious quotient map and  $i$  to be the obvious inclusion. As shown before,  $q$  is surjective and  $i$  is injective. We define  $g$  to be the assignment which takes an equivalence class  $[x]$  to the element  $f(x) \in B$ . Note that  $g$  is well-defined, since if  $[x] = [y]$  then  $x \sim_f y$  and therefore, by the definition of  $\sim_f$ , we have  $f(x) = f(y)$ . We now show that  $g$  is a bijection.  $g$  is injective since for every  $x, y \in A$ , if  $g([x]) = g([y])$  then  $f(x) = f(y)$  and therefore,  $[x] = [y]$ .

## Proof.


We have to construct the sets  $C$ ,  $D$  and a surjection  $q$ , a bijection  $g$  and an injection  $i$ . Now we define  $C$  to be  $A / \sim_f$ , and we define  $D$  to be the image  $f_*(A)$  of  $A$  under  $f$ . We also define  $q$  to be the obvious quotient map and  $i$  to be the obvious inclusion. As shown before,  $q$  is surjective and  $i$  is injective. We define  $g$  to be the assignment which takes an equivalence class  $[x]$  to the element  $f(x) \in B$ . Note that  $g$  is well-defined, since if  $[x] = [y]$  then  $x \sim_f y$  and therefore, by the definition of  $\sim_f$ , we have  $f(x) = f(y)$ . We now show that  $g$  is a bijection.  $g$  is injective since for every  $x, y \in A$ , if  $g([x]) = g([y])$  then  $f(x) = f(y)$  and therefore,  $[x] = [y]$ . Also,  $g$  is surjective: given  $b$  in  $f_*(A)$  there is some  $a \in A$  such that  $b = f(a) = g([a])$ .




## Proof.

We have to construct the sets  $C$ ,  $D$  and a surjection  $q$ , a bijection  $g$  and an injection  $i$ . Now we define  $C$  to be  $A / \sim_f$ , and we define  $D$  to be the image  $f_*(A)$  of  $A$  under  $f$ . We also define  $q$  to be the obvious quotient map and  $i$  to be the obvious inclusion. As shown before,  $q$  is surjective and  $i$  is injective. We define  $g$  to be the assignment which takes an equivalence class  $[x]$  to the element  $f(x) \in B$ . Note that  $g$  is well-defined, since if  $[x] = [y]$  then  $x \sim_f y$  and therefore, by the definition of  $\sim_f$ , we have  $f(x) = f(y)$ . We now show that  $g$  is a bijection.  $g$  is injective since for every  $x, y \in A$ , if  $g([x]) = g([y])$  then  $f(x) = f(y)$  and therefore,  $[x] = [y]$ . Also,  $g$  is surjective: given  $b$  in  $f_*(A)$  there is some  $a \in A$  such that  $b = f(a) = g([a])$ . □

Our factorization diagram becomes



Our factorization diagram becomes



In fact,  $g \circ q = p: X \rightarrow \mathbf{Im}(f)$ .

## Definition

For a function  $f: X \rightarrow X$ , define

$$\text{Fix}(f) =_{\text{def}} \{x \in X \mid f(x) = x\}.$$

We call  $\text{Fix}(f)$  the set of *fix-points* of  $f$ .

## Definition

For a function  $f: X \rightarrow X$ , define

$$\text{Fix}(f) =_{\text{def}} \{x \in X \mid f(x) = x\}.$$

We call  $\text{Fix}(f)$  the set of *fix-points* of  $f$ .

## Definition

A function  $r$  is called an *idempotent* if  $r \circ r = r$ .

## Definition

For a function  $f: X \rightarrow X$ , define

$$\text{Fix}(f) =_{\text{def}} \{x \in X \mid f(x) = x\}.$$

We call  $\text{Fix}(f)$  the set of *fix-points* of  $f$ .

## Definition

A function  $r$  is called an *idempotent* if  $r \circ r = r$ .

## Exercise

Show that if  $f: X \rightarrow X$  is idempotent, then  $\text{Fix}(f) \cong \text{Im}(f)$ .

## Definition

For a function  $f: X \rightarrow X$ , define

$$\text{Fix}(f) =_{\text{def}} \{x \in X \mid f(x) = x\}.$$

We call  $\text{Fix}(f)$  the set of **fix-points** of  $f$ .

## Definition

A function  $r$  is called an **idempotent** if  $r \circ r = r$ .

## Exercise

Show that if  $f: X \rightarrow X$  is idempotent, then  $\text{Fix}(f) \cong \text{Im}(f)$ .

## Exercise

For an idempotent function  $f: X \rightarrow X$ , show that

$$X / \sim_f \cong \text{Fix}(f) \cong \text{Im}(f)$$

## Exercise

*Construct an idempotent  $f: \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N} \times \mathbb{N}$  such that  $\text{Fix}(f)$  is in bijection with the set of integers.*



## Exercise

Suppose  $r: A \rightarrow B$  is a retraction. Show that

$$B \cong A / \sim_r$$

## Exercise

Suppose  $r: A \rightarrow B$  is a retraction. Show that

$$B \cong A / \sim_r$$

## Induction for integers

We identify a natural number  $n$  with the corresponding non-negative integer, i.e. with the image of  $(n, 0) \in \mathbb{N} \times \mathbb{N}$  in  $\mathbb{Z}$ .

# Induction for integers

We identify a natural number  $n$  with the corresponding non-negative integer, i.e. with the image of  $(n, 0) \in \mathbb{N} \times \mathbb{N}$  in  $\mathbb{Z}$ .

## Lemma

*Suppose  $P: \mathbb{Z} \rightarrow \mathbb{Prop}$  is a predicate over integers, and*

- *$P(0)$  holds,*
- *$\forall n : \mathbb{N}, P(n) \Rightarrow P(\text{succ}(n))$ , and*
- *$\forall n : \mathbb{N}, P(-n) \rightarrow P(-\text{succ}(n))$ .*

*Then we have  $\forall z : \mathbb{Z}, P(z)$ .*

We constructed the integers  $\mathbb{Z}$  as a quotient of  $\mathbb{N} \times \mathbb{N}$ , and observed that this quotient is generated by an idempotent.

We constructed the integers  $\mathbb{Z}$  as a quotient of  $\mathbb{N} \times \mathbb{N}$ , and observed that this quotient is generated by an idempotent.

We construct the *field* of rationals  $\mathbb{Q}$  along the same lines as well, namely as the quotient

$$\mathbb{Q} =_{\text{def}} (\mathbb{Z} \times \mathbb{N}) / \approx$$

where

$$(u, a) \approx (v, b) =_{\text{def}} (u(b+1) = v(a+1)).$$

We constructed the integers  $\mathbb{Z}$  as a quotient of  $\mathbb{N} \times \mathbb{N}$ , and observed that this quotient is generated by an idempotent.

We construct the *field* of rationals  $\mathbb{Q}$  along the same lines as well, namely as the quotient

$$\mathbb{Q} =_{\text{def}} (\mathbb{Z} \times \mathbb{N}) / \approx$$

where

$$(u, a) \approx (v, b) =_{\text{def}} (u(b+1) = v(a+1)).$$

In other words, a pair  $(u, a)$  represents the rational number  $u/(1+a)$ .

We constructed the integers  $\mathbb{Z}$  as a quotient of  $\mathbb{N} \times \mathbb{N}$ , and observed that this quotient is generated by an idempotent.

We construct the *field* of rationals  $\mathbb{Q}$  along the same lines as well, namely as the quotient

$$\mathbb{Q} =_{\text{def}} (\mathbb{Z} \times \mathbb{N}) / \approx$$

where

$$(u, a) \approx (v, b) =_{\text{def}} (u(b+1) = v(a+1)).$$

In other words, a pair  $(u, a)$  represents the rational number  $u/(1+a)$ .

Here too we have a canonical choice of representatives, namely fractions in lowest terms.



# The arithmetic of rational numbers

We write down the arithmetical operations on  $\mathbb{Q}$  so that we can compute with fractions.

# The order on rational numbers

We equip  $\mathbb{Q}$  with a total order.

# Real numbers

## Questions

Thanks for your attention!