

# MATH 301

## INTRODUCTION TO PROOFS

Sina Hazratpour

Johns Hopkins University

Fall 2021

- Recursion

## Relevant sections of the textbook

- Chapter 4

# Overview

- 1 Recursion
- 2 Applications of recursion theorem
- 3 Recursion in Lean

Recall that in the last lecture, we defined the set of **natural numbers**  $\mathbb{N}$  to be a set *generated* by the by the number 0 and the **successor function**  $\text{succ}: \mathbb{N} \rightarrow \mathbb{N}$ .

Recall that in the last lecture, we defined the set of **natural numbers**  $\mathbb{N}$  to be a set *generated* by the by the number 0 and the **successor function**  $\text{succ}: \mathbb{N} \rightarrow \mathbb{N}$ .

This means that the only way to construct a natural number is to start with 0 and apply the successor function finitely many times.

Recall that in the last lecture, we defined the set of **natural numbers**  $\mathbb{N}$  to be a set *generated* by the by the number 0 and the **successor function**  $\text{succ}: \mathbb{N} \rightarrow \mathbb{N}$ .

This means that the only way to construct a natural number is to start with 0 and apply the successor function finitely many times. Therefore, natural numbers are

$$0, \text{succ}(0), \text{succ}(\text{succ}(0)), \dots$$

Recall that in the last lecture, we defined the set of **natural numbers**  $\mathbb{N}$  to be a set *generated* by the by the number 0 and the **successor function**  $\text{succ}: \mathbb{N} \rightarrow \mathbb{N}$ .

This means that the only way to construct a natural number is to start with 0 and apply the successor function finitely many times. Therefore, natural numbers are

$$0, \text{succ}(0), \text{succ}(\text{succ}(0)), \dots$$

We also postulated the principle of induction on natural numbers.

## Predicates and subsets

Recall that we can think of a predicate  $P$  on natural numbers as a function  $P: \mathbb{N} \rightarrow \mathbf{2}$  where the set  $\mathbf{2}$  consists of truth values  $\perp$  and  $\top$ .



## Predicates and subsets

Recall that we can think of a predicate  $P$  on natural numbers as a function  $P: \mathbb{N} \rightarrow \mathbf{2}$  where the set  $\mathbf{2}$  consists of truth values  $\perp$  and  $\top$ .

Note that the set  $\mathbb{N} \rightarrow \mathbf{2}$  is in bijection with  $\mathcal{P}(\mathbb{N})$ .

## Predicates and subsets

Recall that we can think of a predicate  $P$  on natural numbers as a function  $P: \mathbb{N} \rightarrow \mathbf{2}$  where the set  $\mathbf{2}$  consists of truth values  $\perp$  and  $\top$ .

Note that the set  $\mathbb{N} \rightarrow \mathbf{2}$  is in bijection with  $\mathcal{P}(\mathbb{N})$ .

In one way, we construct a function

$$\eta(-): (\mathbb{N} \rightarrow \mathbf{2}) \rightarrow \mathcal{P}(\mathbb{N})$$

whose value at a predicate  $P$  is the set consisting of all  $n \in \mathbb{N}$  such that  $P(n)$  is true, i.e.

$$\eta(P) =_{\text{def}} \{n \in \mathbb{N} \mid P(n)\}$$

## Predicates and subsets

Recall that we can think of a predicate  $P$  on natural numbers as a function  $P: \mathbb{N} \rightarrow \mathbf{2}$  where the set  $\mathbf{2}$  consists of truth values  $\perp$  and  $\top$ .

Note that the set  $\mathbb{N} \rightarrow \mathbf{2}$  is in bijection with  $\mathcal{P}(\mathbb{N})$ .

In one way, we construct a function

$$\eta(-): (\mathbb{N} \rightarrow \mathbf{2}) \rightarrow \mathcal{P}(\mathbb{N})$$

whose value at a predicate  $P$  is the set consisting of all  $n \in \mathbb{N}$  such that  $P(n)$  is true, i.e.

$$\eta(P) =_{\text{def}} \{n \in \mathbb{N} \mid P(n)\}$$

In the other direction, we take a subset  $S$  of  $\mathbb{N}$  to the characteristic function  $\chi_S: \mathbb{N} \rightarrow \mathbf{2}$ .

# The principle of induction

The principle of induction says that for any property  $P: \mathbb{N} \rightarrow \mathbf{2}$  of natural numbers, if

- 1  $P(0)$  holds, and
  - 2 whenever  $P(n)$  holds then  $P(n + 1)$  holds,
- we have that  $P$  holds of every natural number.

## The principle of induction reformulated

Let  $S \subseteq \mathbb{N}$  be any set of natural numbers that contains 0 and is closed under the successor operation. Then  $S = \mathbb{N}$ .

## Proofs vs computation

We saw that the principle of induction is a very powerful tool in **proving** universally quantified statements about natural numbers.

## Proofs vs computation

We saw that the principle of induction is a very powerful tool in proving universally quantified statements about natural numbers.

### Example

- *For any finite set  $S$ , if  $S$  has  $n$  elements, then there are  $2^n$  subsets of  $S$ .*
- *For every  $n \in \mathbb{N}$ , we have  $0^2 + 1^2 + 2^2 + \dots + n^2 = \frac{1}{6} n (1 + n) (1 + 2n)$ .*

## Proofs vs computation

We saw that the principle of induction is a very powerful tool in proving universally quantified statements about natural numbers.

But, we also need to **compute** with natural numbers. At the very least, we should be able to define the arithmetic operations  $+$ ,  $\times$ , etc.



## Proofs vs computation

We saw that the principle of induction is a very powerful tool in proving universally quantified statements about natural numbers.

That is why we need another principle to help us with computation of natural numbers. This is the so-called principle of **recursion** which in fact can be proved from the principle of induction!

# Recursion theorem

## Theorem

*Let  $A$  be a set. For all  $a \in A$  and all  $g: \mathbb{N} \times A \rightarrow A$ , there is a unique function  $f: \mathbb{N} \rightarrow A$  such that*

- 1  $f(0) = a$
- 2  $f(\text{succ}(n)) = g(n, f(n))$  for all  $n \in \mathbb{N}$ .

# Recursion theorem

## Theorem

*Let  $A$  be a set. For all  $a \in A$  and all  $g: \mathbb{N} \times A \rightarrow A$ , there is a unique function  $f: \mathbb{N} \rightarrow A$  such that*

- ①  $f(0) = a$
- ②  $f(\text{succ}(n)) = g(n, f(n))$  for all  $n \in \mathbb{N}$ .

## Proof.

Theorem 4.1.2 (Recursion theorem) Page 145.



# Recursion theorem

## Theorem

*Let  $A$  be a set. For all  $a \in A$  and all  $g: \mathbb{N} \times A \rightarrow A$ , there is a unique function  $f: \mathbb{N} \rightarrow A$  such that*

- ①  $f(0) = a$
- ②  $f(\text{succ}(n)) = g(n, f(n))$  for all  $n \in \mathbb{N}$ .

## Proof.

Theorem 4.1.2 (Recursion theorem) Page 145.



Since for every function  $g$  such function  $f$  is uniquely determined, we write  $\text{rec}(g)$  for it.

# Recursion theorem

## Theorem

*Let  $A$  be a set. For all  $a \in A$  and all  $g: \mathbb{N} \times A \rightarrow A$ , there is a unique function  $f: \mathbb{N} \rightarrow A$  such that*

- 1  $f(0) = a$
- 2  $f(\text{succ}(n)) = g(n, f(n))$  for all  $n \in \mathbb{N}$ .

We have

$$\text{rec}(g)(0) = a$$

## Recursion theorem

We have

$$\text{rec}(g)(0) = a$$

$$\text{rec}(g)(1) = \text{rec}(g)(\text{succ}(0)) = g(0, \text{rec}(g)(0)) = g(0, a)$$

# Recursion theorem

We have

$$\text{rec}(g)(0) = a$$

$$\text{rec}(g)(1) = \text{rec}(g)(\text{succ}(0)) = g(0, \text{rec}(g)(0)) = g(0, a)$$

$$\text{rec}(g)(2) = \text{rec}(g)(\text{succ}(1)) = g(1, \text{rec}(g)(1)) = g(1, g(0, a))$$

# Recursion theorem

We have

$$\text{rec}(g)(0) = a$$

$$\text{rec}(g)(1) = \text{rec}(g)(\text{succ}(0)) = g(0, \text{rec}(g)(0)) = g(0, a)$$

$$\text{rec}(g)(2) = \text{rec}(g)(\text{succ}(1)) = g(1, \text{rec}(g)(1)) = g(1, g(0, a))$$

$\vdots$



## Recursion, practically!

In order to specify a function  $f: \mathbb{N} \rightarrow A$ , it suffices to define  $f(0)$  and, for given  $n \in \mathbb{N}$ , assume that  $f(n)$  has been defined, and define  $f(\text{succ}(n))$  in terms of  $n$  and  $f(n)$ .

# Overview

- 1 Recursion
- 2 Applications of recursion theorem**
- 3 Recursion in Lean

## Addition by recursion

We define additions of natural numbers as a function  $+: \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ .

## Addition by recursion

We define additions of natural numbers as a function  $+: \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ .

This means for every  $m \in \mathbb{N}$ , we have to define a function  $m + (-): \mathbb{N} \rightarrow \mathbb{N}$ .

## Addition by recursion

We define additions of natural numbers as a function  $+: \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ .

This means for every  $m \in \mathbb{N}$ , we have to define a function  $m + (-): \mathbb{N} \rightarrow \mathbb{N}$ .

We define the latter by recursion: consider the function  $g: \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$  given by the assignment  $g(i, j) = \text{succ}(j)$ .

## Addition by recursion

We define additions of natural numbers as a function  $+: \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ .

This means for every  $m \in \mathbb{N}$ , we have to define a function  $m + (-): \mathbb{N} \rightarrow \mathbb{N}$ .

We define the latter by recursion: consider the function  $g: \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$  given by the assignment  $g(i, j) = \text{succ}(j)$ .

Choose  $a = m$  in the recursion theorem. Therefore,

## Addition by recursion

We define additions of natural numbers as a function  $+: \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ .

This means for every  $m \in \mathbb{N}$ , we have to define a function  $m + (-): \mathbb{N} \rightarrow \mathbb{N}$ .

We define the latter by recursion: consider the function  $g: \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$  given by the assignment  $g(i, j) = \text{succ}(j)$ .

Choose  $a = m$  in the recursion theorem. Therefore,

$$\text{rec}(g)(0) = m$$

$$\text{rec}(g)(1) = \text{rec}(g)(\text{succ}(0)) = g(0, \text{rec}(g)(0)) = g(0, m) = \text{succ}(m)$$

$$\text{rec}(g)(2) = \text{rec}(g)(\text{succ}(1)) = g(1, \text{rec}(g)(1)) = \text{succ}(\text{succ}(m))$$

$$\text{rec}(g)(\text{succ}(n)) = g(n, \text{rec}(g)(n))$$

## Addition by recursion

We now define  $m + (-): \mathbb{N} \rightarrow \mathbb{N}$  to be  $\text{rec}(g): \mathbb{N} \rightarrow \mathbb{N}$ .



## Addition by recursion

We now define  $m + (-): \mathbb{N} \rightarrow \mathbb{N}$  to be  $\text{rec}(g): \mathbb{N} \rightarrow \mathbb{N}$ .

$$m + 0 = m \tag{1}$$

$$m + \text{succ}(n) = \text{succ}(m + n) \tag{2}$$

## Addition by recursion

We now define  $m + (-) : \mathbb{N} \rightarrow \mathbb{N}$  to be  $\text{rec}(g) : \mathbb{N} \rightarrow \mathbb{N}$ .

$$m + 0 = m \tag{1}$$

$$m + \text{succ}(n) = \text{succ}(m + n) \tag{2}$$

Therefore,

$$m + 1 = m + \text{succ}(0) = \text{succ}(m + 0) = \text{succ}(m) \tag{3}$$

## Addition by recursion

We now define  $m + (-) : \mathbb{N} \rightarrow \mathbb{N}$  to be  $\text{rec}(g) : \mathbb{N} \rightarrow \mathbb{N}$ .

$$m + 0 = m \tag{1}$$

$$m + \text{succ}(n) = \text{succ}(m + n) \tag{2}$$

Therefore,

$$m + 1 = m + \text{succ}(0) = \text{succ}(m + 0) = \text{succ}(m) \tag{3}$$

In particular,

$$1 + 1 = \text{succ}(1) = \text{succ}(\text{succ}(1)) = 2$$

## Combining recursion and induction

### Proposition

*For every natural numbers  $m$  and  $n$ , we have  $m + n = n + m$ .*

### Proof.

Let  $m$  to be an arbitrary natural number. we use induction on  $n$  to prove that  $1 + n = n + 1$  for all  $n \in \mathbb{N}$ .

## Combining recursion and induction

### Proposition

*For every natural numbers  $m$  and  $n$ , we have  $m + n = n + m$ .*

### Proof.

Let  $m$  to be an arbitrary natural number. we use induction on  $n$  to prove that  $1 + n = n + 1$  for all  $n \in \mathbb{N}$ .

When  $n = 0$ , by equations (1) and (2), we have

$$1 + 0 = 1 = \text{succ}(0) = \text{succ}(0 + 0) = 0 + \text{succ}(0) = 0 + 1.$$

## Combining recursion and induction

### Proposition

*For every natural numbers  $m$  and  $n$ , we have  $m + n = n + m$ .*

### Proof.

Let  $m$  to be an arbitrary natural number. we use induction on  $n$  to prove that  $1 + n = n + 1$  for all  $n \in \mathbb{N}$ .

When  $n = 0$ , by equations (1) and (2), we have

$$1 + 0 = 1 = \text{succ}(0) = \text{succ}(0 + 0) = 0 + \text{succ}(0) = 0 + 1.$$

Suppose that  $1 + n = n + 1$ . We want to show that  $1 + \text{succ}(n) = \text{succ}(n) + 1$ .

# Combining recursion and induction

## Proposition

*For every natural numbers  $m$  and  $n$ , we have  $m + n = n + m$ .*

## Proof.

Let  $m$  to be an arbitrary natural number. we use induction on  $n$  to prove that  $1 + n = n + 1$  for all  $n \in \mathbb{N}$ .

When  $n = 0$ , by equations (1) and (2), we have

$$1 + 0 = 1 = \text{succ}(0) = \text{succ}(0 + 0) = 0 + \text{succ}(0) = 0 + 1.$$

Suppose that  $1 + n = n + 1$ . We want to show that  $1 + \text{succ}(n) = \text{succ}(n) + 1$ .

But, by definition of function  $m + -$  for  $m = 1$ ,

$$1 + \text{succ}(n) = \text{succ}(1 + n) = \text{succ}(n + 1) = \text{succ}(\text{succ}(n)) = \text{succ}(n) + 1$$

The last two equations above follow from equation (3).



# Overview

- 1 Recursion
- 2 Applications of recursion theorem
- 3 Recursion in Lean**