

Introductory Statistics Homework Analysis

Overview

The file name with student responses and scores is Topico6.AB.csv. 50 students submitted the homework assignment.

Full credit for this homework assignment is 25. Each student was given 25 questions and this homework assignment contains a total of 89 questions.

The mean score of the 50 students who submitted the homework assignment is 6.24. The lowest and highest scores are 0 and 12, respectively.

The histogram and summary statistics for the scores are given by Figure 1 and Table 1.

```
## Warning: Removed 2 rows containing missing values (geom_bar).
```

Mean	Std.dev	Min	Q1	Median	Q3	Max
6.24	2.38	0.00	5.00	6.00	8.00	12.00

Table 1: Summary statistics of the scores




Figure 1: Histogram of scores.

Topic Learning Outcomes

- A. Use standardizing to determine how many standard deviations an observation is away from the mean value.
- B. Use z-scores to compare observations for different quantitative variables.
- C. Explain how standardizing affects the shape, center, and variability of the distribution of a quantitative variable.
- D. Determine which quantitative variables could be modeled using the normal distribution by interpreting graphical representations of the variable.
- E. Apply the 68-95-99.
- F. Find percentile or area values for any given observation from a normal distribution.
- G. Find the value of an observation when given a percentile or area value from the normal distribution.

Table 2 and Figure 2 show the summary of percentage scores by learning outcome. Among all learning outcomes, Outcome D has the highest correct percentage, while Outcome E has the lowest.

	Mean	Std.dev	Min	Median	Max
D	61.00	45.50	0.00	100.00	100.00
B	38.00	37.20	0.00	50.00	100.00
A	23.00	20.73	0.00	25.00	50.00
C	22.00	27.03	0.00	0.00	100.00
G	21.50	18.22	0.00	25.00	50.00
F	20.80	19.78	0.00	20.00	60.00
E	16.67	12.14	0.00	16.67	33.33

Table 2: Summary statistics of the student percentage correct scores on the topic learning outcomes. The table is sorted from the highest mean to the lowest.

To analyze the students' performance on different learning outcomes, we consider the generalized mixed effects model:

$$g(E[Y_{ij}|u_j]) = \tau_i + u_j$$

where $i = 1, \dots, 7$ learning outcomes; $j = 1, \dots, 50$ students. Y_{ij} is the score (scaled in $[0, 1]$) of the j th student in the i th learning outcome. τ_i is the fixed effect, which represents the actual level of the learning outcome i . u_j is the random effect from the students with $u_j \sim N(0, \sigma_u^2)$.

Since $Y_{ij} \in [0, 1]$, and most of y_{ij} take the value 1 (see Figure 3), it is not easy to find an appropriate model. Beta distribution is possible if 0 and 1 are not in the range of Y_{ij} . Binomial distribution is another possibility if we do not scale the scores. Hence we assume that Y is binomial with the logit link function, then the results are in Table 3 and Figure 4.

	est.	lower	upper
(Intercept)	0.44	0.10	0.79
ObjectiveB	-0.93	-1.40	-0.47
ObjectiveA	-1.64	-2.14	-1.14
ObjectiveC	-1.70	-2.20	-1.19
ObjectiveG	-1.72	-2.23	-1.22
ObjectiveF	-1.76	-2.28	-1.25
ObjectiveE	-2.03	-2.56	-1.49

Table 3: 95% confidence intervals of the fixed effects




Figure 2: Side-by-side boxplots of the correct percentages by learning outcome.




Figure 3: Histogram of the scaled scores by learning outcome.

It shows that D is the best understood learning outcome. Outcomes E,F,G,C,A,B are significantly worse than D.

Nevertheless, the assumption is probably incorrect, as we could check the residuals in Figure 5.





Figure 5: QQ-plot of the residuals by the generalized mixed effects model

Figure 4: 95% confidence intervals of the fixed effects coefficients

Question Sets

Table 4 and Figure ?? shows the summary statistics of the percentage correct score by question for each question set. Percentage correct scores for questions in the same question set should have small variability.

Qset	LO	#	Mean	Std.dev	Min	Median	Max
S	G	5	23.39	19.03	0.00	27.27	50.00
Q	F	5	28.71	18.83	0.00	28.57	50.00
B	A	4	46.13	17.26	25.00	46.43	66.67
J	D	4	62.08	14.74	50.00	57.50	83.33
N	F	5	18.95	14.61	0.00	20.00	37.50
L	E	9	23.02	14.35	5.56	26.32	42.11
U	G	5	20.00	13.94	0.00	16.67	33.33
R	G	5	16.71	13.21	0.00	20.00	33.33
M	F	5	31.24	12.06	25.00	25.00	52.63
O	F	5	11.65	12.03	0.00	14.29	28.57
T	G	5	14.98	11.41	0.00	14.29	27.27
K	E	9	9.56	10.68	0.00	5.88	31.25
A	A	4	42.27	9.74	30.00	43.08	52.94
P	F	5	6.51	6.46	0.00	7.14	14.29
I	D	2	64.16	1.18	63.33	64.16	65.00
C	A	4	0.00	0.00	0.00	0.00	0.00
D	A	4	0.00	0.00	0.00	0.00	0.00
E	B	1	40.00	0.00	40.00	40.00	40.00
F	B	1	36.00	0.00	36.00	36.00	36.00
G	C	1	20.00	0.00	20.00	20.00	20.00
H	C	1	24.00	0.00	24.00	24.00	24.00

Table 4: Summary statistics of the percentage correct score by question for each question sets. The table is sorted from the largest standard deviation to the smallest.




Figure 6: Side-by-side boxplots of the question correct percentage score for each question set

Questions

Table 5 compares the performance on each question.

ID	LO	Qset	Name	Type	FullPt	QinSet	N	CrtPct	Count	NA's	Mean	Std	Flag
1	A	A	1	MC	1	1	4	40.00	10	40	0.40	0.52	
2	A	A	2	MC	1	1	4	46.15	13	37	0.46	0.52	*
3	A	A	3	MC	1	1	4	30.00	10	40	0.30	0.48	*
4	A	A	4	MC	1	1	4	52.94	17	33	0.53	0.51	*
5	A	B	1	MC	1	1	4	42.86	14	36	0.43	0.51	*
6	A	B	2	MC	1	1	4	50.00	16	34	0.50	0.52	*
7	A	B	3	MC	1	1	4	25.00	8	42	0.25	0.46	*
8	A	B	4	MC	1	1	4	66.67	12	38	0.67	0.49	*
9	A	C	1	FB	1	1	4	0.00	11	39	0.00	0.00	
10	A	C	2	FB	1	1	4	0.00	11	39	0.00	0.00	
11	A	C	3	FB	1	1	4	0.00	14	36	0.00	0.00	
12	A	C	4	FB	1	1	4	0.00	14	36	0.00	0.00	
13	A	D	1	FB	1	1	4	0.00	10	40	0.00	0.00	
14	A	D	2	FB	1	1	4	0.00	13	37	0.00	0.00	
15	A	D	3	FB	1	1	4	0.00	11	39	0.00	0.00	
16	A	D	4	FB	1	1	4	0.00	16	34	0.00	0.00	
17	B	E	compare1	MC	1	1	1	40.00	50	0	0.40	0.49	
18	B	F	compare2	MC	1	1	1	36.00	50	0	0.36	0.48	
19	C	G	1	MC	1	1	1	20.00	50	0	0.20	0.40	
20	C	H	1	MC	1	1	1	24.00	50	0	0.24	0.43	
21	D	I	feet	TF	1	1	2	65.00	20	30	0.65	0.49	
22	D	I	lowtemp	TF	1	1	2	63.33	30	20	0.63	0.49	
23	D	J	gain	TF	1	1	4	60.00	10	40	0.60	0.52	*
24	D	J	mpg	TF	1	1	4	83.33	6	44	0.83	0.41	*
25	D	J	blowhole	TF	1	1	4	50.00	14	36	0.50	0.52	*
26	D	J	CDs	TF	1	1	4	55.00	20	30	0.55	0.51	*
27	E	K	whale1	FB	1	3	9	21.05	19	31	0.21	0.42	*
28	E	K	whale2	FB	1	3	9	0.00	18	32	0.00	0.00	*
29	E	K	whale3	FB	1	3	9	0.00	14	36	0.00	0.00	*
30	E	K	cow1	FB	1	3	9	0.00	12	38	0.00	0.00	*

31	E	K	cow2	FB	1	3	9	5.56	18	32	0.06	0.24	*
32	E	K	cow3	FB	1	3	9	11.76	17	33	0.12	0.33	*
33	E	K	bulbs1	FB	1	3	9	31.25	16	34	0.31	0.48	*
34	E	K	bulbs2	FB	1	3	9	5.88	17	33	0.06	0.24	*
35	E	K	bulbs3	FB	1	3	9	10.53	19	31	0.11	0.32	*
36	E	L	heights1	MC	1	3	9	15.79	19	31	0.16	0.37	*
37	E	L	heights2	MC	1	3	9	5.56	18	32	0.06	0.24	*
38	E	L	heights3	MC	1	3	9	42.11	19	31	0.42	0.51	*
39	E	L	mm1	MC	1	3	9	7.69	13	37	0.08	0.28	*
40	E	L	mm2	MC	1	3	9	31.25	16	34	0.31	0.48	*
41	E	L	mm3	MC	1	3	9	26.32	19	31	0.26	0.45	*
42	E	L	IQ1	MC	1	3	9	33.33	18	32	0.33	0.49	*
43	E	L	IQ2	MC	1	3	9	38.46	13	37	0.38	0.51	*
44	E	L	IQ3	MC	1	3	9	6.67	15	35	0.07	0.26	*
45	F	M	mm1	MC	1	1	5	52.63	19	31	0.53	0.51	*
46	F	M	Bulb1	MC	1	1	5	25.00	8	42	0.25	0.46	*
47	F	M	IQ1	MC	1	1	5	25.00	8	42	0.25	0.46	*
48	F	M	Whale1	MC	1	1	5	28.57	7	43	0.29	0.49	*
49	F	M	Cow1	MC	1	1	5	25.00	8	42	0.25	0.46	*
50	F	N	mm2	MC	1	1	5	37.50	8	42	0.38	0.52	*
51	F	N	Bulb2	MC	1	1	5	0.00	11	39	0.00	0.00	*
52	F	N	IQ2	MC	1	1	5	27.27	11	39	0.27	0.47	*
53	F	N	Whale2	MC	1	1	5	10.00	10	40	0.10	0.32	*
54	F	N	Cow2	MC	1	1	5	20.00	10	40	0.20	0.42	*
55	F	O	mm3	MC	1	1	5	14.29	7	43	0.14	0.38	
56	F	O	Bulb3	MC	1	1	5	0.00	8	42	0.00	0.00	*
57	F	O	IQ3	MC	1	1	5	0.00	8	42	0.00	0.00	*
58	F	O	Whale3	MC	1	1	5	28.57	14	36	0.29	0.47	*
59	F	O	Cow3	MC	1	1	5	15.38	13	37	0.15	0.38	*
60	F	P	mm4	MC	1	1	5	7.14	14	36	0.07	0.27	
61	F	P	Bulb4	MC	1	1	5	0.00	7	43	0.00	0.00	
62	F	P	IQ4	MC	1	1	5	14.29	7	43	0.14	0.38	
63	F	P	Whale4	MC	1	1	5	0.00	13	37	0.00	0.00	
64	F	P	Cow4	MC	1	1	5	11.11	9	41	0.11	0.33	
65	F	Q	mm5	MC	1	1	5	50.00	4	46	0.50	0.58	*
66	F	Q	Bulb5	MC	1	1	5	25.00	12	38	0.25	0.45	*
67	F	Q	IQ5	MC	1	1	5	0.00	7	43	0.00	0.00	*
68	F	Q	Whale5	MC	1	1	5	28.57	7	43	0.29	0.49	*

69	F	Q	Cow5	MC	1	1	5	40.00	20	30	0.40	0.50	*
70	G	R	mm1	MC	1	1	5	20.00	10	40	0.20	0.42	*
71	G	R	Bulb1	MC	1	1	5	7.14	14	36	0.07	0.27	*
72	G	R	IQ1	MC	1	1	5	0.00	4	46	0.00	0.00	*
73	G	R	Whale1	MC	1	1	5	23.08	13	37	0.23	0.44	*
74	G	R	Cow1	MC	1	1	5	33.33	9	41	0.33	0.50	*
75	G	S	mm2	MC	1	1	5	50.00	14	36	0.50	0.52	*
76	G	S	Bulb2	MC	1	1	5	27.27	11	39	0.27	0.47	*
77	G	S	IQ2	MC	1	1	5	0.00	9	41	0.00	0.00	*
78	G	S	Whale2	MC	1	1	5	11.11	9	41	0.11	0.33	*
79	G	S	Cow2	MC	1	1	5	28.57	7	43	0.29	0.49	*
80	G	T	mm3	MC	1	1	5	27.27	11	39	0.27	0.47	*
81	G	T	Bulb3	MC	1	1	5	25.00	8	42	0.25	0.46	*
82	G	T	IQ3	MC	1	1	5	14.29	14	36	0.14	0.36	*
83	G	T	Whale3	MC	1	1	5	0.00	5	45	0.00	0.00	*
84	G	T	Cow3	MC	1	1	5	8.33	12	38	0.08	0.29	*
85	G	U	mm4	MC	1	1	5	0.00	5	45	0.00	0.00	*
86	G	U	Bulb4	MC	1	1	5	33.33	12	38	0.33	0.49	*
87	G	U	IQ4	MC	1	1	5	33.33	21	29	0.33	0.48	*
88	G	U	Whale4	MC	1	1	5	16.67	6	44	0.17	0.41	*
89	G	U	Cow4	MC	1	1	5	16.67	6	44	0.17	0.41	*

Table 5: Summary statistics of each question

Students

The percentages correct by learning outcome and total for each student, sorted by the highest to lowest total are displayed in Table 6. And the ranks in percentage are given in Table 7. Figure 7 shows the percent rank of the students.

	Total	E	F	G	C	A	B	D
w.introstat40	48.00	33.33	60.00	50.00	0.00	50.00	50.00	100.00
w.introStat50	48.00	33.33	60.00	50.00	0.00	50.00	50.00	100.00
w.introstat26	36.00	16.67	60.00	25.00	50.00	0.00	50.00	100.00
w.introstat39	36.00	16.67	20.00	50.00	50.00	0.00	100.00	100.00
w.introstat42	36.00	33.33	0.00	25.00	50.00	25.00	100.00	100.00
w.introStat52	36.00	33.33	0.00	25.00	50.00	25.00	100.00	100.00
w.introstat65	36.00	16.67	20.00	50.00	50.00	25.00	50.00	100.00
w.introstat33	32.00	16.67	40.00	50.00	50.00	25.00	50.00	0.00
w.introstat34	32.00	33.33	0.00	25.00	50.00	25.00	50.00	100.00
w.introstat37	32.00	16.67	40.00	0.00	0.00	50.00	50.00	100.00
w.introstat45	32.00	33.33	40.00	50.00	0.00	50.00	0.00	0.00
w.introstat49	32.00	0.00	40.00	25.00	50.00	0.00	100.00	100.00
w.introStat55	32.00	33.33	40.00	50.00	0.00	50.00	0.00	0.00
w.introStat59	32.00	0.00	40.00	25.00	50.00	0.00	100.00	100.00
w.introstat69	32.00	16.67	20.00	25.00	50.00	25.00	50.00	100.00
w.introstat31	28.00	16.67	40.00	25.00	50.00	25.00	0.00	50.00
w.introstat35	28.00	33.33	40.00	0.00	0.00	25.00	0.00	100.00
w.introstat44	28.00	33.33	0.00	25.00	0.00	0.00	100.00	100.00
w.introstat48	28.00	0.00	20.00	25.00	50.00	50.00	100.00	0.00
w.introStat54	28.00	33.33	0.00	25.00	0.00	0.00	100.00	100.00
w.introStat58	28.00	0.00	20.00	25.00	50.00	50.00	100.00	0.00
w.introstat61	28.00	16.67	20.00	50.00	0.00	0.00	50.00	100.00
w.introstat62	28.00	16.67	40.00	50.00	0.00	0.00	50.00	50.00
w.introstat63	28.00	16.67	20.00	50.00	0.00	25.00	0.00	100.00
w.introstat32	24.00	0.00	40.00	0.00	50.00	0.00	50.00	100.00
w.introstat41	24.00	16.67	0.00	0.00	50.00	50.00	0.00	100.00
w.introstat43	24.00	0.00	0.00	25.00	50.00	50.00	0.00	100.00
w.introStat51	24.00	16.67	0.00	0.00	50.00	50.00	0.00	100.00
w.introStat53	24.00	0.00	0.00	25.00	50.00	50.00	0.00	100.00
w.introstat66	24.00	16.67	60.00	0.00	0.00	0.00	50.00	50.00
w.introstat67	24.00	0.00	0.00	25.00	0.00	50.00	50.00	100.00
w.introstat70	24.00	16.67	20.00	25.00	100.00	0.00	50.00	0.00
w.introstat21	20.00	16.67	20.00	25.00	0.00	25.00	0.00	50.00
w.introstat28	20.00	33.33	20.00	25.00	0.00	0.00	50.00	0.00
w.introstat38	20.00	0.00	20.00	25.00	0.00	0.00	50.00	100.00

w.introstat47	20.00	16.67	20.00	0.00	0.00	50.00	0.00	50.00
w.introStat57	20.00	16.67	20.00	0.00	0.00	50.00	0.00	50.00
w.introstat60	20.00	33.33	0.00	25.00	0.00	25.00	50.00	0.00
w.introstat22	16.00	16.67	20.00	0.00	0.00	0.00	50.00	50.00
w.introstat23	16.00	16.67	60.00	0.00	0.00	0.00	0.00	0.00
w.introstat29	16.00	33.33	0.00	0.00	0.00	0.00	0.00	100.00
w.introstat36	16.00	0.00	20.00	0.00	0.00	25.00	0.00	100.00
w.introstat46	16.00	16.67	20.00	25.00	0.00	25.00	0.00	0.00
w.introStat56	16.00	16.67	20.00	25.00	0.00	25.00	0.00	0.00
w.introstat68	16.00	0.00	0.00	0.00	0.00	50.00	0.00	100.00
w.introstat24	12.00	16.67	0.00	0.00	50.00	25.00	0.00	0.00
w.introstat27	12.00	0.00	0.00	25.00	0.00	25.00	50.00	0.00
w.introstat30	12.00	16.67	0.00	0.00	50.00	0.00	50.00	0.00
w.introstat64	4.00	16.67	0.00	0.00	0.00	0.00	0.00	0.00
w.introstat25	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 6: Sorted learning outcomes sets and total correct percentages

	Total	E	F	G	C	A	B	D
w.introstat40	0.00	0.00	0.00	0.00	42.00	0.00	18.00	0.00
w.introStat50	0.00	0.00	0.00	0.00	42.00	0.00	18.00	0.00
w.introstat26	4.00	26.00	0.00	20.00	2.00	62.00	18.00	0.00
w.introstat39	4.00	26.00	30.00	0.00	2.00	62.00	0.00	0.00
w.introstat42	4.00	0.00	64.00	20.00	2.00	30.00	0.00	0.00
w.introStat52	4.00	0.00	64.00	20.00	2.00	30.00	0.00	0.00
w.introstat65	4.00	26.00	30.00	0.00	2.00	30.00	18.00	0.00
w.introstat33	14.00	26.00	10.00	0.00	2.00	30.00	18.00	68.00
w.introstat34	14.00	0.00	64.00	20.00	2.00	30.00	18.00	0.00
w.introstat37	14.00	26.00	10.00	66.00	42.00	0.00	18.00	0.00
w.introstat45	14.00	0.00	10.00	0.00	42.00	0.00	58.00	68.00
w.introstat49	14.00	74.00	10.00	20.00	2.00	62.00	0.00	0.00
w.introStat55	14.00	0.00	10.00	0.00	42.00	0.00	58.00	68.00
w.introStat59	14.00	74.00	10.00	20.00	2.00	62.00	0.00	0.00
w.introstat69	14.00	26.00	30.00	20.00	2.00	30.00	18.00	0.00
w.introstat31	30.00	26.00	10.00	20.00	2.00	30.00	58.00	54.00
w.introstat35	30.00	0.00	10.00	66.00	42.00	30.00	58.00	0.00
w.introstat44	30.00	0.00	64.00	20.00	42.00	62.00	0.00	0.00
w.introstat48	30.00	74.00	30.00	20.00	2.00	0.00	0.00	68.00
w.introStat54	30.00	0.00	64.00	20.00	42.00	62.00	0.00	0.00
w.introStat58	30.00	74.00	30.00	20.00	2.00	0.00	0.00	68.00
w.introstat61	30.00	26.00	30.00	0.00	42.00	62.00	18.00	0.00
w.introstat62	30.00	26.00	10.00	0.00	42.00	62.00	18.00	54.00

w.introstat63	30.00	26.00	30.00	0.00	42.00	30.00	58.00	0.00
w.introstat32	48.00	74.00	10.00	66.00	2.00	62.00	18.00	0.00
w.introstat41	48.00	26.00	64.00	66.00	2.00	0.00	58.00	0.00
w.introstat43	48.00	74.00	64.00	20.00	2.00	0.00	58.00	0.00
w.introStat51	48.00	26.00	64.00	66.00	2.00	0.00	58.00	0.00
w.introStat53	48.00	74.00	64.00	20.00	2.00	0.00	58.00	0.00
w.introstat66	48.00	26.00	0.00	66.00	42.00	62.00	18.00	54.00
w.introstat67	48.00	74.00	64.00	20.00	42.00	0.00	18.00	0.00
w.introstat70	48.00	26.00	30.00	20.00	0.00	62.00	18.00	68.00
w.introstat21	64.00	26.00	30.00	20.00	42.00	30.00	58.00	54.00
w.introstat28	64.00	0.00	30.00	20.00	42.00	62.00	18.00	68.00
w.introstat38	64.00	74.00	30.00	20.00	42.00	62.00	18.00	0.00
w.introstat47	64.00	26.00	30.00	66.00	42.00	0.00	58.00	54.00
w.introStat57	64.00	26.00	30.00	66.00	42.00	0.00	58.00	54.00
w.introstat60	64.00	0.00	64.00	20.00	42.00	30.00	18.00	68.00
w.introstat22	76.00	26.00	30.00	66.00	42.00	62.00	18.00	54.00
w.introstat23	76.00	26.00	0.00	66.00	42.00	62.00	58.00	68.00
w.introstat29	76.00	0.00	64.00	66.00	42.00	62.00	58.00	0.00
w.introstat36	76.00	74.00	30.00	66.00	42.00	30.00	58.00	0.00
w.introstat46	76.00	26.00	30.00	20.00	42.00	30.00	58.00	68.00
w.introStat56	76.00	26.00	30.00	20.00	42.00	30.00	58.00	68.00
w.introstat68	76.00	74.00	64.00	66.00	42.00	0.00	58.00	0.00
w.introstat24	90.00	26.00	64.00	66.00	2.00	30.00	58.00	68.00
w.introstat27	90.00	74.00	64.00	20.00	42.00	30.00	18.00	68.00
w.introstat30	90.00	26.00	64.00	66.00	2.00	62.00	18.00	68.00
w.introstat64	96.00	26.00	64.00	66.00	42.00	62.00	58.00	68.00
w.introstat25	98.00	74.00	64.00	66.00	42.00	62.00	58.00	68.00

Table 7: Rank of the students by the total and learning outcome scores. The percentages are the proportion of students in this section who got a higher score in the corresponding column.





Figure 7: Heat map of the student ranks. Blue represents the top rank, while yellow means the bottom.


Summary of Questions

(1) Question "To6.A.A.04.1.1.MC.1" is given on the right. This question was selected from the question set with a frequency of 0.25. The question was administered to 10 out of the total of 50 students. The average score was 0.4 out of 1.

(Back to the question summary Table 5.)



Answer	Count
a	4
b	3
c	2
Unanswered	1




Summary	Value
Mean	0.40
Std.dev	0.52
Min	0.00
Median	0.00
Max	1.00

The z-score for a particular observation is $z = 1.5$. This means the observation is


- *a. 1.5 standard deviations above the mean.
- b. 1.5 standard deviations below the mean.
- c. 1.5 units above the mean.
- d. 1.5 units below the mean.

(2) Question "To6.A.A.04.1.1.MC.2" is given on the right. This question was selected from the question set with a frequency of 0.25. The question was administered to 13 out of the total of 50 students. The average score was 0.46 out of 1.

(Back to the question summary Table 5.)



Answer	Count
a	6
b	5
c	1
d	1




Summary	Value
Mean	0.46
Std.dev	0.52
Min	0.00
Median	0.00
Max	1.00

The z-score for a particular observation is $z = 0.4$. This means the observation is

- *a. 0.4 standard deviations above the mean.
- b. 0.4 standard deviations below the mean.
- c. 0.4 units above the mean.
- d. 0.4 units below the mean.

(3) Question "To6.A.A.04.1.1.MC.3" is given on the right. This question was selected from the question set with a frequency of 0.25. The question was administered to 10 out of the total of 50 students. The average score was 0.3 out of 1.

(Back to the question summary Table 5.)



Answer	Count
a	3
c	3
d	3
Unanswered	1

Summary	Value
Mean	0.30
Std.dev	0.48
Min	0.00
Median	0.00
Max	1.00

The z-score for a particular observation is $z = 2.3$. This means the observation is

*a. 2.3 standard deviations above the mean.


b. 2.3 standard deviations below the mean.

c. 2.3 units above the mean.


d. 2.3 units below the mean.

(4) Question "To6.A.A.04.1.1.MC.4" is given on the right. This question was selected from the question set with a frequency of 0.25. The question was administered to 17 out of the total of 50 students. The average score was 0.53 out of 1.

(Back to the question summary Table 5.)



Answer	Count
a	9
d	4
b	3
c	1





Summary	Value
Mean	0.53
Std.dev	0.51
Min	0.00
Median	1.00
Max	1.00

The z-score for a particular observation is $z = 3.4$. This means the observation is

- *a. 3.4 standard deviations above the mean.
- b. 3.4 standard deviations below the mean.
- c. 3.4 units above the mean.
- d. 3.4 units below the mean.

(5) Question "To6.A.B.04.1.1.MC.1" is given on the right. This question was selected from the question set with a frequency of 0.25. The question was administered to 14 out of the total of 50 students. The average score was 0.43 out of 1.

(Back to the question summary Table 5.)



Answer	Count
b	6
c	4
d	3
a	1

Summary	Value
Mean	0.43
Std.dev	0.51
Min	0.00
Median	0.00
Max	1.00

The z-score for a particular observation is $z = -1.2$. This means the observation is

a. 1.2 standard deviations above the mean.


*b. 1.2 standard deviations below the mean.

c. 1.2 units above the mean.

d. 1.2 units below the mean.

(6) Question "To6.A.B.04.1.1.MC.2" is given on the right. This question was selected from the question set with a frequency of 0.25. The question was administered to 16 out of the total of 50 students. The average score was 0.5 out of 1.

(Back to the question summary Table 5.)



Answer	Count
b	8
c	5
a	1
d	1
Unanswered	1

Summary	Value
Mean	0.50
Std.dev	0.52
Min	0.00
Median	0.50
Max	1.00

The z-score for a particular observation is $z = -0.8$. This means the observation is

a. 0.8 standard deviations above the mean.


*b. 0.8 standard deviations below the mean.

c. 0.8 units above the mean.

d. 0.8 units below the mean.

(7) Question "To6.A.B.04.1.1.MC.3" is given on the right. This question was selected from the question set with a frequency of 0.25. The question was administered to 8 out of the total of 50 students. The average score was 0.25 out of 1.

(Back to the question summary Table 5.)



Answer	Count
a	3
b	2
c	1
d	1
Unanswered	1

Summary	Value
Mean	0.25
Std.dev	0.46
Min	0.00
Median	0.00
Max	1.00

The z-score for a particular observation is $z = -2.7$. This means the observation is

a. 2.7 standard deviations above the mean.



*b. 2.7 standard deviations below the mean.

c. 2.7 units above the mean.

d. 2.7 units below the mean.

(8) Question "To6.A.B.04.1.1.MC.4" is given on the right. This question was selected from the question set with a frequency of 0.25. The question was administered to 12 out of the total of 50 students. The average score was 0.67 out of 1.

(Back to the question summary Table 5.)




Summary		Value
Mean		0.67
Std.dev		0.49
Min		0.00
Median		1.00
Max		1.00

Answer	Count
b	8
d	3
a	1

- The z-score for a particular observation is $z = -3.1$. This means the observation is
- a. 3.1 standard deviations above the mean.
 - *b. 3.1 standard deviations below the mean.
 - c. 3.1 units above the mean.
 - d. 3.1 units below the mean.

(9) Question "To6.A.C.04.1.1.FB.1" is given on the right. This question was selected from the question set with a frequency of 0.25. The question was administered to 11 out of the total of 50 students. The average score was 0 out of 1.

(Back to the question summary Table 5.)



Answer	Count
-3.21	2
1.45	2
2.35	2
1.23	1
3	1
3.1	1
34	1
436	1

Summary	Value
Mean	0.00
Std.dev	0.00
Min	0.00
Median	0.00
Max	0.00

In a sample of 25 male newborns, the mean birth weight was 3.4 kg and the standard deviation was 0.35 kg.


The z-score for a birth weight of 4.5 kg is _____. Round your answer to 2 decimal places.

Correct Answer(s):

- a. 3.14
- b. 3.15
- c. 3.13

(10) Question "To6.A.C.04.1.1.FB.2" is given on the right. This question was selected from the question set with a frequency of 0.25. The question was administered to 11 out of the total of 50 students. The average score was 0 out of 1.

(Back to the question summary Table 5.)



Answer	Count
3.4	2
Unanswered	2
1.35	1
34	1
4.0	1
4.32	1
443	1
45	1
67	1

Summary	Value
Mean	0.00
Std.dev	0.00
Min	0.00
Median	0.00
Max	0.00

In a sample of 25 male newborns, the mean birth weight was 3.4 kg and the standard deviation was 0.35 kg.


The z-score for a birth weight of 4.0 kg is _____. Round your answer to 2 decimal places.

Correct Answer(s):

- a. 1.71
- b. 1.72
- c. 1.70
- d. 1.7

(11) Question "To6.A.C.04.1.1.FB.3" is given on the right. This question was selected from the question set with a frequency of 0.25. The question was administered to 14 out of the total of 50 students. The average score was 0 out of 1.

(Back to the question summary Table 5.)



Answer	Count
34	3
-1.5	2
0.97	2
Unanswered	2
0	1
12	1
24.53	1
3	1
4.5963	1

Summary	Value
Mean	0.00
Std.dev	0.00
Min	0.00
Median	0.00
Max	0.00

In a sample of 25 male newborns, the mean birth weight was 3.4 kg and the standard deviation was 0.35 kg.


The z-score for a birth weight of 4.2 kg is _____. Round your answer to 2 decimal places.

Correct Answer(s):

- a. 2.28
- b. 2.29
- c. 2.27

(12) Question "To6.A.C.04.1.1.FB.4" is given on the right. This question was selected from the question set with a frequency of 0.25. The question was administered to 14 out of the total of 50 students. The average score was 0 out of 1.

(Back to the question summary Table 5.)



Answer	Count
23	3
2.15	2
3.1	2
4.3	2
Unanswered	2
0.00	1
1	1
1.43	1

Summary	Value
Mean	0.00
Std.dev	0.00
Min	0.00
Median	0.00
Max	0.00

In a sample of 25 male newborns, the mean birth weight was 3.4 kg and the standard deviation was 0.35 kg.



The z-score for a birth weight of 4.3 kg is _____. Round your answer to 2 decimal places.

Correct Answer(s):

- a. 2.57
- b. 2.56
- c. 2.58

(13) Question "To6.A.D.04.1.1.FB.1" is given on the right. This question was selected from the question set with a frequency of 0.25. The question was administered to 10 out of the total of 50 students. The average score was 0 out of 1.

(Back to the question summary Table 5.)



Answer	Count
-0.98	2
3.52	2
-1.23	1
-1.2342	1
1.35	1
3	1
36	1
4	1

Summary	Value
Mean	0.00
Std.dev	0.00
Min	0.00
Median	0.00
Max	0.00

In a sample of 25 male newborns, the mean birth weight was 3.4 kg and the standard deviation was 0.35 kg.


The z-score for a birth weight of 2.3 kg is _____. Round your answer to 2 decimal places.

Correct Answer(s):

- a. -3.14
- b. -3.13
- c. -3.15

(14) Question "To6.A.D.04.1.1.FB.2" is given on the right. This question was selected from the question set with a frequency of 0.25. The question was administered to 13 out of the total of 50 students. The average score was 0 out of 1.

(Back to the question summary Table 5.)



Answer	Count
0.00	2
2.32	2
34	2
Unanswered	2
-100.23	1
3	1
36	1
45	1
56234	1

Summary	Value
Mean	0.00
Std.dev	0.00
Min	0.00
Median	0.00
Max	0.00

In a sample of 25 male newborns, the mean birth weight was 3.4 kg and the standard deviation was 0.35 kg.


The z-score for a birth weight of 2.5 kg is _____. Round your answer to 2 decimal places.

Correct Answer(s):

- a. -2.57
- b. -2.58
- c. -2.56

(15) Question "To6.A.D.04.1.1.FB.3" is given on the right. This question was selected from the question set with a frequency of 0.25. The question was administered to 11 out of the total of 50 students. The average score was 0 out of 1.

(Back to the question summary Table 5.)



Answer	Count
-1.2	2
-1.5	2
-1.08	1
1.56	1
2.2	1
2.31	1
25	1
45	1
Unanswered	1

Summary	Value
Mean	0.00
Std.dev	0.00
Min	0.00
Median	0.00
Max	0.00

In a sample of 25 male newborns, the mean birth weight was 3.4 kg and the standard deviation was 0.35 kg.


The z-score for a birth weight of 2.2 kg is _____. Round your answer to 2 decimal places.

Correct Answer(s):

- a. -3.43
- b. -3.44
- c. -3.42

(16) Question "To6.A.D.04.1.1.FB.4" is given on the right. This question was selected from the question set with a frequency of 0.25. The question was administered to 16 out of the total of 50 students. The average score was 0 out of 1.

(Back to the question summary Table 5.)



Answer	Count
Unanswered	5
-0.35	2
-3.43	2
34	2
-0.23	1
1.23	1
13	1
2	1
25	1

Summary	Value
Mean	0.00
Std.dev	0.00
Min	0.00
Median	0.00
Max	0.00

In a sample of 25 male newborns, the mean birth weight was 3.4 kg and the standard deviation was 0.35 kg.



The z-score for a birth weight of 2.8 kg is _____. Round your answer to 2 decimal places.

Correct Answer(s):

- a. -1.71
- b. -1.72
- c. -1.70
- d. -1.7

(17) Question "To6.B.E.01.1.1.MC.compare1" is given on the right. This question was selected from the question set with a frequency of 1. The question was administered to 50 out of the total of 50 students. The average score was 0.4 out of 1.

(Back to the question summary Table 5.)



Answer	Count
a	20
c	17
b	11
Unanswered	2



Summary	Value
Mean	0.40
Std.dev	0.49
Min	0.00
Median	0.00
Max	1.00

The height of adult women is thought to have a mean of 65 inches and a standard deviation of 2.5 inches. The height of adult men is thought to have a mean of 71 inches and a standard deviation of 3 inches. In the same family, the son is 73 inches tall and the daughter is 67 inches tall. Who is taller among their gender, the son or daughter?

- *a. The daughter
- b. The son
- c. The daughter and son are the same height within their gender.

(18) Question "To6.B.F.01.1.1.MC.compare2" is given on the right. This question was selected from the question set with a frequency of 1. The question was administered to 50 out of the total of 50 students. The average score was 0.36 out of 1.

(Back to the question summary Table 5.)



Answer	Count
a	20
b	18
c	10
Unanswered	2


Summary	Value
Mean	0.36
Std.dev	0.48
Min	0.00
Median	0.00
Max	1.00

The height of adult women is thought to have a mean of 65 inches and a standard deviation of 2.5 inches. The height of adult men is thought to have a mean of 71 inches and a standard deviation of 3 inches. In the same family, the son is 67 inches tall and the daughter is 59 inches tall. Who is taller among their gender, the son or daughter?

- a. The daughter
- *b. The son
- c. The daughter and son are the same height within their gender.

(19) Question "To6.C.G.01.1.1.MC.1" is given on the right. This question was selected from the question set with a frequency of 1. The question was administered to 50 out of the total of 50 students. The average score was 0.2 out of 1.

(Back to the question summary Table 5.)



Answer	Count
e	14
b	10
a	9
d	8
c	7
Unanswered	2

Summary	Value
Mean	0.20
Std.dev	0.40
Min	0.00
Median	0.00
Max	1.00

Standardizing makes the following change(s) to a distribution: I. Shifts the distribution by subtracting the mean. II. Rescales the distribution by dividing by the standard deviation. III. Changes the skewness or symmetry of the distribution. IV. Creates outliers.

a. I, II, and III

*b. I and II

c. III and IV

d. II and III


e. I, II and IV

(20) Question "To6.C.H.01.1.1.MC.1" is given on the right. This question was selected from the question set with a frequency of 1. The question was administered to 50 out of the total of 50 students. The average score was 0.24 out of 1.

(Back to the question summary Table 5.)

Which part of the distribution is NOT affected by standardizing?

- *a. shape
- b. median
- c. IQR
- d. range




Answer	Count
d	16
a	12
c	11
b	9
Unanswered	2


Summary	Value
Mean	0.24
Std.dev	0.43
Min	0.00
Median	0.00
Max	1.00

(21) Question "To6.D.I.02.1.1.TF.feet" is given on the right. This question was selected from the question set with a frequency of 0.5. The question was administered to 20 out of the total of 50 students. The average score was 0.65 out of 1.

(Back to the question summary Table 5.)




Answer	Count
a	13
b	6
Unanswered	1



Summary	Value
Mean	0.65
Std.dev	0.49
Min	0.00
Median	1.00
Max	1.00


NBA players tend to have very large feet. The following plots display what these shoe sizes may hypothetically look like. Based on the output, it is reasonable to model the distribution of NBA players shoe sizes with a normal




- *a. True
- b. False

(22) Question "To6.D.I.02.1.1.TF.lowtemp" is given on the right. This question was selected from the question set with a frequency of 0.5. The question was administered to 30 out of the total of 50 students. The average score was 0.63 out of 1.

(Back to the question summary Table 5.)




Answer	Count
a	19
b	10
Unanswered	1



Summary	Value
Mean	0.63
Std.dev	0.49
Min	0.00
Median	1.00
Max	1.00


Below is the distribution of low temperatures (in degrees F) for 52 cities in the U.S. Based on the output, it is reasonable to model the distribution of low temperature for these cities with a normal distribution.



- *a. True
- b. False

(23) Question "To6.D.J.04.1.1.TF.gain" is given on the right. This question was selected from the question set with a frequency of 0.25. The question was administered to 10 out of the total of 50 students. The average score was 0.6 out of 1.

(Back to the question summary Table 5.)




Answer	Count
b	6
a	3
Unanswered	1

Summary	Value
Mean	0.60
Std.dev	0.52
Min	0.00
Median	1.00
Max	1.00

The distribution of the gain of 120 different amplifiers is depicted below.

Based on the output, it is reasonable to model the distribution of amplifier




a. True


*b. False

(24) Question "To6.D.J.04.1.1.TF.mpg" is given on the right. This question was selected from the question set with a frequency of 0.25. The question was administered to 6 out of the total of 50 students. The average score was 0.83 out of 1.

(Back to the question summary Table 5.)




Answer	Count
b	5
a	1



Summary	Value
Mean	0.83
Std.dev	0.41
Min	0.00
Median	1.00
Max	1.00


The distribution of estimated highway miles per gallon (mpg) for various makes and models of cars is given below. Based on the output, it is reasonable to model the distribution of highway mpg values with a normal



a. True
*b. False

(25) Question "To6.D.J.04.1.1.TF.blowhole" is given on the right. This question was selected from the question set with a frequency of 0.25. The question was administered to 14 out of the total of 50 students. The average score was 0.5 out of 1.

(Back to the question summary Table 5.)




Summary		Value
Mean		0.50
Std.dev		0.52
Min		0.00
Median		0.50
Max		1.00

Answer	Count
a	7
b	7

A blowhole is a hole in a cliff that produces eruptions of water when the ocean swell hits the cliff. Below are 40 times (in seconds) between eruptions for the Kiama blowhole in Australia.


Based on the output, it is reasonable to model the distribution of times between eruptions with a normal



a. True
*b. False


(26) Question "To6.D.J.04.1.1.TF.CDs" is given on the right. This question was selected from the question set with a frequency of 0.25. The question was administered to 20 out of the total of 50 students. The average score was 0.55 out of 1.

(Back to the question summary Table 5.)



Summary		Value
Answer	Count	
b	11	
a	7	
Unanswered	2	
Mean	0.55	
Std.dev	0.51	
Min	0.00	
Median	1.00	
Max	1.00	


A random sample of 120 students was selected from those students who completed the Stat 101 survey over the last 2 years. The survey asked the number of music CDs owned by each of these students. The histogram of the number of music CDs owned by the students is shown below. Based on the output, it is reasonable to model the distribution of the number of CDs owned by Stat 101 students with a normal distribution.



- a. True
- *b. False

(27) Question "To6.E.K.09.3.1.FB.whale1" is given on the right. This question was selected from the question set with a frequency of 0.33. The question was administered to 19 out of the total of 50 students. The average score was 0.21 out of 1.

(Back to the question summary Table 5.)



Answer	Count
95	4
a	4
99.7	2
Unanswered	2
23.1	1
3	1
34	1
35	1
68.0	1
95.0	1
99.70	1

Summary	Value
Mean	0.21
Std.dev	0.42
Min	0.00
Median	0.00
Max	1.00


Fill in the blank with the correct number: Assume the length of female humpback whales can be modeled with a normal distribution with a mean of 13.7 meters and a standard deviation of 0.5 meters. According to the Empirical Rule or 68-95-99.7 Rule, _____ percent of female humpback whales will have a length between 13.2 meters and 14.2 meters.

Correct Answer(s):

- a. 68
- b. 68%

(28) Question "To6.E.K.09.3.1.FB.whale2" is given on the right. This question was selected from the question set with a frequency of 0.33. The question was administered to 18 out of the total of 50 students. The average score was 0 out of 1.

(Back to the question summary Table 5.)



Answer	Count
68	5
23	2
99.7	2
Unanswered	2
1	1
1.34	1
10%	1
12	1
3	1
34	1
45	1

Summary	Value
Mean	0.00
Std.dev	0.00
Min	0.00
Median	0.00
Max	0.00


Fill in the blank with the correct number: Assume the length of female humpback whales can be modeled with a normal distribution with a mean of 13.7 meters and a standard deviation of 0.5 meters. According to the Empirical Rule or 68-95-99.7 Rule, _____ percent of female humpback whales will have a length between 12.7 meters and 14.7 meters.

Correct Answer(s):

- a. 95
- b. 95%

(29) Question "To6.E.K.09.3.1.FB.whale3" is given on the right. This question was selected from the question set with a frequency of 0.33. The question was administered to 14 out of the total of 50 students. The average score was 0 out of 1.

(Back to the question summary Table 5.)



Answer	Count
23	2
95	2
95.0	2
Unanswered	2
23.5	1
24	1
25	1
67	1
68	1
92%	1

Summary	Value
Mean	0.00
Std.dev	0.00
Min	0.00
Median	0.00
Max	0.00


Fill in the blank with the correct number: Assume the length of female humpback whales can be modeled with a normal distribution with a mean of 13.7 meters and a standard deviation of 0.5 meters. According to the Empirical Rule or 68-95-99.7 Rule, _____ percent of female humpback whales will have a length between 12.2 meters and 15.2 meters.

Correct Answer(s):

- a. 99.7
- b. 99.7%

(30) Question "To6.E.K.09.3.1.FB.cow1" is given on the right. This question was selected from the question set with a frequency of 0.33. The question was administered to 12 out of the total of 50 students. The average score was 0 out of 1.

(Back to the question summary Table 5.)



Answer	Count
1	1
12	1
2.53	1
24	1
3	1
34	1
4	1
435	1
45	1
54	1
99	1
Unanswered	1

Summary	Value
Mean	0.00
Std.dev	0.00
Min	0.00
Median	0.00
Max	0.00


Fill in the blank with the correct number: Assume the weight of a certain breed of cow can be modeled with a normal distribution with a mean of 750 kg and a standard deviation of 30 kg. According to the Empirical Rule or 68-95-99.7 Rule, _____ percent of cows from this breed will weigh between 720 kg and 780 kg.

Correct Answer(s):

- a. 68
- b. 68%

(31) Question "To6.E.K.09.3.1.FB.cow2" is given on the right. This question was selected from the question set with a frequency of 0.33. The question was administered to 18 out of the total of 50 students. The average score was 0.06 out of 1.

(Back to the question summary Table 5.)



Answer	Count
68.0	3
99.7	3
68	2
Unanswered	2
2	1
23	1
234	1
3	1
45	1
56	1
67	1
a	1

Summary	Value
Mean	0.06
Std.dev	0.24
Min	0.00
Median	0.00
Max	1.00


Fill in the blank with the correct number: Assume the weight of a certain breed of cow can be modeled with a normal distribution with a mean of 750 kg and a standard deviation of 30 kg. According to the Empirical Rule or 68-95-99.7 Rule, _____ percent of cows from this breed will weigh between 690 kg and 810 kg.

Correct Answer(s):

- a. 95
- b. 95%

(32) Question "To6.E.K.09.3.1.FB.cow3" is given on the right. This question was selected from the question set with a frequency of 0.33. The question was administered to 17 out of the total of 50 students. The average score was 0.12 out of 1.

(Back to the question summary Table 5.)



Answer	Count
95	7
45	2
a	2
32	1
34	1
35	1
6	1
85%	1
95.0	1

Summary	Value
Mean	0.12
Std.dev	0.33
Min	0.00
Median	0.00
Max	1.00


Fill in the blank with the correct number: Assume the weight of a certain breed of cow can be modeled with a normal distribution with a mean of 750 kg and a standard deviation of 30 kg. According to the Empirical Rule or 68-95-99.7 Rule, _____ percent of cows from this breed will weigh between 660 kg and 840 kg.

Correct Answer(s):

- a. 99.7
- b. 99.7%

(33) Question "To6.E.K.09.3.1.FB.bulbs1" is given on the right. This question was selected from the question set with a frequency of 0.33. The question was administered to 16 out of the total of 50 students. The average score was 0.31 out of 1.

(Back to the question summary Table 5.)



Answer	Count
99.7	6
a	5
124	1
3	1
68.0	1
95.0	1
Unanswered	1

Summary	Value
Mean	0.31
Std.dev	0.48
Min	0.00
Median	0.00
Max	1.00



Fill in the blank with the correct number: Assume the lifespan of light bulbs manufactured by Bright Inc. can be modeled with a normal distribution with a mean of 300 days and a standard deviation of 40 days. According to the Empirical Rule or 68-95-99.7 Rule, _____ percent of light bulbs made by Bright Inc. will last between 260 and 340 days.

Correct Answer(s):

- a. 68
- b. 68%

(34) Question "To6.E.K.09.3.1.FB.bulbs2" is given on the right. This question was selected from the question set with a frequency of 0.33. The question was administered to 17 out of the total of 50 students. The average score was 0.06 out of 1.

(Back to the question summary Table 5.)



Answer	Count
45	3
99.7	3
68	2
1	1
13	1
23	1
32	1
34	1
35	1
67	1
95.0	1
a	1

Summary	Value
Mean	0.06
Std.dev	0.24
Min	0.00
Median	0.00
Max	1.00


Fill in the blank with the correct number: Assume the lifespan of light bulbs manufactured by Bright Inc. can be modeled with a normal distribution with a mean of 300 days and a standard deviation of 40 days. According to the Empirical Rule or 68-95-99.7 Rule, _____ percent of light bulbs made by Bright Inc. will last between 220 and 380 days.

Correct Answer(s):

- a. 95
- b. 95%

(35) Question "To6.E.K.09.3.1.FB.bulbs3" is given on the right. This question was selected from the question set with a frequency of 0.33. The question was administered to 19 out of the total of 50 students. The average score was 0.11 out of 1.

(Back to the question summary Table 5.)



Answer	Count
95	5
68	4
a	2
Unanswered	2
14	1
23	1
25	1
3	1
5	1
95.0	1

Summary	Value
Mean	0.11
Std.dev	0.32
Min	0.00
Median	0.00
Max	1.00



Fill in the blank with the correct number: Assume the lifespan of light bulbs manufactured by Bright Inc. can be modeled with a normal distribution with a mean of 300 days and a standard deviation of 40 days. According to the Empirical Rule or 68-95-99.7 Rule, _____ percent of light bulbs made by Bright Inc. will last between 180 and 420 days.

Correct Answer(s):

- a. 99.7
- b. 99.7%

(36) Question "To6.E.L.09.3.1.MC.heights1" is given on the right. This question was selected from the question set with a frequency of 0.33. The question was administered to 19 out of the total of 50 students. The average score was 0.16 out of 1.

(Back to the question summary Table 5.)



Answer	Count
d	7
a	6
b	3
c	3


Summary	Value
Mean	0.16
Std.dev	0.37
Min	0.00
Median	0.00
Max	1.00

Assume the distribution of the height of adult females can be modeled with a normal distribution with mean 66 inches and standard deviation 3 inches. According to the Empirical Rule or 68-95-99.7 Rule, the center 99.7% of all women will have heights between _____ and 75 inches.

- a. 60
- *b. 57
- c. 63
- d. 66

(37) Question "To6.E.L.09.3.1.MC.heights2" is given on the right. This question was selected from the question set with a frequency of 0.33. The question was administered to 18 out of the total of 50 students. The average score was 0.06 out of 1.

(Back to the question summary Table 5.)



Answer	Count
a	6
d	6
b	4
c	1
Unanswered	1

Summary	Value
Mean	0.06
Std.dev	0.24
Min	0.00
Median	0.00
Max	1.00

Assume the distribution of the height of adult females can be modeled with a normal distribution with mean 66 inches and standard deviation 3 inches. According to the Empirical Rule or 68-95-99.7 Rule, the center 68% of all women will have heights between _____ and 69 inches.



- a. 60
- b. 57
- *c. 63
- d. 66

(38) Question "To6.E.L.09.3.1.MC.heights3" is given on the right. This question was selected from the question set with a frequency of 0.33. The question was administered to 19 out of the total of 50 students. The average score was 0.42 out of 1.

(Back to the question summary Table 5.)

Assume the distribution of the height of adult females can be modeled with a normal distribution with mean 66 inches and standard deviation 3 inches. According to the Empirical Rule or 68-95-99.7 Rule, the center 95% of all women will have heights between _____ and 72 inches.

- *a. 60
- b. 57
- c. 63
- d. 66




Summary		Value
Mean		0.42
Std.dev		0.51
Min		0.00
Median		0.00
Max		1.00

Answer	Count
d	10
a	8
Unanswered	1

(39) Question "To6.E.L.09.3.1.MC.mm1" is given on the right. This question was selected from the question set with a frequency of 0.33. The question was administered to 13 out of the total of 50 students. The average score was 0.08 out of 1.

(Back to the question summary Table 5.)



Answer	Count
b	4
d	4
c	3
a	1
Unanswered	1


Summary	Value
Mean	0.08
Std.dev	0.28
Min	0.00
Median	0.00
Max	1.00

Assume the weight of bags of M&Ms can be modeled with the normal distribution with mean 50 grams and standard deviation 1 gram. According to the Empirical Rule or 68-95-99.7 rule, 99.7% of all M&M bags will have weights between _____ and 53 grams.


- *a. 47
- b. 48
- c. 49
- d. 50

(40) Question "To6.E.L.09.3.1.MC.mm2" is given on the right. This question was selected from the question set with a frequency of 0.33. The question was administered to 16 out of the total of 50 students. The average score was 0.31 out of 1.

(Back to the question summary Table 5.)



Answer	Count
d	6
c	5
b	4
a	1



Summary	Value
Mean	0.31
Std.dev	0.48
Min	0.00
Median	0.00
Max	1.00

Assume the weight of bags of M&Ms can be modeled with the normal distribution with mean 50 grams and standard deviation 1 gram. According to the Empirical Rule or 68-95-99.7 rule, 68% of all M&M bags will have weights between _____ and 51 grams.



- a. 47
- b. 48
- *c. 49
- d. 50

(41) Question "To6.E.L.09.3.1.MC.mm3" is given on the right. This question was selected from the question set with a frequency of 0.33. The question was administered to 19 out of the total of 50 students. The average score was 0.26 out of 1.

(Back to the question summary Table 5.)

Assume the weight of bags of M&Ms can be modeled with the normal distribution with mean 50 grams and standard deviation 1 gram. According to the Empirical Rule or 68-95-99.7 rule, 95% of all M&M bags will have weights between _____ and 52 grams.

- a. 47
- *b. 48
- c. 49
- d. 50




Answer	Count
d	7
a	5
b	5
c	2

Summary	Value
Mean	0.26
Std.dev	0.45
Min	0.00
Median	0.00
Max	1.00

(42) Question "To6.E.L.09.3.1.MC.IQ1" is given on the right. This question was selected from the question set with a frequency of 0.33. The question was administered to 18 out of the total of 50 students. The average score was 0.33 out of 1.

(Back to the question summary Table 5.)



Answer	Count
c	6
d	5
a	4
b	2
Unanswered	1


Summary	Value
Mean	0.33
Std.dev	0.49
Min	0.00
Median	0.00
Max	1.00

Assume the distribution of IQ scores for adults can be modeled with a normal distribution with a mean score of 100 points and a standard deviation of 10 points. According to the Empirical Rule or 68-95-99.7 Rule, the middle 95% of all adults will have an IQ score between 80 and _____ points.

- a. 110
- b. 100
- *c. 120
- d. 130

(43) Question "To6.E.L.09.3.1.MC.IQ2" is given on the right. This question was selected from the question set with a frequency of 0.33. The question was administered to 13 out of the total of 50 students. The average score was 0.38 out of 1.

(Back to the question summary Table 5.)



Answer	Count
a	5
c	4
b	2
d	1
Unanswered	1



Summary	Value
Mean	0.38
Std.dev	0.51
Min	0.00
Median	0.00
Max	1.00

Assume the distribution of IQ scores for adults can be modeled with a normal distribution with a mean score of 100 points and a standard deviation of 10 points. According to the Empirical Rule or 68-95-99.7 Rule, the middle 68% of all adults will have an IQ score between 90 and _____ points.

- *a. 110
- b. 100
- c. 120
- d. 130

(44) Question "To6.E.L.09.3.1.MC.IQ3" is given on the right. This question was selected from the question set with a frequency of 0.33. The question was administered to 15 out of the total of 50 students. The average score was 0.07 out of 1.

(Back to the question summary Table 5.)



Answer	Count
a	5
c	5
b	4
d	1



Summary	Value
Mean	0.07
Std.dev	0.26
Min	0.00
Median	0.00
Max	1.00

Assume the distribution of IQ scores for adults can be modeled with a normal distribution with a mean score of 100 points and a standard deviation of 10 points. According to the Empirical Rule or 68-95-99.7 Rule, the middle 99.7% of all adults will have an IQ score between 70 and _____ points.

- a. 110
- b. 100
- c. 120
- *d. 130

(45) Question "To6.FM.05.1.1.MC.mm1" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 19 out of the total of 50 students. The average score was 0.53 out of 1.

(Back to the question summary Table 5.)



Answer	Count
a	10
d	4
b	3
c	2



Summary	Value
Mean	0.53
Std.dev	0.51
Min	0.00
Median	1.00
Max	1.00

For the remaining questions, use either a z-table or an applet or both to do the calculations. Depending on the method used, the final answer could be subject to a small amount of rounding error. Assume the weight of bags of M&Ms can be modeled with the normal distribution with mean 50 grams and standard deviation 1 gram. The weight on the label of these M&M bags is 47.9 grams. What percentage of all M&M bags have a weight below the label weight?

- *a. 1.79%
- b. 98.21%
- c. 1.39%
- d. 2.28%
- e. 97.72%

(46) Question "To6.F.M.05.1.1.MC.Bulb1" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 8 out of the total of 50 students. The average score was 0.25 out of 1.

(Back to the question summary Table 5.)



Answer	Count
b	2
c	2
d	2
e	2


Summary	Value
Mean	0.25
Std.dev	0.46
Min	0.00
Median	0.00
Max	1.00

For the remaining questions, use either a z-table or an applet or both to do the calculations. Depending on the method used, the final answer could be subject to a small amount of rounding error. Assume the lifespan of light bulbs manufactured by Bright Inc. can be modeled with a normal distribution with a mean of 300 days and a standard deviation of 40 days. What percentage of light bulbs produced by Bright Inc. will survive less than 200 days?

- a. 99.38%
- *b. 0.62%
- c. 96.49%
- d. 12.3%
- e. 2.02%

(47) Question "To6.F.M.05.1.1.MC.IQ1" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 8 out of the total of 50 students. The average score was 0.25 out of 1.

(Back to the question summary Table 5.)





Answer	Count
e	3
a	2
b	2
Unanswered	1

Summary	Value
Mean	0.25
Std.dev	0.46
Min	0.00
Median	0.00
Max	1.00

For the remaining questions, use either a z-table or an applet or both to do the calculations. Depending on the method used, the final answer could be subject to a small amount of rounding error. Assume the distribution of IQ scores for adults can be modeled with a normal distribution with a mean score of 100 points and a standard deviation of 10 points. What percentage of adults have an IQ score of less than 87 points?

- *a. 9.68%
- b. 15.15%
- c. 4.46%
- d. 90.32%
- e. 84.85%

(48) Question "To6.FM.05.1.1.MC.Whale1" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 7 out of the total of 50 students. The average score was 0.29 out of 1.
(Back to the question summary Table 5.)




Summary		Value
Mean		0.29
Std.dev		0.49
Min		0.00
Median		0.00
Max		1.00

Answer	Count
b	4
a	2
d	1

For the remaining questions, use either a z-table or an applet or both to do the calculations. Depending on the method used, the final answer could be subject to a small amount of rounding error. Assume the length of female humpback whales can be modeled with a normal distribution with a mean of 13.7 meters and a standard deviation of 0.5 meters. What percentage of female humpback whales will be shorter than 13 meters in length?
*a. 8.08%
b. 91.92%
c. 14.92%
d. 85.08%
e. 0.82%

(49) Question "To6.F.M.05.1.1.MC.Cow1" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 8 out of the total of 50 students. The average score was 0.25 out of 1.

(Back to the question summary Table 5.)



Answer	Count
a	2
b	2
c	1
d	1
e	1
Unanswered	1



Summary	Value
Mean	0.25
Std.dev	0.46
Min	0.00
Median	0.00
Max	1.00

For the remaining questions, use either a z-table or an applet or both to do the calculations. Depending on the method used, the final answer could be subject to a small amount of rounding error. Assume the weight of a certain breed of cow can be modeled with a normal distribution with a mean of 750 kg and a standard deviation of 30 kg. What percentage of cows from this breed will weigh less than 680 kg?

- *a. 0.99%
- b. 99.01%
- c. 2.12%
- d. 97.88%
- e. 0.38%

(50) Question "To6.FN.05.1.1.MC.mm2" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 8 out of the total of 50 students. The average score was 0.38 out of 1.

(Back to the question summary Table 5.)



Answer	Count
a	3
e	3
d	2


Summary	Value
Mean	0.38
Std.dev	0.52
Min	0.00
Median	0.00
Max	1.00

Assume the weight of bags of M&Ms can be modeled with the normal distribution with mean 50 grams and standard deviation 1 gram. What percentage of all M&M bags will have a weight below 51.5 grams?

- *a. 93.32%
- b. 6.68%
- c. 85.31%
- d. 14.69%
- e. 90.32%

(51) Question "To6.F.N.05.1.1.MC.Bulb2" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 11 out of the total of 50 students. The average score was 0 out of 1.

(Back to the question summary Table 5.)



Answer	Count
c	5
d	4
b	1
Unanswered	1



Summary	Value
Mean	0.00
Std.dev	0.00
Min	0.00
Median	0.00
Max	0.00

Assume the lifespan of light bulbs manufactured by Bright Inc. can be modeled with a normal distribution with a mean of 300 days and a standard deviation of 40 days. What percentage of light bulbs produced by Bright Inc. will survive less than 405 days?

- *a. 99.57%
- b. 0.43%
- c. 99.09%
- d. 0.91%
- e. 94.84%

(52) Question "To6.FN.05.1.1.MC.IQ2" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 11 out of the total of 50 students. The average score was 0.27 out of 1.

(Back to the question summary Table 5.)



Answer	Count
a	3
e	3
b	2
c	2
d	1



Summary	Value
Mean	0.27
Std.dev	0.47
Min	0.00
Median	0.00
Max	1.00

Assume the distribution of IQ scores for adults can be modeled with a normal distribution with a mean score of 100 points and a standard deviation of 10 points. What percentage of adults have an IQ score of less than 107 points?

- *a. 75.8%
- b. 24.2%
- c. 72.3%
- d. 85.7%
- e. 14.3%

(53) Question "To6.F.N.05.1.1.MC.Whale2" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 10 out of the total of 50 students. The average score was 0.1 out of 1.

(Back to the question summary Table 5.)



Answer	Count
a	3
c	2
d	2
e	2
b	1


Summary	Value
Mean	0.10
Std.dev	0.32
Min	0.00
Median	0.00
Max	1.00

Assume the length of female humpback whales can be modeled with a normal distribution with a mean of 13.7 meters and a standard deviation of 0.5 meters. What percentage of female humpback whales will be shorter than 14 meters in length?

a. 77.04%
*b. 72.57%
c. 27.43%
d. 68.44%
e. 22.96%

(54) Question "To6.F.N.05.1.1.MC.Cow2" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 10 out of the total of 50 students. The average score was 0.2 out of 1.

(Back to the question summary Table 5.)



Answer	Count
b	5
d	2
a	1
e	1
Unanswered	1


Summary	Value
Mean	0.20
Std.dev	0.42
Min	0.00
Median	0.00
Max	1.00

Assume the weight of a certain breed of cow can be modeled with a normal distribution with a mean of 750 kg and a standard deviation of 30 kg. What percentage of cows from this breed will weigh less than 790 kg?


- a. 6.81%
- b. 86.21%
- c. 93.19%
- *d. 90.82%
- e. 9.18%

(55) Question "To6.F.O.05.1.1.MC.mm3" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 7 out of the total of 50 students. The average score was 0.14 out of 1.

(Back to the question summary Table 5.)



Answer	Count
c	4
a	1
b	1
e	1



Summary	Value
Mean	0.14
Std.dev	0.38
Min	0.00
Median	0.00
Max	1.00


- Assume the weight of bags of M&Ms can be modeled with the normal distribution with mean 50 grams and standard deviation 1 gram. What percent of M&M bags will have a weigh more than 48.5 grams?
- a. 6.68%
 - *b. 93.32%
 - c. 86.64%
 - d. 1.5%
 - e. 98.5%

(56) Question "To6.F.O.05.1.1.MC.Bulb3" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 8 out of the total of 50 students. The average score was 0 out of 1.

(Back to the question summary Table 5.)

Assume the lifespan of light bulbs manufactured by Bright Inc. can be modeled with a normal distribution with a mean of 300 days and a standard deviation of 40 days. What percentage of light bulbs produced by Bright Inc. will survive longer than 225 days?



- a. 3.01%
- *b. 96.99%
- c. 14.01%
- d. 85.99%
- e. 98.08%



Answer	Count	Summary	Value
a	4	Mean	0.00
d	2	Std.dev	0.00
e	1	Min	0.00
Unanswered	1	Median	0.00
		Max	0.00

(57) Question "To6.F.O.05.1.1.MC.IQ3" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 8 out of the total of 50 students. The average score was 0 out of 1.

(Back to the question summary Table 5.)



Answer	Count
d	8


Summary	Value
Mean	0.00
Std.dev	0.00
Min	0.00
Median	0.00
Max	0.00

Assume the distribution of IQ scores for adults can be modeled with a normal distribution with a mean score of 100 points and a standard deviation of 10 points. What percentage of adults have an IQ score higher than 88 points?

a. 11.51%
*b. 88.49%
c. 15.39%
d. 84.61%
e. 89.80%

(58) Question "To6.F.O.05.1.1.MC.Whale3" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 14 out of the total of 50 students. The average score was 0.29 out of 1.

(Back to the question summary Table 5.)





Answer	Count	Summary	Value
e	5	Mean	0.29
b	4	Std.dev	0.47
c	2	Min	0.00
d	2	Median	0.00
Unanswered	1	Max	1.00

Assume the length of female humpback whales can be modeled with a normal distribution with a mean of 13.7 meters and a standard deviation of 0.5 meters. What percentage of female humpback whales are longer than 12.1 meters?

- a. 0.07%
- *b. 99.93%
- c. 98.61%
- d. 1.39%
- e. 94.52%

(59) Question "To6.F.O.05.1.1.MC.Cow3" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 13 out of the total of 50 students. The average score was 0.15 out of 1.

(Back to the question summary Table 5.)



Answer	Count
d	6
c	5
a	2


Summary	Value
Mean	0.15
Std.dev	0.38
Min	0.00
Median	0.00
Max	1.00

Assume the weight of a certain breed of cow can be modeled with a normal distribution with a mean of 750 kg and a standard deviation of 30 kg. What percentage of cows from this breed will weigh more than 675 kg?

- *a. 99.38%
- b. 0.62%
- c. 97.98%
- d. 2.02%
- e. 93.32%

(60) Question "To6.F.P.05.1.1.MC.mm4" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 14 out of the total of 50 students. The average score was 0.07 out of 1.

(Back to the question summary Table 5.)



Answer	Count
a	4
d	4
e	3
b	1
c	1
Unanswered	1



Summary	Value
Mean	0.07
Std.dev	0.27
Min	0.00
Median	0.00
Max	1.00

Assume the weight of bags of M&Ms can be modeled with the normal distribution with mean 50 grams and standard deviation 1 gram. What percent of M&M bags will weigh more than 52.3 grams?

- a. 98.93%
- *b. 1.07%
- c. 2.30%
- d. 97.70%
- e. 9.68%

(61) Question "To6.F.P.05.1.1.MC.Bulb4" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 7 out of the total of 50 students. The average score was 0 out of 1.

(Back to the question summary Table 5.)



Answer	Count
a	3
d	2
b	1
e	1


Summary	Value
Mean	0.00
Std.dev	0.00
Min	0.00
Median	0.00
Max	0.00

Assume the lifespan of light bulbs manufactured by Bright Inc. can be modeled with a normal distribution with a mean of 300 days and a standard deviation of 40 days. What percentage of light bulbs produced by Bright Inc. will survive longer than 365 days?

- a. 2.94%
- b. 94.84%
- *c. 5.16%
- d. 9.34%
- e. 90.68%

(62) Question "To6.F.P.05.1.1.MC.IQ4" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 7 out of the total of 50 students. The average score was 0.14 out of 1.

(Back to the question summary Table 5.)



Answer	Count
a	3
e	2
b	1
Unanswered	1

Summary	Value
Mean	0.14
Std.dev	0.38
Min	0.00
Median	0.00
Max	1.00

Assume the distribution of IQ scores for adults can be modeled with a normal distribution with a mean score of 100 points and a standard deviation of 10 points. What percentage of adults have an IQ score higher than 128 points?



- a. 99.74%
- *b. 0.26%
- c. 89.97%
- d. 0.56%
- e. 10.03%

(63) Question "To6.F.P.05.1.1.MC.Whale4" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 13 out of the total of 50 students. The average score was 0 out of 1.

(Back to the question summary Table 5.)

Assume the length of female humpback whales can be modeled with a normal distribution with a mean of 13.7 meters and a standard deviation of 0.5 meters. What percentage of female humpback whales are longer than 14.3 meters?

- a. 14.46%
- *b. 11.51%
- c. 13.14%
- d. 88.49%
- e. 85.54%





Answer	Count
c	6
a	3
d	3
e	1

Summary	Value
Mean	0.00
Std.dev	0.00
Min	0.00
Median	0.00
Max	0.00

(64) Question "To6.F.P.05.1.1.MC.Cow4" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 9 out of the total of 50 students. The average score was 0.11 out of 1.

(Back to the question summary Table 5.)



Answer	Count
c	4
d	3
a	1
b	1

Summary	Value
Mean	0.11
Std.dev	0.33
Min	0.00
Median	0.00
Max	1.00

Assume the weight of a certain breed of cow can be modeled with a normal distribution with a mean of 750 kg and a standard deviation of 30 kg. What percentage of cows from this breed will weigh more than 770 kg?



- *a. 25.14%
- b. 31.21%
- c. 74.86%
- d. 27.09%
- e. 68.79%

(65) Question "To6.F.Q.05.1.1.MC.mm5" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 4 out of the total of 50 students. The average score was 0.5 out of 1.

(Back to the question summary Table 5.)

Assume the weight of bags of M&Ms can be modeled with the normal distribution with mean 50 grams and standard deviation 1 gram. What percent of M&M bags will have a weight between 49.5 and 51.5 grams?

- *a. 62.47%
- b. 97.72%
- c. 24.17%
- d. 84.13%





Answer	Count
a	2
b	1
d	1

Summary	Value
Mean	0.50
Std.dev	0.58
Min	0.00
Median	0.50
Max	1.00

(66) Question "To6.F.Q.05.1.1.MC.Bulb5" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 12 out of the total of 50 students. The average score was 0.25 out of 1.

(Back to the question summary Table 5.)



Answer	Count
c	5
a	3
d	2
b	1
Unanswered	1

Summary	Value
Mean	0.25
Std.dev	0.45
Min	0.00
Median	0.00
Max	1.00

Assume the lifespan of light bulbs manufactured by Bright Inc. can be modeled with a normal distribution with a mean of 300 days and a standard deviation of 40 days. What percentage of light bulbs produced by Bright Inc. will survive for between 230 days and 330 days?

*a. 73.33%



b. 99.38%

c. 62.55%

d. 34.46%

(67) Question "To6.F.Q.05.1.1.MC.IQ5" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 7 out of the total of 50 students. The average score was 0 out of 1.

(Back to the question summary Table 5.)



Summary		Value
Mean		0.00
Std.dev		0.00
Min		0.00
Median		0.00
Max		0.00



Answer	Count
b	3
a	2
d	2

Assume the distribution of IQ scores for adults can be modeled with a normal distribution with a mean score of 100 points and a standard deviation of 10 points. What percentage of adults have an IQ score between 92 and 112 points?

a. 97.72%
b. 77.4%
*c. 67.3%
d. 63.2%

(68) Question "To6.F.Q.05.1.1.MC.Whale5" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 7 out of the total of 50 students. The average score was 0.29 out of 1.
(Back to the question summary Table 5.)

Assume the length of female humpback whales can be modeled with a normal distribution with a mean of 13.7 meters and a standard deviation of 0.5 meters. What percentage of female humpback whales are between 12.5 and 14.7 meters in length?
*a. 96.90%
b. 99.91%
c. 65.54%
d. 91.92%



Summary		Value
Mean		0.29
Std.dev		0.49
Min		0.00
Median		0.00
Max		1.00


Answer	Count
d	4
a	2
b	1

(69) Question "To6.F.Q.05.1.1.MC.Cow5" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 20 out of the total of 50 students. The average score was 0.4 out of 1.

(Back to the question summary Table 5.)

Assume the weight of a certain breed of cow can be modeled with a normal distribution with a mean of 750 kg and a standard deviation of 30 kg. What percentage of cows from this breed will weigh between 725 kg and 785 kg?

- *a. 67.57%
- b. 97.72%
- c. 63.31%
- d. 53.59%





Answer	Count
a	8
b	5
c	3
d	3
Unanswered	1

Summary	Value
Mean	0.40
Std.dev	0.50
Min	0.00
Median	0.00
Max	1.00

(70) Question "To6.G.R.05.1.1.MC.mm1" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 10 out of the total of 50 students. The average score was 0.2 out of 1.

(Back to the question summary Table 5.)



Answer	Count
d	4
c	3
a	2
b	1

Summary	Value
Mean	0.20
Std.dev	0.42
Min	0.00
Median	0.00
Max	1.00

Assume the weight of bags of M&Ms can be modeled with the normal distribution with mean 50 grams and standard deviation 1 gram. The Mars Company that manufactures M&Ms wants to set the label weight so that only 2% of all bags of this size are under the label weight. What should they make the label weight?

- *a. 47.95 grams
- b. 49.16 grams
- c. 50.51 grams
- d. 50.58 grams

(71) Question "To6.G.R.05.1.1.MC.Bulb1" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 14 out of the total of 50 students. The average score was 0.07 out of 1.

(Back to the question summary Table 5.)



Assume the lifespan of light bulbs manufactured by Bright Inc. can be modeled with a normal distribution with a mean of 300 days and a standard deviation of 40 days. 20% of light bulbs produced by Bright Inc. survive less than how many days?

*a. 266.4 days

b. 218.0 days

c. 248.8 days

d. 273.2 days





Answer	Count
d	7
c	3
b	2
a	1
Unanswered	1

Summary	Value
Mean	0.07
Std.dev	0.27
Min	0.00
Median	0.00
Max	1.00

(72) Question "To6.G.R.05.1.1.MC.IQ1" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 4 out of the total of 50 students. The average score was 0 out of 1.

(Back to the question summary Table 5.)



Summary		Value
Mean		0.00
Std.dev		0.00
Min		0.00
Median		0.00
Max		0.00

Answer	Count
a	2
d	2

Assume the distribution of IQ scores for adults can be modeled with a normal distribution with a mean score of 100 points and a standard deviation of 10 points. 25% of adults have an IQ score lower than what value?



a. 100.3 points
b. 89.1 points
*c. 93.3 points
d. 97.8 points

(73) Question "To6.G.R.05.1.1.MC.Whale1" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 13 out of the total of 50 students. The average score was 0.23 out of 1.

(Back to the question summary Table 5.)

Assume the length of female humpback whales can be modeled with a normal distribution with a mean of 13.7 meters and a standard deviation of 0.5 meters. 15% of female humpback whales are shorter than what length?

- a. 14.79 meters
- b. 12.62 meters
- c. 14.22 meters
- *d. 13.18 meters



Answer	Count
b	7
d	3
a	1
c	1
Unanswered	1



Summary	Value
Mean	0.23
Std.dev	0.44
Min	0.00
Median	0.00
Max	1.00

(74) Question "To6.G.R.05.1.1.MC.Cow1" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 9 out of the total of 50 students. The average score was 0.33 out of 1.

(Back to the question summary Table 5.)

Assume the weight of a certain breed of cow can be modeled with a normal distribution with a mean of 750 kg and a standard deviation of 30 kg. 35% of cows from this breed will weigh less than what weight?

a. 703.2 kg
b. 802.3 kg
c. 751.2 kg
*d. 738.3 kg



Answer	Count
a	3
d	3
b	2
c	1



Summary	Value
Mean	0.33
Std.dev	0.50
Min	0.00
Median	0.00
Max	1.00

(75) Question "To6.G.S.05.1.1.MC.mm2" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 14 out of the total of 50 students. The average score was 0.5 out of 1.

(Back to the question summary Table 5.)

Assume the weight of bags of M&Ms can be modeled with the normal distribution with mean 50 grams and standard deviation 1 gram. 80% of all bags will have a weight under what value?

- *a. 50.84 grams
- b. 51.28 grams
- c. 50.67 grams
- d. 51.65 grams



Summary		Value
Mean		0.50
Std.dev		0.52
Min		0.00
Median		0.50
Max		1.00



Answer	Count
a	7
d	4
c	3

(76) Question "To6.G.S.05.1.1.MC.Bulb2" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 11 out of the total of 50 students. The average score was 0.27 out of 1.

(Back to the question summary Table 5.)

Assume the lifespan of light bulbs manufactured by Bright Inc. can be modeled with a normal distribution with a mean of 300 days and a standard deviation of 40 days. 95% of light bulbs produced by Bright Inc. survive less than how many days?

- a. 359.1 days
- b. 331.2 days
- c. 394.4 days
- *d. 365.8 days





Answer	Count
a	6
d	3
b	1
c	1

Summary	Value
Mean	0.27
Std.dev	0.47
Min	0.00
Median	0.00
Max	1.00

(77) Question "To6.G.S.05.1.1.MC.IQ2" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 9 out of the total of 50 students. The average score was 0 out of 1.

(Back to the question summary Table 5.)



Summary		Value
Mean		0.00
Std.dev		0.00
Min		0.00
Median		0.00
Max		0.00

Answer	Count
c	6
b	3

Assume the distribution of IQ scores for adults can be modeled with a normal distribution with a mean score of 100 points and a standard deviation of 10 points. 90% of adults will have an IQ score lower than what value?


a. 126.9 points
b. 109.8 points
c. 115.6 points
*d. 112.8 points

(78) Question "To6.G.S.05.1.1.MC.Whale2" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 9 out of the total of 50 students. The average score was 0.11 out of 1.

(Back to the question summary Table 5.)

Assume the length of female humpback whales can be modeled with a normal distribution with a mean of 13.7 meters and a standard deviation of 0.5 meters. 65% of female humpback whales are shorter than what length?

- a. 14.65 meters
- b. 13.51 meters
- c. 14.07 meters
- *d. 13.90 meters



Answer	Count
a	2
b	2
c	2
Unanswered	2
d	1



Summary	Value
Mean	0.11
Std.dev	0.33
Min	0.00
Median	0.00
Max	1.00

(79) Question "To6.G.S.05.1.1.MC.Cow2" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 7 out of the total of 50 students. The average score was 0.29 out of 1.

(Back to the question summary Table 5.)

Assume the weight of a certain breed of cow can be modeled with a normal distribution with a mean of 750 kg and a standard deviation of 30 kg. 70% of cows from this breed will weigh less than what weight?

- *a. 765.6 kg
- b. 770.1 kg
- c. 775.2 kg
- d. 788.4 kg





Summary		Value
Mean		0.29
Std.dev		0.49
Min		0.00
Median		0.00
Max		1.00

Answer	Count
c	3
a	2
d	2

(80) Question "To6.G.T.05.1.1.MC.mm3" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 11 out of the total of 50 students. The average score was 0.27 out of 1.

(Back to the question summary Table 5.)



Answer	Count
c	5
a	3
b	2
Unanswered	1



Summary	Value
Mean	0.27
Std.dev	0.47
Min	0.00
Median	0.00
Max	1.00

Assume the weight of bags of M&Ms can be modeled with the normal distribution with mean 50 grams and standard deviation 1 gram. 20% of all M&M bags will weigh more than what weight?

- *a. 50.84 grams
- b. 49.16 grams
- c. 50.20 grams
- d. 49.80 grams

(81) Question "To6.G.T.05.1.1.MC.Bulb3" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 8 out of the total of 50 students. The average score was 0.25 out of 1.

(Back to the question summary Table 5.)



Answer	Count
c	3
a	2
b	2
d	1



Summary	Value
Mean	0.25
Std.dev	0.46
Min	0.00
Median	0.00
Max	1.00

Assume the lifespan of light bulbs manufactured by Bright Inc. can be modeled with a normal distribution with a mean of 300 days and a standard deviation of 40 days. 5% of light bulbs produced by Bright Inc. survive more than how many days?

- *a. 366.0 days
- b. 378.4 days
- c. 351.2 days
- d. 341.6 days

(82) Question "To6.G.T.05.1.1.MC.IQ3" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 14 out of the total of 50 students. The average score was 0.14 out of 1.

(Back to the question summary Table 5.)



Answer	Count
b	7
a	5
d	2

Summary	Value
Mean	0.14
Std.dev	0.36
Min	0.00
Median	0.00
Max	1.00

Assume the distribution of IQ scores for adults can be modeled with a normal distribution with a mean score of 100 points and a standard deviation of 10 points. 30% of adults will have an IQ score higher than what value?


- a. 109.1 points
- b. 116.6 points
- c. 98.6 points
- *d. 105.2 points

(83) Question "To6.G.T.05.1.1.MC.Whale3" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 5 out of the total of 50 students. The average score was 0 out of 1.

(Back to the question summary Table 5.)

Assume the length of female humpback whales can be modeled with a normal distribution with a mean of 13.7 meters and a standard deviation of 0.5 meters. 40% of female humpback whales are longer than what length?

- a. 13.90 meters
- b. 14.06 meters
- *c. 13.83 meters
- d. 11.95 meters





Summary		Value
Mean	0.00	
Std.dev	0.00	
Min	0.00	
Median	0.00	
Max	0.00	

Answer	Count
d	3
a	1
Unanswered	1

(84) Question "To6.G.T.05.1.1.MC.Cow3" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 12 out of the total of 50 students. The average score was 0.08 out of 1.

(Back to the question summary Table 5.)



Answer	Count
d	8
c	3
b	1

Summary	Value
Mean	0.08
Std.dev	0.29
Min	0.00
Median	0.00
Max	1.00

Assume the weight of a certain breed of cow can be modeled with a normal distribution with a mean of 750 kg and a standard deviation of 30 kg. 45% of cows from this breed will weigh more than what weight?


- a. 772.0 kg
- *b. 753.9 kg
- c. 761.3 kg
- d. 758.9 kg

(85) Question "To6.G.U.05.1.1.MC.mm4" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 5 out of the total of 50 students. The average score was 0 out of 1.

(Back to the question summary Table 5.)

Assume the weight of bags of M&Ms can be modeled with the normal distribution with mean 50 grams and standard deviation 1 gram. 65% of all M&M bags will weigh more than what weight?

*a. 49.61 grams
b. 50.39 grams
c. 50.35 grams
d. 49.65 grams





Answer		Summary	
Answer	Count		Value
c	3	Mean	0.00
d	1	Std.dev	0.00
Unanswered	1	Min	0.00
		Median	0.00
		Max	0.00

(86) Question "To6.G.U.05.1.1.MC.Bulb4" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 12 out of the total of 50 students. The average score was 0.33 out of 1.

(Back to the question summary Table 5.)

Assume the lifespan of light bulbs manufactured by Bright Inc. can be modeled with a normal distribution with a mean of 300 days and a standard deviation of 40 days. 70% of light bulbs produced by Bright Inc. survive longer than how many days?

- *a. 279.2 days
- b. 303.2 days
- c. 281.0 days
- d. 263.5 days





Answer	Count
b	5
a	4
c	2
Unanswered	1

Summary	Value
Mean	0.33
Std.dev	0.49
Min	0.00
Median	0.00
Max	1.00

(87) Question "To6.G.U.05.1.1.MC.IQ4" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 21 out of the total of 50 students. The average score was 0.33 out of 1.

(Back to the question summary Table 5.)



Answer	Count
a	7
d	7
c	5
b	2

Summary	Value
Mean	0.33
Std.dev	0.48
Min	0.00
Median	0.00
Max	1.00



Assume the distribution of IQ scores for adults can be modeled with a normal distribution with a mean score of 100 points and a standard deviation of 10 points. 90% of adults will have an IQ score higher than what value?

- *a. 87.2 points
- b. 112.8 points
- c. 91.0 points
- d. 109.0 points

(88) Question "To6.G.U.05.1.1.MC.Whale4" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 6 out of the total of 50 students. The average score was 0.17 out of 1.
(Back to the question summary Table 5.)

Assume the length of female humpback whales can be modeled with a normal distribution with a mean of 13.7 meters and a standard deviation of 0.5 meters. 75% of female humpback whales are longer than what length?

- a. 13.90 meters
- b. 14.05 meters
- *c. 13.37 meters
- d. 14.04 meters





Answer	Count
a	2
b	2
c	1
d	1

Summary	Value
Mean	0.17
Std.dev	0.41
Min	0.00
Median	0.00
Max	1.00

(89) Question "To6.G.U.05.1.1.MC.Cow4" is given on the right. This question was selected from the question set with a frequency of 0.2. The question was administered to 6 out of the total of 50 students. The average score was 0.17 out of 1.
(Back to the question summary Table 5.)

Assume the weight of a certain breed of cow can be modeled with a normal distribution with a mean of 750 kg and a standard deviation of 30 kg. 85% of cows from this breed will weigh more than what weight?
*a. 718.8 kg
b. 781.2 kg
c. 775.5 kg
d. 724.5 kg



Answer		Count
b		3
d		2
a		1

Summary	Value
Mean	0.17
Std.dev	0.41
Min	0.00
Median	0.00
Max	1.00