

KMS: $\Omega^2 = 1 + 10\,\phi^2 + 10.02\,\chi^2$, $\tilde{V} = \lambda_\phi(\phi^4 + 2\phi^2\chi^2 + \chi^4/2)/4$, metric formulation
(a) Einstein-frame potential $V = \tilde{V}/\Omega^4$
trajectory 1
trajectory 2
trajectory 3
$V/\lambda_\phi M_{pl}^4$
$\chi/M_{pl}$
$\phi/M_{pl}$
(b) top view: the fields roll from the star to the origin
start at rest, $\phi = 3.1$
plateau
valley $\phi = 0$
$\chi/M_{pl}$
$\phi/M_{pl}$
$V/\lambda_\phi M_{pl}^4$
(c) the ball's view: a flat ridge at $\theta = 0$
ridge
valley
canonical radial distance $\rho/M_{pl}$
angle $\theta = \arctan(\chi/\phi)$ [deg]
(d) off the ridge: $\chi \propto e^{0.078N}$ (thin lines) until the fall
$|\chi|/M_{pl}$
trajectory 1, $\chi_0 = 1.1e-02$
trajectory 2, $\chi_0 = 1.1e-03$
trajectory 3, $\chi_0 = 1.1e-04$
e-folds from the start, $N$