










Postmastectomy Radiation Therapy: An ASTRO-ASCO-SSO Clinical Practice Guideline

Rachel B. Jimenez, MD¹ ; Yara Abdou, MD² ; Penny Anderson, MD³; Parul Barry, MD⁴; Lisa Bradfield, BA⁵; Julie A. Bradley, MD⁶; Lourdes D. Heras, MPH⁷; Atif Khan, MD, MS⁸ ; Cindy Matsen, MD⁹ ; Rachel Rabinovitch, MD¹⁰ ; Chantal Reyna, MD, MHA¹¹; Kilian E. Salerno, MD¹²; Sarah E. Schellhorn, MD¹³ ; Deborah Schofield, PhD¹⁴; Kekoa Tapparra, MD, PhD, MPH¹⁵; Iman Washington, MD¹⁶ ; Jean L. Wright, MD¹⁷ ; Youssef H. Zeidan, MD, PhD¹⁸ ; Richard C. Zellars, MD¹⁹; and Kathleen C. Horst, MD²⁰

DOI <https://doi.org/10.1200/JCO-25-01747>

ABSTRACT

ASCO Guidelines provide recommendations with comprehensive review and analyses of the relevant literature for each recommendation, following the guideline development process as outlined in the *ASCO Guidelines Methodology Manual*. ASCO Guidelines follow the *ASCO Conflict of Interest Policy for Clinical Practice Guidelines*.

Clinical Practice Guidelines and other guidance (“Guidance”) provided by ASCO is not a comprehensive or definitive guide to treatment options. It is intended for voluntary use by clinicians and should be used in conjunction with independent professional judgment. Guidance may not be applicable to all patients, interventions, diseases or stages of diseases. Guidance is based on review and analysis of relevant literature, and is not intended as a statement of the standard of care. ASCO does not endorse third-party drugs, devices, services, or therapies and assumes no responsibility for any harm arising from or related to the use of this information. See complete ASCO and ASTRO disclaimers in [Appendix 1](#) (online only) for more.

PURPOSE This guideline provides recommendations on use of postmastectomy radiation therapy (PMRT) in breast cancer treatment. Updated recommendations detail indications for PMRT in the upfront surgical setting and after neoadjuvant systemic therapy, and provide guidance on appropriate target volumes, dosing, and treatment techniques.

METHODS A multidisciplinary American Society for Radiation Oncology–ASCO–Society of Surgical Oncology task force addressed four key radiation therapy (RT) questions in patients with breast cancer who undergo mastectomy: (1) indications for PMRT after upfront surgery, (2) indications for PMRT after neoadjuvant systemic therapy followed by surgery, (3) appropriate PMRT volumes and dose-fractionation regimens, and (4) treatment techniques. Recommendations were based on a systematic review and created using a predefined consensus-building methodology for quality of evidence grading and strength of recommendation.

RECOMMENDATIONS After upfront mastectomy, PMRT is indicated for most patients with node-positive breast cancer and select patients with node-negative disease. PMRT is also recommended after neoadjuvant systemic therapy for patients presenting with locally advanced disease and for those with residual nodal disease at the time of surgery. PMRT is conditionally recommended for patients with cT1–3N1 or cT3N0 breast cancer with pathologically negative nodes after neoadjuvant systemic therapy (ypNo). When PMRT is delivered, treatment to the ipsilateral chest wall or reconstructed breast and regional lymphatics is recommended, with moderate hypofractionation preferred, but with conventional fractionation approaches acceptable in rare cases. Computed tomography–based volumetric treatment planning with 3-dimensional conformal RT is recommended, with intensity-modulated RT advised when three-dimensional conformal RT is unable to achieve treatment goals. Deep inspiration breath-hold techniques are also recommended for normal tissue sparing. For patients with skin involvement, positive superficial margins, and/or lymphovascular invasion, use of a bolus is recommended, but routine use of tissue-equivalent bolus is not recommended.

Additional information is available at www.asco.org/breast-cancer-guidelines.

ACCOMPANYING CONTENT

 Appendix
 Data Supplement

Accepted July 30, 2025

Published September 16, 2025

ASTRO Board of Directors approval:
May 23, 2025

ASCO Evidence Based Medicine
Committee approval:

May 27, 2025

SSO Executive Committee approval:
June 2, 2025

J Clin Oncol 00:1-20

© 2025 by American Society of
Clinical Oncology



[View Online
Article](#)

TARGET POPULATION AND AUDIENCE

Target Population

Adults (age ≥ 18 years) who received a diagnosis of breast cancer and underwent mastectomy.

Target Audience

Radiation oncologists, surgical oncologists, medical oncologists, oncology nurses, patients, radiologists, and other relevant oncologic professionals.

(eg, 3-dimensional conformal radiation therapy [3-D CRT], intensity-modulated radiation therapy [IMRT], protons, breath hold, bolus) for treating patients who receive PMRT?

METHODS

Guideline Development Process

This systematic review–based guideline product was developed by an ASTRO-ASCO-SSO joint task force that consisted of a multidisciplinary team of radiation, medical, and surgical oncologists; a radiation oncology resident; a medical physicist; a patient representative, and an ASTRO guidelines staff member with health research methodology expertise (Appendix [Table A1](#), online only). This guideline was also developed in collaboration with the European Society for Radiotherapy and Oncology, which provided representatives and peer reviewers. This work was funded by ASTRO.

The joint task force used evidence-based methodologies to develop guideline recommendations in accordance with the National Academy of Medicine standards.^{4,5} The evidence identified from key questions (KQs) is assessed using the Population, Intervention, Comparator, Outcome, Timing, Setting (PICOTS) framework. A systematic review of the KQs is completed, which includes creation of evidence tables that summarize the evidence base task force members use to formulate recommendations. [Table 1](#) describes ASTRO's recommendation grading system. See Data Supplement 1 (online only) for a list of abbreviations used in the guideline.

Consensus Development

Consensus is evaluated using a modified Delphi approach. Task force members confidentially indicate their level of agreement on each recommendation based on a five-point Likert scale, from “strongly agree” to “strongly disagree.” A prespecified threshold of 75% ($\geq 90\%$ for expert opinion recommendations) of raters who select “strongly agree” or “agree” indicates consensus is achieved. Recommendation(s) that do not meet this threshold are removed or revised. Recommendations edited in response to task force or reviewer comments are resurveyed before submitting for approval.

Scope of the Guideline

The scope of this guideline is to define the role of PMRT in the curative-intent management of invasive breast cancer, including the indications for PMRT after upfront surgery and following neoadjuvant systemic therapy, and to discuss the appropriate target volumes and technical specifications for PMRT. Given the rapid adoption of biologically tailored neoadjuvant systemic therapy and the de-escalation of axillary surgery with the use of sentinel lymph node biopsy or targeted axillary dissection, this guideline seeks to

INTRODUCTION

According to the WHO, in 2022, breast cancer was the second most common cancer and the fourth leading cause of cancer mortality worldwide.¹ Although some patients may undergo breast-conservation therapy, others undergo mastectomy either by medical necessity or by choice. For these patients, postmastectomy radiation therapy (PMRT), which delivers radiation therapy (RT) to the residual skin and soft tissue of the ipsilateral chest wall and regional draining lymphatics, can decrease the risk of a locoregional recurrence (LRR) and improve breast cancer mortality.² As the absolute benefit of PMRT can vary according to patient and tumor characteristics, it is important to individualize treatment decision making to balance considerations of LRR and breast cancer mortality with the side effects of treatment.

The American Society for Radiation Oncology (ASTRO), ASCO, and the Society of Surgical Oncology (SSO) sought to jointly develop a new guideline to clarify patient selection criteria and appropriate technical approaches for the delivery of PMRT. This evidence review was completed to replace the 2016 PMRT guideline³ and to reflect the evolving understanding of the benefit of PMRT. With advancements in the management of breast cancer, including improved diagnostic imaging, trends in de-escalation of axillary surgery, newer and more tailored systemic therapy agents, and advances in RT techniques, there is a need to provide updated guidance regarding the appropriate indications for, and approaches to, PMRT in the modern era.

GUIDELINE QUESTIONS

This clinical practice guideline addresses four overarching clinical questions: (1) What are the indications for PMRT in patients who receive mastectomy as their initial treatment for breast cancer? (2) What are the indications for PMRT in patients who receive neoadjuvant systemic therapy before mastectomy? (3) What are the appropriate treatment volumes (eg, chest wall or reconstructed breast, regional nodes, boost) and dose-fractionation regimens for patients who receive PMRT? (4) What are the appropriate techniques

TABLE 1. ASTRO Recommendation Grading Classification System

ASTRO's recommendations are based on evaluation of multiple factors including the QoE and panel consensus which, among other considerations, inform the strength of recommendation. QoE is based on the body of evidence available for a particular key question and includes consideration of number of studies, study design, adequacy of sample sizes, consistency of findings across studies, and generalizability of samples, settings, and treatments.

Strength of Recommendation	Definition	Overall QoE Grade	Recommendation Wording
Strong	Benefits clearly outweigh risks and burden, or risks and burden clearly outweigh benefits. All or almost all informed people would make the recommended choice.	Any (usually high, moderate, or expert opinion)	"Recommend/Should"
Conditional	Benefits are finely balanced with risks and burden, or appreciable uncertainty exists about the magnitude of benefits and risks. Most informed people would choose the recommended course of action, but a substantial number would not. A shared decision-making approach regarding patient values and preferences is particularly important.	Any (usually moderate, low, or expert opinion)	"Conditionally Recommend"
Overall QoE Grade	Type/Quality of Study	Evidence Interpretation	
High	Two or more well-conducted and highly generalizable RCTs or well-conducted meta-analyses of such randomized trials.	The true effect is very likely to lie close to the estimate of the effect based on the body of evidence.	
Moderate	One well-conducted and highly generalizable RCT or a meta-analysis including such a trial OR Two or more RCTs with some weaknesses of procedure or generalizability OR Two or more strong observational studies with consistent findings.	The true effect is likely to be close to the estimate of the effect based on the body of evidence, but it is possible that it is substantially different.	
Low	One RCT with some weaknesses of procedure or generalizability OR One or more RCTs with serious deficiencies of procedure or generalizability or extremely small sample sizes OR Two or more observational studies with inconsistent findings, small sample sizes, or other problems that potentially confound interpretation of data.	The true effect may be substantially different from the estimate of the effect. There is a risk that future research may significantly alter the estimate of the effect size or the interpretation of the results.	
Expert opinion ^a	Consensus of the panel based on clinical judgment and experience, due to the absence of evidence or limitations in evidence.	Strong consensus ($\geq 90\%$) of the panel guides the recommendation despite insufficient evidence to discern the true magnitude and direction of the net effect. Further research may better inform the topic.	

NOTE. ASTRO's methodology allows for the use of implementation remarks meant to convey clinically practical information that may enhance the interpretation and application of the recommendation. While each recommendation is graded according to recommendation strength and QoE, these grades should not be assumed to extend to the implementation remarks.

Abbreviations: ASTRO, American Society for Radiation Oncology; QoE, quality of evidence; RCTs, randomized controlled trials.

^aA lower QoE, including expert opinion, does not imply that the recommendation is conditional. Many important clinical questions addressed in guidelines do not lend themselves to clinical trials, but there is still consensus that the benefits of a treatment or diagnostic test clearly outweigh its risks and burden.

address the indications and approaches for PMRT in the context of these advances in the multidisciplinary care of breast cancer. In this guideline, PMRT refers to treatment of the chest wall and ipsilateral regional nodes, including at-risk axillary, supraclavicular or infraclavicular, and internal mammary nodes (IMN). Specific situations where treatment volumes may be less comprehensive are noted in the text.

Evidence Review: KQs, Key Outcomes, and Inclusion and Exclusion Criteria

KQs were developed by the ASTRO guideline subcommittee in conjunction with the guideline chairs and then reviewed by the full task force. Using the PICOTS framework (Table 2), a systematic search of human participant studies retrieved from the Ovid MEDLINE and Embase databases was conducted for English-language publications between January 1,

2005, through October 2023, and then the search was updated through October 15, 2024.

The key outcomes of interest include LRR, disease-free survival (DFS), breast cancer mortality, distant metastasis-free survival, and overall survival (OS). Other key outcomes of interest include appropriate dose-fractionation regimens, nodal volumes considered for treatment, and optimal RT techniques to minimize toxicities. This guideline addresses only the subjects specified in the KQs (Table 2). There are several important questions in the management of patients with breast cancer that are outside the scope of this guideline, including inflammatory breast cancer, management of ductal carcinoma in situ after mastectomy, management of locally or regionally recurrent disease, and detailed discussions of chemotherapy regimens and surgical approaches, including axillary management. This guideline also does not encompass recommendations on reirradiation, RT in the setting of

TABLE 2. Key Questions in PICO Format

KQ	Population	Intervention	Comparator	Outcome
1. What are the indications for PMRT in patients who receive mastectomy as their initial treatment for breast cancer?				
	Adult patients with breast cancer	PMRT	No PMRT	Local recurrence Regional recurrence Locoregional recurrence Disease-free survival Breast cancer mortality Distant metastasis-free survival Overall survival
2. What are the indications for PMRT in patients who receive neoadjuvant systemic therapy before mastectomy?				
	Same as KQ1	PMRT after neoadjuvant systemic therapy	No PMRT after neoadjuvant systemic therapy	Local recurrence Regional recurrence Locoregional recurrence Disease-free survival Breast cancer mortality Distant metastasis-free survival Overall survival
3. What are the appropriate treatment volumes (eg, chest wall/reconstructed breast, regional nodes, boost) and dose-fractionation regimens for patients who receive PMRT?				
	Same as KQ1	Hypofractionation Chest wall/reconstructed breast without RNI RNI including IMNs Boost	Conventional fractionation Chest wall/reconstructed breast with RNI RNI without IMNs No boost	Local recurrence Regional recurrence Locoregional recurrence Disease-free survival Breast cancer mortality Distant metastasis-free survival Toxicity and adverse effects
4. What are the appropriate techniques (eg, 3-D CRT, IMRT, protons, breath hold, bolus) for treating patients who receive PMRT?				
	Same as KQ1	IMRT (including VMAT) Electrons Protons Set-up verification, image guidance/surface guidance Respiratory management, gating, breath hold Bolus	3-D CRT PMRT with photons No bolus	Local recurrence Regional recurrence Locoregional recurrence Disease-free survival Breast cancer mortality Distant metastasis-free survival Toxicity and adverse effects

Abbreviations: 3-DCRT, 3-dimensional conformal radiation therapy; IMN, internal mammary nodes; IMRT, intensity-modulated radiation therapy; KQs, key questions; PICO, Population, Intervention, Comparator, Outcome; PMRT, postmastectomy radiation therapy; RNI, regional nodal irradiation; RT, radiation therapy; VMAT, volumetric modulated arc therapy.

oligometastatic or palliative disease, phyllodes tumors, or sarcomas of the breast.

Allowable publication types included prospective randomized controlled trials (RCTs), prospective nonrandomized studies, meta-analyses, and retrospective studies. The population of interest was adults (age ≥ 18 years) who received a diagnosis of breast cancer and underwent mastectomy. Trial size required for inclusion was ≥ 50 patients for RCTs and meta-analyses, and ≥ 100 patients for prospective nonrandomized and retrospective studies. KQ1 addresses indications for PMRT in patients who receive mastectomy as their initial treatment. Retrospective studies were excluded for KQ1 given the strength of the prospective data available for this question. Universal exclusion criteria included the following: preclinical and nonhuman studies; publication types such as abstract only, review articles, case reports, comments, or editorials; and study types such as dosimetric or contouring studies, health economics or cost analysis studies, or large registry database studies. For specific subquestions where limited data were available,

expert opinion was relied on to support recommendations. Full-text articles were assessed by the task force to determine the final included study list resulting in 104 studies (see the [Preferred Reporting Items for Systematic Reviews and Meta-Analyses](#)^{6,7} flow diagram showing the number of articles screened, excluded, and included in the evidence review, and Data Supplement 2 for the literature search strategy, which includes the evidence search parameters and inclusion and exclusion criteria).

Guideline Review and Approval

The guideline was reviewed by 17 official peer reviewers (Data Supplement 3) and revised accordingly. The modified guideline was posted on the ASTRO website for public comment from September to October 2024. The final guideline was approved by the ASTRO Board of Directors, the ASCO Evidence Based Medicine Committee, and SSO Executive Committee; and endorsed by the American Society of Breast Surgeons and the Royal Australian and New Zealand College of Radiologists.

Guideline Updating

Based on periodic formal review of the emerging literature, the respective organizations will determine the need to update. The ASCO Guidelines Methodology Manual (available at www.asco.org/guideline-methodology) provides additional information about the guideline update process. This is the most recent information as of the publication date.

RESULTS

The data used by the task force to formulate recommendations are summarized in evidence tables available in Data Supplement 4. References selected and published in this document are representative and not all-inclusive. Additional ancillary articles not in the evidence tables are included in the text; these were not used to support the evidence-based recommendations but may have informed expert opinion.

KQs AND RECOMMENDATIONS

KQ1: Indications for PMRT With Mastectomy as Initial Treatment (Table 3)

See evidence tables in Data Supplement 4 for the data supporting the recommendations for KQ1, and Figure 1.

What Are the Indications for PMRT in Patients Who Receive Mastectomy as Their Initial Treatment for Breast Cancer?

Over the last 4 decades, multiple RCTs and pooled analyses have shown a significant reduction in LRR and improved DFS

and OS in women with pT3-4 and/or node-positive breast cancer who receive PMRT.^{2,8-10,15-17} Support for the use of PMRT in patients with nodal involvement comes from the Early Breast Cancer Trialists' Collaborative Group (EBCTCG) meta-analysis.^{2,11} This analysis included women who underwent mastectomy and axillary dissection, and were enrolled in trials evaluating PMRT to the chest wall and regional lymph nodes. PMRT significantly reduced breast cancer recurrence, breast cancer mortality, and all-cause mortality in patients with positive lymph nodes.^{2,11} Among these patients, the risk of LRR and the benefit of PMRT increased with nodal burden, with the greatest absolute reduction of LRR and improvement in DFS and OS observed in patients with ≥ 4 positive nodes (pN2), but still with significant benefits for those with 1-3 positive nodes (pN1). Notably, there was no differentiation between patients with pN1 or pN1mic status after axillary dissection in these trials. However, among patients with pN1mic disease, the magnitude of benefit of PMRT is often considered to be lower than in those with higher nodal burden, and therefore, requires assessment of other clinicopathologic features, as noted in the discussion of patients with node-negative disease to follow.

It should also be acknowledged that the EBCTCG meta-analysis was limited to trials initiated by 1995,^{2,18} so while the majority of the included studies reflected the receipt of appropriate systemic therapies for the time period, most did not use current evidence-based systemic regimens (eg, immunotherapy, human epidermal growth factor receptor 2 [HER2]-directed therapy), which have been recognized to further confer a locoregional control and DFS benefit.^{8,10,17} In this context, the benefit of PMRT for low-volume, node-positive disease (pN1) has been questioned. The Selective Use of Postoperative Radiotherapy after Mastectomy

TABLE 3. Indications for PMRT With Mastectomy as Initial Treatment

KQ1 Recommendation	Strength of Recommendation	Quality of Evidence (refs)
1. For patients with node-positive (pN+) breast cancer, PMRT is recommended. <i>Implementation remarks:</i> Omission of PMRT may be appropriate for select patients with pN1mic or low nodal burden pN1a disease following ALND who have favorable clinicopathologic features. Favorable clinicopathologic features include pT1-2 disease, low-to-intermediate grade HR-positive/HER2-negative subtype, postmenopausal status, absence of LVI, and a low 21-gene recurrence score.	Strong	High ^{2,8-12}
2. For patients with any pT4 breast cancer, PMRT is recommended even in the absence of any other risk factors.	Strong	High ^{2,8}
3. For patients with pT3N0 breast cancer, PMRT is conditionally recommended. <i>Implementation remark:</i> PMRT may be omitted or treatment volumes reduced (eg, chest wall alone) for patients with favorable clinicopathologic features including low-to-intermediate grade, HR-positive/HER2-negative subtype, postmenopausal status, absence of LVI, and a low 21-gene recurrence score.	Conditional	High ^{2,8,10}
4. For patients with pT1-2N0 breast cancer, PMRT is not recommended. <i>Implementation remark:</i> Select patients with pT1-2N0 breast cancer who have multiple unfavorable clinicopathologic features (eg, triple-negative, high histologic grade, LVI, young age, and/or central/medially located tumors) may benefit from PMRT.	Strong	Low ^{2,13,14}
5. For patients with positive surgical margins after mastectomy and no other indication for PMRT, RT to the chest wall/reconstructed breast alone is conditionally recommended.	Conditional	Expert opinion

Abbreviations: ALND, axillary lymph node dissection; HR/HER2, hormone receptor/human epidermal growth factor receptor 2; KQ, key question; LVI, lymphovascular invasion; PMRT, postmastectomy radiation therapy; RT, radiation therapy.




FIG 1. Indications for PMRT. ^aSee implementation remarks in Table 3 for details. ^bSee implementation remarks in Table 4 for details. ^cPMRT may be omitted in the setting of complete pathological response in the breast and lymph nodes (ypT0N0). PMRT, postmastectomy radiation therapy.

(SUPREMO; ClinicalTrials.gov identifier: [NCT00966888](https://clinicaltrials.gov/ct2/show/study/NCT00966888)) trial evaluated the impact of PMRT on OS for patients with limited nodal disease in the upfront surgical setting after axillary lymph node dissection with at least eight lymph nodes removed. Final results from this study will provide additional insights regarding the value of PMRT in this favorable-risk population.¹⁹ Additionally, in an era where the biology of breast cancer guides systemic therapy, questions arise as to whether biology should also inform RT recommendations. Indeed, MA.39/TAILOR-RT (A Randomized Trial of Regional Radiotherapy in Biomarker Low-Risk Node-Positive Breast Cancer, ClinicalTrials.gov identifier: [NCT03488693](https://clinicaltrials.gov/ct2/show/study/NCT03488693)) randomizes patients with estrogen receptor (ER)-positive, HER2-negative pT1-2N1a disease and a non-high-risk recurrence score (recurrence score ≤ 25) to PMRT or no PMRT, with a primary end point of recurrence-free interval. The results from this trial will also inform recommendations for PMRT for patients receiving upfront surgery with limited axillary nodal disease including pN1mic and favorable ER-positive tumor biology. Notably, in this study, axillary lymph node dissection is not mandatory; however, there can be no more than two positive lymph nodes present if sentinel lymph node biopsy alone is performed.²⁰ While this study evaluates selective omission of PMRT in favorable-risk ER-positive, HER2-negative breast cancer, it should be noted that in historical studies evaluating the mortality benefit of PMRT, the magnitude of

benefit was higher for patients with ER-positive biology, despite a comparatively lower local recurrence risk,²¹ largely due to the competing risk for distant failure. Therefore, while LRR is an important end point, it need not be the sole consideration in recommendations for PMRT.

In the node-negative setting, data support the use of PMRT in patients with high-risk features. Larger tumor size (≥ 5 cm), younger age (< 40 years), and hormone receptor-negative disease have all independently been associated with a greater benefit of PMRT in node-negative patients.^{17,22,23} Although specific RCTs directly focusing on T4N0 breast cancer are limited, there are data supporting the benefits of PMRT in reducing LRR and improving survival outcomes in this patient population.^{2,8-10,24,25} Invasion of the skin and pectoralis muscle has also been associated with higher rates of LRR,⁸ and were considered high-risk criteria for eligibility in both the Danish 82b/c trials.^{24,25} For patients with pT3N0 breast cancer, who were included in these RCTs, there was a $> 50\%$ reduction in LRR with PMRT.^{24,25} However, this group comprised $< 10\%$ of the study cohorts, modern systemic regimens known to reduce LRR were not used, and neither trial demonstrated a significant improvement in breast cancer-specific or OS in patients with pT3N0 breast cancer.²⁶ Multiple population data set analyses have demonstrated no breast cancer-specific survival benefit of PMRT across unselected patients with pT3N0 disease, even for patients

<50 years of age.²⁷⁻²⁹ Patients with pT3No disease were included in the European Organisation for Research and Treatment of Cancer (EORTC) 22922 trial, which demonstrated a benefit of regional nodal irradiation (RNI) in terms of any breast cancer recurrence and breast cancer mortality, with no significant difference in OS. However, only 3.5% of the patients had pT3No disease. Given the demonstrated local regional control benefit and uncertain survival benefit of PMRT for patients with pT3No breast cancer, PMRT is conditionally recommended and when employed, smaller treatment volumes (eg, chest wall RT alone) may be used at the discretion of the provider.^{2,8} PMRT may be omitted for patients with favorable clinicopathological features including low-to-intermediate grade, ER-positive, HER2-negative subtype, postmenopausal status, absence of lymphovascular invasion (LVI), and low 21-gene recurrence score. These patients are included in both the SUPREMO (ClinicalTrials.gov identifier: [NCT00966888](#)) and the TAILOR-RT (ClinicalTrials.gov identifier: [NCT03488693](#)) trials, and these results may better define the impact of PMRT in this patient population.

Few RCTs have evaluated PMRT in the pT1-2No setting.¹³ A single study in patients with stage I or II triple-negative breast cancer demonstrated a relapse-free survival and OS benefit with PMRT following total mastectomy, partial axillary dissection, and adjuvant chemotherapy; however, the systemic therapy regimens used are no longer considered standard of care.¹³ Additionally, 19% of patients had node-positive disease and no subset analysis was performed to determine if the benefit of PMRT was primarily in the node-positive subgroup.¹³ EORTC 22922 also included patients with stage I and II breast cancer with lymph node-negative, central or medially located tumors, and identified a breast cancer recurrence and breast cancer mortality benefit with

the addition of chest wall and RNI, although mastectomy patients comprised only approximately 25% of participants.¹⁴ Overall, meta-analyses and retrospective studies of patients with pT1-2No breast cancer demonstrate excellent outcomes without PMRT for most patients, with reported 10-year LRR rates between 2.1% and 12.8%, and the majority reporting rates of 3% to 7%.^{2,30} However, these data also suggest that LVI, young age, high histologic grade disease, and positive margins increase the risk of LRR such that PMRT may be beneficial, particularly for patients with multiple high-risk features.³⁰

Finally, there are no RCTs evaluating the role of RT in patients with positive margins following mastectomy. Positive margins, however, are consistently associated with a greater risk of local recurrence.³¹ Recognizing the consistent reduction in local recurrence of approximately 50% with the use of PMRT, PMRT is conditionally recommended in the setting of positive margins when re-excision is not feasible.⁸ The extent and location of positive margins, tumor biology, consideration of other high-risk features (eg, LVI, young age, tumor grade), and plan for adjuvant therapies should be weighed together to determine the value of PMRT for an individual patient.

KQ2: Indications for PMRT With Neoadjuvant Systemic Therapy (Table 4)

See evidence tables in Data Supplement 4 for the data supporting the recommendations for KQ2 and Figure 1.

What Are the Indications for PMRT in Patients Who Receive Neoadjuvant Systemic Therapy Before Mastectomy?

Over the past decade, the use of neoadjuvant systemic therapy has increased for specific subsets of patients with

TABLE 4. Indications for PMRT With Neoadjuvant Systemic Therapy

KQ2 Recommendation	Strength of Recommendation	Quality of Evidence (refs)
1. For patients with initial cT4 or cN2-3 breast cancer who receive neoadjuvant systemic therapy, PMRT is recommended regardless of pathologic response.	Strong	Moderate ³²⁻³⁶
2. For patients with positive lymph nodes after neoadjuvant systemic therapy (ypN+), PMRT is recommended.	Strong	Moderate ³⁷⁻⁴¹
3. For patients with cT1-3N1 or cT3N0 breast cancer with pathologic negative nodes after neoadjuvant systemic therapy (ypN0), PMRT is conditionally recommended <i>Implementation remarks:</i> Patients with high-risk features (eg, young age, LVI, high residual cancer burden in the breast) may derive a greater benefit from PMRT. PMRT may be omitted in the setting of complete pathologic response in the breast and lymph nodes (ypTON0).	Conditional	Moderate ^{37,38,40-50}
4. For patients with cT1-2N0 breast cancer with pathologic negative nodes after neoadjuvant systemic therapy (ypN0), PMRT is not recommended. <i>Implementation remark:</i> Patients with multiple high-risk features (eg, young age, LVI, high residual cancer burden in the breast) may benefit from PMRT.	Strong	Moderate ^{37,38,43-45,47-49}
5. For patients with positive surgical margins after neoadjuvant systemic therapy, PMRT is recommended	Strong	Expert opinion

Abbreviations: KQ, key question; LVI, lymphovascular invasion; PMRT, postmastectomy radiation therapy.

breast cancer, notably those with cT2 or greater or clinically node-positive disease to downstage the breast and axilla, and in those with HER2-positive or triple-negative biology.^{42,51} Several studies have shown that patients with initial cT4 or cN2-3 (also defined by the American Joint Committee on Cancer 6th edition as stage III) breast cancer who receive neoadjuvant systemic therapy have improved LRR with PMRT regardless of their response to neoadjuvant therapy.³²⁻³⁶ Some studies have also shown an improvement in OS, but these were small retrospective evaluations.^{32,33} Based on the current evidence, PMRT is recommended for patients with initial presentation of cT4 or cN2-3 disease who receive neoadjuvant systemic therapy, regardless of pathological response.³²⁻³⁶ In addition, several studies have demonstrated that residual nodal disease after neoadjuvant systemic therapy (ypN+) is associated with an increased risk of LRR.³⁷⁻³⁹

The extent of axillary nodal disease after neoadjuvant systemic therapy (ie, ypN1 v ypN2-3) is also an important risk factor.^{37,40} This risk is further elevated in patients with cT3 tumors.³⁸ The addition of PMRT in patients with ypN+ improves locoregional control with incremental benefit noted in patients with increased axillary burden.^{39,40} An OS benefit for PMRT has been reported for patients with ypN2-3 disease.⁴⁰ It is worth noting that the benefit of PMRT for residual nodal disease in these studies was evaluated in the setting of axillary nodal dissection. Results from the Alliance A011202 trial (ClinicalTrials.gov identifier: [NCT01901094](#)), evaluating whether RT to the undissected axilla and other regional lymph nodes after sentinel lymph node biopsy is noninferior to axillary lymph node dissection (with RT only to the undissected regional lymph nodes), will further clarify the value of extensive axillary surgery after neoadjuvant systemic therapy and provide guidance regarding the appropriate RT treatment volumes needed in this patient population.

In patients who begin treatment with clinically involved axillary lymph nodes (cN1) and convert to pathologically node-negative after neoadjuvant systemic therapy (ypNo), the full reporting of the NSABP B-51/Radiation Therapy Oncology Group 1304 trial (ClinicalTrials.gov identifier: [NCT01872975](#)), which randomly assigned these patients to PMRT or no RT, will help to resolve the clinical equipoise that exists on the use of PMRT in this setting. On this protocol, patients were eligible if they had clinical axillary nodal involvement (cN1) as assessed before neoadjuvant chemotherapy by palpation, ultrasound, computed tomography (CT) scan, magnetic resonance imaging, positron emission tomography (PET) scan, or PET/CT scan, and patients with N2-3 disease detected clinically or by imaging were ineligible.⁵² Data from a 2022 prospective Dutch registry, in which patients with cT1-2N1 breast cancer (defined as 1-3 suspicious nodes on imaging before neoadjuvant chemotherapy), and had negative nodes at surgery (ypNo) and did not receive PMRT,

demonstrated a low LRR rate of 2.1% at 5 years, supporting de-escalation of PMRT in patients with ypNo disease after neoadjuvant systemic therapy.⁵³ Another pooled analysis showed a 5-year LRR rate of 3% after mastectomy without PMRT in patients with HER2-positive disease achieving ypNo.³⁹ Although several retrospective studies have shown similar LRR-free survival rates with and without PMRT after achieving ypNo,^{43,44} a meta-analysis including 12 studies of over 17,000 patients who achieved a pathological complete response in the lymph nodes (ypNo) demonstrated a significant benefit with PMRT in all stages, with the greatest benefit in stage III disease.³⁵ For patients who achieve a pathological complete response in the nodes, certain features appear to increase the risk of LRR and may suggest a continued benefit with PMRT. For example, several reports have suggested that baseline clinicopathological factors including young age, cT3-4 disease, triple-negative subtype, and LVI may predict higher rates of LRR, so PMRT is conditionally recommended in patients with multiple high-risk factors.^{33,37,38,43,45,46,54} Similarly, other pathological features after neoadjuvant systemic therapy are associated with demonstrably higher risks of LRR (eg, high-volume residual invasive disease in the breast, persistent LVI, residual HER2-positive and triple-negative disease, close margins) and may be indications for PMRT after neoadjuvant systemic therapy.^{37,40,41,44,45,47,48,55,56}

The benefits of PMRT may be higher in younger women compared with older women.^{32,43,57} In a retrospective study of young women (age <35 years) who received neoadjuvant anthracycline-based chemotherapy, the use of PMRT reduced LRR and improved OS.³² This finding is consistent with a study from Korea that found age ≤40 years to be an independent predictor of LRR.⁴³ Treatment decision making regarding the role for PMRT should include a discussion of risks and benefits, particularly for young patients. For those who have residual invasive disease in the breast, the advances in adjuvant systemic therapy (eg, CDK 4/6 inhibitors, capecitabine, ado-trastuzumab emtansine-1, pembrolizumab) may further impact the risk-benefit ratio of PMRT.⁵⁸⁻⁶⁰

Although neoadjuvant systemic therapy is most often used for larger tumors and those with nodal involvement, there may be some patients with cT1-2N0 disease who receive neoadjuvant treatment, particularly those with HER2+ and triple-negative biological subtype. For these patients, PMRT is not recommended if the nodes are pathologically negative (ypNo) as the risk of an LRR after mastectomy alone is low.³⁸ However, the presence of multiple clinical and pathological risk factors (eg, young age, LVI, high residual cancer burden in the breast) increases the risk of an LRR such that PMRT may be an option.^{37,38,43-45,47-49}

Finally, there are limited data to inform PMRT recommendations for patients with positive surgical margins after

neoadjuvant therapy. However, given that positive margins are a conditional indication for PMRT in the upfront surgery setting,³¹ PMRT is recommended for positive margins after neoadjuvant systemic therapy when re-excision is not feasible based on expert opinion.

KQ3: PMRT Treatment Volumes and Dose-Fractionation Regimens (Table 5)

See evidence tables in Data Supplement 4 for the data supporting the recommendations for KQ3.

What Are the Appropriate Treatment Volumes (eg, chest wall or reconstructed breast, regional nodes, boost) and Dose-Fractionation Regimens for Patients Who Receive PMRT?

In the EBCTCG meta-analysis of 8,135 women pooled from trials comparing no PMRT with PMRT, inclusive of the chest wall and regional lymph nodes, PMRT significantly reduced both LRR, overall recurrence, and breast cancer mortality, with the chest wall being the most common site of LRR.² The meta-analysis also included eight trials that did not include the chest wall in the treatment fields (ie, only treated the regional lymph node basins) and found that RT in those studies did not have a significant impact on overall recurrence or breast cancer mortality. As 50%–80% of all local recurrences identified in RCTs were located in the chest wall,^{9,17} inclusion of the chest wall as a PMRT target structure is recommended regardless of surgical margins, although

direct comparisons of RT with versus without chest wall volumes are limited.

Several large RCTs have evaluated the value of RNI in patients with medially or centrally located tumors, positive lymph nodes, or in patients with high-risk node-negative breast cancer.^{14,61,76} The EORTC 22922 trial randomly assigned patients who had centrally or medially located primary tumors, irrespective of axillary involvement, or laterally located tumors with axillary involvement, to either whole breast/chest wall irradiation and RNI (inclusive of IMNs) or whole breast, chest wall irradiation alone.⁶¹ Approximately one quarter of these patients were treated with mastectomy. At 10 years, the addition of RNI resulted in a significantly improved breast cancer mortality rate, improved DFS, and a trend toward improved OS. The 15-year results continued to demonstrate a significant reduction in breast cancer mortality and any breast cancer recurrence with the addition of IMN or supraclavicular irradiation in patients with stage I to III breast cancer.¹⁴ The Canadian Cancer Trials Group MA.20 trial also evaluated the addition of RT to the supraclavicular lymph nodes, axillary apical lymph nodes, and the IMNs for patients with node-positive disease or high-risk node-negative disease.⁷⁶ Although it did not include patients treated with mastectomy, it did demonstrate that the addition of RNI reduced the rate of any breast cancer recurrence, further supporting the use of RNI when defining target coverage for patients with node-positive or high-risk node-negative breast cancer. For those patients who have undergone an axillary

TABLE 5. PMRT Treatment Volumes and Dose-Fractionation Regimens

KQ3 Recommendation	Strength of Recommendation	Quality of Evidence (refs)
1. For patients receiving PMRT, treatment to the ipsilateral chest wall/reconstructed breast and regional lymphatics (ie, at-risk axillary nodes, supra/infraclavicular nodes, and IMNs) is recommended. <i>Implementation remarks:</i> Treatment to the chest wall/reconstructed breast alone may be used in select patients (eg, pT3N0). Coverage of the IMNs may be individually determined based on tumor location (medial/central), tumor size, and extent of nodal involvement.	Strong	High ^{2,12,61-65}
2. For patients <i>without</i> breast reconstruction receiving PMRT, moderate hypofractionation is recommended. <i>Implementation remarks:</i> Moderate hypofractionation is preferred, given equivalent oncologic outcomes and reduced toxicity. Conventional fractionation may be an option in rare circumstances.	Strong	High ⁶⁶⁻⁷³
3. For patients <i>with</i> breast reconstruction receiving PMRT, moderate hypofractionation (preferred) or conventional fractionation is recommended.	Strong	Moderate ^{65,74} (moderate hypofx) High ^{14,61,62,66,67,71-73,75} (conventional fx)
4. For patients with T4 breast cancer or close/positive margins receiving PMRT, a boost to the chest wall/scar is conditionally recommended.	Conditional	Low ^{66,67,74,76-80}
5. For patients with nodal disease not surgically addressed and at risk of harboring residual disease, a nodal boost is recommended.	Strong	Expert opinion

NOTE. Moderate hypofractionation is most frequently defined as 266 to 267 cGy per fraction for 15-16 fractions. Conventional fractionation is most frequently defined as 180 to 200 cGy per fraction for 25 to 28 fractions.⁶⁶⁻⁷³

Abbreviations: fx, fractionation; hypofx, hypofractionation; IMNs, internal mammary nodes; KQ, key question; PMRT, postmastectomy radiation therapy.

dissection and receive PMRT, data do not support a benefit to including the dissected stations of the axilla, typically axillary levels I and II; however, an increasing number of studies support the omission of axillary lymph node dissection after a positive sentinel lymph node biopsy and in these circumstances, coverage of all axillary nodal basins is advised.^{2,10,81} Additionally, among patients who undergo an inadequate axillary dissection or whose pathological specimens demonstrate tumor deposits and/or emboli into the axillary fat, coverage of the dissected axilla is indicated. Although it is a departure from traditional PMRT to irradiate the chest wall without inclusion of the regional lymph node stations, this approach may be considered in select patients (eg, positive surgical chest wall margins as the only indication for PMRT or pT3N0 tumors in the absence of other high-risk factors), given the concern for local over regional recurrence risk.^{2,17}

Although RNI in the EORTC 22922 and MA.20 trials included treatment of the IMNs, there is debate as to which patients might benefit most from IMN irradiation, particularly with the higher cardiopulmonary exposure associated with this approach and the potential for increased toxicity.^{14,76} The benefit of IMN RT was specifically evaluated in studies from Denmark, France, and South Korea in which patients with breast cancer were treated with whole breast or chest wall RT, supraclavicular, and axillary apex irradiation with or without IMN RT.⁶²⁻⁶⁴ The DBCG trial was a prospective, nonrandomized population-based cohort study that assigned IMN irradiation only to patients with right-sided disease to mitigate concerns for cardiac RT exposure among patients with left-sided cancer.^{63,82} This study demonstrated a significant improvement in distant recurrence, death from breast cancer, and a 4.7% improvement in OS at 15 years among right-sided patients who received IMN RT. A French RCT enrolled patients with positive axillary lymph nodes or central, medial tumors with or without positive axillary lymph nodes and randomly assigned patients to receive RT to the chest wall and supraclavicular nodes with or without IMN RT.⁶⁴ This study did not demonstrate an OS benefit for IMN RT. In patients with positive axillary lymph nodes, a small but nonsignificant benefit was observed in favor of IMN RT. This study was underpowered and was performed in the 2-dimensional era of treatment planning, limiting its applicability.⁶⁴ Finally, the Korean Radiation Oncology Group 08-06 trial randomly assigned patients with pathologically confirmed, node-positive disease after mastectomy or breast-conservation surgery and axillary lymph node dissection to RNI with or without IMN RT.⁶² The study demonstrated a nonstatistically significant 2.6% absolute decrease in distant metastases without a significant improvement in DFS. However, in an ad hoc subgroup analysis of patients with medial or centrally located tumors, both DFS and breast cancer-specific mortality at 7 years were significantly improved with the addition of IMN RT, suggesting that IMN RT in this subgroup of patients is beneficial.⁶² Importantly, none of these trials, or the aforementioned RNI studies, demonstrated an increased risk of cardiac

toxicity with treatment of the IMNs within the reported follow-up periods, lending support for the routine inclusion of IMN RT for patients with clinically or radiographically detected IMN nodes and those with central or medially located breast tumors, particularly when axillary lymph nodes are positive.^{61-64,76}

Most of the studies evaluating PMRT have used conventional fractionation with doses approximating 5,000 cGy, EQD2.9. However, a number of retrospective analyses have suggested that moderately hypofractionated PMRT regimens result in reduced acute and late toxicity compared with conventional regimens, with comparable survival outcomes.^{66,67,74,83-85} There is also precedent from RCTs to support the use of moderately hypofractionated regimens. In the landmark British Columbia study, 3,750 cGy in 16 fractions was used to deliver PMRT.¹² In the United Kingdom Standardization of Breast Radiotherapy A trial, enrolling 2,236 women with breast cancer, 15% underwent PMRT, and hypofractionated schedules resulted in similar locoregional failure rates, and lower adverse events, compared with conventional fractionation.⁶⁸ Additionally, the United Kingdom Standardization of Breast Radiotherapy B trial involved 2,215 women with breast cancer, with approximately 8% receiving PMRT.⁶⁹ At a median follow-up of 10 years, they found that 4,005 cGy in 15 daily fractions yielded comparable outcomes to 5,000 cGy in 25 daily fractions in terms of locoregional tumor control and lower late normal tissue effects, as assessed by both patient- and physician-reported photographs, and arm and shoulder symptoms.⁶⁹ In China, a noninferiority study randomly assigned 820 patients with at least four positive axillary nodes or T3-4 disease, excluding those with internal mammary or supraclavicular nodal involvement, to moderate hypofractionation (4,350 cGy in 15 fractions) or conventional fractionation (5,000 cGy in 25 fractions). At a median follow-up of 58.4 months, locoregional failure was deemed noninferior between arms (8.3% hypofractionation v 8.1% conventional fractionation), and there was a lower rate of grade 3 skin toxicity in the hypofractionation arm.⁷⁰ An additional RCT confirmed that there were no discernible differences in toxicities, LRR, distant failure rate, or DFS between PMRT regimens of 4,005 cGy in 15 fractions and 5,000 cGy in 25 fractions.⁷¹ Given equivalent oncological outcomes and reduced toxicity, moderate hypofractionation is recommended for patients without breast reconstruction who are receiving PMRT, with careful consideration of dose selection for those with more advanced disease (eg, T4 and cN3 disease) or those with limited response to neoadjuvant systemic therapy.

None of these trials, however, were specifically designed to evaluate the impact of hypofractionation on cosmetic outcomes in the setting of breast reconstruction. As such, there has been hesitancy to transition to shorter treatment schedules for patients who opt for breast reconstruction, but there are increasing data to support its use.^{74,83} The phase III Fractionation on Patient Outcomes After Breast

REConstruction trial randomly assigned 400 patients with stage 0–III breast cancer, excluding T4 disease, after mastectomy with implant-based reconstruction to hypofractionated RT (4,256 cGy in 16 fractions) or conventional RT (5,000 cGy in 25 fractions).⁶⁵ The primary end point was improvement in the Physical Well-Being domain of Functional Assessment of Cancer Therapy–Breast at 6 months. Results showed a significant reduction in patients requiring a treatment break with hypofractionation compared with conventional fractionation (2.7% v 7.7%). There was no difference in chest wall toxicity between the two groups at a median follow-up of 40.4 months.^{65,86} Based on these data, the use of moderate hypofractionation is recommended as the preferred PMRT approach in the setting of implant-based reconstruction.⁶⁵

Another completed RCT, Alliance A221505 (RT CHARM: Hypofractionated Post Mastectomy Radiation with Breast Reconstruction; ClinicalTrials.gov identifier: [NCT03414970](https://clinicaltrials.gov/ct2/show/study/NCT03414970))⁸⁷ randomly assigned nearly 900 patients with T1–3N1–2 or T3N0 disease undergoing mastectomy with immediate or delayed reconstruction (implant-based or autologous) to hypofractionated PMRT (4,256 cGy in 16 fractions) or conventional PMRT (5,000 cGy in 25 fractions) with a primary end point of reconstruction complication rate. In this trial, patients with T4 and N3 disease, including IMN involvement, were excluded.⁸⁷ Final published results from this study will provide additional data on the clinical outcomes and toxicity of hypofractionated PMRT with reconstruction.⁸⁷ Until then, conventional fractionation is also recommended as an option.

It is important to note the variability in dose regimens and eligibility criteria used in each of the previously mentioned trials,^{65,70,72,86,87} reflecting uncertainties regarding biologically effective dosing between conventional fractionation and moderate hypofractionation. Because of the evolving understanding of both the alpha/beta ratio of breast cancer and the effect of shorter treatment regimens on repopulation, care should be taken when selecting hypofractionated regimens, particularly for patients with high-risk features (eg, T4 or N3 disease), to ensure that definitive RT doses are used. In these scenarios, a separate boost to suspected residual disease, as could be employed in the conventional fractionation setting, may also be appropriate (see the subsequent discussion of a boost).

One limitation of these trials is the relatively small number of Black, Hispanic, or Asian patients enrolled, which limits the understanding of potential cosmetic differences in these populations. Prior studies have demonstrated Asian, Black, and Hispanic patients experience worse acute and long-term skin quality-of-life outcomes after breast RT than White patients.^{88,89} Therefore, extra consideration in treatment planning and supportive care is advised in these patient populations, recognizing that their relative lack of representation on the available trials should not unduly limit their access to shorter, more convenient treatment

schedules, particularly given recognized disparities in the receipt of PMRT among Black and Hispanic patients with stage III breast cancer.^{90,91}

Finally, there is increasing interest in the use of ultrahypofractionated treatment regimens (ie, 2,600 cGy in five fractions) in breast cancer, although there are limited data in patients receiving PMRT. Early reports suggest comparable outcomes with ultrahypofractionation to the chest wall and nodal regions,⁹² and additional trials are underway to further evaluate these abbreviated treatment regimens for patients requiring PMRT.^{93,94}

Evidence supporting the administration of a chest wall scar boost to improve local control rates is limited and has never been established prospectively. Although the majority of LRRs after mastectomy occur on the chest wall,⁹ only retrospective studies have examined the use of chest wall boosts for high-risk patients and have provided some support for doses up to 6,600 cGy using conventional fractionation.^{95–98} Despite this, a survey among breast radiation oncologists demonstrated that 55% routinely use a chest wall boost following PMRT and an additional 18% prescribe a boost depending on margin status.⁹⁹ Pragmatically, the administration of a chest wall boost is conditionally recommended in cases of T4 disease and positive margins where concern for residual disease is enhanced. Of note, an evaluation of women who had undergone PMRT from the California Cancer Registry identified disparities in the receipt of a chest wall boost, with poor and Hispanic women more commonly receiving a chest wall boost than affluent and non-Hispanic women of similar cancer stage and biology.¹⁰⁰ This suggests that objective criteria for using a chest wall boost may not be uniformly applied and care should be taken, whenever possible, not only to follow consistent criteria, as detailed here, but to ensure representative enrollment of diverse patient populations in prospective studies evaluating treatment techniques.

Similarly, there are no randomized studies examining the use of a boost to gross disease in undissected nodal basins, such as the supraclavicular fossa or internal mammary chain, despite recognition that involvement of these nodes is a poor prognostic factor in breast cancer.^{63,64} Institutional retrospective analyses suggested that an additional boost to involved supraclavicular and internal mammary chain nodes can be delivered safely and may improve local control rates, but these data are limited by small sample sizes.^{101,102} However, if adding a boost to an undissected node, doses of 6,000 cGy EQD2 should be considered for microscopic disease and at least 6,600 cGy EQD2 for gross or residual disease.

KQ4: Appropriate PMRT Delivery Techniques (Table 6)

See evidence tables in Data Supplement 4 for the data supporting the recommendations for KQ4.

TABLE 6. Appropriate PMRT Delivery Techniques

KQ4 Recommendation	Strength of Recommendation	Quality of Evidence (refs)
1. For patients receiving PMRT, CT-based volumetric treatment planning with 3-D CRT is recommended.	Strong	Moderate ^{14,62-64,82,103-105}
2. For patients receiving PMRT, IMRT (including VMAT) is recommended when 3-D CRT is unable to achieve treatment goals (ie, target coverage and normal tissue avoidance). <i>Implementation remark:</i> Use of IMRT (including VMAT) may increase OAR low-dose exposure compared with 3-D CRT.	Strong	Moderate ¹⁰⁶⁻¹¹⁰
3. For patients receiving PMRT, DIBH is recommended when lower doses to normal tissues, including the heart and lungs, can be achieved compared with free breathing. <i>Implementation remarks:</i> Other normal tissue sparing techniques may be used. For DIBH, use of a real-time monitoring device (eg, SGRT, spirometry-based systems, chest wall monitoring system) and image-guided verification are advised.	Strong	Moderate ^{109,111,112}
4. For patients receiving PMRT treated with IMRT (including VMAT), daily image guidance, in conjunction with regular 3-D assessments (eg, CBCT, SGRT), is recommended.	Strong	Low ¹¹³
5. For patients with cT1-3 breast cancer receiving PMRT, the routine use of tissue-equivalent bolus is not recommended. <i>Implementation remark:</i> Bolus may be used in circumstances where improved dosimetric coverage of the skin is needed.	Strong	Moderate ¹¹⁴⁻¹¹⁹
6. For patients with skin involvement, positive superficial margins, and those with dermal lymphatic involvement and/or extensive LVI, the use of tissue-equivalent bolus is recommended.	Strong	Expert opinion

Abbreviations: 3-D CRT, 3-dimensional conformal radiation therapy; CBCT, cone beam computed tomography; CT, computed tomography; DIBH, deep inspiration breath hold; IMRT, intensity-modulated radiation therapy; KQ, key question; LVI, lymphovascular invasion; OAR, organ at risk; PMRT, postmastectomy radiation therapy; SGRT, surface-guided radiation therapy; VMAT, volumetric modulated arc therapy.

What Are the Appropriate Techniques (eg, 3-D CRT, IMRT, protons, breath hold, bolus) for Treating Patients Who Receive PMRT?

High-quality evidence from RCTs directly evaluating various RT techniques for PMRT is limited, and most foundational studies used 2-dimensional or 3-D photon therapy, with or without an electron component.^{8,12,14,62-64,82,103,120} Modern RT design is based on contouring of the target areas (chest wall and nodal basins as indicated) and the adjacent relevant organs at risk (OARs) as appropriate (ie, heart, left ventricle, left anterior descending [LAD] artery and/or right coronary artery, bilateral lungs, contralateral breast, spinal cord, thyroid, esophagus, humeral head, stomach, liver, and/or brachial plexus).^{121,122} Use of contouring guidelines, such as those provided by the Radiation Therapy Oncology Group atlas, RADCOMP (Radiotherapy Comparative Effectiveness),¹²¹ and European atlases,^{122,123} may be used to assist with accurate target and OAR delineation. The goal of volumetric treatment planning is to use CT information to adequately cover the target volumes while minimizing dose to normal tissues, taking individual anatomic variation into account. While this approach has historically been underutilized in RT treatment planning for breast cancer compared with other disease sites, CT-based volumes should be used for individualized RT planning for breast cancer. The task force acknowledges that in many cases, more stringent planning parameters can be achieved than what is detailed in Table 7 and the concept of as low as reasonably achievable should prevail for all RT treatment plans. However, it is also recognized that the guidance provided may not be uniformly achievable for all

patients' plans, given anatomic concerns. When intensity-modulated planning is employed, attention to low doses delivered to OARs that do not typically receive dose exposure with 3-D planning is advised (eg, spinal cord, stomach, liver).^{14,62-64,82,103-105} Finally, given the current state of the data, specific dose constraints are not provided for all relevant vulnerable normal tissues (eg, LAD artery or right coronary artery); however, contouring of these structures is still advised to rationally constrain unnecessary exposure during treatment planning.

For PMRT field design, 3-D CRT treatment planning can use a variety of techniques, for example, partially wide tangent fields to include the IMN contour, a medial electron field matched to narrow photon tangents, or electrons to the chest wall alone with a match to a photon supraclavicular field with or without a posterior axillary field.¹²⁴ Advanced modulated planning techniques (eg, IMRT including volumetric modulated arc therapy [VMAT]) can be used to improve high-dose conformality and target coverage. Studies evaluating the treatment of patients with breast cancer using tomotherapy have also shown feasibility.^{125,126} Studies comparing various techniques have shown low LRR rates regardless of technique.^{66,103,105,106}

Treatment with inverse planned IMRT can decrease the high-dose exposure of OARs compared with 3-DCRT, and in some cases decrease the risk of toxicity.^{66,107,108,127} A retrospective study of patients receiving PMRT comparing 3-D CRT with VMAT reported a reduction in RT pneumonitis in the cohort treated with VMAT.⁶⁶ Another study

TABLE 7. Guidance on Target Coverage

Structure	Goal	5,000-5,040 cGy in 25-28 fx	4,000-4,256 cGy in 15-16 fx
Ipsilateral chest wall ^{52,87}	Ideal	D95 ≥95% PTV	D95 ≥95% PTV
	Acceptable	D90 ≥90% PTV	D90 ≥90% PTV
	Ideal	D0.1 cc ≤110%	D0.1 cc ≤107% ^b
	Acceptable	D0.1 cc ≤115%Rx	D0.1 cc 115% Rx
Axilla ^{a,42,79}	Ideal	D95 ≥95% PTV	D95 ≥95% PTV
	Acceptable	D90 ≥90% PTV	D90 ≥90% PTV
	Ideal	D0.1 cc 110% Rx	D0.1 cc 107% Rx ^b
	Acceptable	D0.1 cc ≤115%Rx	D0.1 cc 115% Rx
Supraclavicular fossa ^{52,87}	Ideal	D95 ≥95% PTV	D95 ≥95% PTV
	Acceptable	D90 ≥90% PTV	D90 ≥90% PTV
	Ideal	D0.1 cc 115% Rx	D0.1 cc 112% Rx ^b
	Acceptable	D0.1 cc ≤120%Rx	D0.1 cc 115% Rx
Internal mammary nodes ^{52,87}	Ideal	D95 ≥90% PTV	D95 ≥90% PTV
	Acceptable	D90 ≥80% PTV	D90 ≥80% PTV

NOTE. This table is a combination of evidence-based constraints and expert opinion.

Abbreviations: PTV, planning target volume; Rx, prescription dose.

^aIf patient has undergone a completion axillary dissection, coverage goals apply only to the targeted axilla.

^bExtrapolated from conventionally fractionated data.

demonstrated that adequate target coverage was achieved with both 3-D CRT and IMRT, with a decrease in moist desquamation in the cohort treated with IMRT (14.3% v 3.8%, respectively).¹⁰⁸ A third study described a decrease in moderate- and high-dose exposure to the shoulder in patients undergoing RNI with IMRT compared with 3-D CRT.¹²⁷ One tradeoff of reduced high-dose exposure to OARs with IMRT is an increase in low-dose OAR exposure. For example, one study described acute radiation-induced nausea associated with low-dose exposure of the upper abdominal structures,¹²⁸ side effects that are uncommon with 3-D CRT. Therefore, the use of IMRT (including VMAT) is recommended when 3-D CRT is unable to achieve treatment goals, with close attention to increased low-dose OAR exposure (see Table 8 for guidance on OARs).

Historically, a key cause of noncancer-related morbidity and mortality from PMRT came from undue cardiac exposure. Therefore, numerous studies comparing treatment planning techniques have been done with the goal of improving cardiac sparing.^{134,135} Although a dose-dependent relationship between cardiac exposure to RT and heart disease has been demonstrated in several landmark studies,¹³⁶⁻¹⁴⁰ no safe threshold has been established to prevent major cardiovascular events. Therefore, it is generally accepted that mean heart dose should be as low as reasonably achievable (Table 8). Special consideration should be given to minimizing RT exposure to the heart for patients with preexisting heart disease and certain risk factors (eg, diabetes, hypertension, and smoking), as these have been shown to be synergistic with cardiac RT exposure in increasing the risk of cardiac disease development.^{141,142}

A deep inspiration breath hold (DIBH) technique is one strategy for reducing dose to normal tissues, including the heart and lungs. Suitability for DIBH should be evaluated based on a patient's ability to maintain the breath hold and individual cardiac anatomy.^{111,112} Among patients for whom DIBH can be successfully implemented, cardiopulmonary dose can be reduced compared with a free-breathing 3-D CRT technique.^{111,143} Notably, there is an understanding that dose exposure to cardiac substructures including the left ventricle and the LAD artery does not correlate with mean heart dose. Both have been implicated in RT-associated cardiac toxicity in patients receiving RT for breast cancer, so particular consideration should be given to these substructures.^{139,144} An RCT comparing IMRT-DIBH with free-breathing 3-D CRT for patients with node-positive breast cancer showed lower mean doses for the ipsilateral lung, heart, and LAD artery, suggesting that patients receiving IMRT can also benefit from DIBH.¹⁰⁹ Although there was no difference in single-photon emission CT perfusion defects in the LAD territory or lung perfusion or function between groups, most patients in the IMRT-DIBH arm had stable or improved left ventricular ejection fraction at 1 year compared with a slightly declining left ventricular ejection fraction in the free-breathing cohort.¹⁰⁹ When DIBH is employed, use of a real-time monitoring device (eg, surface-guided radiation therapy [SGRT], spirometry-based or chest wall monitoring systems) and image-guided RT verification is advised to ensure the fidelity of respiratory displacement throughout treatment delivery.^{109,111}

The use of proton therapy remains under investigation at the time of guideline development. Single-institution series,

TABLE 8. Guidance on Organs at Risk

Structure	Goal	5,000-5,040 cGy in 25-28 fx	4,000-4,256 cGy in 15-16 fx
Ipsilateral lung ^{52,87}		V5 Gy ≤75%	V4 Gy ≤65% ^b
	Ideal	V10 Gy ≤65%	V8 Gy ≤55% ^b
		V20 Gy ≤35% ^a	V17 Gy ≤25% ^b
	Acceptable	V20 Gy ≤40%	V17 Gy ≤35%
Contralateral lung ⁸⁷	Ideal	V5 Gy ≤10%	V4 Gy ≤10% ^b
	Acceptable	V5 Gy ≤15%	V4 Gy ≤15% ^b
Heart ^{c,87}	Ideal (left-sided)	Dmean ≤3 Gy	Dmean ≤2.4 Gy ^b
	Acceptable (left-sided)	Dmean ≤5 Gy	Dmean ≤4Gy ^b
	Ideal (right-sided)	Dmean <2 Gy	Dmean ≤1.6 Gy ^b
	Acceptable (right-sided)	Dmean ≤3 Gy	Dmean ≤2.4 Gy ^b
Contralateral breast/chest wall ^{52,87}	Ideal	V3 Gy ≤10%	V3 Gy ≤10%
	Acceptable	V5 Gy ≤10%	V5 Gy ≤10%
Additional considerations			
Brachial plexus ⁹²	Suggested	D0.1cc ≤105%Rx	D0.1 cc ≤105% Rx
Esophagus	Suggested	V10 Gy <30%/V20 Gy <15% ¹²⁹	V8 Gy <30%/V17 Gy <15% ^{b,130}
Left ventricle ¹³¹	Suggested	V2 Gy <36%	V1.6 Gy <36% ^b
Spinal cord ^a	Suggested	D0.1 cc 45 Gy	D0.1 cc 38.54 Gy
Thyroid ^{132,133}	Suggested	Dmean <21 Gy	Dmean <21 Gy
Humeral head	Suggested	Dmean <20 Gy	Dmean <17 Gy
Stomach (left-sided)	Suggested	Dmean <3 Gy	Dmean <2.4 Gy
Liver (right-sided)	Suggested	Dmean <7 Gy	Dmean <5.6 Gy

NOTE. Where dose constraints differed by protocol, the more conservative guidelines were used. This table is a combination of evidence-based constraints and expert opinion and reflects guidance for routine treatments that do not employ a boost for gross or residual nodal disease. Abbreviations: Dmean, mean dose received by an organ; HYPOG-1, Hypofractionated Radiotherapy for Breast Cancer Nodal Irradiation-1; Rx, prescription dose.

^aBased on the HYPOG-1 trial protocol (ClinicalTrials.gov identifier: [NCT03127995](#)).

^bExtrapolated from conventionally fractionated data.

^cCardiac dose should be constrained as low as reasonably achievable.

prospective registry reports, and retrospective studies have demonstrated improved dosimetric target coverage, alongside preservation of cardiac function, compared with 3-D CRT and IMRT, particularly in the setting of RNI, including IMN irradiation.¹⁴⁵⁻¹⁴⁹ The RAD-COMP trial as well as the PARABLE and Danish Breast Proton trials are all evaluating major cardiac events between patients treated with proton versus photon RT and it is anticipated that these studies will provide more data on the appropriate role of proton PMRT in the future.^{121,130}

There is currently a lack of evidence to support a single optimal strategy for image guidance in the PMRT setting. Minimally, daily planar imaging, in conjunction with regular 3-D assessments (eg, cone beam CT [CBCT], SGRT), is recommended for patient localization for complex planning and multifield techniques, such as IMRT (including VMAT).^{150,151} Volumetric imaging (eg, CBCT) is valuable under these conditions to assess for evolving anatomic changes or setup variability that may adversely affect treatment accuracy. However, the planning target volume margins should account for setup variability and the type and frequency of image guidance used during treatment.¹⁵²

Alternatively, SGRT using the patient's external surface and nonionizing radiation can assist in PMRT patient setup,^{111,153} monitor intrafraction motion,^{154,155} and verify breath hold position.^{112,153} However, in addition to training and workflow issues,¹⁵⁵ significant tissue deformations and limitations in the technology to detect darker skin tones have been identified as potential drawbacks.¹⁵⁶ Currently, data are lacking to support the use of SGRT alone for daily PMRT treatment delivery. When SGRT is employed, it is advised to use it in conjunction with image-guided RT for setup verification. Guidance for the use of SGRT with image guidance, including common challenges and potential errors, has been published.¹⁵⁵

Finally, tissue-equivalent bolus has historically been used in PMRT with the recognition that most chest wall recurrences occur superficially or just under the skin. The skin and most superficial layer of chest wall tissue are key components of the RT target, and depending on the RT technique and beam energy used, surface dose may only reach 70%– to 80% of the prescribed dose. Tissue-equivalent bolus can be used to bring the skin dose closer to prescription dose. However, the application of tissue-equivalent bolus over the chest wall in

PMRT can vary with respect to frequency and thickness, and several clinical trials have permitted bolus at the discretion of the treating physician,^{52,86,157} thereby limiting the ability to formally evaluate the impact of bolus on clinical outcomes to help guide recommendations for the use of bolus with PMRT.

Multiple studies have identified a relationship between the use of bolus and increased skin toxicity.^{114-118,158} At the same time, despite the historical assumption of benefit, the impact of bolus on local control has been questioned, including three small retrospective studies that did not identify a local control benefit with bolus.¹¹⁶⁻¹¹⁸ One RCT of 59 patients, employing a risk-stratified bolus strategy with thicker and more frequent use of bolus in patients with frank skin involvement and no bolus versus 5-mm bolus on alternate days in standard-risk patients without skin involvement, found no decrement in chest wall local control within risk groups, although all patients in the high-risk group were treated with bolus.¹¹⁴ Although these analyses are limited by patient and treatment heterogeneity, they suggest insufficient evidence for a local control benefit with the routine use of bolus for patients with cT1-3 disease without a high risk of skin involvement.^{117,118} Understanding the value of bolus among patients with darker skin tones may be particularly critical, given the higher likelihood of skin toxicity and late skin effects from RT among non-White patients, although no studies to date have specifically evaluated the impact of bolus across different skin tones.^{90,91} Therefore, the routine use of bolus is not advised for all patients, but may be used in circumstances where improved dosimetric coverage of the skin is needed. In addition, for those patients with an increased risk of skin recurrence, including patients who present with skin involvement, positive anterior surgical margins, dermal lymphatic invasion, or extensive LVI, the use of bolus is recommended based on expert opinion.¹¹⁴

CONCLUSIONS AND FUTURE DIRECTIONS

Multiple RCTs and the EBCTCG meta-analysis have confirmed that PMRT reduces the risk of LRR and improves breast cancer mortality. However, the absolute risk reduction varies across individuals. There are ongoing efforts to try to better characterize risk according to tumor biology, and in the era of tailored systemic therapy, to further personalize treatment recommendations. Unfortunately, there are few data from available clinical trials to guide tailored management recommendations for patients based on sociodemographic characteristics, including race and access to health care. It is critical that future trials of PMRT ensure diverse trial enrollment and participation.

In addition, there are several potentially practice-changing trials that remain in active accrual or have not yet been published at the time of this guideline including trials related

to PMRT in favorable-risk disease (SUPREMO, MA.39/TAILOR-RT [ClinicalTrials.gov identifier: [NCT03488693](#)]), hypofractionation (RT CHARM [ClinicalTrials.gov identifier: [NCT03414970](#)],⁸⁷ HYPOG-01 [ClinicalTrials.gov identifier: [NCT03127995](#)], FAST FORWARD nodal substudy,⁹² HYPOR-Adjuvant study¹⁵⁹), PMRT after neoadjuvant chemotherapy (NSABP B-51 [ClinicalTrials.gov identifier: [NCT01872975](#)]),⁵² particle therapy (RADCOMP [ClinicalTrials.gov identifier: [NCT02603341](#)], PARABLE (United Kingdom),¹³⁰ Danish Breast Proton Trial [ClinicalTrials.gov identifier: [NCT04291378](#)]),¹⁶⁰ and the role of axillary surgery (Alliance A011202 [ClinicalTrials.gov identifier: [NCT01901094](#)]) that will impact clinical decision making and future clinical practice.

ASCO believes that cancer clinical trials are vital to inform medical decisions and improve cancer care, and that all patients should have the opportunity to participate.


ADDITIONAL RESOURCES

For current information, including selected updates, supplements, and clinical tools and resources, visit www.asco.org/breast-cancer-guidelines. The Data Supplement for this guideline includes a list of abbreviations used in the guideline, the PICOTS and literature search strategies, a list of guideline peer reviewers, and data supporting the recommendations. Guideline recommendations are also available in the free ASCO Guidelines app (available for download in the [Apple App Store](#) and [Google Play Store](#)). Listen to key recommendations and insights from panel members on the [ASCO Guidelines podcast](#). The Methodology Manual (available at www.asco.org/guideline-methodology) provides additional information about the methods used to develop this guideline. Patient information is available at www.cancer.org.

The task force welcomes your comments on this guideline, including implementation challenges, new evidence, and how this guideline impacts you. To provide feedback, contact us at guidelines@asco.org. Comments may be incorporated into a future guideline update. To submit new evidence or suggest a topic for guideline development, complete the form available at www.asco.org/guidelines.

INCLUSIVE LANGUAGE

ASCO is committed to promoting the health and well-being of all patients. ASCO guidelines are intended to apply to, and be discussed clearly and compassionately with, all patients. For this reason, guideline authors use appropriately inclusive language. In instances in which the guideline draws upon data based on research in a specified population (eg, studies regarding women with ovarian cancer), the guideline authors describe the characteristics and results of the research as reported.



PRISMA 2020 study selection diagram. PRISMA, Preferred Reporting Items for Systematic Reviews and Meta-Analyses.

AFFILIATIONS

¹Department of Radiation Oncology, Massachusetts General Hospital, Boston, MA

²Department of Medical Oncology, University of North Carolina, Chapel Hill, NC

³Department of Radiation Oncology, Fox Chase Cancer Center, Philadelphia, PA

⁴Department of Radiation Oncology, UPMC Hillman Cancer Center, University of Pittsburgh, Pittsburgh, PA

⁵American Society for Radiation Oncology, Arlington, VA

⁶Department of Radiation Oncology, University of Florida, Jacksonville, FL

⁷Patient Representative, Surviving Breast Cancer, Gilbert, AZ

⁸Department of Radiation Oncology, Memorial Sloan Kettering Cancer Center, New York, NY

⁹Department of Surgery, Huntsman Cancer Institute at the University of Utah, Salt Lake City, UT

¹⁰Department of Radiation Oncology, University of Colorado Denver, Aurora, CO

¹¹Department of Surgery, Loyola University Medical Center, Chicago, IL

¹²Radiation Oncology Branch, National Cancer Institute, Bethesda, MD

¹³Department of Medicine, Yale School of Medicine, New Haven, CT

¹⁴Department of Radiation Physics, University of Texas MD Anderson Cancer Center, Houston, TX

¹⁵Department of Radiation Oncology, David Geffen School of Medicine, University of California Los Angeles, Los Angeles, CA

¹⁶Department of Radiation Oncology, Moffitt Cancer Center, Tampa, FL

¹⁷Department of Radiation Oncology, University of North Carolina, Chapel Hill, NC

¹⁸Department of Radiation Oncology, Florida International University and Lynn Cancer Institute, Boca Raton, FL

¹⁹Department of Radiation Oncology, Indiana University School of Medicine, Indianapolis, IN

²⁰Department of Radiation Oncology, Stanford University, Stanford, CA

CORRESPONDING AUTHOR

American Society of Clinical Oncology; e-mail: guidelines@asco.org.

EDITOR'S NOTE

This ASCO Clinical Practice Guideline provides recommendations, with comprehensive review and analyses of the relevant literature for each recommendation. Additional information, including a supplement with additional evidence tables, clinical tools and resources, and links to patient information at www.cancer.org, is available at www.asco.org/breast-cancer-guidelines. These evidence-based recommendations guide clinical practice on the use of PMRT in patients with breast cancer.

EQUAL CONTRIBUTION

R.B.J. and K.C.H. were Expert Panel vice chair and chair, respectively.

SUPPORT

Supported by the American Society for Radiation Oncology.

AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

Disclosures provided by the authors are available with this article at DOI <https://doi.org/10.1200/JCO-25-01747>.

AUTHOR CONTRIBUTIONS

Conception and design: Rachel B. Jimenez, Yara Abdou, Penny Anderson, Parul Barry, Lisa Bradfield, Atif Khan, Chantal Reyna, Kilian E. Salerno, Youssef H. Zeidan, Richard C. Zellars, Kathleen C. Horst

Administrative support: Lisa Bradfield

Provision of study materials or patients: Youssef H. Zeidan

Collection and assembly of data: Rachel B. Jimenez, Yara Abdou, Penny Anderson, Parul Barry, Lisa Bradfield, Julie A. Bradley, Cindy Matsen, Kilian E. Salerno, Sarah E. Schellhorn, Kekoa Tapparra, Youssef H. Zeidan, Richard C. Zellars, Kathleen C. Horst

Data analysis and interpretation: Rachel B. Jimenez, Yara Abdou, Penny Anderson, Parul Barry, Julie A. Bradley, Lourdes D. Heras, Atif Khan, Cindy Matsen, Rachel Rabinovitch, Chantal Reyna, Kilian E. Salerno, Sarah E. Schellhorn, Deborah Schofield, Kekoa Tapparra, Iman

Washington, Jean L. Wright, Youssef H. Zeidan, Richard C. Zellars, Kathleen C. Horst

Manuscript writing: All authors

Final approval of manuscript: All authors

Accountable for all aspects of the work: All authors

ACKNOWLEDGMENT

ASCO wishes to thank Tiffinay Traina, MD, Rachel Yung, MD, and the Evidence Based Medicine Committee for their thoughtful reviews and insightful comments on this guideline. The task force thanks the National Institutes of Health (NIH) Library biomedical librarians, Tracy C. Shields, MSIS, and Alicia A. Livinski, MPH, MA, for conducting the literature searches for this guideline; and Kekoa Tapparra, MD, PhD, MPH (lead resident), Claire Baniel, MD, Katie Hwang, MD, Osagie Igiebor, MD, and Kaidi Wang, MD, for literature review assistance. The task force also thanks the peer reviewers for their comments and time spent reviewing the guideline. See Data Supplement for their names and disclosures. The authors also thank task force participant Angel Montero, MD, PhD, for serving as ESTRO's representative but who withdrew from authorship.

REFERENCES

1. Ferlay JEM, Lam F, Laversanne M, et al: Global Cancer Observatory: Cancer Today. Lyon, France, International Agency for Research on Cancer, 2024
2. EBCTCG Early Breast Cancer Trialists' Collaborative Group, McGale P, Taylor C, et al: Effect of radiotherapy after mastectomy and axillary surgery on 10-year recurrence and 20-year breast cancer mortality: meta-analysis of individual patient data for 8135 women in 22 randomised trials. *Lancet* 383:2127-2135, 2014
3. Recht A, Comen EA, Fine RE, et al: Postmastectomy radiotherapy: An American Society of Clinical Oncology, American Society for Radiation Oncology, and Society of Surgical Oncology focused guideline update. *Pract Radiat Oncol* 6:e219-e234, 2016
4. Institute of Medicine (US) Committee on Standards for Developing Trustworthy Clinical Practice Guidelines, Graham RMM, Miller WD, Steinberg E (eds): *Clinical Practice Guidelines We Can Trust*. Washington, DC, National Academies Press, 2011
5. Institute of Medicine (US) Committee on Standards for Systematic Reviews of Comparative Effectiveness, Eden JLL, Berg A, Morton S (eds): *Finding What Works In Health Care: Standards for Systematic Reviews*. Washington, DC, National Academies Press, 2011
6. Page MJ, McKenzie JE, Bossuyt PM, et al: The PRISMA 2020 statement: An updated guideline for reporting systematic reviews. *Rev Esp Cardiol* 74:790-799, 2021
7. Page MJ, Moher D, Bossuyt PM, et al: PRISMA 2020 explanation and elaboration: Updated guidance and exemplars for reporting systematic reviews. *BMJ* 372:n160, 2021
8. Overgaard M, Nielsen HM, Tramm T, et al: Postmastectomy radiotherapy in high-risk breast cancer patients given adjuvant systemic therapy. A 30-year long-term report from the Danish Breast Cancer Cooperative Group DBCG 82bc trial. *Radiother Oncol* 170:4-13, 2022
9. Danish Breast Cancer Cooperative Group, Nielsen HM, Overgaard M, et al: Study of failure pattern among high-risk breast cancer patients with or without postmastectomy radiotherapy in addition to adjuvant systemic therapy: Long-term results from the Danish Breast Cancer Cooperative Group DBCG 82 b and c randomized studies. *J Clin Oncol* 24:2268-2275, 2006
10. Nielsen HM, Overgaard M, Grau C, et al: Loco-regional recurrence after mastectomy in high-risk breast cancer—risk and prognosis. An analysis of patients from the DBCG 82 b&c randomization trials. *Radiother Oncol* 79:147-155, 2006
11. Early Breast Cancer Trialists' Collaborative Group EBCTCG: Radiotherapy to regional nodes in early breast cancer: An individual patient data meta-analysis of 14 324 women in 16 trials. *Lancet* 402:1991-2003, 2023
12. Ragaz J, Olivetto IA, Spinelli JJ, et al: Locoregional radiation therapy in patients with high-risk breast cancer receiving adjuvant chemotherapy: 20-year results of the British Columbia randomized trial. *J Natl Cancer Inst* 97:116-126, 2005
13. Wang J, Shi M, Ling R, et al: Adjuvant chemotherapy and radiotherapy in triple-negative breast carcinoma: A prospective randomized controlled multi-center trial. *Radiother Oncol* 100:200-204, 2011
14. Poortmans PM, Weltens C, Fortpied C, et al: Internal mammary and medial supraclavicular lymph node chain irradiation in stage I-III breast cancer (EORTC 22922/10925): 15-year results of a randomised, phase 3 trial. *Lancet Oncol* 21:1602-1610, 2020
15. Li Y, Moran MS, Huo Q, et al: Post-mastectomy radiotherapy for breast cancer patients with t1-t2 and 1-3 positive lymph nodes: A meta-analysis. *PLoS One* 8:e81765, 2013
16. Sittenfeld SMC, Zabor EC, Hamilton SN, et al: A multi-institutional prediction model to estimate the risk of recurrence and mortality after mastectomy for T1-2N1 breast cancer. *Cancer* 128:3057-3066, 2022
17. Karlsson P, Cole BF, Chua BH, et al: Patterns and risk factors for locoregional failures after mastectomy for breast cancer: An International Breast Cancer Study Group report. *Ann Oncol* 23:2852-2858, 2012
18. Clarke M, Collins R, Darby S, et al: Effects of radiotherapy and of differences in the extent of surgery for early breast cancer on local recurrence and 15-year survival: An overview of the randomised trials. *Lancet* 366:2087-2106, 2005
19. Kunkler I: Does postmastectomy radiotherapy in 'intermediate-risk' breast cancer impact overall survival? 10-year results of the BIG 2-04 MRC SUPREMO randomised trial. GS2-03; SESS-3537. Presented at the San Antonio Breast Cancer Symposium. San Antonio, TX, 2024
20. Lim SZ, Kusumawidjaja G, Mohd Ishak HM, et al: Outcomes of stage I and II breast cancer with nodal micrometastases treated with mastectomy without axillary therapy. *Breast Cancer Res Treat* 189:837-843, 2021
21. Kyndi M, Overgaard M, Nielsen HM, et al: High local recurrence risk is not associated with large survival reduction after postmastectomy radiotherapy in high-risk breast cancer: A subgroup analysis of DBCG 82 b&c. *Radiother Oncol* 90:74-79, 2009
22. Rao X, Wang X, Jin K, et al: Outcomes with and without postmastectomy radiotherapy for pT3N0-1M0 breast cancer: An institutional experience. *Cancer Med* 13:e6927, 2024
23. Zhao R, He W, Guan X: Benefits of post-mastectomy radiation for T4N0M0 breast cancer patients: A SEER database study. *Cancer Treat Res Commun* 32:100586, 2022
24. Overgaard M, Hansen PS, Overgaard J, et al: Postoperative radiotherapy in high-risk premenopausal women with breast cancer who receive adjuvant chemotherapy. Danish Breast Cancer Cooperative Group 82b Trial. *N Engl J Med* 337:949-955, 1997
25. Overgaard M, Jensen MB, Overgaard J, et al: Postoperative radiotherapy in high-risk postmenopausal breast-cancer patients given adjuvant tamoxifen: Danish Breast Cancer Cooperative Group DBCG 82c randomised trial. *Lancet* 353:1641-1648, 1999
26. Taghian AG, Jeong JH, Mamounas EP, et al: Low locoregional recurrence rate among node-negative breast cancer patients with tumors 5 cm or larger treated by mastectomy, with or without adjuvant systemic therapy and without radiotherapy: Results from five national surgical adjuvant breast and bowel project randomized clinical trials. *J Clin Oncol* 24:3927-3932, 2006
27. McCammon R, Finlayson C, Schwer A, et al: Impact of postmastectomy radiotherapy in T3N0 invasive carcinoma of the breast: A Surveillance, Epidemiology, and End Results database analysis. *Cancer* 113:683-689, 2008
28. Yan W, Christos P, Nori D, et al: Is there a cause-specific survival benefit of postmastectomy radiation therapy in women younger than age 50 with T3N0 invasive breast cancer? A SEER database analysis: Outcomes by receptor status/race/age: Analysis using the NCI Surveillance, Epidemiology, and End Results (SEER) database. *Am J Clin Oncol* 36:552-557, 2013

29. Taghian A, Jeong JH, Mamounas E, et al: Patterns of locoregional failure in patients with operable breast cancer treated by mastectomy and adjuvant chemotherapy with or without tamoxifen and without radiotherapy: Results from five National Surgical Adjuvant Breast and Bowel Project randomized clinical trials. *J Clin Oncol* 22:4247-4254, 2004
30. Peng G, Zhou Z, Jiang M, et al: Can a subgroup at high risk for LRR be identified from T1-2 breast cancer with negative lymph nodes after mastectomy? A meta-analysis. *Biosci Rep* 39: BSR20181853, 2019
31. Bundred J, Michael S, Bowers S, et al: Do surgical margins matter after mastectomy? A systematic review. *Eur J Surg Oncol* 46:P2185-P2194, 2020
32. Garg AK, Oh JL, Oswald MJ, et al: Effect of postmastectomy radiotherapy in patients <35 years old with stage II-III breast cancer treated with doxorubicin-based neoadjuvant chemotherapy and mastectomy. *Int J Radiat Oncol Biol Phys* 69:1478-1483, 2007
33. McGuire SE, Gonzalez-Angulo AM, Huang EH, et al: Postmastectomy radiation improves the outcome of patients with locally advanced breast cancer who achieve a pathologic complete response to neoadjuvant chemotherapy. *Int J Radiat Oncol Biol Phys* 68:1004-1009, 2007
34. Krug D, Lederer B, Seither F, et al: Post-mastectomy radiotherapy after neoadjuvant chemotherapy in breast cancer: A pooled retrospective analysis of three prospective randomized trials. *Ann Surg Oncol* 26:3892-3901, 2019
35. Wang K, Jin X, Wang W, et al: The role of postmastectomy radiation in patients with ypN0 breast cancer after neoadjuvant chemotherapy: A meta-analysis. *BMC Cancer* 21:728, 2021
36. Murchison S, Nichol A, Speers C, et al: Locoregional recurrence and survival outcomes in breast cancer treated with modern neoadjuvant chemotherapy: A contemporary population-based analysis. *Clin Breast Cancer* 22:e773-e787, 2022
37. Arsenault D, Hurley J, Takita C, et al: Predictors of locoregional outcome in HER2-overexpressing breast cancer treated with neoadjuvant chemotherapy. *Am J Clin Oncol* 38:348-352, 2015
38. Mamounas EP, Anderson SJ, Dignam JJ, et al: Predictors of locoregional recurrence after neoadjuvant chemotherapy: Results from combined analysis of national surgical adjuvant breast and bowel project B-18 and B-27. *J Clin Oncol* 30:3960-3966, 2012
39. Safi O, Bachir B, Panoff J, et al: Post-mastectomy radiation therapy in HER-2 positive breast cancer after primary systemic therapy: Pooled analysis of TRYPHAENA and NeoSphere trials. *Radiother Oncol* 184:109668, 2023
40. Huang Z, Zhu L, Huang XB, et al: Postmastectomy radiation therapy based on pathologic nodal status in clinical node-positive stage II to III breast cancer treated with neoadjuvant chemotherapy. *Int J Radiat Oncol Biol Phys* 108:1030-1039, 2020
41. Haffty BG, McCall LM, Ballman KV, et al: Impact of radiation on locoregional control in women with node-positive breast cancer treated with neoadjuvant chemotherapy and axillary lymph node dissection: Results from ACOSOG Z1071 clinical trial. *Int J Radiat Oncol Biol Phys* 105:174-182, 2019
42. Schmid P, Cortes J, Pusztai L, et al: Pembrolizumab for early triple-negative breast cancer. *N Engl J Med* 382:810-821, 2020
43. Shim SJ, Park W, Huh SJ, et al: The role of postmastectomy radiation therapy after neoadjuvant chemotherapy in clinical stage II-III breast cancer patients with pN0: A multicenter, retrospective study (KROG 12-05). *Int J Radiat Oncol Biol Phys* 88:65-72, 2014
44. Le Scodan R, Selz J, Stevens D, et al: Radiotherapy for stage II and stage III breast cancer patients with negative lymph nodes after preoperative chemotherapy and mastectomy. *Int J Radiat Oncol Biol Phys* 82:e1-e7, 2012
45. Kim D, Kim JH, Kim IA, et al: Impact of postmastectomy radiation therapy on breast cancer patients according to pathologic nodal status after modern neoadjuvant chemotherapy. *Cancer Res Treat* 55:592-602, 2023
46. Nagar H, Mittendorf EA, Strom EA, et al: Local-regional recurrence with and without radiation therapy after neoadjuvant chemotherapy and mastectomy for clinically staged T3N0 breast cancer. *Int J Radiat Oncol Biol Phys* 81:782-787, 2011
47. Crown A, Gonen M, Le T, et al: Does failure to achieve pathologic complete response with neoadjuvant chemotherapy identify node-negative patients who would benefit from postmastectomy radiation or regional nodal irradiation? *Ann Surg Oncol* 28:1328-1335, 2021
48. Nagar H, Boothe D, Ginter PS, et al: Disease-free survival according to the use of postmastectomy radiation therapy after neoadjuvant chemotherapy. *Clin Breast Cancer* 15:128-134, 2015
49. Mailhot Vega RB, Wang S, Brooks ED, et al: Evaluating regional nodal irradiation allocation and association with oncologic outcomes in NSABP B-18, B-27, B-40, and B-41. *Int J Radiat Oncol Biol Phys* 113:542-551, 2022
50. Haffty BG, McCall LM, Ballman KV, et al: Patterns of local-regional management following neoadjuvant chemotherapy in breast cancer: Results from ACOSOG Z1071 (Alliance). *Int J Radiat Oncol Biol Phys* 94:493-502, 2016
51. Gianni L, Pienkowski T, Im YH, et al: Efficacy and safety of neoadjuvant pertuzumab and trastuzumab in women with locally advanced, inflammatory, or early HER2-positive breast cancer (NeoSphere): A randomised multicentre, open-label, phase 2 trial. *Lancet Oncol* 13:25-32, 2012
52. Mamounas E, Bandos H, White J, et al: Abstract GS02-07: Loco-regional irradiation in patients with biopsy-proven axillary node involvement at presentation who become pathologically node-negative after neoadjuvant chemotherapy: Primary outcomes of NRG oncology/NSABP B-51/RTOG 1304. *Cancer Res* 84:GS02-07, 2024 (suppl 9)
53. de Wild SR, de Munck L, Simons JM, et al: De-escalation of radiotherapy after primary chemotherapy in cT1-2N1 breast cancer (RAPCHEM; BOOG 2010-03): 5-year follow-up results of a Dutch, prospective, registry study. *Lancet Oncol* 23:1201-1210, 2022
54. Meattini I, Cecchini S, Di Cataldo V, et al: Postmastectomy radiotherapy for locally advanced breast cancer receiving neoadjuvant chemotherapy. *Biomed Res Int* 2014:719175, 2014
55. Gillon P, Touati N, Breton-Callu C, et al: Factors predictive of locoregional recurrence following neoadjuvant chemotherapy in patients with large operable or locally advanced breast cancer: An analysis of the EORTC 10994/BIG 1-00 study. *Eur J Cancer* 79:226-234, 2017
56. Yang TJ, Morrow M, Modi S, et al: The effect of molecular subtype and residual disease on locoregional recurrence in breast cancer patients treated with neoadjuvant chemotherapy and postmastectomy radiation. *Ann Surg Oncol* 22:S495-S501, 2015 (suppl 3)
57. Werutsky G, Untch M, Hanusch C, et al: Locoregional recurrence risk after neoadjuvant chemotherapy: A pooled analysis of nine prospective neoadjuvant breast cancer trials. *Eur J Cancer* 130: 92-101, 2020
58. Masuda N, Lee SJ, Ohtani S, et al: Adjuvant capecitabine for breast cancer after preoperative chemotherapy. *N Engl J Med* 376:2147-2159, 2017
59. von Minckwitz G, Huang CS, Mano MS, et al: Trastuzumab emtansine for residual invasive HER2-positive breast cancer. *N Engl J Med* 380:617-628, 2019
60. Schmid P, Cortes J, Dent R, et al: Overall survival with pembrolizumab in early-stage triple negative breast cancer. *N Engl J Med* 391:1981-1991, 2024
61. Poortmans PM, Collette S, Kirkove C, et al: Internal mammary and medial supraclavicular irradiation in breast cancer. *N Engl J Med* 373:317-327, 2015
62. Kim YB, Byun HK, Kim DY, et al: Effect of elective internal mammary node irradiation on disease-free survival in women with node-positive breast cancer: A randomized phase 3 clinical trial. *JAMA Oncol* 8:96-105, 2022
63. Thorsen LB, Offersen BV, Danø H, et al: DBCG-IMN: A population-based cohort study on the effect of internal mammary node irradiation in early node-positive breast cancer. *J Clin Oncol* 34: 314-320, 2016
64. Hennequin C, Bossard N, Servagi-Vernat S, et al: Ten-year survival results of a randomized trial of irradiation of internal mammary nodes after mastectomy. *Int J Radiat Oncol Biol Phys* 86: 860-866, 2013
65. Wong JS, Uno H, Tramontano AC, et al: Hypofractionated vs conventionally fractionated postmastectomy radiation after implant-based reconstruction: A randomized clinical trial. *JAMA Oncol* 10:1370-1378, 2024
66. Kim N, Chang JS, Shah C, et al: Hypofractionated volumetric-modulated arc therapy for breast cancer: A propensity-score-weighted comparison of radiation-related toxicity. *Int J Cancer* 149: 149-157, 2021
67. Chitapanarux I, Klunklin P, Pinitpatcharalert A, et al: Conventional versus hypofractionated postmastectomy radiotherapy: A report on long-term outcomes and late toxicity. *Radiat Oncol* 14:175, 2019
68. START Trialists' Group, Bentzen SM, Agrawal RK, et al: The UK standardisation of breast radiotherapy (START) trial A of radiotherapy hypofractionation for treatment of early breast cancer: A randomised trial. *Lancet Oncol* 9:331-341, 2008
69. Haviland JS, Mannino M, Griffin C, et al: Late normal tissue effects in the arm and shoulder following lymphatic radiotherapy: Results from the UK START (Standardisation of Breast Radiotherapy) trials. *Radiother Oncol* 126:155-162, 2018
70. Wang SL, Fang H, Song YW, et al: Hypofractionated versus conventional fractionated postmastectomy radiotherapy for patients with high-risk breast cancer: A randomised, non-inferiority, open-label, phase 3 trial. *Lancet Oncol* 20:352-360, 2019
71. Maiti S, Meyur S, Mandal BC, et al: Comparison of conventional and hypofractionated radiation after mastectomy in locally advanced breast cancer: A prospective randomised study on dosimetric evaluation and treatment outcome. *J Radiother Pract* 20:30-38, 2021
72. Zhao XB, Ren GS: Analysis of radiotherapy optimization regimens after modified radical mastectomy. *Eur Rev Med Pharmacol Sci* 20:4705-4709, 2016
73. Zhang HJ, Qu BL, Meng LL, et al: Clinical observation of short-term efficacies of different hypofractionated radiation therapies after modified radical mastectomy for breast cancer. *Eur J Gynaecol Oncol* 39:524-532, 2018
74. Kim DY, Park E, Heo CY, et al: Influence of hypofractionated versus conventional fractionated postmastectomy radiation therapy in breast cancer patients with reconstruction. *Int J Radiat Oncol Biol Phys* 112:445-456, 2022

75. Matzinger O, Heimsoth I, Poortmans P, et al: Toxicity at three years with and without irradiation of the internal mammary and medial supraclavicular lymph node chain in stage I to III breast cancer (EORTC trial 22922/10925). *Acta Oncol* 49:24-34, 2010
76. Whelan TJ, Olivetto IA, Parulekar WR, et al: Regional nodal irradiation in early-stage breast cancer. *N Engl J Med* 373:307-316, 2015
77. Abdel-Rahman O: Impact of regional nodal irradiation on the outcomes of N1 breast cancer patients referred for adjuvant treatment: A patient-level pooled analysis of 2 clinical trials. *Clin Breast Cancer* 18:504-510, 2018
78. Panoff JE, Takita C, Hurley J, et al: Higher chest wall dose results in improved locoregional outcome in patients receiving postmastectomy radiation. *Int J Radiat Oncol Biol Phys* 82:1192-1199, 2012
79. Marhin W, Wai E, Tyldesley S: Impact of fraction size on cardiac mortality in women treated with tangential radiotherapy for localized breast cancer. *Int J Radiat Oncol Biol Phys* 69:483-489, 2007
80. Albert A, Mangana S, Nittala MR, et al: The impact of a postmastectomy chest wall scar boost on local recurrence-free survival in high-risk patients. *Clin Breast Cancer* 19:363-369, 2019
81. Bartels SAL, Donker M, Poncet C, et al: Radiotherapy or surgery of the axilla after a positive sentinel node in breast cancer: 10-Year results of the randomized controlled EORTC 10981-22023 AMAROS trial. *J Clin Oncol* 41:2159-2165, 2023
82. Thorsen LBJ, Overgaard J, Matthiessen LW, et al: Internal mammary node irradiation in patients with node-positive early breast cancer: Fifteen-year results from the Danish breast cancer group internal mammary node study. *J Clin Oncol* 40:4198-4206, 2022
83. Chung SY, Chang JS, Shin KH, et al: Impact of radiation dose on complications among women with breast cancer who underwent breast reconstruction and post-mastectomy radiotherapy: A multi-institutional validation study. *Breast* 56:7-13, 2021
84. Tovnanubra C, Katanyoo K, Uber P, et al: Comparison of treatment outcome between hypofractionated radiotherapy and conventional radiotherapy in postmastectomy breast cancer. *Asian Pac J Cancer Prev* 21:119-125, 2020
85. Stokes EL, Tyldesley S, Woods R, et al: Effect of nodal irradiation and fraction size on cardiac and cerebrovascular mortality in women with breast cancer treated with local and locoregional radiotherapy. *Int J Radiat Oncol Biol Phys* 80:403-409, 2011
86. Wong JS, Uno H, Tramountano A, et al: Patient-reported and toxicity results from the FABREC study: A multicenter randomized trial of hypofractionated vs. conventionally-fractionated postmastectomy radiation therapy after implant-based reconstruction. *IJROBP* 117:e3-e4, 2023
87. Poppe MM, Le-Rademacher J, Haffty BG, et al: A randomized trial of hypofractionated post-mastectomy radiation therapy (PMRT) in women with breast reconstruction (RT CHARM, Alliance A221505). *Int J Radiat Oncol Biol Phys* 120:S11, 2024
88. Diao K, Lei X, He W, et al: Racial and ethnic differences in long-term adverse radiation therapy effects among breast cancer survivors. *Int J Radiat Oncol Biol Phys* 118:626-631, 2024
89. Wright JL, Takita C, Reis IM, et al: Racial variations in radiation-induced skin toxicity severity: Data from a prospective cohort receiving postmastectomy radiation. *Int J Radiat Oncol Biol Phys* 90:335-343, 2014
90. Fang P, He W, Gomez D, et al: Racial disparities in guideline-concordant cancer care and mortality in the United States. *Adv Radiat Oncol* 3:221-229, 2018
91. Ohri N, Moshier E, Ho A, et al: Postmastectomy radiation in breast cancer patients with pathologically positive lymph nodes after neoadjuvant chemotherapy: Usage rates and survival trends. *Int J Radiat Oncol Biol Phys* 99:549-559, 2017
92. Murray Brunt A, Haviland JS, Wheatley DA, et al: Hypofractionated breast radiotherapy for 1 week versus 3 weeks (FAST-Forward): 5-year efficacy and late normal tissue effects results from a multicentre, non-inferiority, randomised, phase 3 trial. *Lancet* 395:1613-1626, 2020
93. Chakraborty S, Chatterjee S, Hypoort Adjuvant Author Group: HYPORADT adjuvant acute toxicity and patient dosimetry quality assurance results - Interim analysis. *Radiother Oncol* 174:59-68, 2022
94. ClinicalTrials.gov: Hypofractionated LocoRegional Radiotherapy in Breast Cancer (RHEAL). <https://clinicaltrials.gov/study/NCT04228991>
95. Liao Z, Strom EA, Buzdar AU, et al: Locoregional irradiation for inflammatory breast cancer: Effectiveness of dose escalation in decreasing recurrence. *Int J Radiat Oncol Biol Phys* 47:1191-1200, 2000
96. Bristol LJ, Woodward WA, Strom EA, et al: Locoregional treatment outcomes after multimodality management of inflammatory breast cancer. *Int J Radiat Oncol Biol Phys* 72:474-484, 2008
97. Purswani JM, Oh C, Teruel JR, et al: Definitive radiation with nodal boost for patients with locally advanced breast cancer. *Pract Radiat Oncol* 13:e103-e114, 2023
98. Klusen ST, Peiler A, Schmidt GP, et al: Simultaneous integrated boost within the lymphatic drainage system in breast cancer: A single center study on toxicity and oncologic outcome. *Front Oncol* 13:989466, 2023
99. Mayadev J, Einck J, Elson S, et al: Practice patterns in the delivery of radiation therapy after mastectomy among the University of California Athena Breast Health Network. *Clin Breast Cancer* 15:43-47, 2015
100. Hess C, Lee A, Fish K, et al: Socioeconomic and racial disparities in the selection of chest wall boost radiation therapy in Californian women after mastectomy. *Clin Breast Cancer* 15:212-218, 2015
101. Sachdev S, Goodman CR, Neuschler E, et al: Radiotherapy of MRI-detected involved internal mammary lymph nodes in breast cancer. *Radiat Oncol* 12:199, 2017
102. Diao K, Andring LM, Barcenas CH, et al: Contemporary outcomes after multimodality therapy in patients with breast cancer presenting with ipsilateral supraclavicular node involvement. *Int J Radiat Oncol Biol Phys* 112:66-74, 2022
103. Wang SL, Li YX, Song YW, et al: Postmastectomy chest wall radiotherapy with single low-energy electron beam: An assessment of outcome and prognostic factors. *Pract Radiat Oncol* 2:106-113, 2012
104. Osei E, Dang S, Darko J, et al: Dosimetric evaluation of 3 and/or 4 field radiation therapy of breast cancers: Clinical experience. *J Radiother Pract* 20:380-394, 2021
105. Grellier Adedjouma N, Chevrier M, Fourquet A, et al: Long-term results of a highly performing conformal electron therapy technique for chest wall irradiation after mastectomy. *Int J Radiat Oncol Biol Phys* 98:206-214, 2017
106. Van Parijs H, Miedema G, Vinh-Hung V, et al: Short course radiotherapy with simultaneous integrated boost for stage I-II breast cancer, early toxicities of a randomized clinical trial. *Radiat Oncol* 7:80, 2012
107. Rastogi K, Sharma S, Gupta S, et al: Dosimetric comparison of IMRT versus 3DCRT for post-mastectomy chest wall irradiation. *Radiat Oncol* J 36:71-78, 2018
108. Li W, Wang J, Cheng H, et al: IMRT versus 3D-CRT for post-mastectomy irradiation of chest wall and regional nodes: A population-based comparison of normal lung dose and radiation pneumonitis. *Int J Clin Exp Med* 9:22331-22337, 2016
109. Jaggi R, Griffith KA, Moran JM, et al: A randomized comparison of radiation therapy techniques in the management of node-positive breast cancer: Primary outcomes analysis. *Int J Radiat Oncol Biol Phys* 101:1149-1158, 2018
110. Mehta A, Kumar P, Silambarasan NS, et al: Comparison of dosimetric parameters of three-dimensional conformal radiotherapy and intensity-modulated radiotherapy in breast cancer patients undergoing adjuvant radiotherapy after modified radical mastectomy. *Asian J Oncol* 08:1-7, 2022
111. Rice L, Goldsmith C, Green MML, et al: An effective deep-inspiration breath-hold radiotherapy technique for left-breast cancer: Impact of post-mastectomy treatment, nodal coverage, and dose schedule on organs at risk. *Breast Cancer (Dove Med Press)* 9:437-446, 2017
112. Gaál S, Kahan Z, Paczona V, et al: Deep-inspirational breath-hold (DIBH) technique in left-sided breast cancer: Various aspects of clinical utility. *Radiat Oncol* 16:89, 2021
113. Laaksomaa M, Aula A, Sarudis S, et al: Surface-guided radiotherapy systems in locoregional deep inspiration breath hold radiotherapy for breast cancer—A multicenter study on the setup accuracy. *Rep Pract Oncol Radiother* 29:176-186, 2024
114. Sapienza LG, Maia MAC, Gomes MJL, et al: Randomized clinical trial of tissue equivalent bolus prescription in postmastectomy radiotherapy stratified by skin involvement status. *Clin Transl Radiat Oncol* 39:100570, 2023
115. Pignol JP, Vu TT, Mitera G, et al: Prospective evaluation of severe skin toxicity and pain during postmastectomy radiation therapy. *Int J Radiat Oncol Biol Phys* 91:157-164, 2015
116. Yap ML, Tieu M, Sappiatzer J, et al: Outcomes in patients treated with post-mastectomy chest wall radiotherapy without the routine use of bolus. *Clin Oncol (R Coll Radiol)* 30:427-432, 2018
117. Tieu MT, Graham P, Browne L, et al: The effect of adjuvant postmastectomy radiotherapy bolus technique on local recurrence. *Int J Radiat Oncol Biol Phys* 81:e165-e171, 2011
118. Nichol A, Narinesingh D, Raman S, et al: The effect of bolus on local control for patients treated with mastectomy and radiation therapy. *Int J Radiat Oncol Biol Phys* 110:1360-1369, 2021
119. de Sousa CFPM, Neto ES, Chen MJ, et al: Postmastectomy radiation therapy bolus associated complications in patients who underwent 2-stage breast reconstruction. *Adv Radiat Oncol* 7:101010, 2022
120. Thomsen MS, Berg M, Nielsen HM, et al: Post-mastectomy radiotherapy in Denmark: From 2D to 3D treatment planning guidelines of the Danish Breast Cancer Cooperative Group. *Acta Oncol* 47:654-661, 2008
121. Bekelman JE, Lu H, Pugh S, et al: Pragmatic randomised clinical trial of proton versus photon therapy for patients with non-metastatic breast cancer: The Radiotherapy Comparative Effectiveness (RadComp) Consortium trial protocol. *BMJ Open* 9:e025556, 2019
122. Nielsen MH, Berg M, Pedersen AN, et al: Delineation of target volumes and organs at risk in adjuvant radiotherapy of early breast cancer: National guidelines and contouring atlas by the Danish breast cancer cooperative group. *Acta Oncol* 52:703-710, 2013
123. Kaidar-Person O, Vrou Offeren B, Hol S, et al: ESTRO ACROP consensus guideline for target volume delineation in the setting of postmastectomy radiation therapy after implant-based immediate reconstruction for early stage breast cancer. *Radiother Oncol* 137:159-166, 2019
124. Kagkiozis J, Platoni K, Kantzou I, et al: Review of the three-field techniques in breast cancer radiotherapy. *J BUON* 22:599-605, 2017

125. Pasquier D, Le Tinier F, Bennadji R, et al: Intensity-modulated radiation therapy with simultaneous integrated boost for locally advanced breast cancer: A prospective study on toxicity and quality of life. *Sci Rep* 9:2759, 2019
126. Bataille B, Raoudha B, Le Tinier F, et al: Prospective study of intensity-modulated radiation therapy for locally advanced breast cancer. *Cancers* 12:3852, 2020
127. Bazan JG, DiCostanzo D, Hock K, et al: Analysis of radiation dose to the shoulder by treatment technique and correlation with patient reported outcomes in patients receiving regional nodal irradiation. *Front Oncol* 11:617926, 2021
128. Lazzari G, Terlizzi A, Leo MG, et al: VMAT radiation-induced nausea and vomiting in adjuvant breast cancer radiotherapy: The incidental effect of low-dose bath exposure. *Clin Translational Radiat Oncol* 7:43-48, 2017
129. Yaney A, Ayan AS, Pan X, et al: Dosimetric parameters associated with radiation-induced esophagitis in breast cancer patients undergoing regional nodal irradiation. *Radiother Oncol* 155: 167-173, 2021
130. Kirby AM, Haviland JS, Mackenzie M, et al: Proton beam therapy in breast cancer patients: The UK PARABLE trial is recruiting. *Clin Oncol (R Coll Radiol)* 35:347-350, 2023
131. Honaryar MK, Locquet M, Allodji R, et al: Cancer therapy-related cardiac dysfunction after radiation therapy for breast cancer: Results from the BACCARAT cohort study. *Cardiooncology* 10:54, 2024
132. Kanyilmaz G, Aktan M, Koc M, et al: Radiation-induced hypothyroidism in patients with breast cancer: A retrospective analysis of 243 cases. *Med Dosim* 42:190-196, 2017
133. Zhao XR, Fang H, Jing H, et al: Radiation-induced hypothyroidism in patients with breast cancer after hypofractionated radiation therapy: A prospective cohort study. *Int J Radiat Oncol Biol Phys* 115:83-92, 2023
134. Cuzick J, Stewart H, Rutqvist L, et al: Cause-specific mortality in long-term survivors of breast cancer who participated in trials of radiotherapy. *J Clin Oncol* 12:447-453, 1994
135. Favourable and unfavourable effects on long-term survival of radiotherapy for early breast cancer: An overview of the randomised trials. Early breast cancer Trialists' collaborative group. *Lancet* 355:1757-1770, 2000
136. Darby SC, Ewertz M, McGale P, et al: Risk of ischemic heart disease in women after radiotherapy for breast cancer. *N Engl J Med* 368:987-998, 2013
137. Laugaard Lorenzen E, Christian Rehammar J, Jensen MB, et al: Radiation-induced risk of ischemic heart disease following breast cancer radiotherapy in Denmark, 1977-2005. *Radiother Oncol* 152:103-110, 2020
138. Carlson LE, Watt GP, Tonorez ES, et al: Coronary artery disease in young women after radiation therapy for breast cancer: The WECARE study. *JACC CardioOncol* 3:381-392, 2021
139. van den Bogaard VA, Ta BD, van der Schaaf A, et al: Validation and modification of a prediction model for acute cardiac events in patients with breast cancer treated with radiotherapy based on three-dimensional dose distributions to cardiac substructures. *J Clin Oncol* 35:1171-1178, 2017
140. Beaton L, Bergman A, Nichol A, et al: Cardiac death after breast radiotherapy and the QUANTEC cardiac guidelines. *Clin Transl Radiat Oncol* 19:39-45, 2019
141. Lee YC, Chuang JP, Hsieh PC, et al: A higher incidence rate of acute coronary syndrome following radiation therapy in patients with breast cancer and a history of coronary artery diseases. *Breast Cancer Res Treat* 152:429-435, 2015
142. Hooning MJ, Botma A, Aleman BM, et al: Long-term risk of cardiovascular disease in 10-year survivors of breast cancer. *J Natl Cancer Inst* 99:365-375, 2007
143. Aznar MC, Carrasco de Fez P, Corradini S, et al: ESTRO-ACROP guideline: Recommendations on implementation of breath-hold techniques in radiotherapy. *Radiother Oncol* 185:109734, 2023
144. Jacob S, Camilleri J, Derreumaux S, et al: Is mean heart dose a relevant surrogate parameter of left ventricle and coronary arteries exposure during breast cancer radiotherapy: A dosimetric evaluation based on individually-determined radiation dose (BACCARAT study). *Radiat Oncol* 14:29, 2019
145. Mutter RW, Choi JI, Jimenez RB, et al: Proton therapy for breast cancer: A consensus statement from the particle therapy cooperative group breast cancer subcommittee. *Int J Radiat Oncol Biol Phys* 111:337-359, 2021
146. Holt F, Probert J, Darby SC, et al: Proton beam therapy for early breast cancer: A systematic review and meta-analysis of clinical outcomes. *Int J Radiat Oncol Biol Phys* 117:869-882, 2023
147. MacDonald SM, Patel SA, Hickey S, et al: Proton therapy for breast cancer after mastectomy: Early outcomes of a prospective clinical trial. *Int J Radiat Oncol Biol Phys* 86:484-490, 2013
148. Jimenez RB, Hickey S, DePauw N, et al: Phase II study of proton beam radiation therapy for patients with breast cancer requiring regional nodal irradiation. *J Clin Oncol* 37:2778-2785, 2019
149. Hassan MZO, Awadalla M, Tan TC, et al: Serial measurement of global longitudinal strain among women with breast cancer treated with proton radiation therapy: A prospective trial for 70 patients. *Int J Radiat Oncol Biol Phys* 115:398-406, 2023
150. Jaffray D, Kupelian P, Djemil T, et al: Review of image-guided radiation therapy. *Expert Rev Anticancer Ther* 7:89-103, 2007
151. Hartford AC, Palisca MG, Eichler TJ, et al: American Society for Therapeutic Radiology and Oncology (ASTRO) and American College of Radiology (ACR) practice guidelines for intensity-modulated radiation therapy (IMRT). *Int J Radiat Oncol Biol Phys* 73:9-14, 2009
152. Jaffray DA, Langen KM, Mageras G, et al: Safety considerations for IGRT: Executive summary. *Pract Radiat Oncol* 3:167-170, 2013
153. Pazos M, Walter F, Reitz D, et al: Impact of surface-guided positioning on the use of portal imaging and initial set-up duration in breast cancer patients. *Strahlenther Onkol* 195:964-971, 2019
154. Reitz D, Carl G, Schönecker S, et al: Real-time intra-fraction motion management in breast cancer radiotherapy: Analysis of 2028 treatment sessions. *Radiat Oncol* 13:128, 2018
155. Freisleder P, Batista V, Öllers M, et al: ESTRO-ACROP guideline on surface guided radiation therapy. *Radiother Oncol* 173:188-196, 2022
156. Al-Hallaq HA, Cerviño L, Gutierrez AN, et al: AAPM task group report 302: Surface-guided radiotherapy. *Med Phys* 49:e82-e112, 2022
157. Blitzblau RC, Horton JK: Treatment planning technique in patients receiving postmastectomy radiation therapy. *Pract Radiat Oncol* 3:241-248, 2013
158. Dahn HM, Boersma LJ, de Ruyscher D, et al: The use of bolus in postmastectomy radiation therapy for breast cancer: A systematic review. *Crit Rev Oncol Hematol* 163:103391, 2021
159. Chatterjee S, Chakraborty S, HYPORT Adjuvant Author Group: Hypofractionated radiation therapy comparing a standard radiotherapy schedule (over 3 weeks) with a novel 1-week schedule in adjuvant breast cancer: An open-label randomized controlled study (HYPORT-Adjuvant)-study protocol for a multicentre, randomized phase III trial. *Trials* 21:819, 2020
160. Fuglsang Jensen M, Stick LB, Høyer M, et al: Proton therapy for early breast cancer patients in the DBCG proton trial: Planning, adaptation, and clinical experience from the first 43 patients. *Acta Oncol* 61:223-230, 2022

AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

Postmastectomy Radiation Therapy: An ASTRO-ASCO-SSO Clinical Practice Guideline

The following represents disclosure information provided by authors of this manuscript. All relationships are considered compensated unless otherwise noted. Relationships are self-held unless noted. I = Immediate Family Member, Inst = My Institution. Relationships may not relate to the subject matter of this manuscript. For more information about ASCO's conflict of interest policy, please refer to www.asco.org/rwc or ascopubs.org/jco/authors/author-center.

Open Payments is a public database containing information reported by companies about payments made to US-licensed physicians (Open Payments).

Rachel B. Jimenez

Consulting or Advisory Role: Alexion Pharmaceuticals

Yara Abdou

Honoraria: WebMD/Medscape, MJH Associates, Clinical Care Options, Cancer Expert Now

Consulting or Advisory Role: AstraZeneca, Pfizer

Research Funding: Natera (Inst), Haystack Oncology (Inst)

Uncompensated Relationships: Exact Sciences

Parul Barry

Employment: UPMC Hillman Cancer Center/University of Pittsburgh School of Medicine

Honoraria: Osler Institute, OncLive/MJH Life Sciences, Varian Medical Systems

Consulting or Advisory Role: El-Sevier Pathways Consultant

Research Funding: Breast Cancer Research Foundation (Inst), ACRO New Practitioner Grant, NCTN (Inst), Beckwith Institute (Inst), Pittsburgh Foundation (Inst), Silverlon (Inst), Philanthropy (Inst)

Travel, Accommodations, Expenses: ACRO, American Society for Radiation Oncology

Other Relationship: ACRO Board Member, ARS Board, WIMS Scientific Program Chair, Rare Tumors—Senior Associate Editor, ASTRO HEDI Chair

Julie A. Bradley

Patents, Royalties, Other Intellectual Property: 2021: Provisional patent for Breast Tissue Expander ReDesign 2023: U.S. non-provisional application for Breast Tissue Expander ReDesign

Uncompensated Relationships: PTCOG-NA Vice President

Lourdes D. Heras

Employment: Gila River Health Care

Uncompensated Relationships: Surviving Breast Cancer.org

Atif Khan

Employment: Memorial Sloan-Kettering Cancer Center

Stock and Other Ownership Interests: Novavax, Xtrava Health, IgniteData

Consulting or Advisory Role: IgniteData

Research Funding: Clovis Oncology, Merck KGaA, Varian Medical Systems

Patents, Royalties, Other Intellectual Property: Use Patent for the Drug Riluzole Awarded to Rutgers University With Me as Inventor

Uncompensated Relationships: AI MindSystems Foundation

Rachel Rabinovitch

Stock and Other Ownership Interests: Bristol Myers Squibb, Intuitive Surgical

Sarah E. Schellhorn

Stock and Other Ownership Interests: Gilead Sciences, Coronado Biosciences, Roche

Honoraria: Cardinal Health

Consulting or Advisory Role: Eisai, Puma Biotechnology, Celgene, Cardinal Health

Research Funding: Genentech, Pfizer

Travel, Accommodations, Expenses: Cardinal Health

Open Payments Link: <https://openpaymentsdata.cms.gov/physician/1076063>

Deborah Schofield

Uncompensated Relationships: American College of Radiology, American Association of Physicists in Medicine

Kekoa Tapparra

Employment: Stanford Health Care

Honoraria: MD Anderson Cancer Center, Stanford Health Care, American Association for Cancer Research, American Society of Hematology, Genentech

Research Funding: Stanford Cancer Institute, Conquer Cancer Foundation

Travel, Accommodations, Expenses: ASTRO-Breast Cancer Research Foundation

Jean L. Wright

Employment: Johns Hopkins Hospital, University of North Carolina at Chapel Hill

Honoraria: American Society for Radiation Oncology, Physicians' Education Resource

Research Funding: Oncospace, Prostate Cancer Foundation-Pfizer Health Equity Challenge (Inst), Prostate Cancer Foundation-Pfizer Health Equity Challenge (Inst)

Patents, Royalties, Other Intellectual Property: Spouse receives royalties from Up To Date (I)

Expert Testimony: Wicker Smith

Travel, Accommodations, Expenses: Physicians' Education Resource

Uncompensated Relationships: American Society for Radiation Oncology, International Journal of Radiation Oncology Biology Physics

Richard C. Zellars

Employment: Veracyte

Stock and Other Ownership Interests: Lilly, Bristol Myers Squibb Co

No other potential conflicts of interest were reported.

APPENDIX 1

ASCO GUIDELINE DISCLAIMER

The Clinical Practice Guidelines and other guidance published herein are provided by ASCO to assist clinicians in clinical decision making. The information herein should not be relied upon as being complete or accurate, nor should it be considered as inclusive of all proper treatments or methods of care or as a statement of the standard of care. With the rapid development of scientific knowledge, new evidence may emerge between the time information is developed and when it is published or read. The information is not continually updated and may not reflect the most recent evidence. The information addresses only the topics specifically identified therein and is not applicable to other interventions, diseases, or stages of diseases. This information does not mandate any particular course of medical care. Further, the information is not intended to substitute for the independent professional judgment of the treating clinician, as the information does not account for individual variation among patients. Recommendations specify the level of confidence that the recommendation reflects the net effect of a given course of action. The use of words like "must," "must not," "should," and "should not" indicates that a course of action is recommended or not recommended for either most or many patients, but there is latitude for the treating physician to select other courses of action in individual cases. In all cases, the selected course of action should be considered by the treating clinician in the context of treating the individual patient. Use of the information is voluntary. ASCO does not endorse third party drugs, devices, services, or therapies used to diagnose, treat, monitor, manage, or alleviate health conditions. Any use of a brand or trade name is for identification purposes only. ASCO provides this information on an "as is" basis and makes no warranty, express or implied, regarding the information. ASCO specifically disclaims any warranties of merchantability or fitness for a particular use or purpose. ASCO assumes no responsibility for any injury or damage to persons or property arising out of or related to any use of this information, or for any errors or omissions.

ASCO GUIDELINE AND CONFLICTS OF INTEREST POLICY

The Expert Panel was assembled in accordance with ASCO's Conflict of Interest Policy Implementation for Clinical Practice Guidelines ("Policy," found at

www.asco.org/guideline-methodology). All members of the Expert Panel completed ASCO's disclosure form, which requires disclosure of financial and other interests, including relationships with commercial entities that are reasonably likely to experience direct regulatory or commercial impact as a result of promulgation of the guideline. Categories for disclosure include employment; leadership; stock or other ownership; honoraria, consulting or advisory role; speaker's bureau; research funding; patents, royalties, other intellectual property; expert testimony; travel, accommodations, expenses; and other relationships. In accordance with the Policy, the majority of the members of the Expert Panel did not disclose any relationships constituting a conflict under the Policy.

ASTRO DISCLOSURE POLICY

As a leading organization in radiation oncology, the American Society for Radiation Oncology (ASTRO) is dedicated to improving quality of care and patient outcomes. A cornerstone of this goal is the development and dissemination of clinical practice guidelines based on systematic methods to evaluate and classify evidence, combined with a focus on patient-centric care and shared decision making. ASTRO develops and publishes guidelines without commercial support, and members volunteer their time.

ASTRO has detailed policies and procedures related to disclosure and management of industry relationships to avoid actual, potential, or perceived conflicts of interest. All task force members are required to disclose industry relationships and personal interests from 12 months before the initiation of the writing effort. Disclosures for the chair and vice chair go through a review process with final approval by ASTRO's Conflict of Interest Review Committee. For the purposes of full transparency, task force members' comprehensive disclosure information is included in this publication. Peer reviewer disclosures are also reviewed and included (Data Supplement). The complete disclosure policy for Formal Papers is online.

Selection of Task Force Members

ASTRO strives to avoid bias and is committed to creating a task force that includes a diverse and inclusive multidisciplinary group of experts considering race, ethnicity, gender, experience, practice setting, and geographic location. Representatives from organizations and professional societies with related interests and expertise are also invited to serve on the task force.

TABLE A1. ASTRO-ASCO-SSO Joint PMRT Task Force Membership

Name	Affiliation	Role or Area of Expertise
Rachel B. Jimenez, MD (vice chair)	Department of Radiation Oncology, Massachusetts General Hospital, Boston, MA	Radiation oncology
Kathleen C. Horst, MD (chair)	Department of Radiation Oncology, Stanford University, Stanford, CA	Radiation oncology
Yara Abdou, MD	Department of Medical Oncology, University of North Carolina, Chapel Hill, NC	Medical oncology (ASCO representative)
Penny Anderson, MD	Department of Radiation Oncology, Fox Chase Cancer Center, Philadelphia, PA	Radiation oncology
Parul Barry, MD	Department of Radiation Oncology, UPMC Hillman Cancer Center, University of Pittsburgh, Pittsburgh, PA	Radiation oncology
Julie A. Bradley, MD	Department of Radiation Oncology, University of Florida, Jacksonville, FL	Radiation oncology
Lourdes D. Heras, MPH	Gilbert, AZ	Patient representative
Atif Khan, MD, MS	Department of Radiation Oncology, Memorial Sloan Kettering Cancer Center, New York, NY	Radiation oncology
Cindy Matsen, MD	Department of Surgery, Huntsman Cancer Institute at the University of Utah, Salt Lake City, UT	Surgical oncology (SSO representative)
Rachel Rabinovitch, MD	Department of Radiation Oncology, University of Colorado Denver, Aurora, CO	Radiation oncology
Chantal Reyna, MD, MHA	Department of Surgery, Loyola University Medical Center, Chicago, IL	Surgical oncology (SSO representative)
Kilian E. Salerno, MD	Radiation Oncology Branch, National Cancer Institute, Bethesda, MD	Radiation oncology
Sarah E. Schellhorn, MD	Department of Medicine, Yale School of Medicine, New Haven, CT	Medical oncology (ASCO representative)
Deborah Schofield, PhD	Department of Radiation Physics, University of Texas MD Anderson Cancer Center, Houston, TX	Radiation oncology
Kekoa Taparra, MD, PhD, MPH	Department of Radiation Oncology, Stanford University, Stanford, CA	Radiation oncology
Iman Washington, MD	Department of Radiation Oncology, Moffitt Cancer Center, Tampa, FL	Radiation oncology
Jean L. Wright, MD	Department of Radiation Oncology, University of North Carolina, Chapel Hill, NC	Radiation oncology
Youssef H. Zeidan, MD, PhD	Department of Radiation Oncology, Florida International University and Lynn Cancer Institute, Boca Raton, FL	Radiation oncology
Richard C. Zellars, MD	Department of Radiation Oncology, Indiana University School of Medicine, Indianapolis, IN	Radiation oncology
Lisa Bradfield, BA	American Society for Radiation Oncology, Arlington, VA	ASTRO Practice Guideline Staff

Abbreviations: ASTRO, American Society for Radiation Oncology; SSO, Society of Surgical Oncology.