

# Analytic and iterative reconstructions in SPECT

J.-P. Guillement and R.G. Novikov

# Sommaire

Abstract

SPECT

Problem

Novikov formula

Chang formula

OAR

Numerical examples

References

Analytic  
and  
iterative  
recon-  
struc-  
tions in  
SPECT

J.-P.  
Guille-  
ment  
and  
R.G.  
Novikov

Abstract

SPECT

Problem

Novikov  
formula

Chang  
formula

OAR

Numerica  
examples

Reference

Abstract

SPECT

Problem

Novikov formula

Chang formula

OAR

Numerical examples

References

Analytic  
and  
iterative  
recon-  
struc-  
tions in  
SPECT

J.-P.  
Guille-  
ment  
and  
R.G.  
Novikov

Abstract

SPECT

Problem

Novikov  
formula

Chang  
formula

OAR

Numerica  
examples

Reference

## Abstract

We consider analytic and iterative reconstructions in the single-photon emission computed tomography (SPECT).

As analytic techniques we use, in particular, Chang's approximate inversion formula and Novikov's exact inversion formula for the attenuated ray transform, on one hand, and Wiener type filters for data with strong Poisson noise, on other hand.

As iterative techniques we consider the least square and expectation maximization iterative reconstructions.

Different comparisons are given.

## Abstract

SPECT

## Problem

## Novikov formula


## Chang formula

OAR

## Numerical examples

## References

# SPECT (Single Photon Emission Computed Tomography)



In the single-photon emission computed tomography (SPECT) one considers a body containing radioactive isotopes emitting photons. The emission data  $p$  in SPECT consist in the radiation measured outside the body by a family of detectors during some fixed time. The basic problem of SPECT consists in finding the distribution  $f$  of these isotopes in the body from the emission data  $p$  and some a priori information concerning the body. Usually this a priori information consists in the photon attenuation coefficient  $a$  in the points of body, where this coefficient is found in advance by the methods of the transmission computed tomography.

$f(x)$  - density of radioactive isotopes

$a(x)$  - photon attenuation coefficient

$x$  - point of (the space containing the) body

$p(\gamma)$  - emission data

$\gamma$  - point of detector set  $\Gamma$

$\Gamma$  - discrete subset of the set  $T$  of all oriented straight lines in the space containing the body

More precisely,  $p(\gamma)$  is the number of photons coming from (the domain containing) the body along oriented straight line  $\gamma$  to the detector associated with  $\gamma$ .





We remind that  $T \approx \mathbb{R} \times \mathbb{S}^1$ , where  $T$  is the set of all oriented straight lines in  $\mathbb{R}^2$ . If  $\gamma = (s, \theta) \in \mathbb{R} \times \mathbb{S}^1$ , then  $\gamma = \{x \in \mathbb{R}^2 : x = s\theta + t\theta^\perp, t \in \mathbb{R}\}$  (modulo orientation) and  $\theta$  gives the orientation of  $\gamma$ , where  $\theta^\perp = (-\theta_2, \theta_1)$  for  $\theta = (\theta_1, \theta_2) \in \mathbb{S}^1$ .

In practice,  $d = 2$  (after restriction to 2D plane),

$a \geq 0, f \geq 0, \text{supp } a \subset \mathcal{B}_R, \text{supp } f \subset \mathcal{B}_R,$

$\mathcal{B}_R = \{x \in \mathbb{R}^2 : |x| \leq R\},$

$R$ - radius of image support,  $\Gamma$  - is a uniform  $n \times n$  sampling of

$$T_R = \{\gamma \in T : \gamma \cap B_R \neq \emptyset\} = \{(s, \theta) \in \mathbb{R} \times \mathbb{S}^1 : |s| \leq R\}.$$

In addition, the standard value for  $n$  is 128.

## Abstract

SPECT

## Problem

## Novikov formula

## Chang formula

OAR

## Numerical examples

## References

## Analytic and iterative reconstructions in SPECT

J.-P.  
Guille-  
ment  
and  
R.G.  
Novikov

## Problem

Chang  
formula

OAR

## Numerical examples

## Reference

**Problem 1.** Find (as well as possible)  $Cf$  from  $p$  and  $a$ .

Our analytic approach to Problem 1 is based on the scheme

$$Cf \approx P_a^{-1} \mathcal{W}p, \quad (4)$$

where  $\mathcal{W}$  is a space-variant Wiener-type filter developed in [GN 2008] (or, more generally, some analytic method for approximate finding the noiseless data  $g$  of (1) from  $p$ ),  $P_a^{-1}$  is a reconstruction from  $\tilde{p} = \mathcal{W}p$ , based on some optimal combination of the Novikov exact (see [N 2002]) and Chang approximate (see [Ch 1978], [N 2011]) inversion formulas for  $P_a$ . In addition, the aforementioned optimal combination is constructed via a Morozov-type discrepancy principle.

## Abstract

SPECT

## Problem

## Novikov formula

## Chang formula

OAR

## Numerical examples

## References

## Analytic and iterative reconstructions in SPECT

J.-P.  
Guille-  
ment  
and  
R.G.  
Novikov

SPECT

Novikov  
formula

Chang  
formula

OAR

## Numerical examples

## Reference

# Novikov formula (see [N 2002])

$$Cf = \mathcal{N}_a g, \quad \text{where} \quad (5)$$

$$\mathcal{N}_a g(x) = \frac{1}{4\pi} \int_{\mathbb{S}^1} \theta^\perp \nabla_x K(x, \theta) d\theta,$$

$$\begin{aligned} K(x, \theta) &= \exp[-\mathcal{D}a(x, -\theta)] \tilde{g}(x\theta^\perp), \\ \tilde{g}(s) &= \exp[A_\theta(s)] \cos(B_\theta(s)) H(\exp[A_\theta] \cos(B_\theta) g_\theta)(s) + \\ &\quad \exp[A_\theta(s)] \sin(B_\theta(s)) H(\exp(A_\theta) \sin(B_\theta) g_\theta)(s), \end{aligned}$$

$$\begin{aligned} A_\theta(s) &= \frac{1}{2} P_0 a(s, \theta), \quad B_\theta(s) = H A_\theta(s), \quad g_\theta(s) = g(s, \theta), \\ g &= CP_a f \text{ (see (1) and (2))}, \end{aligned}$$

$$Hu(s) = \frac{1}{\pi} p.v. \int_{\mathbb{R}} \frac{u(t)}{s-t} dt \quad (\text{Hilbert transform}) \quad (6)$$

$x = (x_1, x_2) \in \mathbb{R}^2$ ,  $\theta = (\theta_1, \theta_2) \in \mathbb{S}^1$ ,  $\theta^\perp = (-\theta_2, \theta_1)$ ,  $s \in \mathbb{R}$ ,  $\mathcal{D}a$  is defined by (3).

Formula (5) is exact (for continuous case) but is not very stable for reconstruction from discrete and noisy data  $p$  of (1).

## Abstract

SPECT

## Problem

## Novikov formula

## Chang formula

OAR

## Numerical examples

## References

# Chang formula (see [Ch 1978], [N 2011])

$$Cf \simeq Ch_a g, \quad \text{where} \tag{7}$$

$$\begin{aligned} Ch_a g(x) &= \frac{1}{4\pi w_0(x)} \int_{\mathbb{S}^1} \theta^\perp \nabla_x H g_\theta(x \theta^\perp) d\theta, \\ w_0(x) &= \frac{1}{2\pi} \int_{\mathbb{S}^1} \exp[-\mathcal{D}a(x, \theta)] d\theta \end{aligned}$$

$$g = CP_a f \text{ (see (1) and (2)), } g_\theta(s) = g(s, \theta),$$

where  $H$  is defined by (6),  $\mathcal{D}a$  is defined by (3),  
 $x \in \mathbb{R}^2$ ,  $\theta = (\theta_1, \theta_2) \in \mathbb{S}^1$ ,  $\theta^\perp = (-\theta_2, \theta_1)$ ,  $s \in \mathbb{R}$ .

Formula (7) is approximate (for continuous case) but its result is sufficiently stable for reconstruction from discrete and noisy data  $p$  of (1).



## Abstract

SPECT

## Problem

## Novikov formula

## Chang formula

OAR

## Numerical examples

## References

# Optimized analytic reconstruction (OAR)

$$Cf \simeq Cf_\alpha = \mathcal{N}_{a_\alpha}(\mathcal{W}p)_\alpha + \mathcal{C}h_a(\mathcal{W}p - (\mathcal{W}p)_\alpha), \quad \text{where} \quad (8)$$

$\alpha$  is optimization parameter,  $\mathcal{W}$  is space-variant Wiener-type filter of [GN 2008],

$(\mathcal{W}p)_\alpha$  and  $a_\alpha$  are the low-frequency parts of  $\mathcal{W}p$  and  $a$  (respectively) obtained via some standard 2D space-invariant filtering dependent on  $\alpha$ .

$\mathcal{N}_a$  and  $\mathcal{C}h_a$  are the inversion operators of formulas (5) and (7).

In addition, we choose  $\alpha$  as a parameter minimizing the discrepancy  $\|P_a Cf - \mathcal{W}p\|_{L^2(\Gamma)}$ .

The ansatz  $Cf_\alpha$  of (8) is motivated by the facts that  $\mathcal{N}_a p$  of the exact formula (5) is sufficiently stable on sufficiently low frequency part of  $p$  and  $a$ , whereas  $\mathcal{C}h_a p$  of the approximate formula (7) is sufficiently stable on reasonably high frequency part of  $p$  and  $a$ .

## Abstract

SPECT

## Problem

## Novikov formula

## Chang formula

OAR


## Numerical examples

## References


# Numerical example 1



Attenuation map  $a$ , ( $128 \times 128$ )




Emission activity  $f$ , ( $128 \times 128$ )




Emission profile


# Noiseless and noisy projections




Projections  $g$  ( $128 \times 128$ )



Spectrum  $|\hat{g}|$




Projections  $\|p - g\|_2 / \|g\|_2 = 30\%$




Spectrum  $|\hat{p}|$


# Space-variant Wiener filter $A_{8,8}^{sym}$ and O.A. reconstruction




Projections  $\|\tilde{p} - g\|_2 / \|g\|_2 = 11\%$



Spectrum




OAR :  $\|r - r_0\|_2 / \|r_0\|_2 = 36\%$




Profile


# Iterative reconstructions (60 It)




Gradient :  $\|r - r_0\|_2 / \|r_0\|_2 = 43\%$



Profile



Em :  $\|r - r_0\|_2 / \|r_0\|_2 = 42\%$




Profile


## Numerical example 2



Attenuation map  $a$ ,  $(128 \times 128)$



Emission activity  $f$ ,  $(128 \times 128)$




Emission profile




# Noiseless and noisy projections




Projections  $g$  ( $128 \times 128$ )



Spectrum  $|\hat{g}|$




Projections  $\|p - g\|_2 / \|g\|_2 = 30\%$




Spectrum  $|\hat{p}|$


# Space-variant Wiener filter $A_{8,8}^{sym}$ and O.A. reconstruction




Projections  $\|\tilde{p} - g\|_2 / \|g\|_2 = 10\%$



Spectrum



OAR :  $\|r - r_0\|_2 / \|r_0\|_2 = 22\%$



Profile

Analytic  
and  
iterative  
recon-  
struc-  
tions in  
SPECT

J.-P.  
Guille-  
ment  
and  
R.G.  
Novikov

Abstract

SPECT

Problem

Novikov  
formula


Chang  
formula

OAR


Numerica  
examples

Reference


# Iterative reconstructions (60 it)




Gradient :  $\|r - r_0\|_2 / \|r_0\|_2 = 24\%$



Profile



Em :  $\|r - r_0\|_2 / \|r_0\|_2 = 52\%$



Profile

## Abstract

SPECT

## Problem

## Novikov formula

## Chang formula

OAR

## Numerical examples

## References

## Analytic and iterative reconstructions in SPECT

J.-P.  
Guille-  
ment  
and  
R.G.  
Novikov

SPECT

Chang  
formula

OAR

## Numerical examples

## Reference

# References

- [Ch 1978] Chang L T 1978 *A method for attenuation correction in radionuclide computed tomography. IEEE Trans. Nucl. Sci.* **25** 638-643
- [GN 2008] Guillement J-P. and Novikov R.G. 2008 *On Wiener type filters in SPECT. Inverse Problems* 24 025001 (26 pp)
- [N 2002] Novikov R G 2002 *An inversion formula for the attenuated x-ray transformation. Ark. Mat.* **40** 145-167
- [N 2011] Novikov R G 2011 *Weighted Radon transforms for which Chang's approximate inversion formula is exact. Uspekhi Mat. Nauk* **66** (2) 237-238