

# Fokas Methods applied to a Boundary Valued Problem for Conjugate Conductivity Equations

Joint work with Slah Chaabi and Franck Wielonsky

I2M, UMR 7373 CNRS (Aix-Marseille University)

Moscow, the 18th of february, 2021

## Different motivations.

- ▶  $\Omega$  is a smooth domain in  $\mathbb{C}$ .

## Different motivations.

- ▶  $\Omega$  is a smooth domain in  $\mathbb{C}$ .
- ▶  $L$  an elliptic differential operator in  $\overline{\Omega}$

$$L = a(x, y) \frac{\partial^2}{\partial x^2} + b(x, y) \frac{\partial^2}{\partial x \partial y} + c(x, y) \frac{\partial^2}{\partial y^2} +$$
$$+ d(x, y) \frac{\partial}{\partial x} + e(x, y) \frac{\partial}{\partial y} + f(x, y)$$

## Different motivations.

- ▶  $\Omega$  is a smooth domain in  $\mathbb{C}$ .
- ▶  $L$  an elliptic differential operator in  $\overline{\Omega}$

$$L = a(x, y) \frac{\partial^2}{\partial x^2} + b(x, y) \frac{\partial^2}{\partial x \partial y} + c(x, y) \frac{\partial^2}{\partial y^2} + d(x, y) \frac{\partial}{\partial x} + e(x, y) \frac{\partial}{\partial y} + f(x, y)$$

- ▶ Is it possible to solve explicitly  $Lu = 0$  in  $\overline{\Omega}$  with the knowledge of  $u$  on the boundary  $\partial\Omega$  ?

## Different motivations.

- ▶  $\Omega$  is a smooth domain in  $\mathbb{C}$ .
- ▶  $L$  an elliptic differential operator in  $\overline{\Omega}$

$$L = a(x, y) \frac{\partial^2}{\partial x^2} + b(x, y) \frac{\partial^2}{\partial x \partial y} + c(x, y) \frac{\partial^2}{\partial y^2} + d(x, y) \frac{\partial}{\partial x} + e(x, y) \frac{\partial}{\partial y} + f(x, y)$$

- ▶ Is it possible to solve explicitly  $Lu = 0$  in  $\overline{\Omega}$  with the knowledge of  $u$  on the boundary  $\partial\Omega$  ?
- ▶ Lax-Milgram theorem gives the existence and unicity of the solution if  $u$  is given on  $\partial\Omega$  (and continuous)

## Different motivations.

- ▶  $\Omega$  is a smooth domain in  $\mathbb{C}$ .
- ▶  $L$  an elliptic differential operator in  $\overline{\Omega}$

$$L = a(x, y) \frac{\partial^2}{\partial x^2} + b(x, y) \frac{\partial^2}{\partial x \partial y} + c(x, y) \frac{\partial^2}{\partial y^2} + d(x, y) \frac{\partial}{\partial x} + e(x, y) \frac{\partial}{\partial y} + f(x, y)$$

- ▶ Is it possible to solve explicitly  $Lu = 0$  in  $\overline{\Omega}$  with the knowledge of  $u$  on the boundary  $\partial\Omega$  ?
- ▶ Lax-Milgram theorem gives the existence and unicity of the solution if  $u$  is given on  $\partial\Omega$  (and continuous)
- ▶ Construction of fundamental solutions gives a representation formula en terms of  $u$  and the normal derivative  $\partial_{\bar{n}}u$  on the boundary  $\partial\Omega$  : we need to much informations.



## Different motivations.

- ▶  $\Omega$  is a smooth domain in  $\mathbb{C}$ .
- ▶  $L$  an elliptic differential operator in  $\overline{\Omega}$

$$L = a(x, y) \frac{\partial^2}{\partial x^2} + b(x, y) \frac{\partial^2}{\partial x \partial y} + c(x, y) \frac{\partial^2}{\partial y^2} + d(x, y) \frac{\partial}{\partial x} + e(x, y) \frac{\partial}{\partial y} + f(x, y)$$


- ▶ Is it possible to solve explicitly  $Lu = 0$  in  $\overline{\Omega}$  with the knowledge of  $u$  on the boundary  $\partial\Omega$  ?
- ▶ Lax-Milgram theorem gives the existence and unicity of the solution if  $u$  is given on  $\partial\Omega$  (and continuous)
- ▶ Construction of fundamental solutions gives a representation formula en terms of  $u$  and the normal derivative  $\partial_{\bar{n}}u$  on the boundary  $\partial\Omega$  : we need to much informations.
- ▶ Is it possible to obtain a relation between  $u$  and  $\partial_{\bar{n}}u$  on  $\partial\Omega$  without computing  $u$  in  $\Omega$  ?

# One other motivation : What is a Tokamak ?






# Mathematical Formulation




# Mathematical Formulation



- Poloidal field  $\psi$  and its normal derivative  $\partial_{\vec{n}}\psi$  are known on  $\partial D(a, r)$ ,  $\psi = C$  on the boundary  $\partial P$  of the plasma and

$$\operatorname{div} \left( \frac{1}{x} \nabla \psi \right) = 0 \quad \text{in} \quad D(a, r) \setminus P$$

# Mathematical Formulation



- Poloidal field  $\psi$  and its normal derivative  $\partial_{\vec{n}}\psi$  are known on  $\partial D(a, r)$ ,  $\psi = C$  on the boundary  $\partial P$  of the plasma and

$$\operatorname{div}\left(\frac{1}{x}\nabla\psi\right)=0 \quad \text{in} \quad D(a, r) \setminus P$$

- Generalized axisymetrical potential for  $\alpha \in \mathbb{R}$  :

$$\operatorname{div}(x^{\alpha}\nabla u)=0 \quad \Leftrightarrow \quad \Delta u + \frac{\alpha}{x}\frac{\partial u}{\partial x}=0.$$

# The Fokas method Fokas (1997) : a very simple example.



$$q_t - q_{xx} = 0 \quad \text{sur} \quad x \geq 0, \quad t \geq 0 \quad (H)$$

# The Fokas method Fokas (1997) : a very simple example.



$$q_t - q_{xx} = 0 \quad \text{sur} \quad x \geq 0, \quad t \geq 0 \quad (H)$$

- ▶ Formulation in terms of Lax pair : splitting of this equation into two ordinary differential equations, compatible if and only if  $q$  is a solution to  $(H)$

$$\begin{cases} \mu_x - ik\mu = q \\ \mu_t - \mu_{xx} = 0 \end{cases}$$

# The Fokas method Fokas (1997) : a very simple example.



$$q_t - q_{xx} = 0 \quad \text{sur} \quad x \geq 0, \quad t \geq 0 \quad (H)$$

- ▶ Formulation in terms of Lax pair : splitting of this equation into two ordinary differential equations, compatible if and only if  $q$  is a solution to  $(H)$

$$\begin{cases} \mu_x - ik\mu = q \\ \mu_t - \mu_{xx} = 0 \end{cases}$$



$$\begin{cases} \mu_x - ik\mu = q \\ \mu_t + k^2\mu = q_x + ikq \end{cases}$$

# The Fokas method Fokas (1997) : a very simple example.



$$q_t - q_{xx} = 0 \quad \text{sur} \quad x \geq 0, \quad t \geq 0 \quad (H)$$

- Formulation in terms of Lax pair : splitting of this equation into two ordinary differential equations, compatible if and only if  $q$  is a solution to  $(H)$

$$\begin{cases} \mu_x - ik\mu = q \\ \mu_t - \mu_{xx} = 0 \end{cases}$$



$$\begin{cases} \mu_x - ik\mu = q \\ \mu_t + k^2\mu = q_x + ikq \end{cases}$$



$$\begin{cases} \left( \mu e^{-ikx+k^2t} \right)_x = q e^{-ikx+k^2t} \\ \left( \mu e^{-ikx+k^2t} \right)_t = (q_x + ikq) e^{-ikx+k^2t} \end{cases}$$

# The Fokas method Fokas (1997) : a very simple example.



$$q_t - q_{xx} = 0 \quad \text{sur} \quad x \geq 0, \quad t \geq 0 \quad (H)$$

- Formulation in terms of Lax pair : splitting of this equation into two ordinary differential equations, compatible if and only if  $q$  is a solution to  $(H)$

$$\begin{cases} \mu_x - ik\mu = q \\ \mu_t - \mu_{xx} = 0 \end{cases}$$



$$\begin{cases} \mu_x - ik\mu = q \\ \mu_t + k^2\mu = q_x + ikq \end{cases}$$



$$\begin{cases} \left( \mu e^{-ikx+k^2t} \right)_x = q e^{-ikx+k^2t} \\ \left( \mu e^{-ikx+k^2t} \right)_t = (q_x + ikq) e^{-ikx+k^2t} \end{cases}$$



$$d \left( \mu e^{-ikx+k^2t} \right) = e^{-ikx+k^2t} (q dx + (q_x + ikq) dt)$$



- ▶  $\nu = e^{-ikx+k^2t} (qdx + (q_x + ikq)dt)$  is closed

- ▶  $\nu = e^{-ikx+k^2t} (qdx + (q_x + ikq)dt)$  is closed
- ▶  $q(x, t)$  and  $q_x(x, t)$  decay to 0 at  $\infty$  sufficiently fast.

- ▶  $\nu = e^{-ikx+k^2t} (qdx + (q_x + ikq)dt)$  is closed
- ▶  $q(x, t)$  and  $q_x(x, t)$  decay to 0 at  $\infty$  sufficiently fast.
- ▶  $\arg k \in \left[\frac{5\pi}{4}, \frac{7\pi}{4}\right]$

$$\int_{x=0}^{\infty} e^{-ikx} q(x, 0) dx = \int_{t=0}^{\infty} e^{k^2t} (q_x(0, t) + ikq(0, t)) dt \quad (GR)$$

- ▶  $\nu = e^{-ikx+k^2t} (qdx + (q_x + ikq)dt)$  is closed
- ▶  $q(x, t)$  and  $q_x(x, t)$  decay to 0 at  $\infty$  sufficiently fast.
- ▶  $\arg k \in \left[\frac{5\pi}{4}, \frac{7\pi}{4}\right]$

$$\int_{x=0}^{\infty} e^{-ikx} q(x, 0) dx = \int_{t=0}^{\infty} e^{k^2 t} (q_x(0, t) + ikq(0, t)) dt \quad (GR)$$

- ▶ Let us assume first that  $q(x, 0)$ ,  $q_x(0, t)$ ,  $q(0, t)$  are known. Reconstruction of  $q$  in  $\mathbb{R}^+ \times \mathbb{R}^+$  is equivalent to the reconstruction of  $\mu$  since

$$d(\mu e^{-ikx+k^2t}) = \nu.$$


- ▶  $\nu = e^{-ikx+k^2t} (qdx + (q_x + ikq)dt)$  is closed
- ▶  $q(x, t)$  and  $q_x(x, t)$  decay to 0 at  $\infty$  sufficiently fast.
- ▶  $\arg k \in \left[\frac{5\pi}{4}, \frac{7\pi}{4}\right]$

$$\int_{x=0}^{\infty} e^{-ikx} q(x, 0) dx = \int_{t=0}^{\infty} e^{k^2t} (q_x(0, t) + ikq(0, t)) dt \quad (GR)$$


- ▶ Let us assume first that  $q(x, 0)$ ,  $q_x(0, t)$ ,  $q(0, t)$  are known. Reconstruction of  $q$  in  $\mathbb{R}^+ \times \mathbb{R}^+$  is equivalent to the reconstruction of  $\mu$  since

$$d(\mu e^{-ikx+k^2t}) = \nu.$$

- ▶ Integration of  $\nu$  between points of the boundary and  $(x, t)$ .




$$\mu_{46}(x, t, k) = e^{ikx - k^2 t} \int_{\gamma_1} \nu$$




$$\mu_{46}(x, t, k) = e^{ikx - k^2 t} \int_{\gamma_1} \nu$$

$$\mu_5(x, t, k) = e^{ikx - k^2 t} \int_{\gamma_2} \nu$$




$$\mu_{46}(x, t, k) = e^{ikx - k^2 t} \int_{\gamma_1} \nu$$

$$\mu_5(x, t, k) = e^{ikx - k^2 t} \int_{\gamma_2} \nu$$









$$\mu_{46}(x, t, k) = e^{ikx - k^2 t} \int_{\gamma_{46}} \nu$$

$$\mu_5(x, t, k) = e^{ikx - k^2 t} \int_{\gamma_5} \nu$$


$$\mu_{123}(x, t, k) = e^{ikx - k^2 t} \int_{\gamma_{123}} \nu$$



# Riemann-Hilbert Problems




# Riemann-Hilbert Problems



►  $\mu(x, t, k) = \mathcal{O}\left(\frac{1}{k}\right)$


# Riemann-Hilbert Problems



►  $\mu(x, t, k) = \mathcal{O}\left(\frac{1}{k}\right)$

►  $\mu^+ - \mu^- = \phi$

# Riemann-Hilbert Problems



- ▶  $\mu(x, t, k) = \mathcal{O}\left(\frac{1}{k}\right)$
- ▶  $\mu^+ - \mu^- = \phi$
- ▶  $\mu = \frac{1}{2\pi i} \int_L \frac{\phi(k') dk'}{k' - k}$  (Plemelj Formula)

# Lax Pairs and closed differential form for the GASP equation.

►  $L_\alpha(u) = \Delta u + \frac{\alpha}{x} \frac{\partial u}{\partial x} = 0$  in  $\mathbb{H} = \{z, \operatorname{Re} z > 0\}$ .

# Lax Pairs and closed differential form for the GASP equation.

- ▶  $L_\alpha(u) = \Delta u + \frac{\alpha}{x} \frac{\partial u}{\partial x} = 0$  in  $\mathbb{H} = \{z, \operatorname{Re} z > 0\}$ .
- ▶  $u_{z\bar{z}} + \frac{\alpha}{2(z + \bar{z})}(u_z + u_{\bar{z}}) = 0.$  (GASP)

# Lax Pairs and closed differential form for the GASP equation.

- ▶  $L_\alpha(u) = \Delta u + \frac{\alpha}{x} \frac{\partial u}{\partial x} = 0$  in  $\mathbb{H} = \{z, \operatorname{Re} z > 0\}$ .
- ▶  $u_{z\bar{z}} + \frac{\alpha}{2(z + \bar{z})}(u_z + u_{\bar{z}}) = 0.$  (GASP)
- ▶  $\begin{cases} \phi_z(z, k) = (k + \bar{z})^{\alpha/2}(k - z)^{\alpha/2-1}u_z(z), \\ \phi_{\bar{z}}(z, k) = (k + \bar{z})^{\alpha/2-1}(k - z)^{\alpha/2}u_{\bar{z}}(z) \end{cases}$



# Lax Pairs and closed differential form for the GASP equation.

- ▶  $L_\alpha(u) = \Delta u + \frac{\alpha}{x} \frac{\partial u}{\partial x} = 0$  in  $\mathbb{H} = \{z, \operatorname{Re} z > 0\}$ .
- ▶  $u_{z\bar{z}} + \frac{\alpha}{2(z + \bar{z})}(u_z + u_{\bar{z}}) = 0. \quad (\text{GASP})$
- ▶ 
$$\begin{cases} \phi_z(z, k) = (k + \bar{z})^{\alpha/2}(k - z)^{\alpha/2-1}u_z(z), \\ \phi_{\bar{z}}(z, k) = (k + \bar{z})^{\alpha/2-1}(k - z)^{\alpha/2}u_{\bar{z}}(z) \end{cases}$$
- ▶  $W(z, k) = [(k - z)(k + \bar{z})]^{\alpha/2-1}[(k + \bar{z})u_z(z)dz + (k - z)u_{\bar{z}}(z)d\bar{z}]$ 

Note that, when  $\alpha \in 2\mathbb{N}^*$ , the differential form has no singularity in  $\Omega$  and  $k$  may be any complex number. Otherwise, for  $\alpha \in \mathbb{R} \setminus 2\mathbb{N}^*$ ,  $W(z, k)$  has a pole or a branching point in  $k$  or  $-\bar{k}$  if one of this point is in  $\Omega$ .

# Lax Pairs and closed differential form for the GASP equation.

- ▶  $L_\alpha(u) = \Delta u + \frac{\alpha}{x} \frac{\partial u}{\partial x} = 0$  in  $\mathbb{H} = \{z, \operatorname{Re} z > 0\}$ .
- ▶  $u_{z\bar{z}} + \frac{\alpha}{2(z + \bar{z})}(u_z + u_{\bar{z}}) = 0. \quad (\text{GASP})$
- ▶ 
$$\begin{cases} \phi_z(z, k) = (k + \bar{z})^{\alpha/2}(k - z)^{\alpha/2-1}u_z(z), \\ \phi_{\bar{z}}(z, k) = (k + \bar{z})^{\alpha/2-1}(k - z)^{\alpha/2}u_{\bar{z}}(z) \end{cases}$$
- ▶  $W(z, k) = [(k - z)(k + \bar{z})]^{\alpha/2-1}[(k + \bar{z})u_z(z)dz + (k - z)u_{\bar{z}}(z)d\bar{z}]$ 


Note that, when  $\alpha \in 2\mathbb{N}^*$ , the differential form has no singularity in  $\Omega$  and  $k$  may be any complex number. Otherwise, for  $\alpha \in \mathbb{R} \setminus 2\mathbb{N}^*$ ,  $W(z, k)$  has a pole or a branching point in  $k$  or  $-\bar{k}$  if one of this point is in  $\Omega$ .
- ▶  $L_\alpha(u) = 0 \Leftrightarrow L_{2-\alpha}(x^{\alpha-1}u) = 0.$

$$\alpha = -2m, \quad m \in \mathbb{N}.$$

► 
$$W(z, k) = \frac{(k + \bar{z})u_z dz + (k - z)u_{\bar{z}} d\bar{z}}{[(k - z)(k + \bar{z})]^{m+1}}$$

$$\alpha = -2m, \quad m \in \mathbb{N}.$$

$$\triangleright W(z, k) = \frac{(k + \bar{z})u_z dz + (k - z)u_{\bar{z}}d\bar{z}}{[(k - z)(k + \bar{z})]^{m+1}}$$




$\triangleright$

$$\phi(z, k) = \int W(z', k)$$

$$\alpha = -2m, \quad m \in \mathbb{N}.$$

$$\triangleright W(z, k) = \frac{(k + \bar{z})u_z dz + (k - z)u_{\bar{z}}d\bar{z}}{[(k - z)(k + \bar{z})]^{m+1}}$$



►


$$\phi(z, k) = \int W(z', k)$$

► poles of order  $m$  in  $-\bar{z}_r, -\bar{z}, z, z_r$  if  $m > 0$ .



$$\alpha = -2m, \quad m \in \mathbb{N}.$$

$$\triangleright W(z, k) = \frac{(k + \bar{z})u_z dz + (k - z)u_{\bar{z}}d\bar{z}}{[(k - z)(k + \bar{z})]^{m+1}}$$



▶

$$\phi(z, k) = \int W(z', k)$$


▶ poles of order  $m$  in  $-\bar{z}_r, -\bar{z}, z, z_r$  if  $m > 0$ .

$$\triangleright \phi(z, -\bar{k}) = -\overline{\phi(z, k)}$$

▶ Jump on  $(z, z_r) \cup (-\bar{z}_r, -\bar{z})$  equal to  $J(k) = -\int_C W(z', k)$

$$\alpha = -2m, \quad m \in \mathbb{N}.$$

$$\triangleright W(z, k) = \frac{(k + \bar{z})u_z dz + (k - z)u_{\bar{z}}d\bar{z}}{[(k - z)(k + \bar{z})]^{m+1}}$$



►

$$\phi(z, k) = \int W(z', k)$$

► poles of order  $m$  in  $-\bar{z}_r, -\bar{z}, z, z_r$  if  $m > 0$ .

$$\triangleright \phi(z, -\bar{k}) = -\overline{\phi(z, k)}$$


► Jump on  $(z, z_r) \cup (-\bar{z}_r, -\bar{z})$  equal to  $J(k) = - \int_C W(z', k)$

►  $J$  has no singularity in  $z$  and  $z_r$ .



$$\alpha = -2m, \quad m \in \mathbb{N}.$$

$$\triangleright W(z, k) = \frac{(k + \bar{z})u_z dz + (k - z)u_{\bar{z}} d\bar{z}}{[(k - z)(k + \bar{z})]^{m+1}}$$



$\triangleright$

$$\phi(z, k) = \int W(z', k)$$

$\triangleright$  poles of order  $m$  in  $-\bar{z}_r, -\bar{z}, z, z_r$  if  $m > 0$ .

$$\triangleright \phi(z, -\bar{k}) = -\overline{\phi(z, k)}$$

$\triangleright$  Jump on  $(z, z_r) \cup (-\bar{z}_r, -\bar{z})$  equal to  $J(k) = - \int_C W(z', k)$

$\triangleright J$  has no singularity in  $z$  and  $z_r$ .

$$\triangleright \phi \sim_{k \rightarrow +\infty} \frac{u(z) - u(z_r)}{k^{2m-1}}$$

- ▶ Renormalization :  $\tilde{\phi}(z, k) = ((k - z)(k + \bar{z}))^m \phi(z, k)$   
 $\tilde{J}(z, k) = ((k - z)(k + \bar{z}))^m J(z, k)$

- ▶ Renormalization :  $\tilde{\phi}(z, k) = ((k - z)(k + \bar{z}))^m \phi(z, k)$

$$\tilde{J}(z, k) = ((k - z)(k + \bar{z}))^m J(z, k)$$

- ▶  $\tilde{\phi}(z, k) \sim_{k \rightarrow +\infty} \frac{u(z) - u(z_r)}{k}$

- ▶ Renormalization :  $\tilde{\phi}(z, k) = ((k - z)(k + \bar{z}))^m \phi(z, k)$   
 $\tilde{J}(z, k) = ((k - z)(k + \bar{z}))^m J(z, k)$
- ▶  $\tilde{\phi}(z, k) \sim_{k \rightarrow +\infty} \frac{u(z) - u(z_r)}{k}$
- ▶  $\tilde{\phi}$  regular in  $z$  and  $-\bar{z}$ , polar singularities in  $z_r$  et  $-\bar{z}_r$ .

- ▶ Renormalization :  $\tilde{\phi}(z, k) = ((k - z)(k + \bar{z}))^m \phi(z, k)$   
 $\tilde{J}(z, k) = ((k - z)(k + \bar{z}))^m J(z, k)$
- ▶  $\tilde{\phi}(z, k) \sim_{k \rightarrow +\infty} \frac{u(z) - u(z_r)}{k}$
- ▶  $\tilde{\phi}$  regular in  $z$  and  $-\bar{z}$ , polar singularities in  $z_r$  et  $-\bar{z}_r$ .
- ▶  $\tilde{\phi}_{z_r, -\bar{z}_r}(z, k)$  polar part in this point

- ▶ Renormalization :  $\tilde{\phi}(z, k) = ((k - z)(k + \bar{z}))^m \phi(z, k)$   
 $\tilde{J}(z, k) = ((k - z)(k + \bar{z}))^m J(z, k)$
- ▶  $\tilde{\phi}(z, k) \sim_{k \rightarrow +\infty} \frac{u(z) - u(z_r)}{k}$
- ▶  $\tilde{\phi}$  regular in  $z$  and  $-\bar{z}$ , polar singularities in  $z_r$  et  $-\bar{z}_r$ .
- ▶  $\tilde{\phi}_{z_r, -\bar{z}_r}(z, k)$  polar part in this point
- ▶  $\tilde{\phi} - \tilde{\phi}_{z_r, -\bar{z}_r}$  analytic outside  $(z, z_r) \cup (-\bar{z}_r, -\bar{z})$  and has jump equal to  $\tilde{J}(z, k)$

- ▶ Renormalization :  $\tilde{\phi}(z, k) = ((k - z)(k + \bar{z}))^m \phi(z, k)$   
 $\tilde{J}(z, k) = ((k - z)(k + \bar{z}))^m J(z, k)$
- ▶  $\tilde{\phi}(z, k) \sim_{k \rightarrow +\infty} \frac{u(z) - u(z_r)}{k}$
- ▶  $\tilde{\phi}$  regular in  $z$  and  $-\bar{z}$ , polar singularities in  $z_r$  et  $-\bar{z}_r$ .
- ▶  $\tilde{\phi}_{z_r, -\bar{z}_r}(z, k)$  polar part in this point
- ▶  $\tilde{\phi} - \tilde{\phi}_{z_r, -\bar{z}_r}$  analytic outside  $(z, z_r) \cup (-\bar{z}_r, -\bar{z})$  and has jump equal to  $\tilde{J}(z, k)$
- ▶ Plemelj formula

$$\tilde{\phi}(z, k) - \tilde{\phi}_{z_r, -\bar{z}_r}(z, k) = \frac{1}{2\pi i} \int_{(-\bar{z}_r, -\bar{z}) \cup (z, z_r)} \frac{\tilde{J}(z, k') dk'}{k' - k}$$

- ▶ Renormalization :  $\tilde{\phi}(z, k) = ((k - z)(k + \bar{z}))^m \phi(z, k)$   
 $\tilde{J}(z, k) = ((k - z)(k + \bar{z}))^m J(z, k)$
- ▶  $\tilde{\phi}(z, k) \sim_{k \rightarrow +\infty} \frac{u(z) - u(z_r)}{k}$
- ▶  $\tilde{\phi}$  regular in  $z$  and  $-\bar{z}$ , polar singularities in  $z_r$  et  $-\bar{z}_r$ .
- ▶  $\tilde{\phi}_{z_r, -\bar{z}_r}(z, k)$  polar part in this point
- ▶  $\tilde{\phi} - \tilde{\phi}_{z_r, -\bar{z}_r}$  analytic outside  $(z, z_r) \cup (-\bar{z}_r, -\bar{z})$  and has jump equal to  $\tilde{J}(z, k)$
- ▶ Plemelj formula

$$\tilde{\phi}(z, k) - \tilde{\phi}_{z_r, -\bar{z}_r}(z, k) = \frac{1}{2\pi i} \int_{(-\bar{z}_r, -\bar{z}) \cup (z, z_r)} \frac{\tilde{J}(z, k') dk'}{k' - k}$$

▶

$$u(z) - u(z_r) = 2 \operatorname{Re} a_r - \frac{1}{\pi} \operatorname{Im} \int_{(z, z_r)} \tilde{J}(z, k') dk'$$



# Computation of the residue $a_r$ of $\tilde{\phi}$ in $z_r$

►  $u_z dz = \frac{1}{2}(u_t + iu_n)ds, \quad u_{\bar{z}} d\bar{z} = \frac{1}{2}(u_t - iu_n)ds.$

# Computation of the residue $a_r$ of $\widetilde{\phi}$ in $z_r$

- ▶  $u_z dz = \frac{1}{2}(u_t + iu_n)ds, \quad u_{\bar{z}} d\bar{z} = \frac{1}{2}(u_t - iu_n)ds.$
- ▶  $\phi(z, k) = \int W(z', k) = \int \frac{[(k - iy')u_t + ix'u_n]ds}{[(k - z')(k + \bar{z}')]^{m+1}}$

## Computation of the residue $a_r$ of $\tilde{\phi}$ in $z_r$

- ▶  $u_z dz = \frac{1}{2}(u_t + iu_n)ds, \quad u_{\bar{z}} d\bar{z} = \frac{1}{2}(u_t - iu_n)ds.$
- ▶  $\phi(z, k) = \int W(z', k) = \int \frac{[(k - iy')u_t + ix'u_n]ds}{[(k - z')(k + \bar{z}')]^{m+1}}$
- ▶  $\phi(z, k) = \int (k - z')^{-m-1} w(z', k) dz'$  where

$$w(z', k) = (k + \bar{z}')^{-m-1} ((k - iy')u_t + ix'u_n) \tau^{-1}(z') dz'$$

$\tau(z')$  is the unit tangent vector to  $\mathcal{C}$  in  $z'$ .

# Computation of the residue $a_r$ of $\widetilde{\phi}$ in $z_r$

- ▶  $u_z dz = \frac{1}{2}(u_t + iu_n)ds, \quad u_{\bar{z}} d\bar{z} = \frac{1}{2}(u_t - iu_n)ds.$
- ▶  $\phi(z, k) = \int W(z', k) = \int \frac{[(k - iy')u_t + ix'u_n]ds}{[(k - z')(k + \bar{z}')]^{m+1}}$
- ▶  $\phi(z, k) = \int (k - z')^{-m-1} w(z', k) dz'$  where

$$w(z', k) = (k + \bar{z}')^{-m-1} ((k - iy')u_t + ix'u_n) \tau^{-1}(z') dz'$$

$\tau(z')$  is the unit tangent vector to  $\mathcal{C}$  in  $z'$ .

- ▶  $m$  integrations by parts give  $\widetilde{\phi}_{z_r, -\bar{z}_r}$ , then  $a_r$ .

Let  $u$  be a solution to the equation  $\Delta u + \alpha x^{-1} \partial_x u = 0$ ,  $\alpha = -2m$ ,  $m \in \mathbb{N}$ , in the domain  $\mathcal{D}$  with smooth tangential derivatives  $u_t$  and normal derivatives  $u_n$  on the boundary  $\mathcal{C}$ .

$$u(z) = -\frac{1}{\pi} \operatorname{Im} \int_{(z, z_r)} ((k-z)(k+\bar{z}))^m J(z, k) dk + 2\operatorname{Re} a_r + u(z_r), \quad (1)$$

where  $a_r$  can be explicitly computed in terms of the tangential derivative along  $\mathcal{C}$  of  $u_t$  and  $u_n$ , up to the order  $m-1$ , in  $z_r$ . Function  $J(z, k)$  is given by

$$J(z, k) = - \int_{\mathcal{C}} W(z', k),$$

where  $W(z, k)$  is the differential form

$$W(z, k) = ((k-z)(k+\bar{z}))^{-m-1} ((k+\bar{z})u_z(z)dz + (k-z)u_{\bar{z}}(z)d\bar{z}) \quad (2)$$

$$= ((k-z)(k+\bar{z}))^{-m-1} ((k-iy)u_t(z) + ixu_n(z)) ds, \quad (3)$$

with  $z = x + iy$  and  $ds$  the unit length element on  $\mathcal{C}$ .

## The case $\alpha = -1$ .


► 
$$W(z, k) = \frac{(k + \bar{z})u_z dz + (k - z)u_{\bar{z}} d\bar{z}}{\sqrt{(k - z)(k + \bar{z})}}$$

## The case $\alpha = -1$ .

- ▶  $W(z, k) = \frac{(k + \bar{z})u_z dz + (k - z)u_{\bar{z}} d\bar{z}}{\sqrt{(k - z)(k + \bar{z})}}$
- ▶ Riemann surface  $\mathbb{S}_z$  : two copies of  $\mathbb{C}$ ,  $\mathbb{S}_{z,1}$  et  $\mathbb{S}_{z,2}$  glued together along the branching cut  $[-\bar{z}, z]$ .

## The case $\alpha = -1$ .


- ▶  $W(z, k) = \frac{(k + \bar{z})u_z dz + (k - z)u_{\bar{z}} d\bar{z}}{\sqrt{(k - z)(k + \bar{z})}}$
- ▶ Riemann surface  $\mathbb{S}_z$  : two copies of  $\mathbb{C}$ ,  $\mathbb{S}_{z,1}$  et  $\mathbb{S}_{z,2}$  glued together along the branching cut  $[-\bar{z}, z]$ .





## The case $\alpha = -1$ .

- ▶  $W(z, k) = \frac{(k + \bar{z})u_z dz + (k - z)u_{\bar{z}} d\bar{z}}{\sqrt{(k - z)(k + \bar{z})}}$
- ▶ Riemann surface  $\mathbb{S}_z$  : two copies of  $\mathbb{C}$ ,  $\mathbb{S}_{z,1}$  et  $\mathbb{S}_{z,2}$  glued together along the branching cut  $[-\bar{z}, z]$ .



▶

- ▶  $\lambda_1(z, k) \sim k$  when  $k \rightarrow \infty_1$  on the sheet above  $\mathbb{S}_{z,1}$   
 $\lambda_2(z, k) \sim -k$  when  $k \rightarrow \infty_2$  on the sheet below  $\mathbb{S}_{z,2}$


## Integration of the form.

► 
$$\phi(z, k) = \int \frac{(k + \bar{z}')u_z dz' + (k - z')u_{\bar{z}} d\bar{z}'}{\lambda(z', k)}$$

# Integration of the form.

► 
$$\phi(z, k) = \int \frac{(k + \bar{z}')u_z dz' + (k - z')u_{\bar{z}} d\bar{z}'}{\lambda(z', k)}$$


► Sheet 1




# Integration of the form.

► 
$$\phi(z, k) = \int \frac{(k + \bar{z}')u_z dz' + (k - z')u_{\bar{z}} d\bar{z}'}{\lambda(z', k)}$$

► Sheet 1




► Sheet 2




# Integration of the form.

► 
$$\phi(z, k) = \int \frac{(k + \bar{z}')u_z dz' + (k - z')u_{\bar{z}} d\bar{z}'}{\lambda(z', k)}$$



► Sheet 1





► Sheet 2



► 
$$\phi(z, \infty_1) = -\phi(z, \infty_2).$$





$$\begin{aligned} \phi(z, k) = & \frac{1}{4i\pi} \int_{C_{up} \cup -\bar{C}_{up}} J(z, k') \left( \frac{\lambda(z, k)}{\lambda_1(z, k')} + 1 \right) \frac{dk'}{k' - k} \\ & + \frac{1}{4i\pi} \int_{C_{low} \cup -\bar{C}_{low}} J(z, k') \left( \frac{\lambda(z, k)}{\lambda_2(z, k')} + 1 \right) \frac{dk'}{k' - k}, \quad (4) \end{aligned}$$

# The notion of good Lax pair.

- ▶  $\int_{\partial\Omega} W(u_t, u_n, k) ds = 0$  for all  $k \in K \subset \mathbb{C}$ .



## The notion of good Lax pair.

- ▶  $\int_{\partial\Omega} W(u_t, u_n, k) ds = 0$  for all  $k \in K \subset \mathbb{C}$ .
- ▶ Is this relation useful ? In other words, if  $u = 0 \Rightarrow u_n = 0$  on  $\partial\Omega$  ? If  $u$  is known on  $\partial\Omega$ , is it possible to obtain  $u_n$  on  $\partial\Omega$  ?

## The notion of good Lax pair.

- ▶  $\int_{\partial\Omega} W(u_t, u_n, k) ds = 0$  for all  $k \in K \subset \mathbb{C}$ .
- ▶ Is this relation useful ? In other words, if  $u = 0 \Rightarrow u_n = 0$  on  $\partial\Omega$  ? If  $u$  is known on  $\partial\Omega$ , is it possible to obtain  $u_n$  on  $\partial\Omega$  ?
- ▶ In full generality, no

## The notion of good Lax pair.

- ▶  $\int_{\partial\Omega} W(u_t, u_n, k) ds = 0$  for all  $k \in K \subset \mathbb{C}$ .
- ▶ Is this relation useful ? In other words, if  $u = 0 \Rightarrow u_n = 0$  on  $\partial\Omega$  ? If  $u$  is known on  $\partial\Omega$ , is it possible to obtain  $u_n$  on  $\partial\Omega$  ?
- ▶ In full generality, no
- ▶ Example :  $\Omega = \mathbb{D}$ ,  $W = \frac{u_z}{k-z} dz + \frac{u_{\bar{z}}}{k+\bar{z}} d\bar{z}$ .  $\Delta u = 0$ .

## The notion of good Lax pair.

- ▶  $\int_{\partial\Omega} W(u_t, u_n, k) ds = 0$  for all  $k \in K \subset \mathbb{C}$ .
- ▶ Is this relation useful ? In other words, if  $u = 0 \Rightarrow u_n = 0$  on  $\partial\Omega$  ? If  $u$  is known on  $\partial\Omega$ , is it possible to obtain  $u_n$  on  $\partial\Omega$  ?
- ▶ In full generality, no
- ▶ Example :  $\Omega = \mathbb{D}$ ,  $W = \frac{u_z}{k-z} dz + \frac{u_{\bar{z}}}{k+\bar{z}} d\bar{z}$ .  $\Delta u = 0$ .
- ▶  $u_z dz = \frac{1}{2}(u_t + iu_n)ds$ ,  $u_{\bar{z}} d\bar{z} = \frac{1}{2}(u_t - iu_n)ds$ .

## The notion of good Lax pair.

- ▶  $\int_{\partial\Omega} W(u_t, u_n, k) ds = 0$  for all  $k \in K \subset \mathbb{C}$ .
- ▶ Is this relation useful ? In other words, if  $u = 0 \Rightarrow u_n = 0$  on  $\partial\Omega$  ? If  $u$  is known on  $\partial\Omega$ , is it possible to obtain  $u_n$  on  $\partial\Omega$  ?
- ▶ In full generality, no
- ▶ Example :  $\Omega = \mathbb{D}$ ,  $W = \frac{u_z}{k-z} dz + \frac{u_{\bar{z}}}{k+\bar{z}} d\bar{z}$ .  $\Delta u = 0$ .
- ▶  $u_z dz = \frac{1}{2}(u_t + iu_n) ds$ ,  $u_{\bar{z}} d\bar{z} = \frac{1}{2}(u_t - iu_n) ds$ .
- ▶ 
$$\int_{\partial\mathbb{D}} u_n(z) \left( \frac{1}{k-z} - \frac{1}{k+\bar{z}} \right) ds(z) =$$
$$i \int_{\partial\mathbb{D}} u_t(z) \left( \frac{1}{k-z} + \frac{1}{k+\bar{z}} \right) ds(z)$$

## The notion of good Lax pair.

- ▶  $\int_{\partial\Omega} W(u_t, u_n, k) ds = 0$  for all  $k \in K \subset \mathbb{C}$ .
- ▶ Is this relation useful ? In other words, if  $u = 0 \Rightarrow u_n = 0$  on  $\partial\Omega$  ? If  $u$  is known on  $\partial\Omega$ , is it possible to obtain  $u_n$  on  $\partial\Omega$  ?
- ▶ In full generality, no
- ▶ Example :  $\Omega = \mathbb{D}$ ,  $W = \frac{u_z}{k-z} dz + \frac{u_{\bar{z}}}{k+\bar{z}} d\bar{z}$ .  $\Delta u = 0$ .
- ▶  $u_z dz = \frac{1}{2}(u_t + iu_n) ds$ ,  $u_{\bar{z}} d\bar{z} = \frac{1}{2}(u_t - iu_n) ds$ .
- ▶ 
$$\int_{\partial\mathbb{D}} u_n(z) \left( \frac{1}{k-z} - \frac{1}{k+\bar{z}} \right) ds(z) =$$
$$i \int_{\partial\mathbb{D}} u_t(z) \left( \frac{1}{k-z} + \frac{1}{k+\bar{z}} \right) ds(z)$$
- ▶  $u_t \equiv 0$  et  $\frac{1}{k-z} = \sum_{n=1}^{\infty} \frac{z^{n-1}}{k^n}$ ,  $\frac{1}{\bar{z}+k} = \sum_{n=1}^{\infty} \frac{(-\bar{z})^{n-1}}{k^n}$

# The notion of good Lax pair.

- ▶  $\int_{\partial\Omega} W(u_t, u_n, k) ds = 0$  for all  $k \in K \subset \mathbb{C}$ .
- ▶ Is this relation useful ? In other words, if  $u = 0 \Rightarrow u_n = 0$  on  $\partial\Omega$  ? If  $u$  is known on  $\partial\Omega$ , is it possible to obtain  $u_n$  on  $\partial\Omega$  ?
- ▶ In full generality, no
- ▶ Example :  $\Omega = \mathbb{D}$ ,  $W = \frac{u_z}{k-z} dz + \frac{u_{\bar{z}}}{k+\bar{z}} d\bar{z}$ .  $\Delta u = 0$ .
- ▶  $u_z dz = \frac{1}{2}(u_t + iu_n) ds$ ,  $u_{\bar{z}} = \frac{1}{2}(u_t - iu_n) ds$ .
- ▶ 
$$\int_{\partial\mathbb{D}} u_n(z) \left( \frac{1}{k-z} - \frac{1}{k+\bar{z}} \right) ds(z) =$$
$$i \int_{\partial\mathbb{D}} u_t(z) \left( \frac{1}{k-z} + \frac{1}{k+\bar{z}} \right) ds(z)$$
- ▶  $u_t \equiv 0$  et  $\frac{1}{k-z} = \sum_{n=1}^{\infty} \frac{z^{n-1}}{k^n}$ ,  $\frac{1}{\bar{z}+k} = \sum_{n=1}^{\infty} \frac{(-\bar{z})^{n-1}}{k^n}$

▶

$$u_n(z) = \sum_{n=-\infty}^{\infty} b_n z^n, \quad \sum_{n=1}^{+\infty} \frac{\overline{b_{n-1}} + (-1)^n b_{n-1}}{k^n} = 0.$$

## Good Lax pair for $\alpha = -2(m-1)$ .

- ▶  $u_t$  known.



## Good Lax pair for $\alpha = -2(m-1)$ .

- ▶  $u_t$  known.
- ▶  $\forall k \in \mathbb{C} \setminus (D_a \cup D_{-a})$

$$\int_{\partial D(a,1)} \frac{x u_n(z) ds(z)}{[(k-z)(k+\bar{z})]^m} = \int_{\partial D(a,1)} \frac{(y+ik) u_t(z) ds(z)}{[(k-z)(k+\bar{z})]^m}$$

## Good Lax pair for $\alpha = -2(m-1)$ .

- ▶  $u_t$  known.
- ▶  $\forall k \in \mathbb{C} \setminus (D_a \cup D_{-a})$

$$\int_{\partial D(a,1)} \frac{x u_n(z) ds(z)}{[(k-z)(k+\bar{z})]^m} = \int_{\partial D(a,1)} \frac{(y+ik) u_t(z) ds(z)}{[(k-z)(k+\bar{z})]^m}$$

- ▶  $u_t \equiv 0$  on  $C_a \Rightarrow u_n \equiv 0$ .

## Good Lax pair for $\alpha = -2(m-1)$ .

- ▶  $u_t$  known.
- ▶  $\forall k \in \mathbb{C} \setminus (D_a \cup D_{-a})$

$$\int_{\partial D(a,1)} \frac{x u_n(z) ds(z)}{[(k-z)(k+\bar{z})]^m} = \int_{\partial D(a,1)} \frac{(y+ik) u_t(z) ds(z)}{[(k-z)(k+\bar{z})]^m}$$

- ▶  $u_t \equiv 0$  on  $C_a \Rightarrow u_n \equiv 0$ .
- ▶  $f(z) := (x+a)u_n(z+a)$ ,  $z \in \mathbb{T}$  has to be recover from the relation

$$\int_{\mathbb{T}} \frac{z^{m-1} f(z) dz}{(z-(k-a))^m (z+(k+a)^{-1})^m} = 0$$

where  $k \in \mathbb{C} \setminus (D_a \cup D_{-a})$ .

## Good Lax pair for $\alpha = -2(m-1)$ .

- ▶  $u_t$  known.
- ▶  $\forall k \in \mathbb{C} \setminus (D_a \cup D_{-a})$

$$\int_{\partial D(a,1)} \frac{x u_n(z) ds(z)}{[(k-z)(k+\bar{z})]^m} = \int_{\partial D(a,1)} \frac{(y+ik) u_t(z) ds(z)}{[(k-z)(k+\bar{z})]^m}$$

- ▶  $u_t \equiv 0$  on  $C_a \Rightarrow u_n \equiv 0$ .
- ▶  $f(z) := (x+a)u_n(z+a)$ ,  $z \in \mathbb{T}$  has to be recover from the relation

$$\int_{\mathbb{T}} \frac{z^{m-1} f(z) dz}{(z - (k-a))^m (z + (k+a)^{-1})^m} = 0$$

where  $k \in \mathbb{C} \setminus (D_a \cup D_{-a})$ .

- ▶ Since  $f$  is real valued, one has for  $z \in \mathbb{T}$

$$f(z) = g(z) + \bar{g}(1/z)$$

with  $g \in H(\mathbb{D})$  and  $\bar{g}(1/z) \in H(\mathbb{C} \setminus \bar{\mathbb{D}})$ .



$$\int_{\mathbb{T}} \frac{z^{m-1}(g(z) + \bar{g}(1/z))dz}{(z - (k - a))^m \left(z + \frac{1}{k+a}\right)^m} = 0.$$



$$\int_{\mathbb{T}} \frac{z^{m-1}(g(z) + \bar{g}(1/z))dz}{(z - (k - a))^m \left(z + \frac{1}{k+a}\right)^m} = 0.$$



$$\begin{aligned} & \left(\frac{-1}{k+a}\right)^m \left[ \frac{z^{m-1}g(z)}{(z - (k - a))^m} \right]^{(m-1)} \left(\frac{-1}{k+a}\right) + \\ & \left(\frac{1}{k-a}\right)^m \left[ \frac{z^{m-1}\bar{g}(z)}{(z + (k + a))^m} \right]^{(m-1)} \left(\frac{1}{k-a}\right) = 0. \end{aligned}$$



$$\varphi(z) = -\frac{z}{1+2az} \qquad \varphi\left(-\frac{1}{k+a}\right) = \frac{1}{k-a}.$$

$$\varphi(z) = -\frac{z}{1+2az} \quad \varphi\left(-\frac{1}{k+a}\right) = \frac{1}{k-a}.$$


$$\Phi(z) + \bar{\Phi}(\varphi(z)) = 0, \quad z \in \mathbb{A}.$$



$$\varphi(z) = -\frac{z}{1+2az} \quad \varphi\left(-\frac{1}{k+a}\right) = \frac{1}{k-a}.$$

$$\Phi(z) + \bar{\Phi}(\varphi(z)) = 0, \quad z \in \mathbb{A}.$$


$$\mathbb{A} = \mathbb{D} \setminus \overline{D}\left(-\frac{2a}{4a^2-1}, \frac{1}{4a^2-1}\right)$$



$$\varphi(z) = -\frac{z}{1+2az} \quad \varphi\left(-\frac{1}{k+a}\right) = \frac{1}{k-a}.$$

$$\Phi(z) + \bar{\Phi}(\varphi(z)) = 0, \quad z \in \mathbb{A}.$$


$$\mathbb{A} = \mathbb{D} \setminus \overline{D}\left(-\frac{2a}{4a^2-1}, \frac{1}{4a^2-1}\right)$$




► For  $\mu \in \mathbb{A}$

$$\Phi(\mu) = \int_{\mathbb{T}} \frac{z^{m-1} g(z) dz}{(1 - \varphi(\mu)z)^m (z - \mu)^m} = \frac{2\pi i}{(m-1)!} \left( \frac{h(z)}{(1 - \varphi(\mu)z)^m} \right)^{(m-1)} (\mu)$$

with  $h(z) = z^{m-1} g(z)$ .



$\varphi : \mathbb{A} \rightarrow \mathbb{A}$ , and  $D\left(-\frac{2a}{4a^2-1}, \frac{1}{4a^2-1}\right) \rightarrow \mathbb{C} \setminus \overline{\mathbb{D}}$   
 and  $-1/(2a) \rightarrow \infty$ .




$\varphi : \mathbb{A} \rightarrow \mathbb{A}$ , and  $D(-\frac{2a}{4a^2-1}, \frac{1}{4a^2-1}) \rightarrow \mathbb{C} \setminus \overline{\mathbb{D}}$   
and  $-1/(2a) \rightarrow \infty$ .

►  $z_1$  and  $z_2$  the roots of  $z^2 + 2az + 1$ .

$$z_1 = -a + \sqrt{a^2 - 1} \in D(-\frac{2a}{4a^2-1}, \frac{1}{4a^2-1})$$

$$z_2 = \varphi(z_1) = -a - \sqrt{a^2 - 1} \in \mathbb{C} \setminus \mathbb{D}$$



$\varphi : \mathbb{A} \rightarrow \mathbb{A}$ , and  $D(-\frac{2a}{4a^2-1}, \frac{1}{4a^2-1}) \rightarrow \mathbb{C} \setminus \overline{\mathbb{D}}$   
and  $-1/(2a) \rightarrow \infty$ .

- ▶  $z_1$  and  $z_2$  the roots of  $z^2 + 2az + 1$ .

$$z_1 = -a + \sqrt{a^2 - 1} \in D(-\frac{2a}{4a^2-1}, \frac{1}{4a^2-1})$$

$$z_2 = \varphi(z_1) = -a - \sqrt{a^2 - 1} \in \mathbb{C} \setminus \mathbb{D}$$

- ▶  $1 - \varphi(\mu)\mu = \frac{\mu^2 + 2a\mu + 1}{1 + 2a\mu} \Rightarrow \Phi$  has zero of order at least  $m$  at  $-1/(2a)$ .

- Multiplying  $\Phi(z) + \bar{\Phi}(\varphi(z)) = 0$  by

$$S(\mu) = \frac{(\mu - z_1)^{2m-1}(\mu - z_2)^{2m-1}}{(2a\mu + 1)^m}$$

one gets  $S(\mu)\Phi(\mu) = -S(\mu)\bar{\Phi}(\varphi(\mu))$ .

- Multiplying  $\Phi(z) + \bar{\Phi}(\varphi(z)) = 0$  by

$$S(\mu) = \frac{(\mu - z_1)^{2m-1}(\mu - z_2)^{2m-1}}{(2a\mu + 1)^m}$$

one gets  $S(\mu)\Phi(\mu) = -S(\mu)\bar{\Phi}(\varphi(\mu))$ .



$$\Phi(z) = \sum_{p=0}^{m-1} \alpha_p z^{m-1-p} (z^2 + 2az + 1)^p h^{(p)}(z) = P_{2m-2}$$

- ▶ Multiplying  $\Phi(z) + \bar{\Phi}(\varphi(z)) = 0$  by

$$S(\mu) = \frac{(\mu - z_1)^{2m-1}(\mu - z_2)^{2m-1}}{(2a\mu + 1)^m}$$

one gets  $S(\mu)\Phi(\mu) = -S(\mu)\bar{\Phi}(\varphi(\mu))$ .



$$\Phi(z) = \sum_{p=0}^{m-1} \alpha_p z^{m-1-p} (z^2 + 2az + 1)^p h^{(p)}(z) = P_{2m-2}$$

- ▶  $h(z) = \sum_{n=0}^{+\infty} a_n (z - z_1)^n$ .



- ▶ Multiplying  $\Phi(z) + \bar{\Phi}(\varphi(z)) = 0$  by

$$S(\mu) = \frac{(\mu - z_1)^{2m-1}(\mu - z_2)^{2m-1}}{(2a\mu + 1)^m}$$

one gets  $S(\mu)\Phi(\mu) = -S(\mu)\bar{\Phi}(\varphi(\mu))$ .



$$\Phi(z) = \sum_{p=0}^{m-1} \alpha_p z^{m-1-p} (z^2 + 2az + 1)^p h^{(p)}(z) = P_{2m-2}$$

- ▶  $h(z) = \sum_{n=0}^{+\infty} a_n (z - z_1)^n$ .
- ▶  $\sum_{\ell=0}^{m-1} a_{n-\ell} \beta_{\ell}^n = 0, \quad n \geq 2m - 1$ .

- ▶ Multiplying  $\Phi(z) + \bar{\Phi}(\varphi(z)) = 0$  by

$$S(\mu) = \frac{(\mu - z_1)^{2m-1}(\mu - z_2)^{2m-1}}{(2a\mu + 1)^m}$$

one gets  $S(\mu)\Phi(\mu) = -S(\mu)\bar{\Phi}(\varphi(\mu))$ .



$$\Phi(z) = \sum_{p=0}^{m-1} \alpha_p z^{m-1-p} (z^2 + 2az + 1)^p h^{(p)}(z) = P_{2m-2}$$

- ▶  $h(z) = \sum_{n=0}^{+\infty} a_n (z - z_1)^n$ .
- ▶  $\sum_{\ell=0}^{m-1} a_{n-\ell} \beta_{\ell}^n = 0, n \geq 2m - 1$ .
- ▶  $h$  is a polynomial of degree less than  $2m - 2$ .

- ▶ Multiplying  $\Phi(z) + \bar{\Phi}(\varphi(z)) = 0$  by

$$S(\mu) = \frac{(\mu - z_1)^{2m-1}(\mu - z_2)^{2m-1}}{(2a\mu + 1)^m}$$

one gets  $S(\mu)\Phi(\mu) = -S(\mu)\bar{\Phi}(\varphi(\mu))$ .



$$\Phi(z) = \sum_{p=0}^{m-1} \alpha_p z^{m-1-p} (z^2 + 2az + 1)^p h^{(p)}(z) = P_{2m-2}$$

- ▶  $h(z) = \sum_{n=0}^{+\infty} a_n (z - z_1)^n$ .
- ▶  $\sum_{\ell=0}^{m-1} a_{n-\ell} \beta_{\ell}^n = 0, n \geq 2m - 1$ .
- ▶  $h$  is a polynomial of degree less than  $2m - 2$ .
- ▶  $g$  is a polynomial of degree less than  $m - 1$ .



$$\int_{\mathbb{T}} \frac{z^{m-1} f(z)}{(z - (k - a))^m (z + (k + a)^{-1})^m} = 0, \quad \forall k \in \mathbb{C} \setminus \{\overline{\mathcal{D}_a} \cup \overline{-\mathcal{D}_a}\},$$

where  $f(z) = g(z) + \bar{g}\left(\frac{1}{z}\right)$ .



$$\int_{\mathbb{T}} \frac{z^{m-1} f(z)}{(z - (k - a))^m (z + (k + a)^{-1})^m} = 0, \quad \forall k \in \mathbb{C} \setminus \{\overline{\mathcal{D}_a} \cup -\overline{\mathcal{D}_a}\},$$

where  $f(z) = g(z) + \bar{g}\left(\frac{1}{z}\right)$ .

► We put  $\xi = -(k + a)^{-1}$ , we get

$$\int_{\mathbb{T}} \frac{z^{m-1} f(z)}{(\xi z + 2a\xi + 1)^m (z - \xi)^m} = 0$$
$$\forall \xi \in \mathbb{D} \setminus \overline{D\left(-\frac{2a}{4a^2 - 1}, \frac{1}{4a^2 - 1}\right)}.$$



$$\frac{\partial^{m-1}}{\partial z^{m-1}} \left( \frac{z^{m-1} f(z)}{(\xi z + 2a\xi + 1)^m} \right) \Big|_{z=\xi} = 0.$$



$$\frac{\partial^{m-1}}{\partial z^{m-1}} \left( \frac{z^{m-1} f(z)}{(\xi z + 2a\xi + 1)^m} \right) \Big|_{z=\xi} = 0.$$

- Let us denote  $F(z) = z^{m-1} f(z) \in \mathbb{C}_{2m-2}[z]$ .

$$\sum_{k=0}^{m-1} \frac{(-1)^k (m+k-1)!}{k! (m-1-k)!} \frac{\xi^k}{(\xi^2 + 2a\xi + 1)^{m+k}} F^{(m-1-k)}(\xi) = 0.$$



$$\frac{\partial^{m-1}}{\partial z^{m-1}} \left( \frac{z^{m-1} f(z)}{(\xi z + 2a\xi + 1)^m} \right) \Big|_{z=\xi} = 0.$$

- Let us denote  $F(z) = z^{m-1} f(z) \in \mathbb{C}_{2m-2}[z]$ .

$$\sum_{k=0}^{m-1} \frac{(-1)^k (m+k-1)!}{k! (m-1-k)!} \frac{\xi^k}{(\xi^2 + 2a\xi + 1)^{m+k}} F^{(m-1-k)}(\xi) = 0.$$





$$\frac{\partial^{m-1}}{\partial z^{m-1}} \left( \frac{z^{m-1} f(z)}{(\xi z + 2a\xi + 1)^m} \right) \Big|_{z=\xi} = 0.$$

- Let us denote  $F(z) = z^{m-1} f(z) \in \mathbb{C}_{2m-2}[z]$ .

$$\sum_{k=0}^{m-1} \frac{(-1)^k (m+k-1)!}{k! (m-1-k)!} \frac{\xi^k}{(\xi^2 + 2a\xi + 1)^{m+k}} F^{(m-1-k)}(\xi) = 0.$$

- Denotes  $z_1 = -a + \sqrt{a^2 - 1}$  and  $z_2 = -a - \sqrt{a^2 - 1}$

$$\forall k \in \{0, 1, \dots, m-1\}, \begin{cases} F^{(k)}(z_1) = 0, \\ F^{(k)}(z_2) = 0. \end{cases}$$



$$\frac{\partial^{m-1}}{\partial z^{m-1}} \left( \frac{z^{m-1} f(z)}{(\xi z + 2a\xi + 1)^m} \right) \Big|_{z=\xi} = 0.$$

- Let us denote  $F(z) = z^{m-1} f(z) \in \mathbb{C}_{2m-2}[z]$ .

$$\sum_{k=0}^{m-1} \frac{(-1)^k (m+k-1)!}{k! (m-1-k)!} \frac{\xi^k}{(\xi^2 + 2a\xi + 1)^{m+k}} F^{(m-1-k)}(\xi) = 0.$$

- Denotes  $z_1 = -a + \sqrt{a^2 - 1}$  and  $z_2 = -a - \sqrt{a^2 - 1}$

$$\forall k \in \{0, 1, \dots, m-1\}, \begin{cases} F^{(k)}(z_1) = 0, \\ F^{(k)}(z_2) = 0. \end{cases}$$

- $F \equiv 0$ .



$$\frac{\partial^{m-1}}{\partial z^{m-1}} \left( \frac{z^{m-1} f(z)}{(\xi z + 2a\xi + 1)^m} \right) \Big|_{z=\xi} = 0.$$

- Let us denote  $F(z) = z^{m-1} f(z) \in \mathbb{C}_{2m-2}[z]$ .

$$\sum_{k=0}^{m-1} \frac{(-1)^k (m+k-1)!}{k! (m-1-k)!} \frac{\xi^k}{(\xi^2 + 2a\xi + 1)^{m+k}} F^{(m-1-k)}(\xi) = 0.$$

- Denotes  $z_1 = -a + \sqrt{a^2 - 1}$  and  $z_2 = -a - \sqrt{a^2 - 1}$

$$\forall k \in \{0, 1, \dots, m-1\}, \begin{cases} F^{(k)}(z_1) = 0, \\ F^{(k)}(z_2) = 0. \end{cases}$$

- $F \equiv 0$ .

- Thank you very much for your attention !