


α :



$\mathcal{L} \in \vee \in \mathbb{Z}, \neg$:


$$A(\mathcal{C}, \neg \forall)$$

$\mathcal{L} \in \vee \wedge \in \vee$:





|||| (EGIPT)


β :




γ :




δ :




Example:





Tabelă semantică închisă $\xRightarrow{TCC} U$ e tautologie



Tabelă deschisă $\xRightarrow{TCC} U$ nu e tautologie

$$3) U \stackrel{\text{not}}{=} (\exists y)(\forall x)p(x,y)$$

$$\neg U = \neg((\exists y)(\forall x)p(x,y)) \quad (1)$$

| $\mathcal{D}(1)$ a constantă implicită

$$\neg(\forall x)p(x,a) \quad (2)$$

$$\neg(\exists y)(\forall x)p(x,y) \quad (1')$$

| $\mathcal{D}(2)$ b const. nouă

$$\neg p(a,b)$$

| $\mathcal{D}(1')$ b const. existentă

$$\neg(\forall x)p(x,b) \quad (3)$$

$$\neg(\exists y)(\forall x)p(x,y) \quad (1'')$$

| $\mathcal{D}(3)$ c const. nouă

$$\neg p(c,b)$$

| $\mathcal{D}(1'')$

$$\neg(\forall x)p(x,c) \quad (4)$$

$$(\exists x)(\forall y)p(x,y) \quad (1''')$$

$$\neg p(b,c)$$

| ∞