

# New records of rare deep-sea fishes (Teleostei) collected from off north-eastern Brazil, including seamounts and islands of the Fernando de Noronha Ridge

Michael Maia Mincarone<sup>1</sup>  | Leandro Nolé Eduardo<sup>2,3</sup>  | Fabio Di Dario<sup>1</sup>  |  
Thierry Frédou<sup>2</sup>  | Arnaud Bertrand<sup>2,3</sup>  | Flávia Lucena-Frédou<sup>2</sup> 

<sup>1</sup>Instituto de Biodiversidade e Sustentabilidade, Universidade Federal do Rio de Janeiro, Macaé, Brazil

<sup>2</sup>Departamento de Pesca e Aquicultura, Universidade Federal Rural de Pernambuco, Recife, Brazil

<sup>3</sup>MARBECC, Université de Montpellier, CNRS Ifremer, IRD, Montpellier, France

## Correspondence

Michael Maia Mincarone, Instituto de Biodiversidade e Sustentabilidade, Universidade Federal do Rio de Janeiro, Av. São José do Barreto, 764, 27965-045 Macaé, RJ, Brazil.  
Email: [mincarone@macae.ufrj.br](mailto:mincarone@macae.ufrj.br)

## Funding information

Brazilian Biodiversity Fund, Grant/Award Numbers: 104/2016, 011/2019; Conselho Nacional de Desenvolvimento Científico e Tecnológico, Grant/Award Numbers: 314644/2020-2, PROTAX 443302/2020, 307422/2020-8, 308554/2019-1; Coordenação de Aperfeiçoamento de Pessoal de Nível Superior, Grant/Award Number: 88881.142689/2017-01; Fundação Carlos Chagas Filho de Amparo à Pesquisa do Estado do Rio de Janeiro, Grant/Award Number: E-26/210.290/2021; H2020 European Research Council, Grant/Award Number: 817578

## Abstract

Here we report on new data on the occurrence and distribution of rare deep-sea fishes (Teleostei) collected from off north-eastern Brazil, including seamounts and oceanic islands of the Fernando de Noronha Ridge (Rocas Atoll and Fernando de Noronha Archipelago). Collections were made by the French RV *Antea* during the ABRACOS (Acoustics along the BRAZilian COaSt) expeditions between 29 September and 21 October 2015, and 9 April and 10 May 2017. Occurrences of *Photostylus pycnopterus* (Alepocephalidae), *Gigantura chuni* (Giganturidae), *Ahliesaurus berryi* (Notosudidae), *Benthallbella infans*, *Rosenblattichthys hubbsi*, *Scopelarchoides danae* (Scopelarchidae), *Scopelengys tristis* (Neoscopelidae), *Zu cristatus* (Trachipteridae), *Stylephorus chordatus* (Stylephoridae) and *Pseudoscopelus cordilluminatus* (Chiasmodontidae) are reported for the first time or confirmed in the Brazilian Exclusive Economic Zone. *Photostylus pycnopterus* and *G. chuni* are also recorded for the first time in the western South Atlantic, whereas records of *P. cordilluminatus* are the first in the western Atlantic. Other records of rare species of those families are also reported and discussed.

## KEYWORDS

deep-sea fishes, distribution, taxonomy, tropical Atlantic

## 1 | INTRODUCTION

Mesopelagic fishes are one of the most important vertebrate groups of the world's oceans, usually presenting high biodiversity and abundance in deep-sea waters. Despite their relevance on a global scale, there is a major lack of knowledge regarding their regional composition. In the western South Atlantic, a still relatively poorly sampled region, knowledge about the diversity and distribution of deep-sea fishes is mostly based on local and fragmentary studies. In this context, studies conducted along the vast Brazilian coastline (about

7500 km) and its associated Exclusive Economic Zone, including oceanic islands and seamounts, are key for better understanding the marine biodiversity of the region (e.g., Reis *et al.*, 2016).

Major scientific surveys performed in the last decades along the Brazilian coast focused mostly on the demersal fish diversity (e.g., Bernardes *et al.*, 2005; Costa *et al.*, 2007; Costa & Mincarone, 2010; Klautau *et al.*, 2020; Lins Oliveira *et al.*, 2015), while relatively less attention has been given to the meso- and bathypelagic fishes (e.g., Braga *et al.*, 2014; Figueiredo *et al.*, 2002; Lima *et al.*, 2011; Mincarone *et al.*, 2014). In recent years, the team members of the Project

ABRACOS (Acoustics along the BRAZilian COaSt) conducted a multi-disciplinary study off north-eastern Brazil to better understand the degree of connectivity between the continental margin and the oceanic islands and seamounts of the region. The two surveys performed along the study area revealed a rich mesopelagic fish fauna, including many new distributional records in groups of the Teleostei such as the Argentiniformes (Mincarone *et al.*, 2021), Stomiiformes (Eduardo *et al.*, 2020a, 2020b; Villarins *et al.*, 2022), Ceratioidei (Mincarone *et al.*, 2021), Stephanoberycoidei (Afonso *et al.*, 2021), Caristiidae (Mincarone *et al.*, 2019), Howellidae (Eduardo *et al.*, 2019) and Trichiuridae (Eduardo *et al.*, 2018). Here, we provide new distributional records of rare species of the families Alepocephalidae, Chiasmodontidae, Giganturidae, Neoscopelidae, Notosudidae, Scopelarchidae, Stylephoridae and Trachipteridae collected during the ABRACOS expeditions off north-eastern Brazil.

## 2 | MATERIALS AND METHODS

Specimens examined were collected during the ABRACOS expeditions carried out between 29 September and 21 October 2015 (ABRACOS 1: AB1; Bertrand, 2015), and between 9 April and 10 May 2017 (ABRACOS 2: AB2; Bertrand, 2017). Both expeditions were conducted by the French RV *Antea* off north-eastern Brazil, including collections along the Fernando de Noronha Ridge (Rocas Atoll, Fernando de Noronha Archipelago and seamounts off Rio Grande do Norte State). The surveys comprised 82 fishing stations between the surface and 1113 m depth, and resulted in the collection of more than 9000 fish specimens. Further information on sampling methods and fishing gear is provided by Mincarone *et al.* (2021). All specimens are deposited in the Fish Collection of the Instituto de Biodiversidade e Sustentabilidade, Universidade Federal do Rio de Janeiro – NPM (Macaé, Brazil). Other institutional abbreviations follow Sabaj (2020). Diagnostic features of the species reported in the text are composed of a mixture of pertinent characters from different taxonomic levels (family, genus and species) to facilitate the identification of specimens.

**Ethical Statement:** All methods were approved and conducted in accordance with regulations of the Brazilian Ministry of Environment (SISBIO authorization number: 47270–5). Operations of the RV *Antea* were approved by the Brazilian Navy Authority ("Estado-Maior da Armada") under the Ordinances 178 (9 August 2015) and 4 (24 January 2017).

## 3 | RESULTS

### 3.1 | *Photostylus pycnopterus* Beebe 1933 (Alepocephalidae)

#### 3.1.1 | Specimens examined

NPM 5015, 1, 95 mm, sta. AB2/49A, 04°10'38.1"S, 33°16'07.4"W to 04°10'58.0"S, 33°15'03.8"W, 770–1020 m, 30 April 2017, 21:17–

21:52 h; NPM 5018, 1, 75 mm, sta. AB2/52A, 03°43'16.2"S, 33°25'09.8"W to 03°42'14.2"S, 33°24'36.2"W, 822–984 m, 2 May 2017, 11:47–12:18 h (Figures 1a and 2a).

#### 3.1.2 | Diagnostic features

*Photostylus pycnopterus* is distinguished by a scaleless body, covered with photophores on raised stalks; a small head (about 14%–19% SL) with an oblique upper profile; snout length longer than eye length; ventral outline of upper jaw with obtuse angle at end of premaxilla; maxilla extends near to posterior margin of eye; pseudobranch absent; pectoral fins fan like and with 17–20 rays; dorsal-fin base shorter than anal-fin base, with 12–15 rays; 17–19 anal-fin rays (Beebe, 1933; McEachran & Fechhelm, 1998; Sutton *et al.*, 2020; Tsukamoto *et al.*, 1992; Wisner, 1976). The genus is monotypic (Fricke *et al.*, 2022).

#### 3.1.3 | Distribution

*Photostylus pycnopterus* is a meso- to bathypelagic species with a wide distribution in the tropical and temperate waters of the Atlantic, Pacific and Indian oceans, from 840 to 2868 m depth (Beebe, 1933; Fitch & Lavenberg, 1968; Goodyear, 1969; Grey, 1958; Kashkin, 1975; Mundy, 2005; Parin *et al.*, 1976; Quéro, 1974; Rivaton *et al.*, 1990; Sutton *et al.*, 2020; Tsukamoto *et al.*, 1992; Wisner, 1976). In the western Atlantic, the species was reported from Davis Strait (Canada) to the Caribbean Sea, including the Gulf of Mexico (Beebe, 1933; Becker *et al.*, 1975; Carter & Hartel, 2003; McEachran & Fechhelm, 1998; Moore *et al.*, 2003; Nakamura & Okamura, 1995). The two specimens reported here were collected from depths between 770 and 1020 m near the seamounts off eastern Rocas Atoll (Figure 2a). They represent the first records of the species in the western South Atlantic and in Brazilian waters.

#### 3.1.4 | Remarks


This is the only species of the Alepocephalidae collected during the ABRACOS expeditions. About 18 species of that family are currently reported in Brazilian waters (e.g., Melo *et al.*, 2020), but this number is certainly underestimated according to ongoing studies by the authors.

### 3.2 | *Gigantura chuni* Brauer 1901 (Giganturidae)


#### 3.2.1 | Specimens examined

NPM 4241, 1, 167 mm, sta. AB2/44A, 03°52'52.5"S, 32°17'33.3"W to 03°51'13.4"S, 32°16'28.0"W, 850 m, 28 April 2017, 12:44–13:17 h; NPM 5238, 2, 42–? (damaged) mm, sta. AB2/53A, 03°48'58.7"S, 33°59'17.1"W to 03°50'05.8"S, 33°58'46.5"W, 610 m, 2 May 2017, 22:08–22:40 h (Figures 1b and 2b).

**FIGURE 1** (a) *Photostylus pycnopterus*, NPM 5015, 95 mm SL; (b) *Gigantura chuni*, NPM 4241, 167 mm SL; (c) *Ahliesaurus berryi*, NPM 4574, 198 mm SL; (d) *Benthalbella infans*, NPM 3553, 57 mm SL; (e) *Rosenblattichthys hubbsi*, NPM 4554, 85 mm SL; (f) *Scopelarchoides danae*, NPM 4555, 80 mm SL; (g) *Scopelengys tristis*, NPM 4620, 145 mm SL; (h) *Zu cristatus*, NPM 4504, 430 mm SL; (i) *Stylephorus chordatus*, NPM 3178, 255 mm SL; (j) *Pseudoscopelus cordilluminatus*, NPM 4536, 57 mm SL. Scale bar = 10 mm



**FIGURE 2** Records of species of (a) Alepocephalidae and (b) Giganturidae in Brazilian waters made during the ABRACOS expeditions. RA, Rocas Atoll; FN, Fernando de Noronha Archipelago



### 3.2.2 | Diagnostic features

*Gigantura chuni* has a relatively deep and compressed body; snout very short; horizontal tubular eyes directed forward; mouth extends to below middle of pectoral-fin base; jaw teeth moderate to long, depressible, arranged in two or three rows; pelvic fin absent in adults; a deeply forked caudal fin, with a greatly elongated lower lobe. It can be distinguished from its only congener, *Gigantura indica* Brauer 1901, by having 30–33 pectoral-fin rays (vs. 36–42), 8–10 anal-fin rays (vs. 11–14), head length 20.0%–30.0% SL (vs. 12.0%–19.0%), and depth of caudal peduncle 8.1%–11.0% SL (vs. 2.0%–3.6%) (Johnson & Bertelsen, 1991; McEachran & Fechhelm, 1998; Sutton *et al.*, 2020).

### 3.2.3 | Distribution

*Gigantura chuni* is meso- to bathypelagic, with a wide distribution in the tropical waters of the Atlantic, Pacific and Indian oceans, from 500 to 2000 m depth (Johnson & Bertelsen, 1991). Nearly all records are within approximately 20°N and 20°S. Previous records of this species are widespread in the Tropical Atlantic, except off Brazil. The three specimens reported herein were collected off Fernando de Noronha and Rocas Atoll, at 610 and 850 m depth (Figure 2b), and are the first records of *G. chuni* in Brazilian waters and in the western South Atlantic.

### 3.2.4 | Remarks

In addition to *G. chuni*, 30 specimens of *G. indica* (24–190 mm SL) were also collected in 17 collection stations of the ABRACOS expeditions. Most specimens of that species were collected along the Fernando de Noronha Ridge, but some specimens were also collected off states of Rio Grande do Norte, Paraíba and Pernambuco between depths of 50 and 1113 m (Figure 2b). Previous records of *G. indica* in Brazilian

waters are known off Rio Grande do Norte State (as *Rosaura rotunda* (Tucher 1954), junior synonym of *G. indica*), Paraíba State, Fernando de Noronha Archipelago and Trindade Island (Johnson & Bertelsen, 1991). Additional unpublished records of *G. indica* in Brazilian waters are also known off northern Bahia State, based on four specimens (MNRJ 49747, 49748, 49749, 42858; 175–230 mm SL) collected in June 2000 by the RV *Thalassa*, during the REVIZEE Program.


## 3.3 | *Ahliesaurus berryi* Bertelsen, Krefft & Marshall 1976 (Notosudidae)

### 3.3.1 | Specimen examined

NPM 4574, 1, 198 mm, sta. AB2/44A, 03°52'52.5"S, 32°17'33.3"W to 03°52'13.4"S, 32°16'28.0"W, 850 m, 28 April 2017, 12:44–13:17 h (Figures 1c and 3a).

### 3.3.2 | Diagnostic features

Body slender, elongate; snout relatively short (6.1%–7.9% SL), sharply pointed, somewhat depressed; jaw extends close to the rear of eye; jaw teeth small, no enlarged fangs; dorsal fin midway on body; adipose dorsal fin above anal fin; anal fin placed far back on body; body and head covered with large, deciduous, cycloid scales; pelvic fins inserted below or just in front of dorsal-fin origin; posterior infraorbitals as simple half tubes, without posterior expansion; ventral body wall thin and transparent, dark peritoneum clearly visible. It can be distinguished from its only congener, *Ahliesaurus brevis* Bertelsen, Krefft and Marshall 1976 (not recorded in the Atlantic), by having 47–50 vertebrae (vs. 42–46) and eight to 11 pyloric caeca (vs. four to eight) (Bertelsen *et al.*, 1976; Krefft, 1986; Sutton *et al.*, 2020).



**FIGURE 3** Records of species of (a) Notosudidae and (b) Scopelarchidae in Brazilian waters made during the ABRACOS expeditions. RA, Rocas Atoll; FN, Fernando de Noronha Archipelago



### 3.3.3 | Distribution

*Ahliesaurus berryi* is a meso- to bathypelagic species known from the subtropical and tropical waters of the Atlantic, Indian and Pacific oceans (Bertelsen *et al.*, 1976; Krefft, 1986, 1990; Moore *et al.*, 2003; Russell, 2016; Sutton *et al.*, 2020). In the western South Atlantic, the species was previously reported from about 10°S to 40°S, outside the Brazilian EEZ (Bertelsen *et al.*, 1976). The species was included in lists of Brazilian marine fishes based on its presumed occurrence in the region (Figueiredo & Menezes, 2003; Melo *et al.*, 2020), but the single specimen collected near the Fernando de Noronha Archipelago at 850 m depth represents the first confirmed record of *A. berryi* in Brazilian waters (Figure 3a).

### 3.3.4 | Remarks

A single specimen of *Scopelosaurus smithii* Bean 1925, another notosudid, was also collected during the ABRACOS expeditions, from off Pernambuco State, at 680 m depth (NPM 4573, 177 mm SL; Figure 3a). This species was previously recorded in Brazilian waters from off Bahia to Rio de Janeiro states, in addition to the Vitória-Trindade Seamount Chain and Trindade Island (Bertelsen *et al.*, 1976, fig. 30; Bonecker & Castro, 2006). Other species of Notosudidae recorded in the Brazilian EEZ are: *Luciosudis normani* Fraser-Brunner 1931, from off Santa Catarina and Rio Grande do Sul states (Bertelsen *et al.*, 1976, fig. 63; Figueiredo *et al.*, 2002); *Scopelosaurus argenteus* (Maul 1954), from off Fernando de Noronha and Saint Peter and Saint Paul archipelagos (MCZ 83603: 01°02'30"S, 34°58'30"W; MCZ 83604–83605: 01°13'30"S, 34°35'30"W; MCZ 83606: 00°26'30"N 32°06'30"W; Bertelsen *et al.*, 1976, fig. 33); and *Scopelosaurus herwigii* Bertelsen, Krefft & Marshall 1976, from off Espírito Santo (DZUFRJ 8979: 21°12'00"S, 37°46'27"W; Bonecker & Castro, 2006) and Rio Grande do Sul (Figueiredo *et al.*, 2002) states, with additional records outside the Brazilian EEZ between 25°S and 33°S (Bertelsen *et al.*, 1976).

## 3.4 | *Benthalbella infans* Zugmayer 1911 (Scopelarchidae)

### 3.4.1 | Specimen examined

NPM 3553, 1, 57 mm, sta. AB1/26, 06°09'13.5"S, 34°34'34.0"W to 06°09'49.6"S, 34°33'44.5"W, 560 m, 12 October 2015, 12:48–13:08 h (Figures 1d and 3b).

### 3.4.2 | Diagnostic features

*Benthalbella infans* is distinguished by a moderately elongated, compressed body; eyes large, tubular, with glistening white oval lens on side of eye tube; mouth large, extending beyond posterior margin of

eye; adipose dorsal fin present; gill-rakers absent, replaced by tooth plates on anterior arches; 55–62 large lateral line scales, each scale consisting of a bony plate with a distinct bony shelf behind the pore that mostly or completely covers the pore; four luminous organs on the ventral surface of body, the first on the isthmus (IO), the second just anterior to the pelvic fin base (POa), the third just posterior to the pelvic fin (POp) and the fourth immediately anterior to the anal fin (AO); dorsal-fin base short, with nine (rarely eight) rays; 25–28 pectoral-fin rays; 20–25 anal-fin rays (Johnson, 1974a, 1986; Merrett *et al.*, 1971, 1973; Stewart, 2015; Sutton *et al.*, 2020).

### 3.4.3 | Distribution

*Benthalbella infans* is meso- to bathypelagic, with a wide distribution in the tropical and warm-temperate waters of the Atlantic, Pacific and Indian oceans, mainly between 45°N and 35°S (Johnson, 1974a, 1982, 1986; Merrett *et al.* 1973; Russell, 2016; Sutton *et al.*, 2020). The species was reported from the western South Atlantic by Johnson (1974a: ISH 1437–1968, 30°01'S, 42°30'W), Menezes and Figueiredo (2003: MCZ 70342, 30°36'S, 41°56'W), and Judkins and Haedrich (2018: RGB 1423, 12°12'S, 31°06'W; RGB 1431, 23°00'S, 32°12'W). Those records are outside the Brazilian EEZ, but Menezes and Figueiredo (2003) and Melo *et al.* (2020) included *B. infans* in their lists of Brazilian marine fishes based on the likely occurrence of the species in waters of the country, given its wide distribution in the Atlantic. The specimen collected at 560 m depth off Rio Grande do Norte State represents the first confirmed record of *B. infans* in the Brazilian EEZ (Figure 3b).

## 3.5 | *Rosenblattichthys hubbsi* Johnson 1974 (Scopelarchidae)

### 3.5.1 | Specimens examined

NPM 4554, 4, 40–100 mm, sta. AB2/21, 06°50'21.0"S, 34°18'24.0"W to 06°48'24.3"S, 34°20'29.5"W, 800 m, 16 April 2017, 11:25–12:12 h (Figures 1e and 3b).

### 3.5.2 | Diagnosis

*Rosenblattichthys hubbsi* is distinguished by a relatively short, deep and robust body; eyes large, tubular, with glistening white oval lens on side of eye tube; a streak of dense black pigment bordering anterior margin of eye; mouth large, extending beyond posterior margin of eye; adipose dorsal fin present; gill-rakers absent, replaced by tooth plates on anterior arches; 47–53 large lateral line scales, each scale consisting of a bony plate with a distinct bony shelf behind the pore that mostly or completely covers the pore; pectoral and pelvic fins elongated and equal in length, pelvic fins inserting slightly anterior to

dorsal-fin origin; 21–23 pectoral-fin rays; eight or nine dorsal-fin rays; 23–25 anal-fin rays; 49 vertebrae (Johnson 1974a, 1974b, 1982, 1986; McEachran & Fechhelm, 1998; Okiyama & Johnson, 1986; Sutton *et al.*, 2020). Species of *Rosenblattichthys* are also characterized by a unique configuration of accessory pigment areas or spots: *R. hubbsi* has only two pigment areas as its congener *R. nemotoi* Okiyama & Johnson 1986, but its dorsal pigmented area is located over the posterior one-third of the anal-fin base instead of over the anterior one-third of the anal-fin base in *R. nemotoi* (Okiyama & Johnson, 1986).

### 3.5.3 | Distribution

*Rosenblattichthys hubbsi* is meso- to bathypelagic, with a wide distribution in the subtropical and tropical waters of the Atlantic, Pacific and Indian oceans (Johnson, 1974a,b, 1982, 1986). The species was included in previous lists of Brazilian marine fishes (Melo *et al.*, 2020; Menezes & Figueiredo, 2003) based on adult specimens collected outside the country's EEZ (e.g., Bonecker & Castro, 2006; Johnson, 1974b, 1982; Judkins & Haedrich, 2018), and larvae collected off Pará (MCZ 70305: 01°00'N, 45°46'W; Judkins & Haedrich, 2018) and off southern Bahia states (DZUFRJ 3237; Bonecker & Castro, 2006). The four specimens collected off Paraíba State, at 800 m depth (Figure 3b), are the first confirmed records of *R. hubbsi* in Brazilian waters.

## 3.6 | *Scopelarchoides danae* Johnson 1974 (Scopelarchidae)

### 3.6.1 | Specimen examined

NPM 4555, 1, 80 mm, sta. AB2/42A, 03°15'28.1"S, 31°48'29.1"W to 03°15'27.8"S, 31°50'40.6"W, 780 m, 27 April 2017, 12:23–12:26 h (Figures 1f and 3b).

### 3.6.2 | Diagnostic features

*Scopelarchoides danae* is distinguished by a moderately elongated, compressed body; heavy dermal pigmentation forming a dark stripe ventral to lateral line that extends from the pelvic-fin region to the lower caudal-fin lobe, pigment absent or weakly developed dorsal to lateral line and on upper caudal-fin lobe; eyes large, tubular, with glistening white oval lens on side of eye tube; mouth large, extending beyond posterior margin of eye; adipose dorsal fin present; gill-rakers absent, replaced by tooth plates on anterior arches; 50–52 large lateral line scales, each scale consisting of a bony plate with a distinct bony shelf behind the pore that mostly or completely covers the pore; pelvic fin distinctly longer than pectoral fin, origin of pelvic fin under anterior one-third of dorsal-fin base; 20–22 pectoral-fin rays; 6–9 (usually 7 or 8) dorsal-fin rays; 24–27 anal-fin rays; 48–50 vertebrae (McEachran & Fechhelm, 1998; Johnson, 1974a,b, 1986; Stewart, 2015; Sutton *et al.*, 2020).

### 3.6.3 | Distribution

*Scopelarchoides danae* is a mesopelagic species with a wide distribution in the tropical and temperate waters of the Atlantic, Pacific and Indian oceans (Johnson, 1974a,b, 1982, 1990). The species was included in previous lists of Brazilian marine fishes (Melo *et al.*, 2020; Menezes & Figueiredo, 2003) based on specimens collected outside the country's EEZ, off northern and north-eastern Brazil (MCZ 70371, 70380; Johnson, 1982). The single specimen reported here from off Fernando de Noronha Archipelago, at 780 m depth, is the first confirmed record of *S. danae* in Brazilian waters (Figure 3b).

### 3.6.4 | Remarks

In addition to *B. infans*, *R. hubbsi* and *S. danae*, two species of the Scopelarchidae previously recorded in Brazilian waters were also collected during the ABRACOS expeditions: *Scopelarchus analis* (Brauer 1902), from off Saint Peter and Saint Paul Archipelago (MCZ 127130: 00°56'24"N, 29°37'00"W; MCZ 69323: 00°58'S, 27°34'W) and Trindade Island (MCZ 69284: 18°10'S, 29°40'W) (Johnson, 1982), and *Scopelarchus guentheri* Alcock 1896, from off Pará State (MCZ 70953: 01°24'N, 45°24'W). During the ABRACOS expeditions, two specimens of *S. analis* (NPM 3260, 3261; 91–115 mm SL) were collected around the Rocas Atoll between depths of 510 and 525 m. In addition, eight specimens of *S. guentheri* (NPM 4553, 4556, 4557, 4558, 4559; 38–113 mm SL) were collected off Rio Grande do Norte and Paraíba states and in the vicinities of the Rocas Atoll, between depths of 610 and 1030 m (Figure 3b).

*Scopelarchus michaelisarsii* Koefoed 1955 is another scopelarchid reported from Brazil (Melo *et al.*, 2020; Menezes & Figueiredo, 2003). The occurrence of the species in Brazilian waters still require confirmation as all known records of the species are outside the country's EEZ (e.g., MCZ 70969–71; ISH 1259–1968; ISH 1087–1968).

## 3.7 | *Scopelengys tristis* Alcock 1890 (Neoscopelidae)


### 3.7.1 | Specimen examined

NPM 4620, 1, 145 mm, sta. AB2/49A, 04°10'38.1"S, 33°16'07.4"W to 04°10'58.0"S, 33°15'03.8"W, 770–1020 m, 30 April 2017, 21:17–21:52 h; NPM 4621, 1, 98 mm, sta. AB2/42A, 03°15'28.1"S, 31°48'29.1"W to 03°15'27.8"S, 31°50'40.6"W, 780 m, 27 April 2017, 12:23–12:26 h (Figures 1g and 4a).

### 3.7.2 | Diagnostic features

Body elongated, compressed; dorsal-fin origin above pelvic-fin origin; dorsal adipose fin over end of anal fin; anal-fin origin well behind dorsal-fin base; pectoral and pelvic fins long, both reaching to anus; no photophores on body; eye small (11.8%–15.9% HL); upper jaw

**FIGURE 4** Records of species of (a) Neoscopelidae and (b) Trachipteridae in Brazilian waters made during the ABRACOS expeditions. RA, Rocas Atoll; FN, Fernando de Noronha Archipelago



extending beyond vertical through posterior margin of orbit; a patch of small teeth on each side of vomer; mesopterygoid teeth absent; scales cycloid, highly deciduous. It can be distinguished from its only congener, *Scopelengys clarkei* Butler & Ahlstrom 1976, by the number of pectoral-fin rays (14–17 vs. 12–13), number of vertebrae (29–32 vs. 34–35), and depth of caudal peduncle (5.6%–8.3% SL vs. 9.4%–10.2% SL) (Butler & Ahlstrom, 1976; Fujii, 1983; Hartel & Craddock, 2003; Nafpaktitis, 1977; Romero Camarena, 2009).

### 3.7.3 | Distribution

*Scopelengys tristis* is a meso- to batypelagic species with a wide distribution in the tropical and temperate waters of the Atlantic, Pacific and Indian oceans (Butler & Ahlstrom, 1976; Hartel & Craddock, 2003; Milkova et al., 2016; Nafpaktitis, 1977; Nakabo, 2002; Romero Camarena, 2009; Sutton et al., 2020). The species was reported by Menezes et al. (2003a) and Marceniuk et al. (2021) as occurring in Brazilian waters based on one specimen (MCZ 101792), which was actually collected in the western North Atlantic outside the country's EEZ. Therefore, the two specimens of *S. tristis* collected off Rocas Atoll and Fernando de Noronha Archipelago between depths of 770 and 1020 m are the first records of the species in the Brazilian EEZ (Figure 4a).

### 3.7.4 | Remarks

Other species of Neoscopelidae reported from Brazil are *Neoscopelus macrolepidotus* Johnson 1863 and *Neoscopelus microchir* Matsubara 1943. *Neoscopelus macrolepidotus* is recorded along the entire Brazilian coast (e.g., Haimovici et al., 1994; Asano Filho et al., 2005; Bernardes et al., 2005; Braga et al., 2007; Bonecker et al., 2012; Lins Oliveira et al., 2015), while *N. microchir* is known only from two specimens collected off southern Brazil (Figueiredo et al., 2002).

## 3.8 | *Zu cristatus* (Bonelli 1820) (Trachipteridae)

### 3.8.1 | Specimens examined

NPM 3186, 1, 86 mm, sta. AB1/5, 04°05'22.2"S, 32°10'51.6"W to 04°04'32.4"S, 32°11'55.2"W, 85 m, 2 October 2015, 21:18–21:48 h; NPM 3187, 1, 89 mm, sta. AB1/1, 03°46'18.6"S, 32°25'13.2"W to 03°46'03.0"S, 32°23'48.6"W, 150 m, 30 September 2015, 22:32–23:02 h; NPM 3188, 1, 78 mm, sta. AB1/22, 04°07'44.8"S, 33°47'24.5"W to 04°07'00.7"S, 33°48'57.9"W, 525 m, 8 October 2015, 21:32–22:12 h; NPM 3193, 1, 53 mm, sta. AB1/20, 03°45'18.6"S, 33°53'10.8"W to 03°44'32.4"S, 33°54'04.8"W, 60 m, 7 October 2015, 21:13–21:37 h; NPM 3309, 1, 10 mm, sta. AB1/11, 03°44'58.8"S, 33°13'39.6"W to 03°44'58.8"S, 33°12'09.0"W, 650 m, 5 October 2015, 12:03–12:47 h; NPM 3310, 1, 15 mm, sta. AB1/21, 03°39'27.4"S, 33°41'26.6"W to 03°38'33.6"S, 33°42'53.4"W, 75–115 m, 8 October 2015, 10:09–10:49 h; NPM 4504, 1, 430 mm, sta. AB2/48A, 04°25'05.3"S, 32°57'52.1"W to 04°25'24.9"S, 32°56'55.5"W, 505 m, 24 April 2017, 10:30–10:58 h; NPM 5033, 1, 69 mm, sta. AB2/39, 04°52'26.9"S, 34°35'22.9"W to 04°50'52.8"S, 34°51'04.7"W, 650–800 m, 24 April 2017, 21:49–22:37 h (Figures 1h and 4b).

### 3.8.2 | Diagnostic features

*Zu cristatus* is distinguished by an elongated and tapering body, with a wavy or scalloped ventral body margin; upper jaw highly protractile; head short and deep; dorsal fin scarlet, origin over posterior margin of orbit, consisting of about five or six elongated rays followed by about 125–145 shorter rays; posterior portion of lateral line running along ventral edge of tail as a series of sharp spines that point in alternating directions; caudal fin reddish black, darker posteriorly. It can be distinguished from its only congener, *Zu elongatus* Heemstra & Kannemeyer 1984, by the body depth in adults (20%–26% vs. 12%–16% SL), eye

diameter (13%–16% vs. 9%–10% of distance from snout to anus), and number of lateral-line scales (99–106 vs. 126–130) (Angulo & López-Sánchez, 2017; Heemstra & Kannemeyer, 1986; McEachran & Fechhelm, 1998; Olney & Hartel, 2016a; Sutton et al., 2020).

### 3.8.3 | Distribution

*Zu cristatus* is a mesopelagic species, with a wide distribution in the tropical and temperate waters of the Atlantic, Pacific and Indian oceans (Angulo & López-Sánchez, 2017; Olney & Hartel, 2016a; Sutton et al., 2020). The presumed occurrence of the species in Brazilian waters (e.g., Menezes et al., 2003b; Melo et al., 2020; Marcenik et al., 2021) was based on a few records of the species in the western Tropical Atlantic outside the country's EEZ (MCZ 58928; MCZ 84634; NSMT-P 40246) and also on larvae collected off Rio de Janeiro State (Bonecker et al., 2014) and near the Martin Vaz Archipelago (Stocco & Joyeux, 2015). The eight specimens reported here, collected between depths of 40 and 800 m off the Rio Grande do Norte, Rocas Atoll and Fernando de Noronha Archipelago, are the first confirmed records of juvenile and adult specimens of *Zu cristatus* in the Brazilian EEZ (Figure 4b).

### 3.8.4 | Remarks

Other species of Trachipteridae reported from Brazilian waters are *Desmodema polystictum* (Ogilby 1898), recorded off Rio de Janeiro and Rio Grande do Sul states (Costa & Mincarone, 2010; Figueiredo et al., 2002), and *Trachipterus jacksonensis* (Ramsay 1881), reported from off Rio de Janeiro to Rio Grande do Sul states (Figueiredo et al., 2002; Mancini et al., 2003; Menezes, 1971; Mincarone et al., 2001). One additional specimen of *Desmodema polystictum* was also collected during the ABRACOS expeditions (NPM 5055, 74 mm SL), at 850 m depth near the Fernando de Noronha Archipelago (Figure 4b). Additionally, five very small specimens of *Trachipterus* sp. (NPM 3206, 3326, 3331, 3369, 5050; 18–55 mm SL) were collected off the Rocas Atoll, Rio Grande do Norte and Pernambuco states (Figure 4b). They could not be identified at species level due to their small sizes but may represent *T. jacksonensis*, since this is so far the only species of the genus reported from the western South Atlantic (Mincarone et al., 2001).

## 3.9 | *Stylephorus chordatus* Shaw 1791 (Stylephoridae)

### 3.9.1 | Specimens examined

NPM 3178, 2, 170–255 mm, sta. AB1/14, 03°58'58.2"S, 34°03'22.2"W to 03°57'42.0"S, 34°04'52.2"W, 510 m, 6 October 2015, 21:40–22:26 h; NPM 4222, 7, 143–210 mm, sta. AB2/39, 04°52'26.9"S, 34°35'22.9"W to 04°50'52.8"S, 34°51'04.7"W, 650–800 m, 24 April 2017, 21:49–22:37 h; NPM 4100, 4, 112–168 mm,

sta. AB2/42A, 03°15'28.1"S, 31°48'29.1"W to 03°15'26.4"S, 31°48'22.9"W, 780 m, 27 April 2017, 12:23–12:26 h; NPM 4126, 5, 145–176 mm, sta. AB2/53A, 03°48'58.7"S, 33°59'17.1"W to 03°50'05.8"S, 33°58'46.5"W, 610 m, 2 May 2017, 22:08–22:40 h; NPM 4143, 5, 130–220 mm, sta. AB2/44A, 03°52'52.5"S, 32°17'33.3"W to 03°51'13.4"S, 32°16'28.0"W, 850 m, 28 April 2017, 12:44–13:17 h; NPM 4183, 4, 192–242 mm, sta. AB2/35, 04°19'36.6"S, 35°29'51.6"W to 04°18'32.4"S, 35°32'19.8"W, 630 m, 20 April 2017, 22:35–23:15 h; NPM 4212, 1, 220 mm, sta. AB2/52A, 03°43'16.2"S, 33°25'09.8"W to 03°42'14.2"S, 33°24'36.2"W, 822–984 m, 2 May 2017, 11:47–12:18 h; NPM 4213, 2, 139–165 mm, sta. AB2/40A, 03°31'21.0"S, 32°31'39.8"W to 03°31'30.8"S, 32°30'40.6"W, 440 m, 26 April 2017, 10:43–11:06 h; NPM 4244, 10, 90–250 mm, sta. AB2/41A, 03°19'59.1"S, 32°24'42.1"W to 03°19'31.8"S, 32°25'04.6"W, 430 m, 26 April 2017, 21:44–22:06 h; NPM 4433, 1, damaged, sta. AB2/41B, 03°19'15.7"S, 32°25'42.3"W to 03°19'13.8"S, 32°25'10.5"W, 25 m, 26 April 2017, 23:16–23:31 h; NPM 4435, 1, 71 mm, sta. AB2/46A, 04°08'31.3"S, 32°18'14.8"W to 04°08'56.5"S, 32°17'28.9"W, 360 m, 29 April 2017, 11:20–11:45 h; NPM 4454, 1, 111 mm, sta. AB2/46B, 04°10'29.7"S, 32°16'04.2"W to 04°10'56.8"S, 32°15'29.0"W, 440 m, 29 April 2017, 13:08–13:34 h; NPM 4602, 1, 121 mm, sta. AB2/52B, 03°41'56.4"S, 33°23'29.4"W to 03°42'34.2"S, 33°22'36.1"W, 385 m, 2 May 2017, 14:00–14:30 h; NPM 5408, 11, 59–279 mm, sta. AB2/21, 06°50'21.0"S, 34°18'24.0"W to 06°48'24.3"S, 34°20'29.5"W, 800 m, 16 April 2017, 11:25–12:12 h (Figures 1i and 5a).

### 3.9.2 | Diagnosis


*Stylephorus chordatus* is the only species in the Stylephoriformes (e.g., Dornburg & Near, 2021; Hughes et al., 2018; Miya et al., 2007). It can be recognized by a slender, compressed, elongate body; eyes conspicuous, telescopic, directed forward and somewhat upward; jaws highly protrusible, mouth small and tubular, lower jaw extending in front of upper jaw; inside of mouth and gill chamber black; teeth absent or minute; dorsal-fin base long, with more than 100 rays, extending from nape to caudal fin, the first and second rays longer than remaining rays; anus immediately before anal-fin origin; pelvic-fin with just one ray, about below pectoral-fin base; anal fin short; caudal fin with two highly modified externally visible lobes, upper caudal-fin lobe with five or six short rays, lower with two long rays, forming a projection or tapering equal to or exceeding body length when intact (Hulley, 1986; Olney & Hartel, 2016b; Sutton et al., 2020). The genus is monotypic (Fricke et al., 2022).

### 3.9.3 | Distribution

*Stylephorus chordatus* is meso- to bathypelagic, distributed in the tropical and temperate waters of the Atlantic, Pacific and Indian oceans (Olney, 2003; Olney & Hartel, 2016b; Sutton et al., 2020). Records are widespread in the Atlantic Ocean, except in the western South Atlantic. The 53 specimens reported here were collected between



**FIGURE 5** Records of species of (a) Stylephoridae and (b) Chiasmodontidae in Brazilian waters made during the ABRACOS expeditions. RA, Rocas Atoll; FN, Fernando de Noronha Archipelago



depths of 25 and 900 m off the Fernando de Noronha Archipelago, Rocas Atoll, the seamounts of Fernando de Noronha Ridge, and off Rio Grande do Norte and Paraíba states (Figure 5a). They represent the first published records of *S. chordatus* in Brazilian waters, but additional specimens are also known from off Maranhão State (MCZ 147847: 01°12'N, 44°39'W), the Rocas Atoll (MCZ 147900: 01°13'S, 34°35'W; MCZ 147840: 01°02'S, 34°58'W), and the Saint Peter and Saint Paul Archipelago (MCZ 96864: 00°34'N, 30°43'W).

### 3.10 | *Pseudoscopus cordilluminatus* Melo 2010 (Chiasmodontidae)

#### 3.10.1 | Specimens examined

NPM 5027, 1, 31 mm, sta. AB2/6, 08°51'21.0"S, 34°35'40.8"W to 08°53'38.9"S, 34°36'19.0"W, 240 m, 11 April 2017, 10:53–11:55 h; NPM 4536, 1, 57 mm, sta. AB2/44A, 03°52'52.5"S, 32°17'33.3"W to 03°51'13.4"S, 32°16'28.0"W, 850 m, 28 April 2017, 12:44–13:17 h (Figures 1j and 5b).

#### 3.10.2 | Diagnosis

*Pseudoscopus cordilluminatus* is distinguished by an elongated and moderately compressed body; dorsal fin divided into two fins; head and body with photophores; belly moderately distensible; mouth terminal, large, nearly horizontal; premaxillary and dentary teeth curved posteriorly, not flared outwards; a long and broadened premaxilla, with a unique dentition pattern where the mesial series is organized in rows of two to three teeth, the first (outer) and second teeth much smaller than the innermost (oral) tooth; lateral series of premaxillary teeth without hooked teeth (Melo, 2010; Sutton *et al.*, 2020). According to Melo

(2010) *P. cordilluminatus* can be distinguished in the *Pseudoscopus scriptus* species group by the premaxillary dentition pattern (a single row in *P. pierbartus* and *P. obtusifrons*; teeth of the mesial series gradually increasing in size in *P. cephalus*, *P. sagamianus* and *P. scriptus*). Also according to Melo (2010), *P. cordilluminatus* can be further distinguished from *P. scriptus* by the series of anal-fin photophores (*saf*) extending anteriorly to about the anus region (vs. *saf* not extending anteriorly to anus), from *P. cephalus* by the vertebral count (total vertebrae 35–36, precaudal 18–19 vs. 31 and 14, respectively), and from *P. sagamianus* by melanophores inside mouth restricted to the region of tooth insertion (vs. internal area of mouth and gill arches completely black).

#### 3.10.3 | Distribution

*Pseudoscopus cordilluminatus* is a mesopelagic species from the equatorial and tropical eastern Atlantic (Melo, 2010; Prokofiev, 2014; Sutton *et al.*, 2020) and possibly the Indian Ocean (Melo, 2010; Prokofiev, 2014). The holotype was collected in the eastern Atlantic close to the Mid-Atlantic Ridge (MCZ 161017; 0°00'30.0"N, 19°45'30.0"W). The two specimens reported here, collected between depths of 240 and 850 m off the Pernambuco State and Fernando de Noronha Archipelago (Figure 5b), are the first records of the species in the western Atlantic and in Brazilian waters.

#### 3.10.4 | Remarks

In addition to *P. cordilluminatus*, other 13 rare species of the Chiasmodontidae are reported from Brazilian waters (Figueiredo & Menezes, 2003; Melo, 2008, 2009, 2010; Séret & Andreata, 1992). Four of those species were also collected during the ABRACOS expeditions:

*Chiasmodon niger* Johnson 1864, *C. braueri* Weber 1913, *Kali kerberti* (Weber 1913), and *Pseudoscopelus scutatus* Krefft 1971.

*Chiasmodon niger* was previously reported from Brazilian waters based on a single specimen (USU 1364, 134 mm SL; apparently lost) collected off Espírito Santo State (21°24.16'S, 39°56.19'W, 1320–1360 m depth; Melo, 2009; Séret & Andreata, 1992), and another specimen (BMNH 1879.5.14.588, 50.6 mm SL) collected off Saint Peter and Saint Paul Archipelago (01°22'N, 26°36'W; Melo, 2009). The identification of other specimens collected off Saint Peter and Saint Paul Archipelago (ISH 1711–1966: 01°22'N, 26°36'W) and Santa Catarina State (ISH 1817–1968: 29°57'S, 47°35'W) requires further corroboration as they were not reported by Melo (2009). In our survey, a single specimen (NPM 4537, 90 mm SL) of *C. niger* was collected at 850 m depth off eastern Fernando de Noronha Archipelago (Figure 5b).

Four specimens of *C. braueri* (53–171 mm SL), collected between depths of 1209 and 1929 off Bahia and Espírito Santo states, are known in Brazilian waters (Melo, 2009). Two additional specimens (NPM 4487, 70–95 mm SL) were identified in our collections from depths between 830 and 1030 m off western Rocas Atoll (Figure 5b).

*Kali kerberti* is known in Brazilian waters by a single specimen (MNRJ 26712, 133.8 mm SL) collected at 21°07'S, 39°46'W (Melo, 2008). Johnson and Cohen (1974) also reported on a specimen of *Kali normani* (Parr 1931) from north-eastern Brazil. This species is now considered a junior synonym of *K. kerberti* (Melo, 2008) or a subspecies of *K. kerberti* restricted to the East Pacific (Prokofiev, 2014). Six specimens of *K. kerberti* (NPM 4488, 4489, 4490, 4491; 69–170 mm SL) were identified in our collections. They were collected off Rio Grande do Norte State, Rocas Atoll and Fernando de Noronha Archipelago between depths of 650 and 1020 m (Figure 5b).

*Pseudoscopelus scutatus* was described from the tropical Atlantic (between 02°N and 28°S, west of 25°W), including some records off north-eastern Brazil, outside the EEZ (Figueiredo et al., 2003; Krefft, 1971). The species was subsequently recorded in Brazilian waters, based on one specimen (MNRJ 26700, 91 mm SL) collected off Espírito Santo State (21°25'S, 39°43'W) between depths of 1712–1721 m (Melo, 2010). Another specimen (NPM 985, 85 mm SL), collected between depths of 1210 and 1215 m from off Rio de Janeiro State (23°24'01"S, 40°59'00"W), is also known (Mincarone et al., 2017). During the ABRACOS expeditions two additional specimens of *P. scutatus* (NPM 4486, 4570; 67–75 mm SL) were collected between depths of 430 and 984 m off Rocas Atoll and the Fernando de Noronha Archipelago (Figure 5b).

## 4 | DISCUSSION

The United Nations Decade of Ocean Science for Sustainable Development (2021–2030) recognizes the deep sea as a frontier of science and discovery, and calls for research to advance understanding of this environment, its diversity, vulnerabilities and roles in the global ecosystems (Howell et al., 2020). However, there is an uneven capacity to conduct deep-ocean science across nations primarily due to differing

levels of funding capabilities. As a joint scientific collaboration between Brazil and France, the ABRACOS expeditions stand out in the context of international efforts that might help alleviate this situation.

When all published accounts of the ABRACOS expeditions are considered together (Afonso et al., 2021; Eduardo et al., 2018, 2019, 2021; Mincarone et al., 2019, 2020, 2021; Villarins et al., 2022; this study), a total of 50 species of deep-sea fishes are newly reported for Brazilian waters. Some of those records are also new for the western South Atlantic or larger areas, as the Atlantic Ocean. This high number of new records is based on just two collecting campaigns, reinforcing that an ecologically relevant portion of the deep-sea fish diversity of South America is still insufficiently known (e.g., Reis et al., 2016). For example, the first record of the enigmatic *Stylephorus chordatus* presented here for the western South Atlantic was based on 53 specimens collected in relatively few trawls, revealing that the species is likely common in the region. A similar situation was also recently reported for *Winteria telescopa* Brauer 1901 (Argentiniformes: Opisthoproctidae) by Mincarone et al. (2020). Other apparently abundant but still unrecorded species of deep-sea fishes were not collected in the region so far, most likely due to the paucity of scientific expeditions. This huge gap in basic knowledge hinders the proper management and conservation of deep-sea habitats, a situation that is particularly problematic in Brazil since the offshore exploration of oil and gas is one of the main economic activities of the country (e.g., ANP, 2021). Expansion of offshore economic activities worldwide is actually a trend in the following decades, and as a consequence proposals for mitigating anthropogenic impacts on deep-sea habitats will be required at an increasing pace, especially in the emerging context of the Blue Economy (Almada & Bernardino, 2017; Novaglio et al., 2021). Therefore, recent efforts towards increasing the knowledge of the deep-sea fish diversity of the western South Atlantic and off Brazil in particular should be celebrated (e.g., Bauer et al., 2021; Klautau et al., 2020; Melo et al., 2020, 2022; Schwarzhans et al., 2020; our studies). However, offshore oil companies and funding agencies in general should further increase the investment necessary for the proper understanding and management of this complex and still insufficiently known environment.

## ACKNOWLEDGEMENTS

We acknowledge the French oceanographic fleet for funding the at-sea survey ABRACOS, and the officers and crew of the RV *Antea* for their contribution to the success of the operations. For providing literature we thank Artem Prokofiev and Ron Fricke. We thank the CNPq – Brazilian National Council for Scientific and Technological Development, for providing research grants to M. M. Mincarone, F. Di Dario, T. Frédou and F. Lucena-Frédou (grants 314644/2020-2, PROTAX 443302/2020, 307422/2020-8 and 308554/2019-1, respectively). Thanks to CAPES – Coordination for the Improvement of Higher Education Personnel for providing a student scholarship to L. N. Eduardo, who is also supported by FUNBIO – Brazilian Fund for Biodiversity, and HUMANISE under the grant "Programa Bolsas Funbio – Conserando o Futuro 2018" (011/2019). This study is a contribution to the

LMI TAPIOCA, program CAPES/COFECUB (88881.142689/2017-01), and EU H2020 TRIATLAS project (817578). The NPM Fish Collection is supported by FUNBIO (grant 104/2016 – "Pesquisa Marinha e Pesqueira") and FAPERJ – Carlos Chagas Filho Foundation for Research Support of the State of Rio de Janeiro (grant E-26/210.290/2021).

## CONFLICTS OF INTEREST

The authors declare no conflict of interest.

## CONTRIBUTIONS

M.M.M.: data generation, data analysis, manuscript preparation. L.N.E.: data generation, data analysis, manuscript preparation. F.D.D.: data generation, data analysis, manuscript preparation. T.F.: data generation, funding. A.B.: data generation, funding. F.L.F.: data generation, funding. All authors have reviewed and approved the final version of the manuscript.

## ADDITIONAL MATERIAL EXAMINED

**GIGANTURIDAE:** *Gigantura indica* – NPM 4232, 2, 131–165 mm, sta. AB2/16, 07°36'14.4"S, 33°59'33.8"W to 07°36'49.3"S, 33°57'18.7"W, 680 m, 14 April 2017, 21:53–22:39 h; NPM 4233, 2, 175–190 mm, sta. AB2/39, 04°52'26.9"S, 34°35'22.9"W to 04°50'52.8"S, 34°51'04.7"W, 650–800 m, 24 April 2017, 21:49–22:37 h; NPM 4234, 3, 63–146 mm, sta. AB2/48B, 04°26'25.2"S, 32°56'15.4"W to 04°27'16.1"S, 32°55'42.1"W, 70 m, 30 April 2017, 12:15–12:45 h; NPM 4235, 1, 183 mm, sta. AB2/52A, 03°43'16.2"S, 33°25'09.8"W to 03°42'14.2"S, 33°24'36.2"W, 822–984 m, 2 May 2017, 11:47–12:18 h; NPM 4236, 4, 73–176 mm, sta. AB2/49A, 04°10'38.1"S, 33°16'07.4"W to 04°10'58.0"S, 33°15'03.8"W, 770–1020 m, 30 April 2017, 21:17–21:52 h; NPM 4237, 2, 142–170 mm, sta. AB2/40A, 03°31'21.0"S, 32°31'39.8"W to 03°31'30.8"S, 32°30'40.6"W, 440 m, 26 April 2017, 10:43–11:06 h; NPM 4238, 4, 50–164 mm, sta. AB2/44A, 03°52'52.5"S, 32°17'33.3"W to 03°51'13.4"S, 32°16'28.0"W, 850 m, 28 April 2017, 12:44–13:17 h; NPM 4239, 1, 65 mm, sta. AB2/41A, 03°19'59.1"S, 32°24'42.1"W to 03°19'31.8"S, 32°25'04.6"W, 430 m, 26 April 2017, 21:44–22:06 h; NPM 4242, 1, 60 mm, sta. AB2/52B, 03°41'56.4"S, 33°23'29.4"W to 03°42'34.2"S, 33°22'36.1"W, 385 m, 2 May 2017, 14:00–14:30 h; NPM 4405, 1, 74 mm, sta. AB2/53A, 03°48'58.7"S, 33°59'17.1"W to 03°50'05.8"S, 33°58'46.5"W, 610 m, 2 May 2017, 22:08–22:40 h; NPM 4431, 2, 54–61 mm, sta. AB2/42A, 03°15'28.1"S, 31°48'29.1"W to 03°15'26.4"S, 31°48'22.9"W, 780 m, 27 April 2017, 12:23–12:26 h; NPM 4434, 1, 84 mm, sta. AB2/42B, 03°15'43.6"S, 31°48'59.5"W to 03°15'26.4"S, 31°48'22.9"W, 50 m, 27 April 2017, 14:52–15:26 h; NPM 4451, 2, 54–64 mm, sta. AB2/44B, 03°52'18.4"S, 32°17'58.5"W to 03°52'46.4"S, 32°18'19.7"W, 130 m, 28 April 2017, 14:47–15:02 h; NPM 4894, 1, 58 mm, sta. AB2/59A, 03°38'01.6"S, 36°03'10.6"W to 03°38'07.9"S, 36°02'22.6"W, 700–1113 m, 5 May 2017, 21:57–22:37 h; NPM 4895, 1, damaged, sta. AB2/58A, 03°56'53.6"S, 36°06'13.9"W to 03°57'56.9"S, 36°09'27.9"W, 520 m, 5 May 2017, 11:58–12:29 h; NPM 5092, 1, 24 mm, sta. AB2/7, 08°46'29.4"S, 34°45'37.0"W to 08°46'28.3"S, 34°42'57.0"W, 112 m, 11 April

2017, 15:39–16:27 h; NPM 5411, 1, 56 mm, sta. AB2/21, 06°50'21.0"S, 34°18'24.0"W to 06°48'24.3"S, 34°20'29.5"W, 800 m, 16 April 2017, 11:25–12:12 h; MNRJ 49747, 1, 201 mm, RV *Thalassa*, sta. E-0496, 13°17.580'S, 38°17.599'W to 13°12.030'S, 38°14.873'W, 1635–1864 m, 7 June 2000; MNRJ 49748, 1, 230 mm, RV *Thalassa*, sta. E-522, 13°30.495'S, 38°38.977'W to 13°29.472'S, 38°37.943'W, 1044–1275 m, 21 June 2000; MNRJ 49749, 175 mm, RV *Thalassa*, sta. E-520, 13°21.837'S, 38°16.683'W to 13°26.455'S, 38°13.836'W, 1981–2271 m, 20 June 2000; MNRJ 42858, 211 mm, RV *Thalassa*, sta. E-519, 13°19.944'S, 38°19.654'W to 13°22.615'S, 38°21.960'W, 1726–1929 m, 20 June 2000.

**NOTOSUDIDAE:** *Scopelosaurus smithii* – NPM 4573, 1, 177 mm, sta. AB2/16, 07°36'15.0"S, 33°59'30.0"W to 07°36'49.3"S, 33°57'18.7"W, 680 m, 14 April 2017, 21:53–22:39 h.

**SCOPELARCHIDAE:** *Scopelarchus analis* – NPM 3260, 1, 115 mm, sta. AB1/14, 03°58'58.2"S, 34°03'22.2"W to 03°57'42.0"S, 34°04'52.2"W, 510 m, 6 October 2015, 21:40–22:26 h; NPM 3261, 1, 91 mm, sta. AB1/22, 04°07'44.8"S, 33°47'24.5"W to 04°07'00.7"S, 33°48'57.9"W, 525 m, 8 October 2015, 21:32–22:12 h. *Scopelarchus guentheri* – NPM 4553, 2, 89–110 mm, sta. AB2/53A, 03°48'58.7"S, 33°59'17.1"W to 03°50'05.8"S, 33°58'46.5"W, 610 m, 2 May 2017, 22:08–22:40 h; NPM 4556, 1, 87 mm, sta. AB2/52B, 03°41'56.4"S, 33°23'29.4"W to 03°42'34.2"S, 33°22'36.1"W, 385 m, 2 May 2017, 14:00–14:30 h; NPM 4557, 2, 100–113 mm, sta. AB2/35, 04°19'36.6"S, 35°29'51.6"W to 04°18'32.4"S, 35°32'19.8"W, 630 m, 20 April 2017, 22:35–23:15 h; NPM 4558, 1, 90 mm, sta. AB2/54B, 03°45'17.2"S, 34°41'04.0"W to 03°44'39.2"S, 34°40'04.5"W, 830–1030 m, 3 May 2017, 13:11–13:47 h; NPM 4559, 2, 38–48 mm, sta. AB2/21, 06°50'21.0"S, 34°18'24.0"W to 06°48'24.3"S, 34°20'29.5"W, 800 m, 16 April 2017, 11:25–12:12 h.

**TRACHIPTERIDAE:** *Desmodema polystictum* – NPM 5055, 1, 74 mm, sta. AB2/44A, 03°52'52.5"S, 32°17'33.3"W to 03°51'13.4"S, 32°16'28.0"W, 850 m, 28 April 2017, 12:44–13:17 h. *Trachipterus* sp. – NPM 3206, 1, 55 mm, AB1/14, 03°58'58.2"S, 34°03'22.2"W to 03°57'42.0"S, 34°04'52.2"W, 510 m, 6 October 2015, 21:40–22:26 h; NPM 3326, 1, 23 mm, AB1/13, 03°54'35.2"S, 33°50'55.8"W to 03°53'56.3"S, 33°51'56.3"W, 110 m, 6 October 2015, 11:03–11:38 h; NPM 3331, 1, 40 mm, AB1/23, 05°08'35.4"S, 34°42'52.2"W to 05°08'02.4"S, 34°44'40.2"W, 35–100 m, 9 October 2015, 10:35–11:20 h; NPM 3369, 1, 18 mm, AB1/21, 03°39'27.4"S, 33°41'26.6"W to 03°38'33.6"S, 33°42'53.4"W, 75–115 m, 8 October 2015, 10:09–10:49 h; NPM 5050, 1, 48 mm, AB2/7, 08°46'29.4"S, 34°45'37.0"W to 08°46'28.3"S, 34°42'57.0"W, 112 m, 11 April 2017, 15:39–16:27 h.

**CHIASMODONTIDAE:** *Chiasmodon braueri* – NPM 4487, 2, 70–95 mm, sta. AB2/54B, 03°45'17.2"S, 34°41'04.0"W to 03°44'39.2"S, 34°40'04.5"W, 830–1030 m, 3 May 2017, 13:11–13:47 h. *Chiasmodon niger* – NPM 4537, 1, 90 mm, sta. AB2/44A, 03°52'52.5"S, 32°17'33.3"W to 03°51'13.4"S, 32°16'28.0"W, 850 m, 28 April 2017, 12:44–13:17 h. *Kali kerberti* – NPM 4488, 2, 69–156 mm, sta. AB2/44A, 03°52'52.5"S, 32°17'33.3"W to 03°51'13.4"S, 32°16'28.0"W, 850 m, 28 April 2017, 12:44–13:17 h; NPM 4489,

1, 170 mm, sta. AB2/49A, 04°10'38.1"S, 33°16'07.4"W to 04°10'58.0"S, 33°15'03.8"W, 770–1020 m, 30 April 2017, 21:17–21:52 h; NPM 4490, 1, 126 mm, sta. AB2/39, 04°52'26.9"S, 34°35'22.9"W to 04°50'52.8"S, 34°51'04.7"W, 650–800 m, 24 April 2017, 21:49–22:37 h; NPM 4491, 2, 106–164 mm, sta. AB2/42A, 03°15'28.1"S, 31°48'29.1"W to 03°15'26.4"S, 31°48'22.9"W, 780 m, 27 April 2017, 12:23–12:26 h. *Pseudoscopelus scutatus* – NPM 4486, 1, 75 mm, sta. AB2/41A, 03°19'59.1"S, 32°24'42.1"W to 03°19'31.8"S, 32°25'04.6"W, 430 m, 26 April 2017, 21:44–22:06 h; NPM 4570, 2, 67 mm, sta. AB2/52A, 03°43'16.2"S, 33°25'09.8"W to 03°42'14.2"S, 33°24'36.2"W, 822–984 m, 2 May 2017, 11:47–12:18 h.

## ORCID

Michael Maia Mincarone  <https://orcid.org/0000-0003-0780-7854>

Leandro Nolê Eduardo  <https://orcid.org/0000-0003-2369-4175>

Fabio Di Dario  <https://orcid.org/0000-0001-9256-8485>

Thierry Frédou  <https://orcid.org/0000-0002-0510-6424>

Arnaud Bertrand  <https://orcid.org/0000-0003-4723-179X>

Flávia Lucena-Frédou  <https://orcid.org/0000-0001-5492-7205>

## REFERENCES

- Afonso, G. V. F., Di Dario, F., Eduardo, L. N., Lucena-Frédou, F., Bertrand, A., & Mincarone, M. M. (2021). Taxonomy and distribution of deep-sea bigscales and whalefishes (Teleostei: Stephanoberycoidei) collected off northeastern Brazil, including seamounts and oceanic islands. *Ichthyology & Herpetology*, 109, 467–488.
- Almada, G. V. M. B., & Bernardino, A. F. (2017). Conservation of deep-sea ecosystems within offshore oil fields on the Brazilian margin, SW Atlantic. *Biological Conservation*, 206, 92–101.
- Angulo, A., & López-Sánchez, M. I. (2017). New records of lampriform fishes (Teleostei: Lampriformes) from the Pacific coast of lower Central America, with comments on the diversity, taxonomy and distribution of the Lampriformes in the eastern Pacific Ocean. *Zootaxa*, 4236, 573–591.
- ANP (Agência Nacional do Petróleo, Gás Natural e Biocombustíveis). (2021). *Especial ANP 20 anos*. Available at: <https://www.gov.br/anp/pt-br/aceso-a-informacao/institucional/especial-anp-20-anos>
- Asano Filho, M., Holanda, F. C. A. F., Santos, F. J. S., & Cavalcante, T. S., Jr. (2005). *Recursos pesqueiros de grandes profundidades na costa norte do Brasil*. Brasília: Ibama.
- Bauer, A. B., Schwarzhans, W. W., Moura, R. L., Nunes, J. A. C. C., & Mincarone, M. M. (2021). A new species of viviparous brotula genus *Pseudogilbia* (Ophidiiformes: Dinematchthyidae) from Brazilian reefs, with an updated diagnosis of the genus. *Journal of Fish Biology*, 99, 1292–1298.
- Becker, V. E., Shcherbachev, Y. N., & Tchuvasov, V. M. (1975). Deep-sea pelagic fishes of the Caribbean Sea, Gulf of Mexico, and Puerto-Rican trench. *Trudy Instituta Okeanologii im. P. P. Shirshova, Akademiya Nauk SSSR*, 100, 289–336.
- Beebe, W. (1933). Deep-sea isospondylous fishes. Two new genera and four new species. *Zoologica*, 13, 159–167.
- Bernardes, R. A., Figueiredo, J. L., Rodrigues, A. R., Fischer, L. G., Vooren, C. M., Haimović, M., & Rossi-Wongtschowski, C. L. D. B. (2005). *Peixes da Zona Econômica Exclusiva da região sul-sudeste do Brasil. Levantamento com armadilhas, pargueiras e rede de arrasto de fundo*. São Paulo: Edusp.
- Bertelsen, E., Krefft, G., & Marshall, N. B. (1976). The fishes of the family Notosudidae. *Dana Report*, 86, 1–114.
- Bertrand, A. (2015). *ABRACOS cruise, RV Antea*. Available at: <https://doi.org/10.17600/15005600>.
- Bertrand, A. (2017). *ABRACOS 2 cruise, RV Antea*. Available at: <https://doi.org/10.17600/17004100>.
- Bonecker, A. C. T., & Castro, M. S. (2006). *Atlas de larvas de peixes da região central da Zona Econômica Exclusiva Brasileira*. Rio de Janeiro: Museu Nacional.
- Bonecker, A. C. T., Katsuragawa, M., Castro, M. S., Gomes, E. A. P., Namiki, C. A. P., & Zani-Teixeira, M. L. (2012). Larval fish of the Campos Basin, southeastern Brazil. *Check List*, 8, 1280–1291.
- Bonecker, A. C. T., Namiki, C. A. P., Castro, M. S., & Campos, P. N. (2014). Ordem Lampriformes. In *Catálogo dos estágios iniciais de desenvolvimento dos peixes da Bacia de Campos* (pp. 128–130). Curitiba: Sociedade Brasileira de Zoologia.
- Braga, A. C., Costa, P. A. S., Lima, A. T., Nunan, G. W., Olavo, G., & Martins, A. S. (2007). Padrões de distribuição de teleósteos epi- e mesopelágicos na costa central (11–22°S) brasileira. In P. A. S. Costa, G. Olavo, & A. S. Martins (Eds.), *Biodiversidade da fauna marinha profunda na costa central brasileira* (pp. 63–86). Rio de Janeiro: Museu Nacional.
- Braga, A. C., Costa, P. A. S., Martins, A. S., Olavo, G., & Nunan, G. W. A. (2014). Lanternfish (Myctophidae) from eastern Brazil, Southwest Atlantic Ocean. *Latin American Journal of Aquatic Research*, 42, 245–257.
- Butler, J. L., & Ahlstrom, E. H. (1976). Review of the deep-sea fish genus *Scopelogadus* (Neoscopelidae) with a description of a new species, *Scopelogadus clarkii*, from the Central Pacific. *Fishery Bulletin*, 74, 142–150.
- Carter, J. A., & Hartel, K. E. (2003). Alepocephalidae. In K. E. Carpenter (Ed.), *The living marine resources of the Western Central Atlantic. Volume 2: Bony fishes part 1 (Acipenseridae to Grammatidae)* (pp. 874–878). FAO Species Identification Guide for Fishery Purposes and American Society of Ichthyologist and Herpetologists Special Publication No. 5). Rome: FAO.
- Costa, P. A. S., & Mincarone, M. M. (2010). Ictiofauna demersal. In H. P. Lavrado, & A. C. S. Brasil (Eds.), *Biodiversidade da região oceânica profunda da Bacia de Campos: Megafauna e ictiofauna* (pp. 295–373). Rio de Janeiro: SAG Serv.
- Costa, P. A. S., Olavo, G., & Martins, A. S. (Eds.). (2007). *Biodiversidade da fauna marinha profunda na costa central brasileira*. Rio de Janeiro: Museu Nacional.
- Dornburg, A., & Near, T. J. (2021). The emerging phylogenetic perspective on the evolution of actinopterygian fishes. *Annual Review of Ecology, Evolution and Systematics*, 52, 427–452. <https://doi.org/10.1146/annurev-ecolsys-122120-122554>.
- Eduardo, L. N., Bertrand, A., Mincarone, M. M., Martins, J. R., Frédou, T., Vieira Assunção, R., ... Lucena-Frédou, F. (2021). Distribution, vertical migration, and trophic ecology of lanternfishes (Myctophidae) in the southwestern tropical Atlantic. *Progress in Oceanography*, 199, 102695. <https://doi.org/10.1016/j.pocean.2021.102695>.
- Eduardo, L. N., Bertrand, A., Mincarone, M. M., Santos, L. V., Frédou, T., Assunção, R. V., ... Lucena-Frédou, F. (2020a). Hatchetfishes (Stomiiformes: Stenopterychiidae) biodiversity, trophic ecology, vertical niche partitioning and functional roles in the western tropical Atlantic. *Progress in Oceanography*, 187, 102389. <https://doi.org/10.1016/j.pocean.2020.102389>.
- Eduardo, L. N., Lucena-Frédou, F., Mincarone, M. M., Soares, A., Le Loc'h, F., Frédou, T., ... Bertrand, A. (2020b). Trophic ecology, habitat, and migratory behavior of the viperfish *Chauliodus sloani* reveal a key mesopelagic player. *Scientific Reports*, 10, 20996. <https://doi.org/10.1038/s41598-020-77222-8>.
- Eduardo, L. N., Villarins, B. T., Lucena-Frédou, F., Frédou, T., Lira, A. S., Bertrand, A., & Mincarone, M. M. (2018). First record of the intermediate scabbardfish *Aphanopus intermedius* (Scombriformes: Trichiuridae) in the western South Atlantic Ocean. *Journal of Fish Biology*, 93, 992–995.



- Eduardo, L. N., Villarins, B. T., Martins, J. R., Lucena-Frédou, F., Frédou, T., Lira, A. S., ... Mincarone, M. M. (2019). Deep-sea oceanic basslets (Perciformes, Howellidae) from Brazil: New records and range extensions. *Check List*, 15, 965–971.
- Figueiredo, J. L., & Menezes, N. A. (2003). Família Notosudidae. In N. A. Menezes, P. A. Buckup, J. L. Figueiredo, & R. L. Moura (Eds.), *Catálogo das espécies de peixes marinhos do Brasil* (pp. 49). São Paulo: Museu de Zoologia da Universidade de São Paulo.
- Figueiredo, J. L., Santos, A. P., Yamaguti, N., Bernardes, R. A., & Rossi-Wongtschowski, C. L. D. B. (2002). *Peixes da Zona Econômica Exclusiva da região sudeste-sul do Brasil*. São Paulo: Edusp.
- Fitch, J. E., & Lavenberg, R. J. (1968). *Deep-water fishes of California*. Berkeley: University of California Press.
- Fricke, R., Eschmeyer, W. N., & R. van der Laan (Eds.) (2022). *Catalog of fishes: genera species, references*. <http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmain.asp>. Eletronic version accessed 20 June 2022.
- Fujii, E. (1983). Neoscopelidae. In T. Uyeno, K. Matsuura, & E. Fujii (Eds.), *Fishes trawled off Suriname and French Guiana* (pp. 188–189). Tokyo: Japan Marine Fishery Resource Research Center.
- Goodyear, R. H. (1969). Records of the alepocephalid fish *Photostylus pycnopterus* in the Indian and Pacific oceans. *Copeia*, 1969, 398–400.
- Grey, M. (1958). Second specimen of the bathypelagic fish *Photostylus pycnopterus*. *Copeia*, 1958, 56–57.
- Haimovici, M., Martins, A. S., Figueiredo, J. L., & Vieira, P. C. (1994). Demersal bony fish of the outer shelf and upper slope of the southern Brazil subtropical convergence ecosystem. *Marine Ecology Progress Series*, 108, 59–77.
- Hartel, K. E., & Craddock, J. E. (2003). Neoscopelidae. In K. E. Carpenter (Ed.), *The living marine resources of the western Central Atlantic. Volume 2: Bony fishes part 1 (Acipenseridae to Grammatidae)* (pp. 942–943). FAO Species Identification Guide for Fishery Purposes and American Society of Ichthyologists and Herpetologists Species Publication No. 5). Rome: FAO.
- Heemstra, P. C., & Kannemeyer, S. X. (1986). Trachipteridae. In M. M. Smith, & P. C. Heemstra (Eds.), *Smiths' sea fishes* (pp. 399–402). Berlin: Springer-Verlag.
- Howell, K. L., Hilário, A., Allcock, A. L., Bailey, D., Baker, M., Clark, M. R., ... Xavier, J. R. (2020). A decade to study deep-sea life. *Nature Ecology and Evolution*, 5, 265–267.
- Hughes, L. C., Ortí, G., Huang, Y., Sun, Y., Baldwin, C. C., Thompson, A. W., ... Shi, Q. (2018). Comprehensive phylogeny of ray-finned fishes (Actinopterygii) based on transcriptomic and genomic data. *PNAS*, 115, 6249–6254.
- Hulley, P. A. (1986). Stylephoridae. In M. M. Smith & P. C. Heemstra (Eds.), *Smiths' sea fishes* (p. 404). Berlin: Springer-Verlag.
- Johnson, R. K. (1974a). A revision of the alepisauroid family Scopelarchiidae (Pisces: Myctophiformes). *Fieldiana (Zoology)*, 66, 1–249.
- Johnson, R. K. (1974b). Five new species and a new genus of Alepisauroid fishes of the Scopelarchidae (Pisces: Myctophiformes). *Copeia*, 1974, 449–457.
- Johnson, R. K. (1982). Fishes of the families Evermannellidae and Scopelarchidae: Systematics, morphology, interrelationships, and zoogeography. *Fieldiana (Zoology)*, 12, 1–252.
- Johnson, R. K. (1986). Scopelarchidae. In M. M. Smith & P. C. Heemstra (Eds.), *Smiths' sea fishes* (pp. 265–267). Berlin: Springer-Verlag.
- Johnson, R. K. (1990). Scopelarchidae. In J. C. Quero, J. C. Hureau, C. Karrer, A. Post, & L. Saldanha (Eds.), *Check-list of the fishes of the eastern tropical Atlantic (CLOFETA)* (pp. 393–397). Lisbon: JNICT; Paris: SEI & UNESCO.
- Johnson, R. K., & Bertelsen, E. (1991). The fishes of the family Giganturidae: Systematics, development, distribution and aspects of biology. *Dana Reports*, 91, 1–45.
- Johnson, R. K., & Cohen, D. M. (1974). Results of the research cruises of FRV "Walther Herwig" to South America. XXX. Revision of the chiasmodontid fish genera *Dysalotus* and *Kali*, with descriptions of two new species. *Archiv für Fischereiwissenschaft*, 25, 13–46.
- Judkins, D. C., & Haedrich, R. L. (2018). The deep scattering layer micro-nektonic fish faunas of the Atlantic mesopelagic ecoregions with comparison of the corresponding decapod shrimp faunas. *Deep Sea Research Part I: Oceanographic Research Papers*, 136, 1–30.
- Kashkin, N. I. (1975). Finds of *Photostylus pycnopterus* Beebe, 1933 (Pisces, Alepocephalidae) in the northeastern Atlantic Ocean and in the Gulf of Mexico. *Voprosy Ichtiologii*, 15, 911–914.
- Klautau, A. G. C. M., Cintra, I. H. A., Rotundo, M. M., Carvalho-Filho, A., Caires, R. A., & Marceniuk, A. P. (2020). The deep sea teleost fish fauna of the Brazilian north coast. *Neotropical Ichthyology*, 18, e200030. <https://doi.org/10.1590/1982-0224-2020-0030>.
- Krefft, G. (1971). Ergebnisse der Forschungsreisen des FFS "Walther Herwig" nach Südamerika. XVIII. *Pseudoscopelus scutatus* spec. nov. (Pisces, Perciformes, Trachinoidei, Chiasmodontidae), ein neuer Kreuz-zahnbarsch aus dem tropischen Atlantik. *Archiv für Fischereiwissenschaft*, 22, 165–174.
- Krefft, G. (1986). Notosudidae. In M. M. Smith, & P. C. Heemstra (Eds.), *Smiths' sea fishes* (pp. 268–270). Berlin: Springer-Verlag.
- Krefft, G. (1990). Notosudidae. In J.-C. Quéro, J.-C. Hureau, C. Karrer, A. Post, & L. Saldanha (Eds.), *Check-list of the fishes of the eastern tropical Atlantic. CLOFETA* (pp. 361–364). Paris: UNESCO.
- Lima, A. T., Costa, P. A. S., Braga, A. C., Nunan, G. W. A., & Mincarone, M. M. (2011). Fishes of the family Sternoptychidae (Stomiiformes) collected on the Brazilian continental slope between 118 and 238S. *Zootaxa*, 2742, 34–48.
- Lins Oliveira, J. E., Nóbrega, M. F., Garcia Júnior, J., Sampaio, C. L. S., Di Dario, F., Fischer, L. G., & Mincarone, M. M. (2015). *Biodiversidade marinha da Baía Potiguar/RN: Peixes do talude continental*. Rio de Janeiro: Museu Nacional.
- Mancini, P. L., Amorim, A. F., & Arfelli, C. A. (2003). Observações em *Trachipterus jacksonensis* capturados no Brasil. In *Anais do Congresso Brasileiro de Pesquisas Ambientais e Saúde* (pp. 56–58). Santos: Congresso Brasileiro de Pesquisas Ambientais e Saúde.
- Marceniuk, A. P., Caires, R. A., Carvalho-Filho, A., Klautau, A. G. C. M., Santos, W. C. R., Wosiacki, W. B., ... Rotundo, M. M. (2021). Teleostei fishes of the north coast of Brazil. *Revista CEPISUL – Biodiversidade e Conservação Marinha*, 10, e2021006. <https://doi.org/10.37002/revistacepsul.vol10.1969e2021006>.
- McEachran, J. D., & Fechhelm, J. D. (1998). *Fishes of the Gulf of Mexico. Volume 1: Myxiniformes to Gasterosteiformes*. Austin: University of Texas Press.
- Melo, M. R. S. (2008). The genus *Kali* Lloyd (Chiasmodontidae: Teleostei) with description of new two species, and the revalidation of *K. kerberti* Weber. *Zootaxa*, 1747, 1–33.
- Melo, M. R. S. (2009). Revision of the genus *Chiasmodon* (Acanthomorpha: Chiasmodontidae), with the description of two new species. *Copeia*, 2009, 583–608.
- Melo, M. R. S. (2010). A revision of the genus *Pseudoscopelus* Lutken (Chiasmodontidae: Acanthomorpha) with descriptions of three new species. *Zootaxa*, 2710, 1–78.
- Melo, M. R. S., Caires, R. A., & Sutton, T. T. (2020). The scientific explorations of the deep-sea fishes in Brazil: The known knowns, the known unknowns, and the unknown unknowns. In P. Y. G. Sumida, A. F. Bernardino, & F. C. De Léo (Eds.), *Brazilian deep-sea biodiversity* (pp. 153–216). Berlin/Heidelberg: Springer.
- Melo, M. R. S., Gomes, A. A., Møller, P. R., & Nielsen, J. G. (2022). A new species of the rare deep-sea genus *Sciadonus* Garman, 1899 (Teleostei, Bythitidae) from off Brazil, with a discussion of the evolution of troglomorphy and miniaturization in the aphyonid clade. *Deep Sea Research Part I: Oceanographic Research Papers*, 180, 103684. <https://doi.org/10.1016/j.dsr.2021.103684>.
- Menezes, N. A. (1971). First record of *Trachipterus* in the Atlantic coast of South America (Pisces, Trachipteridae). *Papéis Avulsos de Zoologia*, 23, 205–207.

- Menezes, N. A., & Figueiredo, J. L. (2003). Família Scopelarchidae. In N. A. Menezes, P. A. Buckup, J. L. Figueiredo, & R. L. Moura (Eds.), *Catálogo das espécies de peixes marinhos do Brasil* (p. 49). São Paulo: Museu de Zoologia da Universidade de São Paulo.
- Menezes, N. A., Figueiredo, J. L., & Santos, A. P. (2003a). Família Neoscolopelidae. In N. A. Menezes, P. A. Buckup, J. L. Figueiredo, & R. L. Moura (Eds.), *Catálogo das espécies de peixes marinhos do Brasil* (pp. 51–52). São Paulo: Museu de Zoologia da Universidade de São Paulo.
- Menezes, N. A., Figueiredo, J. L., & Santos, A. P. (2003b). Família Trachipteridae. In N. A. Menezes, P. A. Buckup, J. L. Figueiredo, & R. L. Moura (Eds.), *Catálogo das espécies de peixes marinhos do Brasil* (p. 58). São Paulo: Museu de Zoologia da Universidade de São Paulo.
- Merrett, N. R., Badcock, J., & Herring, P. J. (1971). Observations on bioluminescence in a scopelarchid fish, *Benthalbella*. *Deep Sea Research and Oceanographic Abstracts*, 18, 1265–1267.
- Merrett, N. R., Badcock, J., & Herring, P. J. (1973). The status of *Benthalbella infans* (Pisces: Myctophidae), its development, bioluminescence, general biology and distribution in the eastern North Atlantic. *Journal of Zoology*, 170, 1–48.
- Milkova, V., Hanke, G., Gillespie, G., Fong, K., Boutillier, J., & Bedard, J. (2016). Range records for ten species of stomiiform, aulopiform, and myctophiform fishes in British Columbia, Canada. *Northwestern Naturalist*, 97, 113–123.
- Mincarone, M. M., Afonso, G. V. F., Di Dario, F., Eduardo, L. N., Frédou, T., Lucena-Frédou, F., ... Pietsch, T. W. (2021). Deep-sea anglerfishes (Lophiiformes: Ceratioidei) from off northeastern Brazil, with remarks on the ceratioids reported from the Brazilian Exclusive Economic Zone. *Neotropical Ichthyology*, 19, e200151. <https://doi.org/10.1590/1982-0224-2020-0151>.
- Mincarone, M. M., Di Dario, F., & Costa, P. A. S. (2014). Deep-sea bigscales, pricklefishes, gibberfishes and whalefishes (Teleostei: Stephanoberycoidei) off Brazil: New records, range extensions for the southwestern Atlantic Ocean and remarks on the taxonomy of *Poromitra*. *Journal of Fish Biology*, 85, 1546–1570.
- Mincarone, M. M., Lima, A. T., & Soto, J. M. R. (2001). Sobre a ocorrência do peixe-fita *Trachipterus jacksonensis* (Ramsay, 1881) (Lampridiformes, Trachipteridae) na costa brasileira. *Mare Magnum*, 1, 121–124.
- Mincarone, M. M., Martins, A. S., Costa, P. A. S., Braga, A. C., & Haimocivi, M. (2017). Peixes marinhos da Bacia de Campos: uma revisão da diversidade. In M. P. Curbelo-Fernandez & A. C. Braga (Eds.), *Comunidades demersais e bioconstrutores: caracterização ambiental regional da Bacia de Campos, Atlântico Sudoeste* (Vol. 4, pp. 187–216). Rio de Janeiro: Elsevier.
- Mincarone, M. M., Martins, J. R., Di Dario, F., Eduardo, L. N., Frédou, T., Lucena-Frédou, F., & Bertrand, A. (2020). Deep-sea smelts, pencil smelts, and barreleyes (Teleostei: Argentiniformes) from oceanic islands and seamounts off northeastern Brazil. *Marine Biology Research*, 16, 762–773.
- Mincarone, M. M., Villarins, B. T., Eduardo, L. N., Caires, R. A., Lucena-Frédou, F., Frédou, T., ... Bertrand, A. (2019). Deep-sea lanternfishes (Perciformes: Caristiidae) from oceanic islands and seamounts off northeastern Brazil, with comments on the caristiids previously reported in Brazilian waters. *Marine Biology Research*, 15, 297–304.
- Miya, M., Holcroft, N. I., Satoh, T. P., Yamaguchi, M., Nishida, M., & Wiley, E. O. (2007). Mitochondrial genome and a nuclear gene indicate a novel phylogenetic position of deep-sea tube-eye fish (Stylephoridae). *Ichthyological Research*, 54, 323–332.
- Moore, J. A., Hartel, K. E., Craddock, J. E., & Galbraith, J. K. (2003). An annotated list of deepwater fishes from off the New England region, with new area records. *Northeastern Naturalist*, 10, 159–248.
- Mundy, B. C. (2005). Checklist of the fishes of the Hawaiian archipelago. *Bishop Museum Bulletins in Zoology*, 6, 1–703.
- Nafpaktitis, B. G. (1977). Family Neoscolopelidae. In R. H. Gibbs, Jr., F. H. Berry, W. N. Eschmeyer, J. E. Böhlke, G. W. Mead, D. M. Cohen, et al. (Eds.), *Fishes of the western North Atlantic. Part 7: Order Iniomi* (Myctophiformes) (pp. 1–12). New Haven: Memoir, Sears Foundation for Marine Research.
- Nakabo, T. (2002). Neoscolopelidae. In T. Nakabo (Ed.), *Fishes of Japan with pictorial keys to the species* (English ed., p. 377). Tokyo: Tokai University Press.
- Nakamura, K., & Okamura, O. (1995). Alepocephalidae. In O. Okamura, K. Amaoka, M. Takeda, K. Yano, K. Okada, & S. Chikuni (Eds.), *Fishes collected by the R/V Shinkai Maru around Greenland* (pp. 72–80). Tokyo: Japan Marine Fishery Resources Research Center.
- Novaglio, C., Bax, N., Boschetti, F., Emad, G. R., Frusher, S., Fullbrook, L., ... Fulton, E. A. (2021). Deep aspirations: Towards a sustainable offshore blue economy. *Reviews in Fish Biology and Fisheries*, 32, 209–230. <https://doi.org/10.1007/s11160-020-09628-6>.
- Okiyama, M., & Johnson, R. K. (1986). *Rosenblattichthys nemotoi*, a new species of Scopelarchidae, from the South Indian Ocean sub-tropical convergence zone. *Japanese Journal of Ichthyology*, 32, 409–412.
- Olney, J. E. (2003). Stylephoridae. In K. E. Carpenter (Ed.), *The living marine resources of the western Central Atlantic. Volume 2: Bony fishes part 1 (Acipenseridae to Grammatidae)* (p. 953). FAO Species Identification Guide for Fishery Purposes and American Society of Ichthyologists and Herpetologists Species Publication No. 5). Rome: FAO.
- Olney, J. E., & Hartel, K. E. (2016a). Trachipteridae. In K. E. Carpenter & N. De Angelis (Eds.), *The living marine resources of the eastern Central Atlantic. Volume 3: Bony fishes part 1 (Elopiformes to Scorpaeniformes)*. FAO Species Identification Guide for Fishery Purposes. Rome: FAO.
- Olney, J. E., & Hartel, K. E. (2016b). Stylephoridae. In K. E. Carpenter & N. De Angelis (Eds.), *The living marine resources of the eastern Central Atlantic. Volume 3: Bony fishes part 1 (Elopiformes to Scorpaeniformes)*. FAO Species Identification Guide for Fishery Purposes. Rome: FAO.
- Parin, N. V., Pokhilskaia, G. N., Sazonov, Y. I., & Fedoryako, B. I. (1976). Rare and poorly known midwater fishes from the central and eastern equatorial Pacific Ocean. *Trudy Instituta Okeanologii*, 104, 206–236.
- Prokofiev, A. M. (2014). Swallowfishes (Chiasmodontidae) of the East Pacific. *Journal of Ichthyology*, 54, 631–641.
- Quérou, J.-C. (1974). Capture d'une espèce nouvelle pour la faune de l'Atlantique nord-est *Photostylus pycnopterus* Beebe, 1933 Poissons, Clupeiformes, Alepocephalidae. *Revue des Travaux de l'Institut des Pêches Maritimes*, 38, 435–436.
- Reis, R. E., Albert, J. S., Di Dario, F., Mincarone, M. M., Petry, P., & Rocha, L. A. (2016). Fish biodiversity and conservation in South America. *Journal of Fish Biology*, 89, 12–47.
- Rivaton, J., Fourmanoir, P., Bourret, P., & Kulbicki, M. (1990). *Catalogue des poissons de Nouvelle-Calédonie. Rapport provisoire*. Nouméa: Sciences de la Mer: Biologie Marine, ORSTOM.
- Romero Camarena, M. (2009). Neoscolopelidae. In K. Nakauia, M. Yabe, H. Imamura, M. Romero Camarena, & M. Yoshida (Eds.), *Deep-sea fishes of Peru* (p. 143). Tokyo: Japan Deep Sea Trawler Association.
- Russell, B. C. (2016). Scopelarchidae. In K. E. Carpenter, & N. De Angelis (Eds.), *The living marine resources of the eastern Central Atlantic. Volume 3: Bony fishes part 1 (Elopiformes to Scorpaeniformes)*. FAO Species Identification Guide for Fishery Purposes. Rome: FAO.
- Sabaj, M. H. (2020). Codes for natural history collections in ichthyology and herpetology. *Copeia*, 108, 593–669.
- Schwarzans, W. W., Mincarone, M. M., & Villarins, B. T. (2020). A new species of the genus *Verilus* (Teleostei, Percomorpha, Acropomatidae) from Brazil. *Zootaxa*, 4751, 589–596.
- Séret, B., & Andreata, J. V. (1992). Deep-sea fishes collected during the oceanographic cruise MD-55 off Brazil. *Cybium*, 16, 81–100.
- Stewart, A. L. (2015). Family Scopelarchidae. In C. D. Roberts, A. L. Stewart, & C. D. Struthers (Eds.), *The fishes of New Zealand* (pp. 566–574). Wellington: Te Papa Press.
- Stocco, L. B., & Joyeux, J.-C. (2015). Distribution of fish larvae on the Vitória-Trindade Chain, southwestern Atlantic. *Check List*, 11, 1590. <https://doi.org/10.15560/11.2.1590>.

- Sutton, T. T., Hulley, P. A., Wienerroither, R., Zaera-Perez, D., & Paxton, J. R. (2020). *Identification guide to the mesopelagic fishes of the central and Southeast Atlantic Ocean*. FAO species identification guide for fishery purposes. Rome: FAO.
- Tsukamoto, Y., Aizawa, M., & Okiyama, M. (1992). First record of the Alepocephalid fish, *Photostylus pycnopterus*, from Japan. *Japanese Journal of Ichthyology*, 39, 255–258.
- Villarins, B. T., Di Dario, F., Eduardo, L. N., Lucena-Frédou, F., Bertrand, A., Prokofiev, A. M., & Mincarone, M. M. (2022). Deep-sea dragonfishes (Teleostei: Stomiiformes) collected from off northeastern Brazil, with a review of the species reported from the Brazilian Exclusive Economic Zone. *Neotropical Ichthyology*, 20, e220004. <https://doi.org/10.1590/1982-0224-2022-0004>.
- Wisner, R. L. (1976). New data on the rare alepocephalid fish *Photostylus pycnopterus*. *Bulletin of the Southern California Academy of Sciences*, 75, 153–158.

**How to cite this article:** Mincarone, M. M., Eduardo, L. N., Di Dario, F., Frédou, T., Bertrand, A., & Lucena-Frédou, F. (2022). New records of rare deep-sea fishes (Teleostei) collected from off north-eastern Brazil, including seamounts and islands of the Fernando de Noronha Ridge. *Journal of Fish Biology*, 101(4), 945–959. <https://doi.org/10.1111/jfb.15155>