

Analysis of Time Series

Chapter 8: Smoothing in the time and frequency domains

Edward L. Ionides

Outline

- 1 Smoothing a time series
- 2 Seasonal adjustment in the frequency domain
 - The frequency response function
- 3 Loess smoothing
 - The frequency response function for loess
 - Business cycles in detrended economic data

Introduction to smoothing

Objectives

- 1 Estimating a nonparametric trend from a time series is known as smoothing. We will review some standard smoothing methods.
- 2 We can also smooth the periodogram to estimate a spectral density.
- 3 Many smoothers can be represented as linear filters. We will see that the statistical properties of linear filters for dependent (time-domain) stationary models can be conveniently studied in the frequency domain.

A motivating example

- The economy fluctuates between periods of rapid expansion and periods of slower growth or contraction.
- High unemployment is one of the most visible signs of a dysfunctional economy, in which labor is under-utilized, leading to hardships for many individuals and communities.
- Economists, politicians, businesspeople and the general public therefore have an interest in understanding fluctuations in unemployment.
- Economists try to distinguish between fundamental structural changes in the economy and the shorter-term cyclical booms and busts that appear to be a natural part of capitalist business activity.
- Monthly US unemployment figures are published by the [Bureau of Labor Statistics \(BLS\)](#).
- Measuring unemployment has subtleties, but these are not our immediate focus.

```
system("head unadjusted_unemployment.csv",intern=TRUE)
```


```
[1] "# Data extracted on: February 4, 2016 (10:06:56 AM)"
[2] "# from http://data.bls.gov/timeseries/LNU04000000"
[3] "# Labor Force Statistics from the Current Population Survey"
[4] "# Not Seasonally Adjusted"
[5] "# Series title:          (Unadj) Unemployment Rate"
[6] "# Labor force status:    Unemployment rate"
[7] "# Type of data:         Percent or rate"
[8] "# Age:                  16 years and over"
[9] "Year,Jan,Feb,Mar,Apr,May,Jun,Jul,Aug,Sep,Oct,Nov,Dec"
[10] "1948,4.0,4.7,4.5,4.0,3.4,3.9,3.9,3.6,3.4,2.9,3.3,3.6"
```

```
U1 <- read.table(file="unadjusted_unemployment.csv",
  sep="," ,header=TRUE)
head(U1,3)
```

Year	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
1948	4.0	4.7	4.5	4.0	3.4	3.9	3.9	3.6	3.4	2.9	3.3	3.6
1949	5.0	5.8	5.6	5.4	5.7	6.4	7.0	6.3	5.9	6.1	5.7	6.0
1950	7.6	7.9	7.1	6.0	5.3	5.6	5.3	4.1	4.0	3.3	3.8	3.9


Question 8.1. A coding exercise: Explain how the tabulated data in U1 are converted to a time series, below.

```
u1 <- t(as.matrix(U1[2:13]))  
dim(u1) <- NULL  
date <- seq(from=1948,length=length(u1),by=1/12)  
plot(date,u1,type="l",ylab="Unemployment rate (unadjusted)")
```



- We see seasonal variation, and perhaps we see business cycles on top of a slower trend.
- The seasonal variation looks like an additive effect, say an annual fluctuation with amplitude around 1 percentage point.
- Sometimes, we may prefer to look at monthly seasonally adjusted unemployment, [also provided by BLS](#).

Unemployment. Raw (black) and seasonally adjusted (red)




- We can wonder how the Bureau of Labor Statistics adjusts the data, and if this introduces any artifacts that a careful statistician should be aware of.

- To help understand the seasonal adjustment, we look at what it does to the smoothed periodogram.
- Using the `ts` class we can tell R the units of time.

```
u1_ts <- ts(u1,start=1948,frequency=12)
u2_ts <- ts(u2,start=1948,frequency=12)
spectrum(ts.union(u1_ts,u2_ts),spans=c(3,5,3),
  main="Unemployment: raw (black), seasonally adjusted (red)")
```

Unemployment: raw (black), seasonally adjusted (red)



Comments on the smoothed periodogram

- Note: For a report, we should add units to plots. Extra details (like bandwidth in the periodogram plot) should be explained or removed.

Question 8.2. What are the x-axis units?

Question 8.3. Comment on what you learn from comparing these smoothed periodograms.

The frequency response function

- The ratio of the periodograms of the smoothed and unsmoothed time series is the **frequency response** of the smoother.
- The frequency response function tells us how much the smoother contracts (or inflates) the sine and cosine components at each frequency ω .
- A frequency response may involve change in phase as well as magnitude, but here we consider only magnitude.
- Linear, time invariant transformations do not move power between frequencies, so they are characterized by their frequency response function.
- Smoothers are linear and time invariant, at least approximately. If we scale or shift the data, we expect the smoothed estimate to have the same scale or shift. We expect a smooth approximation to the sum of two time series to be approximately the sum of the two smoothed series.

Calculating a frequency response function

- We investigate the frequency response of the smoother used by Bureau of Labor Statistics to deseasonalize the unemployment data.


```
s <- spectrum(ts.union(u1_ts,u2_ts),plot=FALSE)
```

- We find the parts of `s` that we need to plot the frequency response.

```
names(s)
```

```
[1] "freq"      "spec"      "coh"       "phase"     "kernel"
[6] "df"        "bandwidth" "n.used"    "orig.n"    "series"
[11] "snames"    "method"    "taper"     "pad"       "detrend"
[16] "demean"
```

```
plot(s$freq,s$spec[,2]/s$spec[,1],type="l",log="y",
     ylab="frequency ratio", xlab="frequency",
     main="frequency response (red lines at 0.8 and 1.2)")
abline(h=c(0.8,1.2),col="red")
```




Question 8.4. What do you learn from this frequency response plot?

Estimating trend by Loess smoothing

- Loess is a **Local linear regression** approach (perhaps an acronym for L^ocal E^stimation by S^moothing) also known as **Lowess** (perhaps L^ocally W^eighted S^um of S^quares).
- At each point in time, Loess computes a linear regression (a constant, linear or quadratic trend estimate) using only neighboring times.
- We can imagine a moving window of points included in the regression.
- `loess` is an R implementation, with the fraction of points included in the moving window being scaled by the `span` argument.
- We can choose a value of the span that visually separates long term trend from business cycle.

A Loess smooth of unemployment


```
u1_loess <- loess(u1~date,span=0.5)
plot(date,u1,type="l",col="red")
lines(u1_loess$x,u1_loess$fitted,type="l")
```



Now, we compute the frequency response function for what we have done.

```
s2 <- spectrum(ts.union(  
  u1_ts, ts(u1_loess$fitted, start=1948, frequency=12)),  
  plot=FALSE)  
plot(s2$freq, s2$spec[,2]/s$spec[,1], type="l", log="y",  
  ylab="frequency ratio", xlab="frequency", xlim=c(0, 1.5),  
  main="frequency response (red line at 1.0)")  
abline(h=1, lty="dashed", col="red")
```

frequency response (red line at 1.0)




Question 8.5. Describe the frequency domain behavior of this filter.


Extracting business cycles: A band pass filter

- For the unemployment data, high frequency variation might be considered “noise” and low frequency variation might be considered trend.
- A band of mid-range frequencies might be considered to correspond to the business cycle.
- Let's build a smoothing operation in the time domain to extract business cycles, and then look at its frequency response function.

```
u_low <- ts(loess(u1~date,span=0.5)$fitted,  
            start=1948,frequency=12)  
u_hi <- ts(u1 - loess(u1~date,span=0.1)$fitted,  
            start=1948,frequency=12)  
u_cycles <- u1 - u_hi - u_low  
plot(ts.union(u1, u_low,u_hi,u_cycles),  
      main="Decomposition of unemployment as trend + noise + cycles")
```


Decomposition of unemployment as trend + noise + cycles





Question 8.6. Describe the frequencies (and corresponding periods) that this decomposition identifies as business cycles. Note: Usually, we should specify units for frequency and period. Here, the units are omitted to give you an exercise!

To help answer this question, let's add some lines to the previous plot




```
kable(matrix(freq_cycles,nrow=1,  
  dimnames=list("frequency",c("low","hi"))),digits=3)
```

	low	hi
frequency	0.069	0.194

Question 8.7. So far as we have opinions on business cycles, use them to comment on this decomposition.

Looking for business cycles

We plot a smoothed periodogram for the raw unemployment data. We cut out higher frequencies to zoom in on frequencies relevant to our business cycle investigation.



Question 8.8. Comment on the evidence for and against the concept of a business cycle in the above figure.

Common smoothers in R

- Above, we have used the **local regression smoother** `loess` but there are other options.
- Our immediate goal is to get practical experience using a smoother and then statistically assessing what we have done.
- You can learn about alternative smoothers, and try them out, if you like.
- `ksmooth` is a **kernel smoother**. The default periodogram smoother in `spectrum` is also a kernel smoother. See https://en.wikipedia.org/wiki/Kernel_smoother
- `smooth.spline` is a **spline smoother**. https://en.wikipedia.org/wiki/Smoothing_spline

Bandwidth for a smoother

- All these smoothers have some concept of a **bandwidth**, which is a measure of the size of the neighborhood of time points in which data affect the smoothed value at a particular time point.
- The concept of bandwidth is most obvious for kernel smoothers, but exists for other smoothers.
- We usually only interpret bandwidth up to a constant. For a particular smoothing algorithm and software implementation, you learn by experience to interpret the comparative value. Smaller bandwidth means less smoothing.
- Typically, when writing reports, it makes sense to focus on the tuning parameter for the smoother in question, which is not the bandwidth except for a kernel smoother.

Further reading

- Section 2.3 of Shumway and Stoffer (2017) discusses smoothing of time series, in the time domain.
- Section 4.2 of Shumway and Stoffer (2017) presents a frequency response function for linear filters, related to this chapter but in a different context.

License, acknowledgments, and links

- Licensed under the [Creative Commons Attribution-NonCommercial license](#). Please share and remix non-commercially, mentioning its origin.
- The materials builds on [previous courses](#).
- Compiled on February 17, 2021 using R version 4.0.3.



[Back to course homepage](#)

References

Shumway RH, Stoffer DS (2017). *Time Series Analysis and its Applications: With R Examples*. Springer. URL <http://www.stat.pitt.edu/stoffer/tsa4/tsa4.pdf>.