

2020-03-22



# Contents

<b>1</b>	<b>5</b>
<b>2 Introduction</b>	<b>7</b>
2.1 1.1 . . . . .	7
2.2 1.2 . . . . .	7
2.3 1.3 . . . . .	8
2.4 1.4 . . . . .	8
2.5 1.5 . . . . .	9
<b>3 Literature</b>	<b>11</b>
<b>4 Methods</b>	<b>13</b>
<b>5 Applications</b>	<b>15</b>
5.1 Example one . . . . .	15
5.2 Example two . . . . .	15
<b>6 Final Words</b>	<b>17</b>



# Chapter 1

*Life, thin and light-off time and time again*

*Frivolous tireless*



# Chapter 2

## Introduction

### 2.1 1.1

scalar

vector

$xx \qquad X \qquad X_1 \qquad X_2$

matrix

$A$

tensor

$A \quad \text{“A”} \quad A \quad (i,j,k) \quad A_{(i,j,k)}$

$0$

$/ \ /$

### 2.2 1.2

- “ ” “ ”  $n \quad n$  “ ”
- 
- $3 \times 3$
- $1 \times 1 \ 1 \times 3$

## 2.3 1.3

$$m \ n \quad n \quad m$$

## 2.4 1.4

$$\vec{a} = [-5, 6, 8, -10] \quad \vec{x} = (x_1, x_2, \dots, x_N)$$

$$\bullet \quad 1 \quad \vec{a} \ 1 \quad 29$$

$$\|\vec{x}\|_1 = \sum_{i=1}^N |x_i|$$

$$\bullet \quad 2 \quad \vec{a} \ 2 \quad 15$$

$$\|\vec{x}\|_2 = \sqrt{\sum_{i=1}^N |x_i|^2}$$

$$\bullet \quad \vec{a} \quad 5$$

$$\|\vec{x}\|_{-\infty} = \min |x_i|$$

$$\bullet \quad \vec{a} \quad 10$$

$$\|\vec{x}\|_{+\infty} = \max |x_i|$$

$$\bullet \quad p$$

$$L_p = \|\vec{x}\|_p = \sqrt[p]{\sum_{i=1}^N |x_i|^p}$$

$$A = [-1, 2, -3; 4, -6, 6] \quad A_{m \times n} \quad a_{ij}$$

$$\|A\|_p := \sup_{x \neq 0} \frac{\|Ax\|_p}{\|x\|_p}$$

,

$$\bullet \quad 1 \quad , \quad A \ 1 \quad [5, 8, 9] \quad 9$$



- $\mathbf{2} \quad A^T A \quad A \mathbf{2} \quad 10.0623$

$$\|A\|_2 = \sqrt{\lambda_{max}(A^T A)}$$

$\lambda_{max}(A^T A) \quad A^T A \quad - \quad A \mathbf{1} \quad [616] \quad 16$

$$\|A\|_\infty = \max_{1 \leq i \leq n} \sum_{j=1}^n |a_{ij}|$$

- $\text{svd} \quad \text{---} \quad A \quad 10.9287$
- $\mathbf{L0} \quad 0 \quad \mathbf{L0} \quad 0 \quad A \quad 6$
- $\mathbf{L1} \quad \mathbf{L0} \quad A \quad 22$
- $\mathbf{F} \quad \mathbf{L2} \quad A \quad 10.0995$

$$\|A\|_F = \sqrt{(\sum_{i=1}^m \sum_{j=1}^n |a_{ij}|^2)}$$

- $\mathbf{L21} \quad \mathbf{F} \quad 2 \quad \mathbf{L1} \quad 1 \quad \mathbf{L1} \mathbf{L2} \quad A \quad 17.1559$
- $\mathbf{p}$

$$\|A\|_p = \sqrt[p]{(\sum_{i=1}^m \sum_{j=1}^n |a_{ij}|^p)}$$

2.5 1.5

- $0$
- $C \ C^T C$
- $\begin{matrix} n \\ E \end{matrix} \quad E$
- 
- 
-



## Chapter 3

# Literature

Here is a review of existing methods.



## Chapter 4

# Methods

We describe our methods in this chapter.



## Chapter 5

# Applications

Some *significant* applications are demonstrated in this chapter.

### 5.1 Example one

### 5.2 Example two





## Chapter 6

# Final Words

We have finished a nice book.