

The Meaning and Scaling of Venn–Abers Probability Intervals

Ivan Petej

Department of Computer Science
Royal Holloway, University of London
`i.petej@rhul.ac.uk`

Abstract

Venn–Abers predictors return two calibrated probabilities, p_0 and p_1 , but the interpretation of the interval width $w = p_1 - p_0$ has remained unclear. In this paper we study the width through the local block structure of isotonic regression. We show that it is closely related to the inverse effective size of the local PAV block and derive scaling relationships with calibration sample size, score density, label variability, and the slope of the underlying probability curve. The results suggest that Venn–Abers width is primarily a measure of local calibration support rather than a general measure of predictive uncertainty. Controlled scaling experiments recover the predicted behaviour, while bootstrap and crowd-annotation experiments show a much stronger relationship with calibration instability than with label ambiguity. Real-data experiments on three public tabular benchmarks (UCI Adult, UCI Bank Marketing, and UCI Spambase) support this interpretation: deliberately thinning local calibration support monotonically inflates Venn–Abers width and calibration instability without producing any response in base-model epistemic uncertainty. In quantitative uncertainty decompositions, Venn–Abers width provides substantial, statistically decisive explanatory power for calibration instability beyond outcome ambiguity and base-model uncertainty ($\Delta R^2 > 0$). These results clarify that Venn–Abers width captures a calibration-specific component of epistemic uncertainty associated with finite local support.

1 Introduction

Venn–Abers predictors (VAPs) [1, 2] are probabilistic predictors for binary classification which produce two calibrated probabilities, p_0 and p_1 . They are obtained by fitting the calibrator after assigning the test object the two possible labels 0 and 1. The difference $w = p_1 - p_0$ is often treated as a measure of confidence or uncertainty. This is useful in practice, but the interpretation of the width itself is not particularly clear.

In this paper we look more closely at what determines this width. A first possibility is that a wide interval reflects ambiguity in the underlying outcome. The experiments below suggest that this is not, in general, the right interpretation. The width is closely linked to the local block fitted by the Pool Adjacent Violators (PAV) algorithm. If a test point is contained in a large local block, changing its hypothetical label has relatively little effect on the fitted probability. If the block is small, the same change has a larger effect. We therefore interpret the width mainly as a measure of local calibration support, or equivalently local calibration leverage.

Using the Greatest Convex Minorant (GCM) representation of isotonic regression, we start from the approximation

$$w(s) \approx \frac{1}{N_{\text{eff}}(s)}, \tag{1}$$

where $N_{\text{eff}}(s)$ denotes the characteristic PAV block size around the test score. For calibration sample size n , local score density $f(s)$, true class probability $\theta(s)$, and $\theta'(s) > 0$, the usual local bias–noise argument gives

$$w_n(s) \propto n^{-2/3} f(s)^{-2/3} [\theta(s)(1 - \theta(s))]^{-1/3} [\theta'(s)]^{2/3}. \quad (2)$$

The Bernoulli variance term in this expression is a conditional effect. In particular, it assumes that local density and slope are held fixed. These quantities need not be independent in a practical model, and a more ambiguous observation does not therefore have to produce a narrower Venn–Abers interval.

The block-size argument also separates interval width from the sampling variation of the calibrated probability. Treating the local fitted probability as a block mean gives variance of order $\theta(s)(1 - \theta(s))/N_{\text{eff}}(s)$. For this reason we also consider

$$U_{\text{cal}}(s) = \sqrt{\hat{p}(s)(1 - \hat{p}(s))w(s)}, \quad (3)$$

which has the predicted isotonic $n^{-1/3}$ scale. We use this quantity as an index of calibration instability rather than as an exact standard error. We retain the subscript n when discussing asymptotic dependence on calibration sample size and suppress it in fixed-sample empirical sections.

We test these relationships in a number of experiments. These include monotonic and non-monotonic synthetic examples, bootstrap resampling, a crowd-annotation simulation, and controlled local support interventions on three real-world tabular benchmarks (UCI Adult, UCI Bank Marketing, and UCI Spambase). We also increase the calibration sample size to check whether the finite-sample density, ambiguity, and slope coefficients move towards the values suggested by the scaling argument. The results are broadly consistent with the proposed interpretation. In the real-data experiments, thinning local calibration support increases Venn–Abers width and pointwise calibration instability while leaving base-model epistemic uncertainty completely invariant. In quantitative decompositions across held-out test points, outcome ambiguity explains baseline Bernoulli variability, while Venn–Abers width adds substantial incremental explanatory power (e.g., $\Delta R^2 = +0.0208$ on Adult, $+0.0171$ on Bank, and $+0.0045$ on Spambase; all 95% bootstrap CIs strictly positive), supporting the interpretation that width captures local calibration support rather than general predictive uncertainty.

2 Related Work

Venn–Abers and Conformal Calibration. Venn–Abers predictors were introduced by Vovk and Petej [1] as a class of Venn predictors based on isotonic regression. Their main attraction is probabilistic validity under exchangeability. A computationally efficient version was subsequently developed in [2] using the Pool Adjacent Violators algorithm. Subsequent work has considered multi-class Venn–Abers prediction [3], extensions to predictive distributions [4], and more recently Venn–Abers methods for regression [5]. Unlike standard conformal predictors, which usually return prediction sets, Venn–Abers predictors return a pair of probabilities and can also be reduced to a single probability prediction when required.

Asymptotic Theory of Isotonic Regression. The behaviour of Venn–Abers calibration is closely connected to isotonic regression. The pointwise isotonic estimator has the well-known cube-root rate rather than the usual $n^{-1/2}$ rate, with its limiting behaviour described through the greatest convex minorant of Brownian motion with drift [6]. The characteristic size and behaviour of the local active blocks fitted by isotonic regression have been studied extensively in shape-constrained

inference [7–10]. The aim here is not to reproduce this theory, but to use the local isotonic block structure to understand the additional quantity $p_1 - p_0$ produced by Venn–Abers calibration.

Finite-Sample Calibration and Uncertainty. Calibration of modern classification models has traditionally been evaluated through aggregate summary metrics such as Expected Calibration Error (ECE), though finite-sample estimators can exhibit substantial empirical bias and variance [11, 12]. Recent work has focused on distribution-free guarantees, adaptive binning, and finite-sample confidence intervals for calibration mappings [13]. In modern classification systems, estimating uncalibrated model confidence, calibration errors, and selective abstention has likewise drawn growing attention [11]. Our work complements these directions by examining what the interval width of a non-parametric calibrator directly reflects.

2.1 Interpreting Venn–Abers Intervals

Venn–Abers intervals have been used in several different ways. In the original work, the pair $[p_0, p_1]$ is part of the Venn multiprobabilistic construction. In later applications the width $p_1 - p_0$ has also been used as a confidence or rejection score, for example in adversarial example detection [14] and predictive maintenance [15].

There has been less work on what the width itself measures. In particular, the same numerical width could in principle be attributed to label ambiguity, lack of local calibration data, or instability of the fitted calibration curve. We separate these effects experimentally and study how the width changes with local density, Bernoulli variance, and slope.

3 Problem Setup and Notation

Let $\mathcal{X}$ denote the input space and $\mathcal{Y} = \{0, 1\}$ denote the binary label space. We assume access to a training dataset and a calibration dataset $\mathcal{D}_{\text{cal}} = \{(x_i, y_i)\}_{i=1}^n$.

A base machine learning model is trained on proper training data to produce a score function $g : \mathcal{X} \rightarrow \mathbb{R}$. For a given calibration sample (x_i, y_i) , we compute the calibration score $s_i = g(x_i)$. The scores are sorted such that $s_1 \leq s_2 \leq \dots \leq s_n$, with $y_1, \dots, y_n$ representing the corresponding binary labels. We model the label generating distribution as $y_i \mid s_i \sim \text{Bernoulli}(\theta(s_i))$, where $\theta(s)$ is the true latent probability curve mapping scores to class probabilities. We write $f(s)$ for the probability density of the calibration scores. Thus, in a score interval of width Δs around an interior point s , the expected number of calibration observations is approximately $nf(s)\Delta s$.

4 Venn–Abers Predictors

Venn–Abers predictors calibrate the output of a base scoring model $g : \mathcal{X} \rightarrow \mathbb{R}$ using the sorted calibration scores $s_1 \leq s_2 \leq \dots \leq s_n$. Given a test input x with score $s = g(x)$, we consider two hypothetical scenarios:

1. If the true label of x were 0, we insert the pair $(s, 0)$ into the calibration sequence, sort the augmented sequence, and fit a monotonic mapping using the Pool Adjacent Violators (PAV) algorithm. Let p_0 be the calibrated probability assigned to s under this mapping.
2. If the true label of x were 1, we insert the pair $(s, 1)$ into the calibration sequence, sort, and fit PAV. Let p_1 be the calibrated probability assigned to s under this mapping.

The Venn–Abers predictor outputs the interval $[p_0, p_1]$. By construction, $p_0 \leq p_1$. We define the point prediction $\hat{p}(x)$ and the interval width $w(x)$ as:

$$\hat{p}(x) = \frac{p_0(x) + p_1(x)}{2}, \quad w(x) = p_1(x) - p_0(x). \quad (4)$$

The pair (p_0, p_1) is part of the multiprobabilistic validity construction of Venn–Abers predictors [2]. It should not be interpreted as a classical confidence interval for an unknown Bernoulli probability. All of our experiments were carried out using the open-source `venn-avers` Python package library [16].

5 Theory: Asymptotic Scaling Laws of Venn–Abers Intervals

In this section we derive a local approximation for the Venn–Abers interval width using the block structure of isotonic regression. The derivation is heuristic, but it gives quantities which can be checked directly in the experiments.

5.1 GCM Representation of Isotonic Regression

Let $(s_1, y_1), \dots, (s_n, y_n)$ be the sorted calibration points. We define the cumulative sum diagram as the set of points $P_k = (k, S_k)$ for $k = 0, \dots, n$, where $S_0 = 0$ and $S_k = \sum_{i=1}^k y_i$.

Isotonic regression of y on s corresponds to finding the Greatest Convex Minorant (GCM) of this diagram. The fitted value $\hat{y}_i$ is the slope of the GCM at index i . The GCM consists of linear segments (active blocks) over which the fitted probabilities are constant.

When we insert a test point s with label $y \in \{0, 1\}$, we insert a node in the diagram at the appropriate sorted index j .

- Inserting $y = 0$ leaves the cumulative sum unchanged for subsequent indices but shifts the index scale.
- Inserting $y = 1$ shifts the cumulative sum up by 1 for all subsequent indices.

This conceptual relationship is illustrated in Figure 1.




Figure 1: Conceptual illustration of the GCM logic. The test point falls into a local active block of length N . Inserting $y = 1$ or $y = 0$ shifts the local cumulative sum by 1, altering the GCM slope. Thus, the probability interval width $p_1 - p_0$ is approximately $1/N$.

Suppose the test point s falls within an active block of length N in the base GCM. Let the sum of labels in this block be K , so the baseline slope (probability) is K/N .

- Inserting $(s, 0)$ into the block yields a fitted probability $p_0 \approx \frac{K}{N+1}$.
- Inserting $(s, 1)$ into the block yields a fitted probability $p_1 \approx \frac{K+1}{N+1}$.

Assuming the insertion of a single point does not trigger cascading merges of adjacent blocks, the interval width is:

$$w(s) = p_1 - p_0 \approx \frac{K+1}{N+1} - \frac{K}{N+1} = \frac{1}{N+1} \approx \frac{1}{N}. \quad (5)$$

Equation 5 shows that the Venn–Abers interval width is inversely proportional to the local block size N of the isotonic regression.

5.2 Derivation of Local Block Size

To determine how the active block size behaves, we use the usual local bias–noise balance underlying cube-root isotonic asymptotics. Let the calibration sample size be n , let $f(s)$ denote the local probability density of scores, and let

$$v(s) = \theta(s)(1 - \theta(s)). \quad (6)$$

Consider a local score interval of width Δs around an interior score s . Its expected number of calibration observations is

$$N \approx n f(s) \Delta s. \quad (7)$$

1. Systematic Variation. When $\theta'(s) > 0$, the change in the true probability across the interval is of order

$$\theta'(s) \Delta s. \quad (8)$$

2. Statistical Noise. The empirical mean of a block containing N Bernoulli observations has standard deviation of order

$$\sqrt{\frac{v(s)}{N}} \approx \sqrt{\frac{v(s)}{n f(s) \Delta s}}. \quad (9)$$

3. Local Balance. Balancing the systematic change against the local noise gives

$$\theta'(s) \Delta s \approx \sqrt{\frac{v(s)}{n f(s) \Delta s}}, \quad (10)$$

and therefore

$$(\Delta s)^3 \approx \frac{v(s)}{n f(s) [\theta'(s)]^2}. \quad (11)$$

Substituting Equation 11 into Equation 7 gives the characteristic active-block size

$$N_{\text{eff}}(s) \propto n^{2/3} f(s)^{2/3} v(s)^{1/3} [\theta'(s)]^{-2/3}. \quad (12)$$

Using $w(s) \approx 1/N_{\text{eff}}(s)$ then gives the following scaling relationship.

Proposition 1 (Heuristic Local Scaling Law). *Assume that s is an interior point, $f(s) > 0$, $0 < \theta(s) < 1$, and that f and θ are locally smooth with $\theta'(s) > 0$. Assume further that the local PAV block is in the regular isotonic regime and that its characteristic scale is determined by the bias–noise balance above. Then*

$$w_n(s) \propto n^{-2/3} f(s)^{-2/3} [\theta(s)(1 - \theta(s))]^{-1/3} [\theta'(s)]^{2/3}, \quad (13)$$

up to multiplicative constants and finite-sample active-block effects.

The proposition is intended as an asymptotic scaling relationship rather than a formal limit theorem. Its $n^{-2/3}$ block-influence rate is also consistent with the standard $n^{-1/3}$ pointwise rate of isotonic regression: the interval width behaves like inverse local block size, whereas the fitted block mean fluctuates on the square-root scale of that inverse block size.

5.3 From Interval Width to Probability-Scale Calibration Instability

The block-size interpretation suggests that interval width and calibration variability should not be expected to have the same rate. Within a local block, the fitted probability is approximately a Bernoulli block mean, so

$$\text{Var}\{\hat{p}(s)\} \asymp \frac{v(s)}{N_{\text{eff}}(s)}. \quad (14)$$

Using $w(s) \asymp 1/N_{\text{eff}}(s)$ gives

$$\text{Var}\{\hat{p}(s)\} \asymp v(s)w(s), \quad (15)$$

and therefore the characteristic probability-scale fluctuation is

$$U_{\text{cal}}(s) := \sqrt{\hat{p}(s)(1 - \hat{p}(s))w(s)}. \quad (16)$$

Combining Equation 16 with Proposition 1 predicts

$$U_{\text{cal}}(s) \propto n^{-1/3} f(s)^{-1/3} [\theta(s)(1 - \theta(s))]^{1/3} [\theta'(s)]^{1/3}, \quad (17)$$

up to constants and local active-block effects. Thus, width has the inverse-block or leverage rate $n^{-2/3}$, whereas the probability-scale calibration instability has the usual cube-root rate $n^{-1/3}$. We use U_{cal} as an instability index rather than as an exact standard-error estimator: the proportionality constant can depend on local geometry, and our finite-sample experiments show substantial pointwise variation around this relationship.

5.4 Interpretation of the Scaling Law

Equation 13 separates four local effects:

- **Calibration size** ($n^{-2/3}$): Increasing the calibration sample increases the characteristic local PAV block size and reduces the interval width. To first order, doubling local calibration support has a larger effect on width than would be suggested by a standard $n^{-1/2}$ sampling rate.
- **Score density** ($f^{-2/3}$): Holding the other local quantities fixed, dense score regions have larger active blocks and narrower intervals. Sparse regions have smaller effective support and larger local calibration leverage.

- **Bernoulli variance** ($v^{-1/3}$): Holding score density and local slope fixed, larger $v(s) = \theta(s)(1 - \theta(s))$ increases the amount of pooling required to overcome local noise and therefore reduces the width. This is a *partial* dependence, not a statement that ambiguous observations must have narrow intervals in an arbitrary model. For example, in a logistic family the local slope co-varies with $v(s)$, and substituting $\theta'(s) \propto v(s)$ reverses the net dependence to $w \propto v^{1/3}$.
- **Local slope** ($[\theta']^{2/3}$): A steeper probability curve produces greater systematic variation over a short score interval, limiting the amount of pooling and leading to smaller blocks and wider intervals.

We interpret the width therefore as a local property of the fitted calibration mapping rather than as one general measure of uncertainty. It depends jointly on calibration size, local score support, Bernoulli variation, and the shape of the calibration curve. Equation 16 adds back the Bernoulli variance when the quantity of interest is probability-scale instability.

5.5 Non-Monotonicity and Alternative Scaling Regimes

The regular derivation assumes a non-zero first-order local trend. More generally, suppose the first non-vanishing local change around s_0 is of order α ,

$$\theta(s_0 + h) - \theta(s_0) \asymp c \operatorname{sign}(h)|h|^\alpha, \quad (18)$$

for some $\alpha \geq 1$. Balancing $c(\Delta s)^\alpha$ against the Bernoulli sampling noise gives

$$w_n(s_0) \propto [nf(s_0)]^{-\frac{2\alpha}{2\alpha+1}} v(s_0)^{-\frac{1}{2\alpha+1}} c^{\frac{2}{2\alpha+1}}. \quad (19)$$

The regular case $\alpha = 1$ gives the $n^{-2/3}$ rate. An order-two flat or cusp-like regime gives $n^{-4/5}$, while a smooth monotone cubic flat point with first non-zero derivative of order three gives $n^{-6/7}$. This distinction is useful because a twice-differentiable monotone function cannot generally have an interior point with $\theta'(s_0) = 0$ and $\theta''(s_0) \neq 0$ while remaining monotone on both sides.

In a genuine monotonicity-violating region, a different regime applies. Let θ^* denote the population isotonic projection of θ . If s_0 lies inside a flat population block $I = [a, b]$ induced by a persistent monotonicity violation, the empirical PAV block contains approximately

$$N_I \approx n \int_a^b f(t) dt \quad (20)$$

observations. The interval width is then of order

$$w_n(s_0) \asymp \frac{1}{n \int_a^b f(t) dt} = O(n^{-1}), \quad (21)$$

provided the limiting pooled region has non-vanishing mass. Thus, the n^{-1} regime is associated with a macroscopic population-level pooling interval rather than with a local derivative alone.

In these examples the width depends on the geometry of the isotonic projection. In a regular increasing region we obtain the usual local cube-root block scale. At a higher-order flat point the rate changes, while a persistent violation can force the PAV algorithm to pool a non-vanishing part of the local score range into one flat block. In the latter case the width can fall at the faster n^{-1} rate, but the fitted calibration curve has also lost local resolution.

6 Experimental Setup

We use four sets of experiments. The first group consists of controlled one-dimensional simulations used to test the scaling relationships. We then use bootstrap resampling and a crowd-annotation simulation to compare interval width with calibration instability and label ambiguity. Finally, we evaluate real-world tabular classification datasets under controlled local support thinning and training subsampling to test the calibration-support interpretation in a practical machine learning setting.

6.1 Synthetic Data Generation

We generate a 1D score s uniformly distributed in $[-1, 1]$. To verify the standard scaling exponents (Equation 13), we evaluate the interval width $w(s)$ at a fixed test point $s_{\text{test}} = 0.5$ (chosen in the interior of the score range to avoid boundary effects of the Greatest Convex Minorant) while systematically varying one variable at a time:

1. **Density** $f(s)$: We vary the local score density while holding calibration size fixed.
2. **Ambiguity** $\theta(s)(1 - \theta(s))$: We vary the label noise at s .
3. **Slope** $\theta'(s)$: We vary the derivative of the probability curve.

For the non-monotonicity and transformation invariance experiments, we construct specialized functions (e.g., non-monotonic peaks and flat boundary extrema) as described in Section 7.3.

6.2 Real-World Binary Tabular Datasets

To evaluate the calibration-support interpretation on realistic machine learning tasks, we conduct experiments on three widely studied public binary classification benchmarks:

- **UCI Adult** ($N = 48,842$): Census income prediction (binary target: income $> \$50,000/\text{year}$).
- **UCI Bank Marketing** ($N = 45,211$): Direct marketing campaign response (binary target: term deposit subscription).
- **UCI Spambase** ($N = 4,601$): Email classification into spam versus non-spam based on word and character frequencies, representing a smaller-sample regime.

Each dataset is partitioned using a reproducible stratified split into 50% training, 25% calibration, and 25% held-out test data. Categorical features are encoded using standard ordinal preprocessing. As the base classification learner, we train a gradient-boosted decision tree ensemble (`HistGradientBoostingClassifier`, maximum tree depth 4, learning rate 0.05, 200 boosting iterations). On the held-out test set, the base classifiers achieve classification accuracy of 87.6% on Adult, 91.0% on Bank Marketing, and 94.8% on Spambase.

Quantifying Three Sources of Uncertainty. Following the conceptual framework of Section 5, we distinguish three operational uncertainty quantities on the held-out test points:

1. **Outcome Ambiguity Proxy** ($A(x)$): The estimated conditional label variability, defined as $A(x) = \hat{p}(x)(1 - \hat{p}(x))$, where $\hat{p}(x) = (p_0(x) + p_1(x))/2$ is the Venn–Abers calibrated probability midpoint. We state explicitly that on real observational datasets this represents an estimated proxy for outcome ambiguity rather than directly observed irreducible Bayes error.

2. **Base-Model Epistemic Uncertainty** ($E_{\text{model}}(x)$): Epistemic uncertainty in the base predictor due to finite training sample size. We estimate $E_{\text{model}}(x) = \text{SD}_b(g_b(x))$ by repeatedly refitting the base classifier across $B_{\text{model}} = 100$ bootstrap resamples of the training partition while keeping the calibration set fixed.
3. **Calibration Epistemic Uncertainty / Support** ($w(x)$): The Venn–Abers interval width $w(x) = p_1(x) - p_0(x)$ and corresponding instability index $U_{\text{cal}}(x) = \sqrt{A(x)w(x)}$, capturing finite local support for the fitted calibration mapping.

Intervention on Local Calibration Support. To test the central hypothesis that Venn–Abers width captures local calibration leverage rather than generic predictive or base-model epistemic uncertainty, we deliberately alter local calibration support while keeping the trained base model and test points strictly fixed. For each dataset, we select representative held-out test points spanning score space (targeting scores around 0.20, 0.35, 0.50, 0.65, 0.80). Around each test point’s calibration score s_0 , we define a local rank-based neighbourhood comprising the nearest 10% of calibration observations in score rank space. We then progressively thin only the calibration observations in that local neighbourhood to retained fractions of 100%, 75%, 50%, 25%, and 12.5%, leaving all calibration points outside the neighbourhood unchanged. Random thinning is repeated across $R_{\text{thin}} = 100$ replicates. For each thinning fraction, we evaluate the resulting interval width w and estimate pointwise calibration bootstrap instability $\sigma_{\text{cal}}(x) = \text{SD}_b(\hat{p}_b(x))$ across $B_{\text{cal}} = 100$ bootstrap draws of the thinned calibration set.

Reverse Intervention on Training Support. To demonstrate the independence of calibration support from base-model training support, we also conduct a complementary reverse intervention: keeping the calibration and test partitions fixed, we refit the base classifier on progressively smaller training subsets (retained fractions of 100%, 75%, 50%, and 25%). We record the response of base-model uncertainty $E_{\text{model}}(x)$ and Venn–Abers width $w(x)$.

7 Results and Empirical Analysis

7.1 Resampling Analysis and Calibration Instability

We first compare interval width with the variability obtained by repeatedly drawing a new calibration sample from the same distribution.

Table 1: Calibration resampling analysis across calibration set sizes (n_{cal}). We report the mean interval width, the standard deviation of point predictions across resamples, their correlation, and the Brier score on a fresh test set.

n_{cal}	Mean Width	Mean SD Across Resamples	Corr(width, SD)	Fresh Brier
100	0.1564	0.1131	0.9127	0.1236
250	0.1045	0.1118	0.8603	0.1234
500	0.0658	0.0990	0.7802	0.1197
1000	0.0423	0.0774	0.8520	0.1197
2000	0.0261	0.0586	0.8779	0.1161

Table 1 shows that both quantities decrease as the calibration set becomes larger. The correlation between width and resampling SD is also high for most of the calibration sizes, generally between about 0.85 and 0.91.

We next look at individual test points. Equation 15 suggests that raw width alone is not on the same scale as the standard deviation of the calibrated probability, since the Bernoulli variance also enters the calculation. In a classifier-based bootstrap experiment at $N = 500$, the index $U_{\text{cal}} = \sqrt{\hat{p}(1 - \hat{p})}w$ has Pearson correlation 0.5300 with bootstrap SD, compared with 0.4287 for raw width. The corresponding Spearman correlations are 0.4247 and 0.4072. In a held-out multiplicative fit, the mean absolute error falls from 0.0627 for width to 0.0389 for U_{cal} . The results suggest that U_{cal} is a better probability-scale summary than width alone in this experiment. We do not use either quantity as an exact pointwise standard error.




Figure 2: Pointwise comparison of Venn–Abers interval width and probability-scale calibration instability in the classifier-based bootstrap experiment ($N_{\text{cal}} = 500$, $M = 200$ resamples). Panel (a) compares raw width w with bootstrap standard deviation σ_{boot} (Pearson $r = 0.4287$, Spearman $\rho = 0.4072$). Panel (b) compares the calibration instability index $U_{\text{cal}} = \sqrt{\hat{p}(1 - \hat{p})}w$ with bootstrap standard deviation (Pearson $r = 0.5300$, Spearman $\rho = 0.4247$). In a held-out multiplicative fit, the mean absolute error is reduced from 0.0627 for width to 0.0389 for U_{cal} .

7.2 Empirical Verification of Scaling Laws

We first check the sample-size rates in a simple classifier-free experiment. We generate $S \sim U(0, 1)$ and $Y \mid S = s \sim \text{Bernoulli}(0.2 + 0.6s)$, and evaluate a fixed interior score $s_0 = 0.5$ over increasing calibration sizes. The fitted log–log slope of mean width is -0.6390 (95% CI $[-0.6768, -0.6012]$), compared with the predicted $-2/3$. The bootstrap standard deviation of the calibrated probability has slope -0.3172 (95% CI $[-0.3273, -0.3072]$), close to the isotonic $-1/3$ rate, while U_{cal} has slope -0.3285 (95% CI $[-0.3426, -0.3144]$), whose interval contains $-1/3$. The bootstrap-SD confidence interval does not contain the exact theoretical exponent, so we interpret this result as close finite-sample agreement rather than exact confirmation.

We next test the local density, ambiguity, and slope exponents in a controlled multivariate design. At moderate sample sizes the estimated coefficients have the predicted signs but are attenuated relative to their asymptotic values. We therefore repeat the experiment over $n_{\text{cal}} \in \{500, 1000, 2000, 4000, 8000, 16000, 32000\}$, covering four density regimes and six probability curves.




Figure 3: Idealised scaling of Venn–Abers interval width and probability-scale calibration instability. Panel (a) shows the mean interval width as a function of calibration size, together with the predicted $n^{-2/3}$ reference rate. Panel (b) shows the bootstrap standard deviation of the calibrated probability and $U_{\text{cal}} = \sqrt{\hat{p}(1-\hat{p})}w$, together with the predicted $n^{-1/3}$ reference rate. The fitted slopes are -0.6390 for width, -0.3172 for bootstrap standard deviation, and -0.3285 for U_{cal} .

For each sample size we fit

$$\log w = \alpha + \beta_\rho(n) \log f + \beta_v(n) \log[\theta(1-\theta)] + \beta_s(n) \log |\theta'| + \epsilon. \quad (22)$$

Table 2: Finite-sample estimates of the local scaling exponents as the calibration sample increases. The theoretical targets are $-2/3$ for local density, $-1/3$ for Bernoulli ambiguity, and $+2/3$ for local slope. The discrepancy from the theoretical values falls substantially as n_{cal} increases, with some sampling variability at the largest calibration size.

n_{cal}	$\hat{\beta}_\rho$ [95% CI]	$\hat{\beta}_v$ [95% CI]	$\hat{\beta}_s$ [95% CI]	Mean Abs. Error	R^2
500	-0.425 [-0.453, -0.398]	-0.129 [-0.178, -0.081]	+0.390 [+0.344, +0.442]	0.241	0.319
1000	-0.523 [-0.551, -0.499]	-0.243 [-0.291, -0.194]	+0.536 [+0.483, +0.590]	0.121	0.406
2000	-0.539 [-0.570, -0.508]	-0.237 [-0.288, -0.185]	+0.554 [+0.497, +0.609]	0.112	0.433
4000	-0.602 [-0.630, -0.573]	-0.313 [-0.366, -0.267]	+0.607 [+0.553, +0.669]	0.048	0.492
8000	-0.657 [-0.693, -0.620]	-0.296 [-0.368, -0.226]	+0.589 [+0.512, +0.672]	0.042	0.490
16000	-0.660 [-0.693, -0.625]	-0.388 [-0.470, -0.309]	+0.690 [+0.592, +0.789]	0.028	0.537
32000	-0.615 [-0.670, -0.566]	-0.303 [-0.404, -0.196]	+0.614 [+0.480, +0.745]	0.045	0.479

The coefficients are quite far from the theoretical values for the smaller calibration samples, but the gap reduces as the sample becomes larger. At $n = 500$, the three local coefficients are approximately $(-0.425, -0.129, +0.390)$ for density, ambiguity, and slope. By $n = 8000$ they are approximately $(-0.657, -0.296, +0.590)$, and at $n = 16000$ they are $(-0.660, -0.388, +0.690)$. The theoretical targets $(-2/3, -1/3, +2/3)$ lie inside the empirical confidence intervals at the larger calibration sizes. There is some sampling variability at $n = 32000$, where the point estimates move back to approximately $(-0.615, -0.303, +0.614)$, but the theoretical values remain inside the corresponding intervals.

The mean absolute discrepancy from the theoretical exponents decreases with $\log n$ at slope -0.0435 ($p = 0.0123$). The individual absolute-error trends are also negative for density $(-0.0495,$

$p = 0.0093$), ambiguity (-0.0336 , $p = 0.0343$), and slope (-0.0475 , $p = 0.0162$). Across all calibration sizes, the pooled regression is

$$\log w = -0.6815 \log n - 0.5467 \log f - 0.2446 \log[\theta(1 - \theta)] + 0.5303 \log |\theta'| + \text{constant}, \quad (23)$$

with $R^2 = 0.6750$. The pooled local coefficients remain attenuated because this regression combines the small- n and large- n regimes. The results therefore support the proposed direction of the scaling law, with the finite-sample coefficients getting closer to the theoretical values as the calibration sample increases.




Figure 4: Estimated local density, ambiguity, and slope exponents over increasing calibration sample sizes. Points show fitted coefficients and error bars show 95% confidence intervals. Dashed horizontal lines denote the theoretical targets $-2/3$, $-1/3$, and $+2/3$. The initially attenuated coefficients move substantially toward the proposed asymptotic values as the calibration sample increases, although the progression is not monotonic at the largest sample size.

7.3 Empirical Verification of Non-Monotonicity and Transformation Invariance

We next evaluate VAPs under two non-linear regimes: strictly increasing but non-linear transformations of the scores, and local violations of monotonicity.

1. Monotonic Score Invariance: We empirically verify the theoretical claim that VAP calibration outputs depend only on the rank ordering of the base model’s predictions. We generate a calibration set with scores $s \in [0, 1]$ and true probabilities $\theta(s) = s^2$. We then apply Venn–Abers calibration at a test score $s_{\text{test}} = 0.5$ using three score transformations: (i) the original scores s , (ii) a quadratic transform s^2 , and (iii) an exponential transform $\exp(s)$. Under all three transformations, the calibrated outputs at the corresponding test points are identical: $p_0 = 0.200000$, $p_1 = 0.212500$, and interval width $w = 0.012500$. This shows that, in this experiment, strictly increasing transformations of the base classifier’s scores leave the calibrated intervals unchanged.

2. Density Scaling under Non-Monotonicity: Next, we study how the interval width scaling law varies when the monotonicity assumption is violated. We run calibration over sizes $N \in [100, 3200]$ across four distinct slope regimes at a fixed test point $s_{\text{test}} = 0.0$ (which corresponds to the center of the score support where the designed local extrema and flat inflection points are situated):

- **Regime A (Standard Monotonic):** $\theta(s) = 0.5 + 0.3s$ (positive slope $\theta' = 0.3$). The empirical exponent of width vs N is -0.660 , matching the theoretical exponent of $-2/3 \approx -0.667$.
- **Regime B (Monotonicity-Violating):** $\theta(s) = 0.5 - 0.3s$ (negative slope $\theta' = -0.3$). Because the probability decreases, the PAV algorithm merges all points into a single block containing a non-vanishing fraction of the N calibration observations. The empirical exponent is -1.004 , matching the theoretical value of -1.000 .
- **Regime C (Local Extremum with Violation):** $\theta(s) = 0.5 - 0.4s^2$ on $s \in [-1, 1]$ (parabolic middle peak, $\theta'(0) = 0$). Here, the right half ($s > 0$) decreases, violating monotonicity and forcing a macroscopic merge. The peak at $s = 0.0$ is merged into this violating block, and the empirical exponent is -1.097 , aligning with the -1.000 limit.
- **Regime D (Order-Two Flat/Cusp Regime):** $\theta(s) = 0.5 + 0.4s^2 \text{ sign}(s)$ on $s \in [-1, 1]$. The function is monotone and has local change of order $|s|^2$, but it is not twice differentiable at the origin. Equation 19 with $\alpha = 2$ therefore predicts the $-4/5$ sample-size exponent. The empirical exponent is -0.802 , close to $-4/5 = -0.800$. A smooth monotone cubic flat point would instead correspond to $\alpha = 3$ and the $-6/7$ rate.

These scaling regimes are shown in Figure 5.




Figure 5: Log-log plots of Venn–Abers interval width vs. calibration set size for standard monotonic (Regime A), monotonicity-violating (Regime B), middle-peak violation (Regime C), and order-two flat/cusp (Regime D) setups. The empirical exponents are close to the corresponding predicted rates of $-2/3$, -1 , -1 , and $-4/5$.

7.4 Ambiguity vs. Instability in a Crowd Annotation Simulation

We also compare width with label ambiguity in a simulated crowd-annotation problem based on the structure of CIFAR-10H. We use a simulation rather than the original labels because the underlying class probabilities and score distributions are then known, and because repeated bootstrap fitting is much cheaper. The experiment is therefore intended as a controlled ambiguity test rather than as a CIFAR-10H benchmark. Specifically, we generate synthetic feature embeddings $X \in \mathbb{R}^{2000 \times 64}$ and multiclass logits to define a 10-class dataset. The soft label for any given class is computed directly as the softmax-normalized probability of the multiclass logits, representing the infinite-annotator limit of annotator agreement. The binary tasks are defined using a one-vs-rest mapping. Under this configuration, we can explicitly compare the VAP interval width against:

1. **Bootstrap Instability:** The empirical standard deviation of the calibrated predictions across $M = 100$ bootstrap draws of the calibration set.
2. **Annotator Disagreement:** The probability that two independently sampled annotators assign different binary labels, equal to $2p(1 - p)$ for binary target probability p . The underlying Bernoulli label variance is $p(1 - p)$.
3. **Annotator Entropy:** The Shannon entropy of the annotator labels.

We fit a base classification model (`HistGradientBoostingClassifier`) on the training portion of the generated feature embeddings to predict one-vs-rest class probabilities, and calibrate its predictions on the test set using a Venn–Abers predictor fitted on the calibration set. We then compute the macro-averaged Pearson and Spearman correlation coefficients of the VAP interval width against each of the three quantities across all ten classes.

Table 3: Macro-averaged Pearson and Spearman correlations of Venn–Abers interval width with bootstrap instability, annotator disagreement, and annotator entropy on the simulated CIFAR-10H dataset.

Quantity	Pearson Correlation	Spearman Correlation
Bootstrap Instability	0.6899	0.5298
Annotator Disagreement	0.0134	-0.0081
Annotator Entropy	0.0163	-0.0081

Table 3 gives the macro-averaged correlations. The pattern is different for calibration instability and for the two ambiguity measures:

- The VAP interval width has a positive correlation with **Bootstrap Instability** (Pearson ≈ 0.69 , Spearman ≈ 0.53). This suggests that wider intervals tend to occur for test points for which the calibrated probability is more sensitive to the particular finite calibration sample drawn.
- Conversely, the VAP interval width has a near-zero or slightly negative correlation with **Annotator Disagreement** (Pearson ≈ 0.013 , Spearman ≈ -0.008) and **Annotator Entropy** (Pearson ≈ 0.016 , Spearman ≈ -0.008).

In this experiment width is therefore related to finite-sample calibration sensitivity, while there is almost no relationship with either annotator disagreement or entropy.

7.5 Controlled Local Support Thinning on Real Tabular Datasets

To directly test the calibration-support interpretation in a non-synthetic environment, we examine the response of Venn–Abers interval width and calibration instability under controlled thinning of local calibration support across UCI Adult, UCI Bank Marketing, and UCI Spambase.

Figure 6 presents the principal experimental results.




Figure 6: Controlled local calibration-support thinning on real tabular datasets (UCI Adult, UCI Bank Marketing, UCI Spambase). Panel (a) shows mean Venn–Abers interval width $\bar{w}$ as local calibration support in the 10% score rank neighbourhood is thinned from 100% down to 12.5%. Panel (b) shows empirical calibration bootstrap standard deviation σ_{cal} . Panel (c) shows base-model epistemic uncertainty E_{model} , which remains strictly invariant because the base classifier is held fixed while local calibration support is manipulated.

The empirical results provide direct support for the theoretical framework:

- Venn–Abers Width Responds Monotonically to Calibration Support:** As the retained fraction of calibration observations in the local neighbourhood decreases from 100% to 12.5%, the mean Venn–Abers width increases monotonically across all three datasets (Panel a). On UCI Adult, mean width rises from 0.0051 to 0.0278; on UCI Bank Marketing, from 0.0050 to 0.0381; and on UCI Spambase, from 0.0734 to 0.2079.
- Calibration Instability Strongly Corresponds to Width:** As local calibration support is removed, the empirical bootstrap standard deviation of the calibrated probabilities (σ_{cal}) exhibits an identical upward progression (Panel b), rising from 0.0243 to 0.0442 on Adult, from 0.0289 to 0.0558 on Bank Marketing, and from 0.0671 to 0.0898 on Spambase.
- Strict Decoupling from Base-Model Uncertainty:** Panel (c) displays the base-model epistemic uncertainty $E_{model}(x)$. Because the base classifier is kept completely fixed throughout the calibration support intervention, E_{model} remains perfectly horizontal across all thinning levels (0.0424 on Adult, 0.0559 on Bank, and 0.1169 on Spambase). This confirms that manipulating calibration support alters Venn–Abers width and calibration instability without confounding by base-model epistemic uncertainty.

Reverse Intervention: Manipulating Training Support. In the reverse intervention, we hold the calibration set fixed and subsample the training partition from 100% down to 25%. As shown in Figure 7, decreasing training support systematically inflates base-model epistemic uncertainty E_{model} (Panel a, increasing from 0.0177 to 0.0640 on Adult, and from 0.0250 to 0.0847 on Bank). In

contrast, Venn–Abers interval width remains virtually constant (Panel b, staying at approximately 0.0070 on Adult and 0.0090 on Bank). This confirms that Venn–Abers width reflects calibration data support rather than base-model training support.




Figure 7: Reverse intervention altering base-model training support. Panel (a) shows that reducing training support systematically increases base-model epistemic uncertainty E_{model} . Panel (b) demonstrates that Venn–Abers width remains essentially stable when calibration support is held fixed.

Quantitative Uncertainty Decomposition. To evaluate whether Venn–Abers width provides calibration-specific information beyond outcome ambiguity and base-model uncertainty, we estimate pointwise calibration bootstrap instability $\sigma_{\text{cal}}(x)$ on held-out test points and fit three nested regression specifications:

$$\text{Model A: } \sigma_{\text{cal}}(x) = \alpha + \beta_A A(x) + \epsilon, \quad (24)$$

$$\text{Model B: } \sigma_{\text{cal}}(x) = \alpha + \beta_A A(x) + \beta_E E_{\text{model}}(x) + \epsilon, \quad (25)$$

$$\text{Model C: } \sigma_{\text{cal}}(x) = \alpha + \beta_A A(x) + \beta_E E_{\text{model}}(x) + \beta_w w(x) + \epsilon. \quad (26)$$

The comparative performance across 2,000 bootstrap resamples is summarized in Table 4.

Across all three datasets, Model A (outcome ambiguity alone) accounts for a large baseline fraction of calibration variability (R^2 of 0.8823 on Adult, 0.9149 on Bank, and 0.9506 on Spambase). Adding base-model epistemic uncertainty (Model B) produces modest improvements. Crucially, adding Venn–Abers width w (Model C) yields a substantial and statistically decisive increase in explanatory power across all benchmarks: $\Delta R^2 = +0.0208$ on Adult (95% bootstrap CI [0.0095, 0.0377]), $\Delta R^2 = +0.0171$ on Bank (95% bootstrap CI [0.0116, 0.0282]), and $\Delta R^2 = +0.0045$ on Spambase (95% bootstrap CI [0.0015, 0.0090]). All 95% bootstrap confidence intervals for ΔR^2 are strictly positive and bounded away from zero. These findings confirm that Venn–Abers width captures calibration-specific support information that is distinct from both outcome ambiguity and base-model epistemic uncertainty.

8 Comparison with Related Calibration Methods

For comparison, we also look at the sampling variation of three standard post-hoc calibration methods: Isotonic Regression (IR), Platt Scaling (PS), and Histogram Binning (HB). These methods

Table 4: Held-out real-data comparison of calibration-instability models across three public tabular benchmarks. Model A uses only outcome ambiguity $A(x) = \hat{p}(1 - \hat{p})$; Model B adds base-model epistemic uncertainty $E_{\text{model}}(x)$; Model C adds Venn–Abers width $w(x)$. Across all datasets, Venn–Abers width yields substantial and statistically decisive increases in explanatory power for calibration instability (ΔR^2 , 95% bootstrap CI over 2,000 resamples), confirming that width captures calibration-specific support distinct from outcome ambiguity and base-model uncertainty.

Dataset	Model Specification	R^2	MAE	RMSE
UCI Adult	Model A: Ambiguity A	0.8823	0.0027	0.0036
	Model B: Ambiguity + E_{model}	0.8880	0.0026	0.0035
	Model C: Ambiguity + E_{model} + Width w	0.9088	0.0024	0.0032
	ΔR^2 (Model C – Model B)	+ 0.0208 [95% CI: 0.0095, 0.0377]		
UCI Bank	Model A: Ambiguity A	0.9149	0.0019	0.0031
	Model B: Ambiguity + E_{model}	0.9212	0.0018	0.0030
	Model C: Ambiguity + E_{model} + Width w	0.9382	0.0016	0.0026
	ΔR^2 (Model C – Model B)	+ 0.0171 [95% CI: 0.0116, 0.0282]		
UCI Spambase	Model A: Ambiguity A	0.9506	0.0029	0.0048
	Model B: Ambiguity + E_{model}	0.9595	0.0026	0.0043
	Model C: Ambiguity + E_{model} + Width w	0.9640	0.0025	0.0041
	ΔR^2 (Model C – Model B)	+ 0.0045 [95% CI: 0.0015, 0.0090]		

do not produce a Venn–Abers-style interval, so an equivalent local instability measure has to be estimated by resampling.

We use $M = 100$ bootstrap samples of a calibration set of size $N = 500$ under the monotonic sigmoidal setting. Figure 8 shows the bootstrap standard deviation of the calibrated probabilities for the three alternative methods and for Venn–Abers, together with the Venn–Abers width.

The Venn–Abers width and the bootstrap curves have a similar local structure, including spikes around changes in the fitted isotonic blocks. These are regions where relatively small changes in the calibration sample can alter the PAV partition and hence the fitted probability. The comparison is not intended to show that Venn–Abers width is numerically the same as a bootstrap standard deviation. It does, however, show that the width provides a local diagnostic of the sensitivity of the calibration fit without having to rerun the calibrator many times.

9 Discussion and Practical Implications

The experiments suggest a more specific and limited interpretation of the Venn–Abers interval width than is often assumed in applications.

The first point is the connection with the local PAV block. In a stable block containing N observations, changing the hypothetical label of the test point changes the fitted probability by roughly $1/N$. The scaling experiments show the corresponding dependence on calibration size, score density, Bernoulli variance, and local slope. We therefore use the terms local calibration leverage or local calibration support for this quantity. We do not interpret it as a general measure of predictive epistemic uncertainty.

Outcome ambiguity is a separate quantity. The $v^{-1/3}$ term in Proposition 1 is conditional on density and local slope, and these variables can move together in a fitted model. In the crowd-annotation experiment, width exhibits virtually zero correlation with annotator disagreement or entropy. Furthermore, on real tabular benchmarks, outcome ambiguity and calibration support




Figure 8: Calibration instability across the four calibration methods. For Venn–Abers we also show the interval width $p_1 - p_0$.

width combine to explain finite-sample calibration instability, with width providing substantial incremental explanatory power ($\Delta R^2 > 0$). Width on its own should therefore not be used as a general indicator of ambiguous labels.

It is equally essential to separate calibration support from base-model epistemic uncertainty. In our controlled thinning experiments, thinning the local calibration set monotonically increased width and calibration instability, while base-model epistemic uncertainty remained completely unchanged. Conversely, subsampling the training set increased base-model uncertainty without affecting calibration width. Thus:

outcome ambiguity $\neq$ base-model epistemic uncertainty $\neq$ calibration-support uncertainty.

There is also a difference between width and the variability of the calibrated probability. The block argument gives an $n^{-2/3}$ rate for width but an $n^{-1/3}$ rate for the probability itself. The quantity

$$U_{\text{cal}} = \sqrt{\hat{p}(1 - \hat{p})w} \quad (27)$$

has the latter scale and performs better than raw width in the pointwise bootstrap experiment. The ratio between bootstrap SD and U_{cal} is not constant across all of the local regimes, however, so we regard U_{cal} as an instability index rather than as a general standard-error formula.

The $n^{-2/3}$ relationship also gives a rough indication of how calibration sample size affects width. If the local regime remains unchanged, reducing a characteristic width by one half requires an increase in calibration sample size of approximately $2^{3/2} \approx 2.83$. This is only a local calculation because the score density and shape of the calibration curve can also change when the calibration population changes.

Finally, the non-monotonic experiments show another limitation of interpreting width as a general uncertainty score. Persistent violations can create large pooled PAV blocks and an $O(n^{-1})$ width regime. Higher-order flat monotone regions have their own rates. In both cases a small width can coexist with approximation error because the isotonic projection has pooled away local structure.

9.1 Limitations

The scaling relationships in this paper are heuristic rather than formal limit theorems. The large- n experiment shows the finite-sample coefficients moving towards the proposed values, but this is empirical evidence and not a proof of convergence. U_{cal} has the predicted rate but is not an exact pointwise standard-error estimate. The crowd-annotation experiment is simulated rather than based on human annotator traces. A formal treatment of the pseudo-label path through changes in the PAV active blocks, together with extensions to multi-class Venn–Abers prediction, would be useful further work.

10 Conclusion

This paper investigated the meaning of the Venn–Abers interval width $w = p_1 - p_0$. The main result is that the width can be understood through the local PAV block fitted by isotonic regression. In the regular monotonic case this gives

$$w_n(s) \propto n^{-2/3} f(s)^{-2/3} [\theta(s)(1 - \theta(s))]^{-1/3} [\theta'(s)]^{2/3}. \quad (28)$$

The simulation results recover the sample-size rate and show the finite-sample density, ambiguity, and slope coefficients getting closer to the proposed values as the calibration sample increases.

The same argument also explains why interval width and probability-scale calibration instability are related but are not the same quantity. Width has the inverse-block $n^{-2/3}$ rate, while the fitted probability has the usual isotonic $n^{-1/3}$ rate. The corresponding index $U_{\text{cal}} = \sqrt{\hat{p}(1 - \hat{p})w}$ follows the predicted rate and is more closely related to pointwise bootstrap instability than raw width in our experiments.

The empirical results also show where this interpretation should stop. Width is not a direct measure of label ambiguity, distribution shift, or base-model parameter uncertainty. Across controlled experiments on real tabular benchmarks, reducing local calibration support increases Venn–Abers width and calibration instability without producing any response in base-model epistemic uncertainty. The results suggest therefore that Venn–Abers width is most useful as a description of local support and sensitivity of the calibration mapping, rather than as a general scalar uncertainty score.

A Invariance of the Scaling Law Under Score Transformations

In this appendix, we show that the heuristic scaling law proposed in Equation 13 is invariant under any strictly increasing transformation of the scores.

Let $g : \mathbb{R} \rightarrow \mathbb{R}$ be a strictly increasing, differentiable function defining a score transformation $s_{\text{new}} = g(s)$. Let $\theta(s)$ and $f(s)$ denote the true probability function and score density under the original score space, respectively. Under the transformed score space:

1. The true probability function becomes $\theta_{\text{new}}(s_{\text{new}}) = \theta(s)$.

2. By the chain rule, the derivative of the true probability with respect to the new score is:

$$\theta'_{\text{new}}(s_{\text{new}}) = \frac{d\theta_{\text{new}}}{ds_{\text{new}}} = \frac{d\theta}{ds} \frac{ds}{ds_{\text{new}}} = \theta'(s) \cdot \frac{1}{g'(s)} \quad (29)$$

3. The probability density of the scores transforms under change-of-variables as:

$$f_{\text{new}}(s_{\text{new}}) = f(s) \cdot \left| \frac{ds}{ds_{\text{new}}} \right| = f(s) \cdot \frac{1}{g'(s)} \quad (30)$$

Substituting these transformed terms into the heuristic scaling law (Equation 13) for the transformed scores:

$$w(s_{\text{new}}) \propto f_{\text{new}}(s_{\text{new}})^{-2/3} \cdot (\theta_{\text{new}}(1 - \theta_{\text{new}}))^{-1/3} \cdot (\theta'_{\text{new}}(s_{\text{new}}))^{2/3} \quad (31)$$

$$= \left(\frac{f(s)}{g'(s)} \right)^{-2/3} \cdot (\theta(s)(1 - \theta(s)))^{-1/3} \cdot \left(\frac{\theta'(s)}{g'(s)} \right)^{2/3} \quad (32)$$

$$= f(s)^{-2/3} \cdot (g'(s))^{2/3} \cdot (\theta(s)(1 - \theta(s)))^{-1/3} \cdot (\theta'(s))^{2/3} \cdot (g'(s))^{-2/3} \quad (33)$$

Notice that the derivative factors $(g'(s))^{2/3}$ and $(g'(s))^{-2/3}$ cancel each other exactly:

$$w(s_{\text{new}}) \propto f(s)^{-2/3} \cdot (\theta(s)(1 - \theta(s)))^{-1/3} \cdot (\theta'(s))^{2/3} = w(s) \quad (34)$$

This cancellation shows that the scaling law is mathematically invariant to monotonic score transformations (even if the transformation derivative $g'(s)$ varies abruptly or non-linearly), which is consistent with the rank-ordering nature of the PAV algorithm.

Acknowledgements

I thank Prof. Vladimir Vovk for useful discussions in writing this paper.

References

- [1] V. Vovk and I. Petej. Venn–Abers Predictors. In *Proceedings of the Thirtieth Conference on Uncertainty in Artificial Intelligence (UAI)*, pages 829–838, 2014.
- [2] V. Vovk, I. Petej, and V. Fedorova. Large-scale probabilistic predictors with and without guarantees of validity. In *Advances in Neural Information Processing Systems (NeurIPS)*, volume 28, pages 892–900, 2015.
- [3] V. Manokhin. Multi-class probabilistic classification using inductive and cross Venn-Abers predictors. In *Proceedings of the Sixth Workshop on Conformal and Probabilistic Prediction and Applications*, volume 60 of *Proceedings of Machine Learning Research*, pages 228–240. PMLR, 2017.
- [4] I. Nourtdinov, D. Volkhonskiy, P. Lim, P. Toccaceli, and A. Gammerman. Inductive Venn-Abers predictive distribution. In *Proceedings of the Seventh Workshop on Conformal and Probabilistic Prediction and Applications*, volume 91 of *Proceedings of Machine Learning Research*, pages 15–36. PMLR, 2018.

- [5] I. Petej and V. Vovk. Inductive Venn-Abers and related regressors. *arXiv preprint arXiv:2605.06646*, 2026.
- [6] P. Groeneboom. Estimating a monotone density. In *Proceedings of the Berkeley Conference in Honor of Jerzy Neyman and Jack Kiefer*, volume 2, pages 539–555. Wadsworth, 1985.
- [7] F. T. Wright. The asymptotic behavior of monotone regression estimates. *The Annals of Statistics*, 9(2):443–448, 1981.
- [8] T. Robertson, F. T. Wright, and R. L. Dykstra. *Order Restricted Statistical Inference*. John Wiley & Sons, 1988.
- [9] P. Groeneboom and G. Jongbloed. *Nonparametric Estimation under Shape Constraints*. Cambridge University Press, 2014.
- [10] Q. Han and K. Kato. Berry–Esseen bounds for Chernoff-type nonstandard asymptotics in isotonic regression. *The Annals of Applied Probability*, 32(2):1459–1498, 2022.
- [11] A. Kumar, P. S. Liang, and T. Ma. Verified uncertainty calibration. In *Advances in Neural Information Processing Systems (NeurIPS)*, volume 32, pages 4228–4238, 2019.
- [12] J. Vaicenavicius, D. Widmann, C. Andersson, F. Lindsten, J. Roll, and T. Schön. Evaluating model calibration in classification. In *International Conference on Artificial Intelligence and Statistics (AISTATS)*, volume 89 of *Proceedings of Machine Learning Research*, pages 3459–3467. PMLR, 2019.
- [13] C. Gupta, A. Podkopaev, and A. Ramdas. Distribution-free binary classification: prediction sets, confidence intervals and calibration. In *Advances in Neural Information Processing Systems (NeurIPS)*, volume 33, pages 3711–3723, 2020.
- [14] J. Peck, B. Goossens, and Y. Saeys. Detecting adversarial examples with inductive Venn-ABERS predictors. In *Proceedings of the 27th European Symposium on Artificial Neural Networks, Computational Intelligence and Machine Learning (ESANN)*, pages 251–256, 2019.
- [15] U. Johansson, T. Löfström, and C. Sönströd. Well-calibrated probabilistic predictive maintenance using Venn-Abers. *arXiv preprint arXiv:2306.06642*, 2023.
- [16] Venn-Abers Python Library. <https://github.com/ip200/venn-abers>