Exact modular-form certificates for four 2-colour overpartition congruences

Anonymous

12 October 2026

Abstract

For positive integers ℓ, define $\sum_{n\geq 0}\overline{p}_\ell(n)q^n = f_2 f_{2\ell}/(f_1^2 f_\ell^2)$, where $f_k = \prod_{m\geq 1}(1-q^{km})$. We give exact modular-form certificates for $\overline{p}_8(32n+28)\equiv 0 \pmod{32}$, $\overline{p}_6(24n+17)\equiv \overline{p}_6(24n+23)\equiv 0 \pmod{64}$ and $\overline{p}_9(36n+33)\equiv 0 \pmod{128}$. The first statement already occurs in Bharadwaj–Sujatha–Chandankumar's Theorem 4.4, as well as their conjecture list; we provide a complete independent certificate, not new statement priority. At most 541 extracted Fourier coefficients are tested against a Sturm bound. The largest selected index of the underlying form is 19,440, and the direct progression confirmation reaches 19,473.

1 Introduction

Let $f_k = (q^k;q^k)_\infty$. In [1] Bharadwaj, Sujatha and Chandankumar study the functions $\overline{p}_\ell(n)$ with

$$\sum_{n\geq 0}\overline{p}_\ell(n)q^n = \frac{f_2\, f_{2\ell}}{f_1^2\, f_\ell^2},$$

the generating function of pairs of overpartitions in which the second component has all parts divisible by ℓ. Their methods are elementary (q-series dissections), and they deliberately avoid modular forms. In their final section they state the following.

Theorem 1 (Statements in [1]; attribution below)**.** *For all $n \geq 0$,*

$$\begin{aligned}\overline{p}_8(32n+28) &\equiv 0 \pmod{32},\\ \overline{p}_6(24n+17) &\equiv 0 \pmod{64},\\ \overline{p}_6(24n+23) &\equiv 0 \pmod{64}.\end{aligned}$$

Theorem 2 (Second congruence of a further conjecture in [1])**.** *For all $n \geq 0$, $\overline{p}_9(36n+33)\equiv 0 \pmod{128}$.*

The first congruence is the $\alpha = 2$ case of [1, Theorem 4.4, equation (4.51)], and also appears in its Conjecture 2, equation (5.4). The inspected proof does not explicitly derive the full all-α assertion. We record this ambiguity rather than claim either new statement priority or a complete earlier proof. The other three are equations (5.5), (5.6), (5.8) of that source. No exact earlier proof was located in the bounded search documented with this collection; this is not a priority certification. The companion family paper treats the modulo-8 parameter family and extends the dyadic modulo-16 family.

We prove Theorems 1 and 2 by the standard method of Ono and Radu [2, 3]: we multiply the generating function by an eta-quotient in q^M chosen to make it a holomorphic modular form,

apply U_M, and check the finitely many coefficients that Sturm's theorem [4] requires. The only work is finding suitable multipliers. We found them by integer linear programming; the proof itself depends only on checking the certificates in Table 1.

2 Tools

Let $\eta(z) = q^{1/24} f_1$ with $q = e^{2\pi i z}$. An eta-quotient is $F(z) = \prod_{d|N} \eta(dz)^{r_d}$.

Lemma 3 (Gordon–Hughes–Newman, Ligozat; [2, Thms. 1.64, 1.65])**.** *Suppose $k = \frac{1}{2}\sum r_d$ is a positive integer, $\sum_d d\, r_d \equiv 0 \pmod{24}$ and $\sum_d (N/d)\, r_d \equiv 0 \pmod{24}$. Then F satisfies the transformation law of weight k on $\Gamma_0(N)$ with character $\chi(m) = (\frac{(-1)^k s}{m})$, where $s = \prod d^{r_d}$. If moreover for every divisor $c \mid N$*

$$\frac{N}{24} \sum_{d|N} \frac{\gcd(c,d)^2\, r_d}{\gcd(c, N/c)\, c\, d} \;\geq\; 0,$$

then F is holomorphic at all cusps, so $F \in M_k(\Gamma_0(N), \chi)$.

Lemma 4 ([2, Prop. 2.22])**.** *Let χ be a Dirichlet character modulo N. If $M \mid N$, then $U_M : \sum a(n) q^n \mapsto \sum a(Mn) q^n$ maps $M_k(\Gamma_0(N), \chi)$ to itself.*

Lemma 5 (Sturm [4]; [2, Thm. 2.58])**.** *Let $G \in M_k(\Gamma_0(N), \chi)$ have integer coefficients $a(n)$, and let p be a prime. If $a(n) \equiv 0 \pmod{p}$ for all $n \leq S = \frac{k}{12}[\mathrm{SL}_2(\mathbb{Z}) : \Gamma_0(N)]$, then $a(n) \equiv 0 \pmod{p}$ for all n.*

Applying Lemma 5 j times shows that if $2^j \mid a(n)$ for all $n \leq S$ then $2^j \mid G$. After each step, $G/2$ is again in $M_k(\Gamma_0(N), \chi)$ with integer coefficients.

3 The reduction

Fix ℓ, a modulus M and a residue r. Let $A_\ell(z) = \eta(2z)\eta(2\ell z)/(\eta(z)^2 \eta(\ell z)^2)$. Its exponents satisfy $\sum d\, r_d = 2 + 2\ell - 2 - 2\ell = 0$, so $A_\ell = \sum \overline{p}_\ell(n) q^n$ exactly. Choose integers b_d for $M \mid d \mid N$ and put

$$B(z) = \prod_{M|d|N} \eta(dz)^{b_d} = q^c\, g(q^M), \qquad c = \tfrac{1}{24}\textstyle\sum d\, b_d, \quad g \in 1 + q\mathbb{Z}[[q]].$$

Suppose $c \in \mathbb{Z}$ and $c \equiv -r \pmod{M}$, and that $F = A_\ell B$ satisfies the hypotheses of Lemma 3. Then

$$F = q^c g(q^M) \sum_{n \geq 0} \overline{p}_\ell(n) q^n,$$

and the exponents divisible by M are exactly those with $n \equiv r \pmod{M}$. Hence

$$G := F \mid U_M = q^{(c+r)/M}\, g(q) \sum_{n \geq 0} \overline{p}_\ell(Mn + r)\, q^n \in M_k(\Gamma_0(N), \chi)$$

by Lemma 4. Since g is a unit in $\mathbb{Z}[[q]]$, 2^j divides every coefficient of G if and only if $2^j \mid \overline{p}_\ell(Mn+r)$ for all $n \geq 0$. By the remark after Lemma 5, it suffices to check the coefficients of G up to q^S.

4 Certificates

Table 1 lists the multipliers. In each case:

- $\sum(N/d)r_d = 0$ for $F = A_\ell B$;
- the minimum over the cusps of Ligozat's order is 0, so the hypotheses of Lemma 3 hold;
- $M \mid N$ and χ is a character modulo N, so Lemma 4 applies (indeed the conductor of χ, 1 or 3, even divides N/M).

For (A), $s = 2^{156}$ is a square and k is even, so χ is trivial. In the other cases k is odd and the squarefree part of s is 3, so $\chi = (\frac{-3}{\cdot})$.

(ℓ, M, r)	N	k	c	S	j	coeffs
$(8, 32, 28)$	128	18	4	288	5	289
$(6, 24, 17)$	288	11	7	528	6	529
$(6, 24, 23)$	288	11	1	528	6	529
$(9, 36, 33)$	216	15	3	540	7	541

$$
\begin{aligned}
B_{8,28} &= \eta_{32}^{69}\eta_{64}^{-29}\eta_{128}^{-2},\\
B_{6,17} &= \eta_{24}^{61}\eta_{48}^{-31}\eta_{72}^{-16}\eta_{96}^{2}\eta_{144}^{8},\\
B_{6,23} &= \eta_{24}^{59}\eta_{48}^{-25}\eta_{72}^{-16}\eta_{96}^{-2}\eta_{144}^{8},\\
B_{9,33} &= \eta_{36}^{88}\eta_{72}^{-43}\eta_{108}^{-26}\eta_{216}^{13}.
\end{aligned}
$$

Table 1: Certificates; $\eta_d = \eta(dz)$. The indices $[\mathrm{SL}_2(\mathbb{Z}) : \Gamma_0(N)]$ are 192, 576, 432.

Proposition 6. *For each row of Table 1, all coefficients of $q^0, \dots, q^S$ in G are divisible by 2^j.*

Proof. This is an exact integer computation. The failure-sensitive script `verify.py` expands F, applies U_M, tests every coefficient $0 \le m \le S$, and checks the extraction identity and direct progression. A separate standard-library checker reconstructs the overpartition coefficients by theta inversion and the multiplier by a logarithmic-derivative recurrence, without FLINT. The historical auxiliary checker shared FLINT and a pentagonal-product approach and is not described as independent reproduction. A compact receipt records cusp orders, characters, shifts and exact ranges. The universal conclusion also requires the written modular-form argument above. □

Proof of Theorems 1 and 2. Combine Section 3 with Proposition 6. □

Remark 7. Explicit witnesses are $\overline{p}_8(28) = 86{,}368 = 32 \cdot 2699$, $\overline{p}_6(17) = 3648 = 64 \cdot 57$, $\overline{p}_6(23) = 25{,}024 = 64 \cdot 391$ and $\overline{p}_9(69) = 594{,}986{,}112 = 128 \cdot 4{,}648{,}329$. Thus none of the four moduli can be doubled uniformly. Stronger subprogressions are not excluded. The integer-programming discovery search is not a proof prerequisite and no minimality of its multipliers is claimed.

Version and assurance

Version 1.1.0-candidate, DOI `https://doi.org/10.5281/zenodo.23320550`. Unrefereed candidate; internal exact checking and AI-assisted review are not formal verification or external peer review.

References

[1] H. S. Sumanth Bharadwaj, N. Sujatha and S. Chandankumar, Arithmetic properties of the 2-color overpartition function $\overline{p}_\ell(n)$, arXiv:2607.16608 (2026).

[2] K. Ono, *The Web of Modularity: Arithmetic of the Coefficients of Modular Forms and q-series*, CBMS Regional Conf. Ser. Math. 102, Amer. Math. Soc., 2004.

[3] S. Radu, An algorithmic approach to Ramanujan's congruences, *Ramanujan J.* 20 (2009), 215–251.

[4] J. Sturm, On the congruence of modular forms, in *Number Theory (New York, 1984–1985)*, Lecture Notes in Math. 1240, Springer, 1987, 275–280.

Parameter classification and binary-form support for 2-colour overpartitions

Anonymous

12 October 2026

Abstract

Let $\sum_{n\geq 0}\overline{p}_\ell(n)q^n = f_2 f_{2\ell}/(f_1^2 f_\ell^2)$ with $f_k = \prod_{m\geq 1}(1-q^{km})$. Bharadwaj, Sujatha and Chandankumar conjectured that $\overline{p}_\ell(4n+3)\equiv 0 \pmod 8$ for every $\ell \equiv 0,1 \pmod 4$, and that $\overline{p}_9(2^\alpha(8n+7))\equiv 0 \pmod{16}$ for every $\alpha \geq 1$. We prove both statements; the second in fact holds for all $\alpha \geq 0$. We further classify the modulo-8 parameter range exactly, prove a binary-form support theorem for square or twice-square parameters, and extend the dyadic family to every parameter whose odd part is 1 modulo 8. The methods are elementary and classical.

1 Introduction

Write $f_k = (q^k;q^k)_\infty$, $\varphi(q) = \sum_{k\in\mathbb{Z}} q^{k^2}$, and

$$\sum_{n\geq 0}\overline{p}_\ell(n)q^n = \frac{f_2\, f_{2\ell}}{f_1^2\, f_\ell^2}.$$

These functions were studied in [1]. The authors proved many congruences modulo 8, including $\overline{p}_\ell(4n+3)\equiv 0 \pmod 8$ for $\ell \equiv 0 \pmod 8$, and stated the following conjectures in their final section.

Theorem 1. *For a positive integer ℓ, the congruence $\overline{p}_\ell(4n+3)\equiv 0 \pmod 8$ holds for every integer $n\geq 0$ if and only if $\ell \equiv 0,1 \pmod 4$.*

Theorem 2. *For every $\alpha \geq 0$ and $n \geq 0$, $\overline{p}_9\big(2^\alpha(8n+7)\big)\equiv 0 \pmod{16}$.*

The sufficient direction of Theorem 1 is Conjecture 1 of [1], where the case $\ell \equiv 0 \pmod 8$ is proved. Theorem 2 for $\alpha \geq 1$ is the first part of their Conjecture 3; no prior statement of the case $\alpha = 0$ was located in our bounded search. For comparison, [1] state a 2-power family $\overline{p}_8(2^\alpha(8n+7)) \equiv 0 \pmod{32}$ in Theorem 4.4, equation (4.51); the companion certificate paper discusses the source's overlap with its conjecture list. The hypothesis $\ell \equiv 0,1 \pmod 4$ in Theorem 1 cannot be dropped: $\overline{p}_2(3)\equiv 4$, $\overline{p}_3(3)\equiv 2$, $\overline{p}_6(7)\equiv 4$ and $\overline{p}_7(7)\equiv 2 \pmod 8$.

Related work and method. The arguments use classical reciprocal-theta expansions and representation parity, as in Mahlburg [6], Hirschhorn–Sellers [3] and Kim [4]. For $\ell = 1$, the coefficients count unrestricted overpartition pairs: Keister–Sellers–Vary [5, Theorem 9, $k=2$] implies the modulo-8 result. For every $4 \mid \ell$, it follows immediately from [3, Theorem 2, equation (6)] and the convolution

$$\overline{p}_\ell(4n+3) = \sum_{j=0}^{\lfloor (4n+3)/\ell \rfloor} \overline{p}(4n+3-\ell j)\overline{p}(j).$$

Each first argument is 3 modulo 4. The uniform argument and extensions below are applications of established methods, not new techniques. The bounded literature check found no matching full parameter theorem, but does not certify priority. For $\ell = 2$ the object is the overcubic partition function, so known overcubic congruences are relevant special cases, not a new object.

Throughout, $S_j = \sum_{k\geq 1} q^{jk^2}$, so $\varphi(q^j) = 1 + 2S_j$ and $\varphi(-q^j) = 1 + 2\sum_{k\geq 1}(-1)^k q^{jk^2}$. All congruences between power series are coefficientwise.

2 Gauss's identities

Lemma 3. *$\varphi(-q) = f_1^2/f_2$, $\varphi(q) = f_2^5/(f_1^2 f_4^2)$, and hence $\varphi(q)\varphi(-q) = \varphi(-q^2)^2$.*

Proof. The two product formulas are classical (Gauss; see [2, Entry 22]). Then $\varphi(q)\varphi(-q) = f_2^4/f_4^2 = \varphi(-q^2)^2$. □

In particular $\sum \overline{p}_\ell(n)q^n = 1/(\varphi(-q)\varphi(-q^\ell))$.

3 Proof of Theorem 1

Lemma 4. *Let $A = 1/\varphi(-q)$. Then $A \equiv 1 + 2S_1 + 4(S_2 + S_4) \pmod 8$.*

Proof. Applying Lemma 3 with q and with q^2 gives $\varphi(-q) = \varphi(-q^2)^2/\varphi(q)$ and $\varphi(-q^2)^2 = \varphi(-q^4)^4/\varphi(q^2)^2$, so

$$A = \frac{\varphi(q)\,\varphi(q^2)^2}{\varphi(-q^4)^4}.$$

Since $\varphi(-q^4) = 1 + 2X$ with $X \in \mathbb{Z}[[q]]$, we have $\varphi(-q^4)^4 \equiv 1 \pmod 8$, and so its inverse is $\equiv 1 \pmod 8$. Next, $\varphi(q^2)^2 = 1 + 4S_2 + 4S_2^2$, and $S_2^2 \equiv S_4 \pmod 2$ because $g^2 \equiv g(q^2) \pmod 2$ for every $g \in \mathbb{Z}[[q]]$. Hence $\varphi(q^2)^2 \equiv 1 + 4S_2 + 4S_4 \pmod 8$ and $A \equiv (1 + 2S_1)(1 + 4S_2 + 4S_4) \equiv 1 + 2S_1 + 4S_2 + 4S_4 \pmod 8$. □

Proof of Theorem 1. By Lemma 4 write $A(q) \equiv 1 + 2a + 4b$ with $a = S_1$ and $b = S_2 + S_4$. Then

$$A(q)A(q^\ell) \equiv 1 + 2\big(a(q) + a(q^\ell)\big) + 4\big(b(q) + b(q^\ell)\big) + 4a(q)a(q^\ell) \pmod 8.$$

Every exponent occurring in $b(q)$ or $b(q^\ell)$ is even. The exponents in $a(q)$ are squares, hence $\equiv 0, 1 \pmod 4$. Those in $a(q^\ell)$ are ℓk^2, also $\equiv 0, 1 \pmod 4$ when $\ell \equiv 0, 1 \pmod 4$. The exponents in $a(q)a(q^\ell)$ are $j^2 + \ell k^2 \equiv 0, 1, 2 \pmod 4$. So no exponent $\equiv 3 \pmod 4$ occurs on the right, and the coefficient of q^{4n+3} in $A(q)A(q^\ell) = \sum \overline{p}_\ell(n)q^n$ is divisible by 8. □

For completeness, the converse in Theorem 1 has uniform witnesses. If $\ell \equiv 3 \pmod 4$, then $\overline{p}_\ell(\ell) = \overline{p}(\ell) + 2 \equiv 2 \pmod 8$. If $\ell \equiv 2 \pmod 4$, then $\overline{p}_\ell(\ell + 1) = \overline{p}(\ell + 1) + 2\overline{p}(1) \equiv 4 \pmod 8$. Here the ordinary congruence follows from Lemma 4. The witness $\overline{p}_4(3) = 8$ shows that modulus 8 is sharp uniformly over the admissible parameter family.

4 Proof of Theorem 2

Let $N = 2^\alpha m$ with $\alpha \geq 0$ and $m \equiv 7 \pmod 8$. We use three facts.

(F1) N is not a perfect square, since its odd part m is not $\equiv 1 \pmod 8$.

(F2) N is not a sum of two squares: $m \equiv 3 \pmod 4$ has a prime factor $p \equiv 3 \pmod 4$ to an odd power.

(F3) $N \neq X^2 + 2Y^2$ for integers X, Y. A positive integer represented by $X^2 + 2Y^2$ has each prime $p \equiv 5, 7 \pmod 8$ to an even power [7, Ch. 1]. Its odd part is therefore a product of primes $\equiv 1, 3 \pmod 8$ and odd squares, which is $\equiv 1$ or $3 \pmod 8$. Here $m \equiv 7$.

Proof of Theorem 2. Write $\varphi(-q) = 1 + 2s$ and $\varphi(-q^9) = 1 + 2t$, where $s = \sum_{k\geq 1}(-1)^k q^{k^2}$ and $t = s(q^9)$. Then, modulo 16,

$$\frac{1}{\varphi(-q)\varphi(-q^9)} \equiv \sum_{i+j\leq 3} (-2)^{i+j} s^i t^j.$$

We show that the coefficient c of q^N in each term $(-2)^{i+j}s^i t^j$ with $1 \leq i + j \leq 3$ is $\equiv 0 \pmod{16}$.

Degree 1. s and t are supported on squares, so their coefficients at q^N vanish by (F1).

Degree 2. The terms s^2, st and t^2 are supported on numbers a^2+b^2, $a^2+(3b)^2$ and $(3a)^2+(3b)^2$. Their coefficients at q^N vanish by (F2).

Degree 3. These terms carry the factor 8, so it suffices that each of s^3, s^2t, st^2, t^3 has an even coefficient at q^N. Each coefficient is a signed count of ordered triples, so its parity equals that of the number of triples.

- For s^3, count triples (a, b, c) of positive integers with $a^2 + b^2 + c^2 = N$, under the symmetric group. Orbits with distinct entries have size 6. Orbits with exactly two equal entries have size 3 and correspond to $N = 2x^2 + y^2$ with $x \neq y$, which is impossible by (F3). The orbit $a = b = c$ needs $N = 3a^2$, whose odd part is $\equiv 3 \pmod 8$; this is impossible.
- For s^2t, consider $a^2 + b^2 + 9c^2 = N$ and swap a and b. A fixed point gives $N = 2a^2 + (3c)^2$, impossible by (F3).
- For st^2, consider $a^2 + 9b^2 + 9c^2 = N$ and swap b and c. A fixed point gives $N = a^2 + 2(3b)^2$, impossible by (F3).
- For t^3, the coefficient is zero unless $9 \mid N$, in which case $N/9 = 2^\alpha(m/9)$ with $m/9 \equiv 7 \pmod 8$, and the s^3 argument applies to $N/9$.

Hence every coefficient is even, and $\overline{p}_9(N) \equiv 0 \pmod{16}$. □

Remark 5. The public checker `checks/families.py` uses exact modular arithmetic. It tests 60,000 modulo-eight instances, 60 excluded-parameter witnesses, 274,780 dyadic instances through argument 20,000 across 55 parameters, and 10,416 binary-form support instances through argument 1,000. A direct Euler-product expansion cross-checks the inverse-theta coefficients through argument 180. These finite checks complement, but do not replace, the universal proofs.

5 A support theorem and a larger dyadic family

Theorem 6. *Let d be a positive integer and $\ell = d^2$ or $2d^2$. If $N > 0$ is represented by neither $X^2 + Y^2$ nor $X^2 + 2Y^2$, with integral coordinates including zero, then $\overline{p}_\ell(N) \equiv 0 \pmod{16}$.*

Proof. Set $s = \sum_{k\geq 1}(-1)^k q^{k^2}$ and $t = s(q^\ell)$. In $\mathbb{Z}[[q]]$,

$$H_\ell \equiv 1 - 2(s + t) + 4(s^2 + st + t^2) - 8(s^3 + s^2t + st^2 + t^3) \pmod{16}.$$

The linear supports are squares and ℓ times squares. The quadratic supports are x^2+y^2, $x^2+\ell y^2$ and $\ell(x^2+y^2)$. All lie in the union of the two stated representation sets; for twice-square ℓ, use $2d^2(x^2+y^2) = (d(x+y))^2 + (d(x-y))^2$. Modulo 2, Frobenius reduces the cubic supports to

$$x^2+2y^2, \quad 2x^2+\ell y^2, \quad x^2+2\ell y^2, \quad \ell(x^2+2y^2).$$

For square ℓ these embed in X^2+2Y^2. For twice-square ℓ, the middle two embed in sums of two squares, using $2x^2+2d^2y^2 = (x+dy)^2+(x-dy)^2$ and $x^2+4d^2y^2 = x^2+(2dy)^2$. Finally $2d^2(x^2+2y^2) = (2dy)^2+2(dx)^2$. Thus at the stipulated N the linear and quadratic coefficients vanish and the cubic coefficients are even. This proves the result. The constant term explains why $N>0$ is required. □

Theorem 7. *Let $\ell = 2^a u$, where $a \geq 0$ and $u > 0$ is odd with $u \equiv 1 \pmod 8$. Then for every $\alpha, n \geq 0$,*

$$\overline{p}_\ell(2^\alpha(8n+7)) \equiv 0 \pmod{16}.$$

Proof. Call a positive value admissible if its odd part is not 7 (mod 8). Multiplication by a power of two, an odd square, or u preserves admissibility. Every positive value of X^2+uY^2 and X^2+2uY^2 is admissible: remove a common power of two from the coordinates. For the first form, opposite parities give odd residue 1 or 5 modulo 8; both odd give residue 2 modulo 8, hence valuation exactly one and odd part 1 modulo 4. For the second, odd X gives residue 1 or 3 modulo 8. Even X and odd Y give residue 2 or 6 modulo 16, so the odd part is 1 or 3 modulo 8. Zero coordinates are included; both zero give no positive value.

Absorbing even powers into Y shows the same for $X^2+2^b uY^2$, $b \geq 0$, also with $u=1$. Apply the expansion in the preceding proof. All its linear and quadratic supports are admissible. Of the four cubic parity supports displayed there, the first, third and fourth are admissible immediately. For $2x^2+\ell y^2$, if $a \geq 1$ factor out two and use $x^2+2^{a-1}uy^2$. If $a=0$, remove a common power of four from uy^2+2x^2: odd y gives residue 1 or 3 modulo 8; even y and odd x give 2 or 6 modulo 16. The same exclusion follows. At an exponent of odd part 7, the lower-degree terms vanish and the cubic coefficients are even, proving the congruence. □

Theorem 7 includes all square and twice-square parameters and also $\ell = 17, 34$, for example. It is a sufficient condition, not a classification. It does not enlarge Theorem 6: 21 is represented by neither of its two forms, but $\overline{p}_{17}(21) = 9932 \equiv 12 \pmod{16}$. The witness $\overline{p}_9(15) = 1552 = 16 \cdot 97$ forbids doubling the modulus uniformly. The support theorem arose in the supplied review, and the odd-part extension in a fresh targeted AI-assisted check during revision; neither carries a first-proof claim. Their significance lies in explaining the parameter dependence, not in introducing a new method.

Version and assurance

Version 1.1.0-candidate, DOI `https://doi.org/10.5281/zenodo.23320550`. Unrefereed candidate; internal exact checking and AI-assisted review are not formal verification or external peer review.

References

[1] H. S. Sumanth Bharadwaj, N. Sujatha and S. Chandankumar, Arithmetic properties of the 2-color overpartition function $\overline{p}_\ell(n)$, arXiv:2607.16608 (2026).

[2] B. C. Berndt, *Ramanujan's Notebooks, Part III*, Springer, New York, 1991.

[3] M. D. Hirschhorn and J. A. Sellers, Arithmetic relations for overpartitions, *J. Combin. Math. Combin. Comput.* 53 (2005), 65–73.

[4] B. Kim, The overpartition function modulo 128, *Integers* 8 (2008), A38.

[5] D. M. Keister, J. A. Sellers and R. G. Vary, Some arithmetic properties of overpartition k-tuples, *Integers* 9 (2009), 181–190, A17.

[6] K. Mahlburg, The overpartition function modulo small powers of 2, *Discrete Math.* 286 (2004), 263–267.

[7] D. A. Cox, *Primes of the Form* $x^2 + ny^2$, 2nd ed., Wiley, 2013.